

1 **In focus: Stay-at-home strategy brings fitness benefits to migrants**

2 James J. Gilroy, School of Environmental Sciences, University of East Anglia

3 **Abstract:**

4 In this issue, two studies examine the extent to which variation in migratory behaviour
5 influences individual fitness across a population. Lok et al. (2017) examine reproductive
6 success and post-fledging survival in a population of Eurasian spoonbills (*Platalea*
7 *leucorodia*), comparing individuals that winter in south-west Europe against those migrating
8 to sub-Saharan Africa, while Grist et al. (2017) measure reproductive success in a population
9 of European shags (*Phalacrocorax aristotelis*) breeding in Scotland that either remain
10 resident or migrate to surrounding waters. Both studies find that individuals migrating longer
11 distances tend to show later initiation of breeding attempts. In turn, longer migration
12 correlates with lower reproductive success in both populations. In spoonbills, this effect is
13 most pronounced in older male birds, whilst young individuals show little difference in
14 breeding success with respect to migration distance. In shags, fitness benefits of residence
15 were most pronounced when both individuals of a pair were resident, although there was no
16 evidence of assortative mating. Both studies provide fascinating new insights into the role
17 migratory variability can play in shaping population dynamics.

18 **Main text:**

19 Migration ranks among the greatest spectacles of the natural world, and yet it remains one of
20 the most challenging aspects of animal ecology to study. Partly owing to the difficulty of
21 tracking individuals as they traverse the globe, we still understand relatively little about key
22 aspects of migratory life, including the mechanisms that underpin variation in migration
23 within and between individuals, and how this influences migratory evolution.

24 For researchers seeking to understand migratory behaviour, fertile ground can be found
25 among species that express multiple migratory strategies within the same breeding
26 population. In particular, partial-migrants - those with both resident and migratory
27 phenotypes – offer opportunities to directly compare fitness consequences between strategies,
28 as well as tease apart factors influencing why individuals adopt one strategy or another
29 (Adriaensen & Dhondt 1990). Partial migration is relatively widespread in nature (Chapman
30 et al. 2011), but researchers have long struggled to collect the detailed data necessary to
31 elucidate the processes underlying emergent patterns. In particular, research has been
32 hampered by the difficulty of measuring migration at the individual scale, and of following
33 individuals for long enough to quantify their subsequent fitness (Marra et al. 1998;
34 Gunnarsson et al. 2006). Such data can only be collected through exhaustive efforts to mark
35 and track individuals across large spatial and temporal scales.

36 In this issue, two studies report on just such exhaustive efforts, shedding new light on the
37 within-species migratory variability and how it relates to individual fitness. Grist et al. (2017)
38 report on a study of European shags (*Phalacrocorax aristotelis*) breeding on the Isle of May,
39 Scotland, in which the reproductive success of resident individuals is compared against those
40 migrating relatively short distances (up to 100km) to wintering locations in surrounding
41 waters. Lok et al. (2017) use a similar approach to report on fitness differences among
42 Eurasian spoonbills (*Platalea leucorodia*) breeding in the Netherlands, this time comparing
43 shorter-distance migrants (up to 2,000 km) against long-distance migrants that winter in sub-
44 Saharan Africa (c. 4,700 km). Combined, these studies thus provide fascinating insights into
45 migratory variability across a broad swathe of the migratory spectrum.

46 Strikingly, both studies report similar overall patterns within their study populations (Fig. 1):
47 that individuals performing longer migrations tend to have lower reproductive fitness than
48 those wintering closer to their breeding sites. In shags, resident individuals tended to raise

49 more young, of higher condition than their migratory counterparts, and thus had greater
50 chance of subsequently recruiting young into the breeding population. In spoonbills, that
51 pattern was echoed for short-distance migrants relative to those that cross the Sahara each
52 winter. Increased fitness amongst shorter-distance migrants has been reported previously in
53 several other species (e.g. Adriaensen & Dhondt 1990; Gillis et al. 2008; Anderson et al.
54 2015), but the new studies in this issue provide unprecedented novel insights into the
55 mechanisms underpinning the emergent pattern.

56 In both studies, fitness differences among individuals were closely linked to the timing of
57 breeding. Individuals migrating shorter distances (spoonbills) or remaining resident (shags)
58 tended to initiate broods earlier in the season than their more migratory counterparts. As is
59 common throughout the avian world, earlier initiation translates into greater chances of
60 successfully raising young to fledging (Forslund & Part 1995). This could be related to nest
61 site quality – those arriving earliest at the breeding colony have access to the best locations
62 (Kokko 1999) - or optimal timing of breeding in relation to seasonal peaks of resource
63 availability (Both & te Marvelde 2007; Jones & Cresswell 2010). Early fledglings also tend
64 to have greater post-fledging survival chances, often as they have a longer period to feed
65 prior to the onset of more difficult conditions at the end of the breeding season (Naef-
66 Daenzer et al. 2001).

67 It should perhaps come as little surprise that longer-distance migrants have poorer breeding
68 success – migration is, after all, an arduous feat that can impose significant costs on the
69 condition of those undertaking it (McWilliams et al. 2004). However, this poses an
70 evolutionary conundrum: if longer-distance migration is more costly than residence, why
71 would it persist in the population? Evolutionary models suggest that multiple migration
72 strategies (including residence) can persist if 1) overall fitness benefits are balanced by the

73 two strategies, or 2) if the optimal outcome for an individual varies depending on its
74 phenotype (Lundberg 1988; Kaitala et al. 1993; Chapman et al. 2011).

75 In both of the new studies, fitness benefits appear to be unbalanced in favour of shorter
76 migration (unless some unmeasured parameter is having a buffering effect), suggesting that
77 condition 1 is not met in these cases. For condition 2 to be the driver underpinning migratory
78 persistence, migration would need to be the best option for at least some of the individuals
79 within each population. This could arise, for example, if individuals with a particular trait are
80 unlikely to survive if they adopt the resident/short distance strategy, despite incurring
81 subsequent costs for reproduction. What might such a trait be?

82 In the shag population, migratory strategies appear to remain fixed though adult life (Grist et
83 al. 2014), suggesting that migratory ‘decisions’ do not depend on any condition-dependent
84 phenotypic trait. Grist et al. (2017) also found no influence of sex on the likelihood of
85 individuals adopting a given strategy, nor on its consequent fitness benefits. Interestingly,
86 fitness benefits were maximised if both male and female of a pair were resident, and yet there
87 was no evidence of assortative mating in relation to migratory strategy. Costly migration
88 therefore does not appear to persist due to asymmetric sex-dependent benefits in this case,
89 and the role of other phenotypic traits (e.g. body size) remains unclear.

90 In spoonbills, Lok et al. (2017) uncover clearer evidence for a trait-dependent fitness cost of
91 migration, relating to age: older long-distance migrants (particularly males) tend to breed
92 significantly later than short-distance migrants of equivalent age, and consequently have
93 lower likelihood of raising young to recruitment. For younger individuals, however, the
94 fitness benefits of long- and short- migration were more similar. Trans-Saharan migration
95 might therefore remain an optimal (or at least break-even) strategy for younger individuals,
96 allowing long-distance migration to persist. Under such a scenario, however, we might expect

97 individuals to switch from longer to shorter migration as they age, and yet spoonbill
98 migratory strategy again typically remains fixed throughout life, just as it does in shags.
99 Condition-dependence therefore seems unlikely to be the principal mechanism allowing more
100 costly long-distance migration to persist in either population.

101 One possibility is that the relative benefits of different migratory strategies vary over
102 timescales greater than those captured in either study. Both sets of authors consider the
103 possibility that the observed benefits of shorter migration may be counterbalanced by other
104 factors over longer time periods, for example by periodic survival costs incurred among
105 residents when winter conditions are poor (Kaitala et al. 1993).

106 Another intriguing possibility is that the relative benefits of short-migration/residency have
107 emerged only recently, as a result of environmental change. This seems particularly plausible
108 in the case of the spoonbills, as the numbers of individuals overwintering in Europe have
109 increased significantly over the last 20 years (Lok, Overdijk & Piersma 2013). This could
110 reflect improving conditions for overwintering in Europe following climatic amelioration, or
111 deteriorating conditions in the traditional African winter range. Links between environmental
112 change and altered migratory behaviour are seldom documented, although climate-related
113 patterns of migratory change have been observed in European wildfowl (Maclean et al.
114 2008), as well as in some shorebirds (Austin & Rehfisch 2005) and passerines (Visser et al.
115 2009). Grist et al (2017) and Lok et al (2017) both provide compelling evidence that within-
116 population variation in migration distance, and consequent fitness impacts, could be
117 important drivers of population-scale changes in migratory behaviour. Such changes may
118 become increasingly important in future, as migratory species respond to climatic changes
119 occurring in each of the areas they visit during the annual cycle.

120 Overall, these studies demonstrate the tremendous insights that can be gained from long-term
121 mark-resight studies of migratory variability within populations. Whilst they reveal intriguing
122 commonalities, they also highlight the extent to which species differ in the mechanisms
123 underpinning migratory variability, and their effects on sex- and age-specific fitness. Many
124 questions remain to be answered, including the role of density dependence in regulating the
125 optimality of different strategies (Chapman et al. 2011), as well as the determinants of
126 strategy selection at early life stages. Understanding the mechanistic processes underpinning
127 migratory variation will be fundamental if we are to fully understand how migratory systems
128 evolve, or predict how they will respond to conservation actions under ongoing
129 environmental change.

130 **References**

- 131 Adriaensen, F. & Dhondt, A.A. (1990) Population dynamics and partial migration of the
132 European robin (*Erithacus rubecula*) in different habitats. *Journal of Animal Ecology*, 59,
133 1077-1090.
- 134 Anderson, A.M., Novak, S.J., Smith, J.F., Steenhof, K., Heath, J.A. (2015) Nesting
135 phenology, mate choice and genetic divergence within a partially migratory population of
136 American Kestrels. *The Auk*, 133, 99-109.
- 137 Austin, G. E., & Rehfisch, M. M. (2005). Shifting nonbreeding distributions of migratory
138 fauna in relation to climatic change. *Global Change Biology*, 11, 31-38.
- 139 Both, C., & te Marvelde, L. (2007). Climate change and timing of avian breeding and
140 migration throughout Europe. *Climate Research*, 35, 93-105.
- 141 Chapman, B. B., Brönmark, C., Nilsson, J. Å., & Hansson, L. A. (2011). The ecology and
142 evolution of partial migration. *Oikos*, 120, 1764-1775.

143 Forslund, P. & Part, T. (1995) Age and reproduction in birds - hypotheses and tests. Trends in
144 Ecology & Evolution, 10, 374-378

145 Gillis, E.A., Green, D.J., Middleton, H.A. & Morrissey, C.A. (2008) Life history correlates of
146 alternative migratory strategies in American Dippers. Ecology, 89, 1687-1695.

147 Grist, H., Daunt, F., Wanless, S., Nelson, E.J., Harris, M.P., Newell, M., Burthe, S. & Reid,
148 J.M. (2014) Site fidelity and individual variation in winter location in partially migratory
149 European shags. PloS One, 9, e98562.

150 Grist, H., Daunt, F., Wanless, S., Burthe, S. J., Newell, M. A., Harris, M. P., & Reid, J. M.
151 (2017). Reproductive performance of resident and migrant males, females and pairs in a
152 partially migratory bird. Journal of Animal Ecology.

153 Gunnarsson, T.G., Gill, J.A., Atkinson, P.W., Gélinaud, G., Potts, P.M., Croger, R.E.,
154 Gudmundsson, G.A., Appleton, G.F. & Sutherland, W.J. (2006) Population-scale drivers of
155 individual arrival times in migratory birds. Journal of Animal Ecology, 75, 1119-1127.

156 Jones, T., & Cresswell, W. (2010). The phenology mismatch hypothesis: are declines of
157 migrant birds linked to uneven global climate change?. Journal of Animal Ecology, 79, 98-
158 108.

159 Kaitala, A., Kaitala, V., & Lundberg, P. (1993). A theory of partial migration. The American
160 Naturalist, 142, 59-81.

161 Kokko, H. (1999). Competition for early arrival in migratory birds. Journal of Animal
162 Ecology, 68, 940-950.

163 Lok, T., Overdijk, O. & Piersma, T. (2013) Migration tendency delays distributional response
164 to differential survival prospects along a flyway. American Naturalist, 181, 520-531.

165 Lok, T., Veldhoen, L., Overdijk, O., Tinbergen, J. M., & Piersma, T. (2017) An age-
166 dependent fitness cost of migration? Old trans-Saharan migrating spoonbills breed later than
167 those staying in Europe, and late breeders have lower recruitment. *Journal of Animal*
168 *Ecology*.

169 Lundberg, P. (1988). The evolution of partial migration in birds. *Trends in Ecology &*
170 *Evolution*, 3, 172-175.

171 Maclean, I., Austin, G. E., Rehfisch, M. M., Blew, J. A. N., Crowe, O., Delany, S., ... & Van
172 Roomen, M. (2008). Climate change causes rapid changes in the distribution and site
173 abundance of birds in winter. *Global Change Biology*, 14, 2489-2500.

174 Marra, P.P., Hobson, K.A. & Holmes, R.T. (1998) Linking winter and summer events in a
175 migratory bird by using stable-carbon isotopes. *Science*, 282, 1884-1886

176 McWilliams, S. R., Guglielmo, C., Pierce, B., & Klaassen, M. (2004). Flying, fasting, and
177 feeding in birds during migration: a nutritional and physiological ecology perspective.
178 *Journal of Avian Biology*, 35, 377-393.

179 Naef-Daenzer, B., Widmer, F., & Nuber, M. (2001). Differential post-fledging survival of
180 great and coal tits in relation to their condition and fledging date. *Journal of Animal ecology*,
181 70, 730-738.

182 Visser, M. E., Perdeck, A. C., BALEN, V., Johan, H., & Both, C. (2009). Climate change
183 leads to decreasing bird migration distances. *Global Change Biology*, 15, 1859-1865.

184

185 **Figure legend**

186 **Figure 1.** Schematic showing key effects of within-population variation in migration distance
187 on subsequent breeding success, as documented by a) Lok et al. (2017) in a population of
188 Eurasian spoonbills (*Platalea leucorodia*) breeding in the Netherlands and migrating to
189 south-west Europe or sub-Saharan Africa, and b) Grist et al. (2017) in a population of
190 European shags (*Phalacrocorax aristotelis*) breeding in Scotland and remaining either
191 resident or migrating to surrounding waters.