

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Cook, JI; Majeed, S; Ignell, R; Pickett, JA; Birkett, MA; Logan, JG (2011) Enantiomeric selectivity in behavioural and electrophysiological responses of *Aedes aegypti* and *Culex quinquefasciatus* mosquitoes. *Bulletin of entomological research*, 101 (5). pp. 541-550. ISSN 0007-4853 DOI: 10.1017/S0007485311000162

Downloaded from: <http://researchonline.lshtm.ac.uk/18636/>

DOI: [10.1017/S0007485311000162](https://doi.org/10.1017/S0007485311000162)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

Enantiomeric selectivity in behavioural and electrophysiological responses of *Aedes aegypti* and *Culex quinquefasciatus* mosquitoes

J.I. Cook¹, S. Majeed², R. Ignell², J.A. Pickett¹, M.A. Birkett¹
and J.G. Logan^{3*}

¹Centre for Sustainable Pest and Disease Management, Chemical Ecology Group, Biological Chemistry Department, Rothamsted Research, Harpenden, Hertfordshire, AL5 2JQ, UK: ²Division of Chemical Ecology, Department of Plant Protection Biology, PO Box 102, Sundsvägen 14, SLU 230 53, Alnarp, Sweden: ³Department of Disease Control, Faculty of Infectious and Tropical Diseases, London School of Hygiene & Tropical Medicine, Keppel Street, London, WC1E 7HT, UK

Abstract

1-Octen-3-ol is a kairomone for many haematophagous insects including mosquitoes. Numerous studies have examined the effects of racemic 1-octen-3-ol; however, few studies have investigated the role of individual enantiomers in relation to mosquito attraction. In the present study, we investigated the behavioural and electrophysiological responses of two mosquito species, *Aedes aegypti* and *Culex quinquefasciatus*, to individual enantiomers and mixtures of 1-octen-3-ol, employing a laboratory Y-tube olfactometer and single sensillum recordings. The olfactory receptor neurons of both *Ae. aegypti* and *Cx. quinquefasciatus* had a significantly higher response to the (*R*)-1-octen-3-ol enantiomer compared to the (*S*)-1-octen-3-ol enantiomer at 10^{-9} g μl^{-1} to 10^{-6} g μl^{-1} . Behaviourally, *Ae. aegypti* was more responsive to the (*R*)-1-octen-3-ol enantiomer, showing an increase in flight activity and relative attraction compared to *Cx. quinquefasciatus*. The (*R*)-1-octen-3-ol enantiomer caused an increase in activation for *Cx. quinquefasciatus*. However, the most notable effect was from an (*R*:*S*)-1-octen-3-ol mixture (84:16) that caused significantly more mosquitoes to sustain their flight and reach the capture chambers (demonstrated by a reduced non-sustained flight activity), suggesting that it may have a behaviourally excitatory effect. For *Cx. quinquefasciatus*, a reduced relative attraction response was also observed for all treatments containing the (*R*)-1-octen-3-ol enantiomer, either on its own or as part of a mixture, but not with the (*S*)-1-octen-3-ol enantiomer. This is the first time enantiomeric selectivity has been shown for *Ae. aegypti* using electrophysiology *in vivo*. The implications of these results for exploitation in mosquito traps are discussed.

Keywords: mosquito, 1-octen-3-ol, enantiomer, electrophysiology, behaviour

(Accepted 22 February 2011)

*Author for correspondence
Fax: +44 (0)207 927 2918
E-mail: james.logan@lshtm.ac.uk

Introduction

Haematophagous insects, including mosquitoes (Diptera: Culicidae), use vision, heat and humidity, but predominantly volatile chemical cues, known as kairomones, to locate potential vertebrate hosts (Khan *et al.*, 1966; Eiras & Jepson, 1994; Gibson & Torr, 1999). One of the most extensively studied kairomones for haematophagous insects is 1-octen-3-ol. This compound, which exists in two enantiomeric forms, is released by many vertebrates and plant species (Hall *et al.*, 1984; Knudsen *et al.*, 1993; Kline *et al.*, 2007). 1-Octen-3-ol was first described in relation to haematophagous insects as a component of ox odour and has since been shown to be effective at increasing catches of tsetse flies (Diptera: Glossinidae) (Bursell, 1984; Hall *et al.*, 1984; Vale & Hall, 1985a,b). In combination with carbon dioxide (CO₂), a major component of vertebrate breath, 1-octen-3-ol has also been shown to increase behavioural responses and trap catches of mosquitoes, including those of the genera *Aedes* and *Anopheles* (Vythilingam *et al.*, 1992; Kemme *et al.*, 1993; Becker *et al.*, 1995; Kline & Lemire, 1995; Mboera *et al.*, 2000; Burkett *et al.*, 2001; Russell, 2004; Miller *et al.*, 2005; Kline *et al.*, 2007). Interestingly, 1-octen-3-ol was found to have little behavioural effect on most *Culex* spp. mosquitoes, despite the fact that some have olfactory receptors on their antennae and maxillary palps (Takken & Kline, 1989; Hill *et al.*, 2009).

1-Octen-3-ol is found in cattle breath to be predominantly (R)-1-octen-3-ol, varying between 80:20 R:S and 92:8 R:S (Hall *et al.*, 1984). Despite this, most investigations on haematophagous insects have assessed this compound only in its racemic form (1:1, R:S). Furthermore, the racemic form of 1-octen-3-ol is used in commercially available insect traps, such as the Mosquito Magnet and Midgeaters (Vythilingam *et al.*, 1992; Kemme *et al.*, 1993; Becker *et al.*, 1995; Mboera *et al.*, 2000; Burkett *et al.*, 2001; Russell, 2004; Miller *et al.*, 2005). Although no differences in either electroantennogram (EAG) activity, behaviour in a wind tunnel or attractiveness in the field between the (R) and the (S) enantiomers have been found for tsetse flies (Bursell, 1984; Hall *et al.*, 1984; Vale & Hall, 1985a,b), there is evidence that mosquitoes can discriminate between the enantiomers. For example, a study by Kline *et al.* (2007) showed an increase in numbers of *Anopheles crucians* and *Ochlerotatus infirmatus* caught in traps when using the (R)-1-octen-3-ol enantiomer compared to the (S)-1-octen-3-ol enantiomer. In line with this, a recent study showed that an olfactory receptor neuron (ORN) in the maxillary palps of *Cx. quinquefasciatus* mosquitoes displayed 'remarkable sensitivity' to the (R)-1-octen-3-ol enantiomer compared to the (S)-1-octen-3-ol enantiomer (Syed & Leal, 2007). Similarly, an odorant receptor of the yellow fever mosquito, *Ae. aegypti*, has been shown to be enantioselective *in vitro*, responding predominantly to the (R)-1-octen-3-ol enantiomer, when heterologously expressed in *Xenopus* oocytes (Bohbot & Dickens, 2009). Therefore, one might expect to see stronger, as yet undefined, behavioural responses of *Aedes* and *Culex* species mosquitoes to the (R) enantiomer. However, this has not been investigated to date and is further complicated by the fact the *Ae. aegypti* and *Cx. quinquefasciatus* have different host preferences. *Aedes aegypti* is a known anthropophilic mosquito that responds preferentially to human host odours, as well as other mammals, and *Cx. quinquefasciatus* preferentially feeds on birds (Takken & Kline, 1989; Takken, 1991; Zinser *et al.*, 2004). 1-Octen-3-ol has been identified from human odour previously; therefore, *Ae. aegypti* would be expected to show

positive behavioural responses to the compound. To our knowledge, 1-octen-3-ol has not been identified from bird odour, and so its ecological role is difficult to predict for *Cx. quinquefasciatus* mosquitoes.

The aim of this study was to further our understanding on the role of 1-octen-3-ol and to provide possible explanations for the inconsistency in trapping efficacy of previous studies. To achieve this, we investigate the behavioural and electrophysiological effect of pure enantiomers of 1-octen-3-ol and mixtures on the attraction of *Ae. aegypti* and *Cx. quinquefasciatus* mosquitoes using laboratory based olfactometer bioassays and electrophysiology employing single-sensillum recordings (SSRs).

Materials and methods

Behaviour

Insects

Aedes aegypti mosquitoes, refm strain, were originally obtained from the London School of Hygiene and Tropical Medicine and were kept in culture in the laboratory at Rothamsted Research. The culture was maintained in a controlled environment at 27 ± 2°C, 55–60% relative humidity (RH) and a L:D 12:12 h photoperiod. The larvae were fed daily with Tetramin[®] Tropical Fish Flakes according to larval density of rearing trays. Adult insects were kept in BugDorm-1 (MegaView, Taiwan) plastic rearing cages and provided with 10% sucrose solution (Fischer Scientific UK Ltd, UK) on cotton wool. Females were blood-fed weekly on sheep blood in Alsevers (TCS Biosciences, UK) using a Hemotek[®] Membrane Feeding System (Discovery Workshops, UK).

Culex quinquefasciatus mosquitoes were originally obtained from the Thai strain provided by the Liverpool School of Tropical Medicine. Similar to *Ae. aegypti*, this culture was maintained at Rothamsted Research in a controlled environment at 27 ± 2°C, 55–60% RH, although the photoperiod was inverted, D:L 12:12 h. Larvae were fed regularly with Brewer's Yeast Tablets (500 mg) (Holland & Barrett, UK). Adults were kept in BugDorm-1 rearing cages and were fed on defibrinated horse blood (TCS Biosciences, UK) weekly.

For behavioural bioassays, female mosquitoes aged between 5–12 days that had not been previously blood-fed, were collected and stored in plastic pre-release chambers (WHO, Malaysia), without sucrose and water, in the bioassay room for two hours, allowing time for the insects to acclimatize.

Behavioural assays

A Y-tube olfactometer (Logan *et al.*, 2008), comprising a Y-shaped Plexiglass tube (id 62 mm; fig. 1B–D) that allows insects to move towards odours coming from the upwind end, was used for the behavioural bioassays. Air was pumped through an activated charcoal filter then a glass bell jar (5 l), containing distilled water (dH₂O, 1 l; fig. 1G–I) prior to being split between the two arms of the olfactometer. Flow meters regulated the airflow in each arm at 101 min⁻¹ (fig. 1F). Chemicals in solvent (hexane) were introduced onto filter paper (10 µl; Whatman No. 1, 9-cm diameter filter papers, Whatman plc, UK) by micropipette (Drummond, UK) and placed in sintered glass expander/reductor couplings (fig. 1E). A second pump provided a charcoal filtered airflow over the chemical samples, regulated at 20 ml min⁻¹, and this was introduced through polytetrafluoroethylene (PTFE) tubing




Fig. 1. Y-tube olfactometer bioassay design. Female insects were collected in pre-release chambers (A) and allowed to acclimatize for two hours. These chambers were designed to easily attach to the main Y-tube olfactometer (B). Air was pumped (I) through a charcoal filter (H) and humidified in a glass bell jar containing dH₂O (G). The airflow was regulated by flowmeters (F). Chemicals were applied to filter papers and placed in sintered glass vessels (E). Where necessary, CO₂ was introduced from a cylinder located in the bioassay room (J).

adjacent to the main airflow. For dose response experiments, chemicals were introduced in the presence of CO₂ presented via a pressurized cylinder (BOC, UK) and introduced directly into the main airflow before the split, or in the required tubing arm after the split (fig. 1). Airflow was regulated by a flow meter (20 ml min⁻¹) and was balanced in the opposite arm of the Y-tube by the addition of another pump introducing charcoal filtered air at the same flow rate (fig. 1). The Y-tube olfactometer and components were changed after every third treatment and cleaned thoroughly using distilled water and Tepol (Harvey, Waddington, UK) then allowed to air dry. Additionally, the PTFE tubing was rinsed with ethanol, and the sintered glass vessels were cleaned with acetone and distilled water. Both the tubing and vessels were then placed for two hours at 180°C in an oven.

In all experiments, three behavioural parameters were recorded: relative attraction (the proportion of mosquitoes in the arm of the Y-tube containing the odour stimulus out of the total number of mosquitoes in both arms); upwind flight activity (the proportion of mosquitoes that had flown upwind beyond 30 cm in the stem of the Y-tube) and non-sustained flight activity (NSFA; the proportion of mosquitoes that had flown beyond 30 cm in the stem of the Y-tube, but did not reach the capture chambers at the end of the arms).

Dose response experiments

Racemic 1-octen-3-ol, (*R*)-1-octen-3-ol, (*S*)-1-octen-3-ol and an (*R*:*S*)-1-octen-3-ol mixture (84:16) were prepared in hexane and serial diluted ten-fold to provide solutions of 10⁻¹²–10⁻⁴ g μl⁻¹. Aliquots of 10 μl were applied in each replicate, giving doses of 10⁻¹¹–10⁻³ g μl⁻¹ (treatment vs. blank). All treatments were tested with CO₂. In addition to the nine treatment doses, three controls were also included: (i) CO₂ vs. blank, (ii) CO₂ vs. CO₂ (to ensure there was no bias in the Y-tube set up) and (iii) aqueous L-(+)-lactic acid + CO₂

vs. distilled water (dH₂O) (a positive control). Experiments were done using a semi-randomized block design. Treatments were randomized then ordered by ascending dose within the groups that occurred between equipment changes/washes. Twelve replicates were conducted and the randomisation also included orientation of the Y-tube olfactometer, so six replicates were conducted with the test compound in the left arm and six in the right arm.

A pre-release chamber containing approximately 20 female mosquitoes was attached to the release chamber and the door opened to allow the mosquitoes to move. The cover of the visual occlusion frame was replaced, the main airflow turned on and the insects were given five minutes to acclimatize. During this time, the treatments were prepared on filter papers as above, allowing one minute for the solvent to evaporate. After acclimatization, the papers were introduced into the sintered glass vessels, and the secondary airflows were turned on along with the CO₂ and the balancing airflow. Once all flow meters had settled, the insects were released for a duration of 90 s. The swivel doors on the capture chambers and release chambers were closed, and the position of all insects in the stem and arms was recorded. The cover to the visual occlusion frame was removed, and the insects in the capture chambers and release chambers were counted. The CO₂ cylinder, as well as the balancing airflow, was turned off. The insects in the olfactometer were removed by suction. Experiments were conducted in the light with *Ae. aegypti* and under red-light (artificial night) with *Cx. quinquefasciatus*.

Electrophysiology

Insects

Aedes aegypti, Rockefeller strain, and *Culex quinquefasciatus*, Thai strain (obtained as eggs from Rothamsted Research Institute, UK), were used for electrophysiology experiments.

Both mosquito species were cultured at SLU, Alnarp. The cultures were maintained at 27°C, with a relative humidity of 70% at an L:D 12:12 h photoperiod (the photoperiod was inverted for *Cx. quinquefasciatus*, D:L 12:12 h). Adults were maintained in mesh and plastic culture cages with free access to 10% sucrose through a filter paper wick. Larvae were reared in a separate chamber (27°C, 60% RH and 12:12 L:D) in distilled water-filled plastic trays (20 cm × 30 cm × 10 cm). Larvae were kept in groups of ~500 per tray with free access to fish food (Tetramin® Tropical Fish Flakes). Pupae were collected daily in 30 ml containers and transferred to the adult cages. For electrophysiology, non-blood-fed female mosquitoes were chosen at random from a population aged between three and five days.

Insect preparation

To obtain a stable recording, a female mosquito was anesthetized (1.0–1.5 min at –20°C) and mounted over double-sided tape on a microscope slide (76 × 26 mm). To secure the insect, half of the thorax and the entire abdomen were covered by tape fixed to the microscope slide. The cover slip, partially covered with double-sided tape, was positioned abutting the mosquito head, above and at a right angle to the proboscis. Maxillary palps were lifted and extended smoothly onto the cover slip. The scales that cover the basiconic sensilla on the maxillary palps were removed by gentle rubbing. The mounted mosquito was placed in the single sensillum recording (SSR) rig under a Nikon Eclipse (E600FN) microscope. The maxillary palps were visualized at high magnification (750×).

Single sensillum recording

Maxillary palps of mosquitoes house three types of olfactory receptor neurons, and their extracellular spikes can easily be resolved into three distinct populations based on differences in amplitude. In order to determine a comparative olfactory response profile for the olfactory receptor neurons to the three forms of 1-octen-3-ol, SSRs were performed on ten maxillary palp basiconic sensilla from each culicine species, *Ae. aegypti* and *Cx. quinquefasciatus*.

Two tungsten electrodes were electrolytically sharpened (~1-µm tip diameter) by dipping repeatedly in a 10% KNO₂ at 5–10 V. The reference electrode was inserted into the eye of the mosquito under low magnification (75×). The recording electrode was connected to a preamplifier (10×, Syntech, Kirchzarten, Germany) and inserted into the base or shaft of a basiconic sensillum to extracellularly record the activity of the olfactory receptor neurons housed within the sensilla. The preamplifier was connected to an analogue to digital signal converter (IDAC, Syntech), which in turn was connected to a PC computer for data acquisition and storage. The data were analyzed with Autospice v.3.0 (Syntech).

Stimuli and stimulus delivery

Seven concentrations of racemic 1-octen-3-ol, (*R*)-1-octen-3-ol and (*S*)-1-octen-3-ol enantiomers were used to describe the dose-response relationships. Each of the three volatiles was dissolved in hexane to make a stock solution of 10 µg µl⁻¹. Aliquots of 10 µl of each serially diluted compound (10⁻¹² g µl⁻¹ – 10⁻⁶ g µl⁻¹) were loaded onto a piece of filter paper (5 × 10 mm) and placed into the glass Pasteur pipette.

Hexane (10 µl) alone was used as a control. All pipettes were prepared in a fume hood and left for 40 min to ensure the evaporation of the hexane solvent.

A continuous humidified air stream delivered at 1.5 l min⁻¹ through a glass tube (10 mm id) passed over the maxillary palp preparation. Stimuli were applied through a hole in the glass tube, which was 11 cm away from the outlet. A stimulus controller (Syntech, Germany) was used to pass 0.5 l min⁻¹ through the stimulus cartridge for 500 ms into the continuous air stream through the hole in the glass tube. Fresh stimulus cartridges were applied for each new replicate. Throughout this study, synthetic CO₂-free air (Scan Gas, Sweden) was used for the main airflow and stimulus delivery. One-to-two recordings were taken from each mounted insect.

Chemicals

Racemic 1-octen-3-ol was obtained from Alfa-Aeser (Heysham, Lancashire, UK). (*R*)-1-Octen-3-ol and (*S*)-1-octen-3-ol were provided by Botanix Ltd (Paddock Wood, Kent, UK), and Agrisense BCS Ltd (Pontypridd, Gwent, UK). The enantiomeric purity/composition of materials was determined by derivitization to respective butyrate esters using pyridine and butyric anhydride. The esters were then analysed by high resolution gas chromatography (GC), using an Agilent 6890 GC equipped with a β-cyclodextrin column (25 m × 0.32 mm id × 0.25 µm film thickness), cool on-column injector and a flame ionization detector (FID). The following oven temperature programme was used: 40°C per 1 min, then 3°C min⁻¹ to 150°C, hold for 0.1 min, then 5°C min⁻¹ to 180°C, hold for 15 min. The derivatized enantiomers resolved clearly and their composition was determined by analysis of the resultant peak areas. Enantiomeric purity was determined to be: (*R*)-(-)-1-octen-3-ol, 99%; (*S*)-(+)-1-octen-3-ol, 99%; (*R*:*S*)-1-octen-3-ol, 84:16 and racemic 1-octen-3-ol (50:50).

Statistical analysis

For behavioural data, a generalized linear model (GLM) with binomial distribution and logistic regression was used to perform pairwise comparisons between the CO₂ or blank control and the test treatments (Genstat®, 12th edn). The mean proportions of mosquitoes responding to each treatment were compared. Initially, the model blocked by day and layered out each of the controls to determine where effects were occurring. Following this, the model was simplified to only block by day to compare all treatments. Predictions, taking into account total variance, were produced along with standard errors and Least Significant Differences (LSDs). Results are shown with significance levels ($P < 0.05$, $P < 0.01$, $P < 0.001$).

The electrophysiological data were analyzed using a two-way repeated measures ANOVA followed by Bonferroni post test (Graph Pad Prism vs. 5.0a).

Results

Behaviour

Relative attraction

To determine the optimum concentration and composition of 1-octen-3-ol enantiomers to elicit a behavioural response, behavioural bioassays were performed in the laboratory. In all cases, with *Ae. aegypti*, the positive control (L-(+)-lactic




Fig. 2. Relative attraction of *Aedes aegypti* female mosquitoes in a Y-tube olfactometer to 1-octen-3-ol treatments (plus CO₂) and controls: CO₂ vs. blank, CO₂ vs. CO₂ and L-(+)-lactic acid + CO₂ vs. dH₂O. Relative attraction is the proportion of mosquitoes that flew upwind and were caught in the capture chamber of the treated arm of the Y-tube olfactometer. (a) racemic 1-octen-3-ol, (b) (S)-1-octen-3-ol, (c) (R)-1-octen-3-ol and (d) (R:S)-1-octen-3-ol (84:16) mixture. Asterisks show statistically significant differences in response compared to the CO₂ vs. blank control. (GLM, * $P < 0.05$, $n = 12$).

acid + CO₂ vs. dH₂O) produced significantly greater relative attraction (the proportion of mosquitoes in the arm of the Y-tube containing the test odour stimulus) compared with the CO₂ vs. blank control (fig. 2). Relative attraction in response to the (R)-1-octen-3-ol enantiomer was significantly greater than the CO₂ vs. blank control for two doses, 10^{-5} g μl^{-1} ($P = 0.019$) and 10^{-7} g μl^{-1} ($P = 0.023$) (fig. 2c).

No significant differences were found between the positive control and CO₂ vs. blank control for *Cx. quinquefasciatus* mosquitoes (fig. 3). The (R)-1-octen-3-ol enantiomer caused a significantly lower relative attraction than the CO₂ vs. blank control at one dose 10^{-3} g μl^{-1} ($P < 0.001$) (fig. 3c). A similar result was noted with racemic 1-octen-3-ol and the (R:S)-1-octen-3-ol mixture (84:16) in which the relative attraction to the 10^{-3} g μl^{-1} dose was also significantly lower than the CO₂ vs. blank control ($P < 0.001$ and $P = 0.014$, respectively) (fig. 3a,d). No statistically significant differences were found for relative attraction with the (S)-1-octen-3-ol enantiomer (fig. 3b).

Flight activation

For flight activation (the proportion of mosquitoes that had flown upwind beyond 30 cm in the stem of the Y-tube) with *Ae. aegypti*, no significant differences were found for the racemic 1-octen-3-ol or either enantiomer when

compared with the CO₂ vs. blank control ($P > 0.05$) (table 1). The (R:S)-1-octen-3-ol mixture (84:16) showed significantly less activation at 10^{-8} g μl^{-1} ($P = 0.030$) (table 1).

With *Cx. quinquefasciatus*, no significant differences were found for racemic 1-octen-3-ol ($P > 0.05$) (table 1). Activation was significantly greater than the CO₂ vs. blank control for (R)-1-octen-3-ol at 10^{-4} g μl^{-1} ($P = 0.027$) and 10^{-6} g μl^{-1} ($P = 0.048$) (table 1). It was noted, generally, that these mosquitoes elicited a lower activation response compared to *Ae. aegypti*.

No significant differences between the positive control and the CO₂ vs. blank control were noted for either mosquito species in relation to flight activation.

Non-sustained flight activity (NSFA)

In all the experiments with *Ae. aegypti*, the positive control (L-(+)-lactic acid + CO₂ vs. dH₂O) produced significantly lower NSFA (the proportion of mosquitoes that had flown beyond 30 cm in the stem of the Y-tube but did not reach the capture chambers at the end of the arms) compared with the CO₂ vs. blank control (table 2). The (R:S)-1-octen-3-ol mixture (84:16) elicited a significantly greater NSFA than the CO₂ vs. blank control at the 10^{-7} g μl^{-1} dose ($P = 0.011$) (table 2). No significance was found when using (S)-1-octen-3-ol ($P > 0.05$).




Fig. 3. Relative attraction of *Culex quinquefasciatus* female mosquitoes in a Y-tube olfactometer to 1-octen-3-ol treatments (plus CO₂) and controls: CO₂ vs. blank, CO₂ vs. CO₂ and L-(+)-lactic acid + CO₂ vs. dH₂O. Relative attraction is the proportion of mosquitoes that flew upwind and were caught in the capture chamber of the treated arm of the Y-tube olfactometer. (a) racemic 1-octen-3-ol, (b) (S)-1-octen-3-ol, (c) (R)-1-octen-3-ol and (d) (R:S)-1-octen-3-ol (84:16) mixture. Asterisks show statistically significant differences in response compared to the CO₂ vs. blank control. (GLM, **P* < 0.05, ****P* < 0.001, *n* = 12).

Table 1. Flight activation of *Aedes aegypti* and *Culex quinquefasciatus* female mosquitoes in a Y-tube olfactometer to 1-octen-3-ol treatments (plus CO₂) and controls: CO₂ vs. blank, CO₂ vs. CO₂ and L-(+)-lactic acid + CO₂ vs. dH₂O.

Activation	<i>Aedes aegypti</i>				<i>Culex quinquefasciatus</i>			
	Racemic	S	R	R:S	Racemic	S	R	R:S
CO vs. CO ₂	0.72 ± 0.05	0.69 ± 0.05	0.70 ± 0.04	0.69 ± 0.05	0.49 ± 0.05	0.37 ± 0.05	0.49 ± 0.05	0.42 ± 0.05
L-(+)-lactic acid	0.79 ± 0.04	0.78 ± 0.04	0.77 ± 0.04	0.85 ± 0.05	0.68 ± 0.05	0.54 ± 0.05	0.69 ± 0.04	0.54 ± 0.05
CO ₂ vs. blank	0.74 ± 0.05	0.75 ± 0.04	0.68 ± 0.04	0.77 ± 0.05	0.56 ± 0.05	0.41 ± 0.05	0.58 ± 0.05	0.37 ± 0.05
10 ⁻³	0.68 ± 0.05	0.75 ± 0.04	0.66 ± 0.04	0.69 ± 0.05	0.62 ± 0.05	0.49 ± 0.05	0.62 ± 0.05	0.48 ± 0.05
10 ⁻⁴	0.65 ± 0.05	0.75 ± 0.05	0.59 ± 0.04	0.72 ± 0.05	0.57 ± 0.05	0.44 ± 0.05	0.72 ± 0.04*	0.46 ± 0.05
10 ⁻⁵	0.63 ± 0.05	0.83 ± 0.04	0.62 ± 0.04	0.78 ± 0.05	0.62 ± 0.05	0.48 ± 0.05	0.66 ± 0.05	0.41 ± 0.05
10 ⁻⁶	0.73 ± 0.05	0.81 ± 0.04	0.74 ± 0.04	0.70 ± 0.05	0.63 ± 0.05	0.43 ± 0.05	0.71 ± 0.04*	0.37 ± 0.05
10 ⁻⁷	0.74 ± 0.05	0.80 ± 0.04	0.72 ± 0.04	0.70 ± 0.05	0.55 ± 0.05	0.53 ± 0.05	0.55 ± 0.05	0.45 ± 0.05
10 ⁻⁸	0.69 ± 0.05	0.73 ± 0.04	0.71 ± 0.04	0.61 ± 0.06*	0.55 ± 0.05	0.38 ± 0.05	0.59 ± 0.05	0.36 ± 0.05
10 ⁻⁹	0.73 ± 0.05	0.83 ± 0.04	0.76 ± 0.04	0.64 ± 0.05	0.61 ± 0.05	0.51 ± 0.05	0.65 ± 0.05	0.49 ± 0.05
10 ⁻¹⁰	0.69 ± 0.05	0.83 ± 0.04	0.73 ± 0.04	0.63 ± 0.05	0.63 ± 0.05	0.52 ± 0.05	0.60 ± 0.05	0.37 ± 0.05
10 ⁻¹¹	0.74 ± 0.05	0.84 ± 0.04	0.76 ± 0.04	0.64 ± 0.05	0.49 ± 0.05	0.36 ± 0.04	0.55 ± 0.05	0.33 ± 0.05

Flight activation is the proportion of mosquitoes that were activated and flew upwind leaving the release chamber of the Y-tube olfactometer in response to racemic 1-octen-3-ol, (S)-1-octen-3-ol, (R)-1-octen-3-ol and (R:S)-1-octen-3-ol (84:16) mixture. Data are presented as mean ± standard error. Asterisks show statistically significant differences in response compared to the CO₂ vs. blank control. (GLM, * *P* < 0.05, *n* = 12).

(table 2). The (R)-1-octen-3-ol produced significantly lower NSFA at 10⁻⁶ g μl⁻¹ (*P* = 0.017) and 10⁻⁹ g μl⁻¹ (*P* = 0.047) (table 2).

With *Cx. quinquefasciatus* mosquitoes, the positive control (L-(+)-lactic acid + CO₂ vs. dH₂O) produced no significant differences in NSFA compared with the CO₂ vs. blank control

Table 2. Non-sustained flight activity (NSFA) of *Aedes aegypti* and *Culex quinquefasciatus* female mosquitoes in a Y-tube olfactometer to 1-octen-3-ol treatments (plus CO₂) and controls: CO₂ vs. blank, CO₂ vs. CO₂ and L-(+)-lactic acid + CO₂ vs. dH₂O. NSFA is the proportion of mosquitoes that were activated and flew upwind leaving the release chamber, however, failed to reach the capture chambers of the Y-tube olfactometer in response to racemic 1-octen-3-ol, (S)-1-octen-3-ol, (R)-1-octen-3-ol and (R:S)-1-octen-3-ol (84:16) mixture.

Non-sustained flight activity Treatment/dose (g µl ⁻¹)	<i>Aedes aegypti</i>				<i>Culex quinquefasciatus</i>			
	Racemic	S	R	R:S	Racemic	S	R	R:S
CO ₂ vs. CO ₂	0.41 ± 0.05	0.49 ± 0.05	0.51 ± 0.05	0.53 ± 0.05	0.51 ± 0.06	0.65 ± 0.06	0.63 ± 0.06	0.35 ± 0.06
L-(+)-lactic acid	0.36 ± 0.04	0.24 ± 0.04	0.29 ± 0.05	0.30 ± 0.05	0.66 ± 0.04	0.68 ± 0.05	0.67 ± 0.04	0.31 ± 0.05
CO ₂ vs. blank	0.50 ± 0.05	0.58 ± 0.04	0.66 ± 0.05	0.53 ± 0.05	0.70 ± 0.05	0.72 ± 0.05	0.73 ± 0.05	0.58 ± 0.07
10 ⁻³	0.58 ± 0.05	0.58 ± 0.04	0.62 ± 0.05	0.65 ± 0.05	0.75 ± 0.04	0.59 ± 0.05	0.78 ± 0.04	0.31 ± 0.05**
10 ⁻⁴	0.57 ± 0.05	0.58 ± 0.05	0.53 ± 0.06	0.55 ± 0.05	0.72 ± 0.05	0.75 ± 0.05	0.74 ± 0.04	0.39 ± 0.06*
10 ⁻⁵	0.61 ± 0.05	0.60 ± 0.04	0.53 ± 0.06*	0.63 ± 0.05	0.59 ± 0.05	0.64 ± 0.05	0.64 ± 0.05	0.37 ± 0.06*
10 ⁻⁶	0.49 ± 0.05	0.51 ± 0.04	0.48 ± 0.05	0.55 ± 0.05	0.74 ± 0.04	0.71 ± 0.05	0.73 ± 0.04	0.46 ± 0.07
10 ⁻⁷	0.48 ± 0.05	0.61 ± 0.04	0.61 ± 0.05	0.70 ± 0.05*	0.60 ± 0.05	0.69 ± 0.05	0.65 ± 0.05	0.26 ± 0.05***
10 ⁻⁸	0.49 ± 0.05	0.59 ± 0.04	0.54 ± 0.05	0.64 ± 0.05	0.68 ± 0.05	0.75 ± 0.05	0.80 ± 0.04	0.52 ± 0.07
10 ⁻⁹	0.52 ± 0.05	0.63 ± 0.04	0.51 ± 0.05*	0.52 ± 0.05	0.74 ± 0.04	0.64 ± 0.05	0.76 ± 0.04	0.40 ± 0.06*
10 ⁻¹⁰	0.49 ± 0.05	0.52 ± 0.04	0.64 ± 0.05	0.60 ± 0.05	0.74 ± 0.04	0.68 ± 0.05	0.67 ± 0.05	0.36 ± 0.07*
10 ⁻¹¹	0.46 ± 0.05	0.61 ± 0.04	0.70 ± 0.05	0.52 ± 0.05	0.71 ± 0.05	0.71 ± 0.06	0.74 ± 0.05	0.38 ± 0.07*

Data are presented as mean ± standard error. Asterisks show statistically significant differences in response compared to the CO₂ vs. blank control. (GLM, **P* < 0.05, ***P* < 0.01, ****P* < 0.001, *n* = 12).

except for the experiment where the 1-octen-3-ol mixture (84:16 R:S) was tested, in which the response to L-(+)-lactic acid + CO₂ was significantly lower (*P* = 0.002) (table 2). No significant differences in NSFA were found using the racemic 1-octen-3-ol or the (R)-1-octen-3-ol enantiomer (*P* > 0.05). The (R:S)-1-octen-3-ol mixture (84:16) elicited a significant decrease in NSFA compared to the CO₂ vs. blank control at most doses: 10⁻³ g µl⁻¹ (*P* = 0.003); 10⁻⁴ g µl⁻¹ (*P* = 0.035); 10⁻⁵ g µl⁻¹ (*P* = 0.03); 10⁻⁷ g µl⁻¹ (*P* < 0.001); 10⁻⁹ g µl⁻¹ (*P* = 0.05); 10⁻¹⁰ g µl⁻¹ (*P* = 0.026) and 10⁻¹¹ g µl⁻¹ (*P* = 0.047) (table 2).

Electrophysiology

The B cell, i.e. the olfactory receptor neuron with an intermediate amplitude, of the maxillary palp basiconic sensilla of both species responded to racemic 1-octen-3-ol, as well as the two enantiomers in a dose dependent manner (fig. 4).

In both species, the B neuron showed a significant difference in selectivity between the enantiomers, with (R)-1-octen-3-ol eliciting a greater response, followed by racemic 1-octen-3-ol then (S)-1-octen-3-ol (fig. 4). For *Ae. aegypti*, (R)-1-octen-3-ol produced significant differences compared to racemic 1-octen-3-ol at 10⁻⁹ g µl⁻¹, 10⁻⁸ g µl⁻¹ and 10⁻⁷ g µl⁻¹ (*P* = 0.047, *P* = 0.002 and *P* = 0.01); whereas, significant differences were only found at 10⁻⁷ g µl⁻¹ (*P* = 0.032) with *Cx. quinquefasciatus* (fig. 4b, d). In both species, the racemic 1-octen-3-ol elicited a significantly greater response compared to (S)-1-octen-3-ol at 10⁻⁸ g µl⁻¹, 10⁻⁷ g µl⁻¹ and 10⁻⁶ g µl⁻¹ (*Ae. aegypti*: *P* = 0.0005, *P* = 0.0002, *P* = 0.004; *Cx. quinquefasciatus*: *P* = 0.001, *P* = 0.0003 and *P* = 0.004, respectively). Similarly, comparison of responses to the enantiomers showed that (R)-1-octen-3-ol produced significantly greater responses in both species at 10⁻⁹ g µl⁻¹, 10⁻⁸ g µl⁻¹, 10⁻⁷ g µl⁻¹ and 10⁻⁶ g µl⁻¹ (*Ae. aegypti*: *P* = 0.003, *P* = 0.0001, *P* = 0.0001, *P* = 0.0008; *Cx. quinquefasciatus*: *P* = 0.005, *P* = 0.0003, *P* = 0.0001 and *P* = 0.001, respectively). No significant differences were found for any treatments below 10⁻⁹ g µl⁻¹.

Discussion

In this study, we investigated the behavioural and electrophysiological responses of *Ae. aegypti* and *Cx. quinquefasciatus* mosquitoes to (R)-1-octen-3-ol, (S)-1-octen-3-ol and to mixtures of the enantiomers in combination with CO₂. Previous studies have investigated the role of racemic 1-octen-3-ol in mosquito attraction. However, few investigations have examined the effect of the individual enantiomers. In our study, clear electrophysiological and behavioural differences between *Ae. aegypti* and *Cx. quinquefasciatus* were recorded in response to the enantiomers.

Aedes aegypti maxillary palp ORNs detected and discriminated between the (R) and (S) enantiomers at a stimulus concentration of 10⁻⁹ g µl⁻¹ and above and discerned (R)-1-octen-3-ol from the racemic mixture at concentrations of 10⁻⁹ g µl⁻¹, 10⁻⁸ g µl⁻¹ and 10⁻⁷ g µl⁻¹. However, only the (R)-1-octen-3-ol enantiomer elicited a behavioural response in the form of significant increase in relative attraction and a reduction in non-sustained flight activity for *Ae. aegypti*. The (R:S)-1-octen-3-ol mixture (84:16), based on ratios from cattle breath (Hall *et al.*, 1984), failed significantly to attract either species. However, this could be due to the fact that *Ae. aegypti* would not normally feed on cattle; and, therefore, it is possible that a different species showing a greater preference for cattle (e.g. certain *Anopheles* spp.) may elicit significant behavioural results (Takken, 1991). All treatments had little effect in terms of flight activation; however, it was noted that, across all experiments, *Ae. aegypti* showed a greater average flight activation compared with *Cx. quinquefasciatus*. (R)-1-Octen-3-ol, (S)-1-octen-3-ol, racemic 1-octen-3-ol and the mixture of the 1-octen-3-ol enantiomers did not elicit a significant relative attraction by *Cx. quinquefasciatus*. These results suggest that *Ae. aegypti* has greater selectivity of the (R) enantiomer at the electrophysiological level and has the most pronounced effect on behaviour, causing activated mosquitoes to sustain their flight and orientate towards the compound. This is the first time electrophysiological selectivity has been shown for a 1-octen-3-ol enantiomer in *Ae. aegypti* *in vivo*. Bohbot & Dickens (2009) demonstrated that the *in vitro* expression of AaOR8 in *Xenopus* oocytes shows a strong




Fig. 4. Dose response relationships for the basiconic sensillum on the maxillary palp in *Aedes aegypti* and *Culex quinquefasciatus* female mosquito. 1-Octen-3-ol and its enantiomers elicit dose-dependent responses in the B neuron of basiconic sensilla on the maxillary palp of *Ae. aegypti* and *Cx. quinquefasciatus* female mosquitoes. (a), (c) Response of the B neuron to (R)-1-octen-3-ol and the A neuron (open circle) to ambient CO₂ trapped in the stimulus cartridge in *Ae. aegypti* and *Cx. quinquefasciatus*, respectively. Closed circles indicate the smallest spiking neuron, the C neuron. Unmarked intermediate response spikes are from the B neuron. (b), (d) The B neurons in *Ae. aegypti* and *Cx. quinquefasciatus* express significantly higher responses to (R)-1-octen-3-ol as compared to (S)-1-octen-3-ol. (two-way repeated measures ANOVA; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$). Significant differences are also found between the response to racemic 1-octen-3-ol and its enantiomers (see results), $n = 10$; mean \pm SEM (\diamond , S (+)1-octen-3-ol *Aedes*; \square , R (-)1-octen-3-ol *Aedes*; \blacklozenge , S (+)1-octen-3-ol *Culex*; \blacksquare , R (-)1-octen-3-ol *Culex*; \blacktriangle , (R+S)1-octen-3-ol *Culex*).

preference towards the (R)-1-octen-3-ol enantiomer, which supports our study.

To our knowledge, no other investigation has examined the effect of the 1-octen-3-ol enantiomers on *Ae. aegypti* behaviour, and few studies have examined the behavioural effect on other mosquitoes species. In the field, Kline *et al.* (2007) demonstrated that traps baited with (R)-1-octen-3-ol, when tested in combination with CO₂, caught greater numbers of mosquitoes (*Anopheles crucians*, *Culex salinarius*) compared to (S)-1-octen-3-ol, which performed equally or less well than CO₂ alone. No significant increase in trap catch of *Cx. quinquefasciatus* was recorded. Numerous other studies have reported no significant behavioural effects with racemic 1-octen-3-ol and *Culex* spp. For example, *Culex sitiens* and *Culex annulirostris* have been reported to show either no response or a repellent response to racemic 1-octen-3-ol (Kempe *et al.*, 1993; Van Essen *et al.*, 1994). In Tanzania, a study using unlit CDC traps and racemic 1-octen-3-ol found that the number of *Cx. quinquefasciatus* caught did not differ significantly from the unbaited control (Mboera *et al.*, 2000). Similarly, studies of other arthropods have shown a lack of preference for the racemic mixture or enantiomers of 1-octen-3-ol. For example, the bont tick, *Amblyomma variegatum*, has been shown to display the same orientated movements towards the racemic mixture and (R)-1-octen-3-ol using a servosphere bioassay (McMahon *et al.*, 2001). In a pitfall olfactometer, an equivalent increase in catches of grain beetle, *Oryzaephilus mercator*, was

recorded to the separate 1-octen-3-ol enantiomers and the racemic mixture in comparison to the controls (Pierce *et al.*, 1989).

Culex quinquefasciatus mosquitoes possess ORNs that detect 1-octen-3-ol, and these are located in the basiconic sensilla of the maxillary palps (B neuron). In previous studies, the B cells in maxillary palp sensilla have been shown to have remarkable selectivity, being more sensitive to the (R)-1-octen-3-ol enantiomer than the (S) enantiomer (Syed & Leal, 2007). In *An. gambiae* it is also the B cell that responds to 1-octen-3-ol (Lu *et al.*, 2007), which suggests that this olfactory receptor neuron (ORN) could be conserved throughout the mosquito lineage. Indeed, in our electrophysiological experiments with *Aedes*, we found supporting evidence for this and the B cell responded in a similar manner, discriminating between the (R) and (S) enantiomers at a stimulus concentration of 10^{-9} g μl^{-1} and above and discerning (R)-1-octen-3-ol from the racemic mixture at a concentration of 10^{-7} g μl^{-1} .

Although the (R)-1-octen-3-ol enantiomer caused an increase in activation for *Cx. quinquefasciatus* at two doses, the most notable effect was that the (R:S)-1-octen-3-ol mixture (84:16) at seven of the doses tested caused significantly more mosquitoes to sustain their flight and reach the capture chambers (demonstrated by a reduced NSFAs), suggesting that it may have a behaviourally excitatory effect. Additionally, at the highest concentration, the few mosquitoes that did reach the capture chambers moved towards the control chamber

rather than the arm containing the (*R*)-1-octen-3-ol enantiomer. Interestingly, this reduced relative attraction response was observed for all treatments containing the (*R*)-1-octen-3-ol enantiomer either on its own or as part of a mixture, but not with the (*S*)-1-octen-3-ol enantiomer. It is unknown whether this effect is true repellency, as the design of the Y-tube olfactometer bioassay does not allow measurement of such behaviour. Similarly, one laboratory study found that CO₂ plus racemic 1-octen-3-ol caused a repellent effect in *Anopheles quadrimaculatus* (Pates *et al.*, 2005).

The behavioural differences observed here between *Aedes* and *Culex* mosquitoes may be due to feeding preferences as, although both species are known to feed on humans (Takken, 1991; Zinser *et al.*, 2004), *Cx. quinquefasciatus* can feed opportunistically, and it is preferentially an ornithophilic species (Takken & Kline, 1989). These results are supported by previous research which shows differences in levels of attraction between mosquito species. For example, one study, using modified Fay-Prince traps to test the effects of CO₂ and racemic 1-octen-3-ol on catches of *Ae. aegypti*, found that it had no effect on increasing trap catches over CO₂ alone (Canyon & Hii, 1997). In contrast, other studies, using racemic 1-octen-3-ol with encephalitis vector surveillance (EVS) traps, showed that 1-octen-3-ol combined with CO₂ considerably enhanced catches of *Aedes vigilax* and *Aedes funereus* (Kempe *et al.*, 1993; Van Essan *et al.*, 1994). Using CDC light traps in a salt marsh area of North Carolina, USA, it was found that *Aedes sollicitans*, *Aedes taeniorhynchus*, *Anopheles bradleyi* and *Culex salinarius* were significantly attracted to racemic 1-octen-3-ol in combination with CO₂ and light. Additionally, *Anopheles atropos* showed a highly significant response regardless of the presence of light. Interestingly, when they repeated the experiment in the creek flood plain, none of the 24 species collected were significantly attracted to racemic 1-octen-3-ol in combination with CO₂ and light compared to CO₂ and light alone (Rueda *et al.*, 2001). More recently, CDC Fay-Prince traps have been used in Maryland, USA, to test the effects of CO₂ and racemic 1-octen-3-ol on catches of *Aedes albopictus*. It was found that, while traps baited with racemic 1-octen-3-ol and CO₂ generally caught more *Ae. albopictus* mosquitoes, this was not significantly different from CO₂ alone (Shone *et al.*, 2003). The reason for such varied responses in field trapping could be due to the fact that the racemic form of 1-octen-3-ol does not provide an appropriate cue and therefore results in inconsistent trap catches.

Conclusion

This study investigated the effects of enantiomers of 1-octen-3-ol using laboratory behaviour bioassays and SSR. In summary, *Ae. aegypti* mosquitoes differentiate the 1-octen-3-ol enantiomers at the electrophysiological level and make oriented movements towards the (*R*)-1-octen-3-ol enantiomer and do not respond in the same way to either the (*S*)-1-octen-3-ol enantiomer or to any mixture tested. Flight activation is little affected by the treatments, although a negative behavioural effect was observed in response to the 84:16 (*R*:*S*)-1-octen-3-ol mixture. Therefore, the (*R*)-1-octen-3-ol could be a more important kairomone and would be the obvious choice for incorporation into a trap for *Ae. aegypti*. *Culex* mosquitoes also distinguished between the treatments at the electrophysiological level and appeared to be more sensitive to the (*R*) enantiomer. However, the associated

behavioural response was not characteristic of a kairomone. Instead, the mosquitoes made oriented movements away from the (*R*)-1-octen-3-ol enantiomer at high concentrations. Although the mosquitoes were also more activated by this enantiomer than the (*S*)-1-octen-3-ol enantiomer or any mixture tested, this could be due to an excitatory response and is not necessarily conducive to the development of a suitable attractant for traps for this species. The fact that the SSR response of *Culex* mosquitoes to the (*R*)-1-octen-3-ol enantiomer plateaued at the highest concentration suggests that the ORNs became saturated, which may reflect the negative behavioural response observed to this compound at the high concentrations. However, it is worth noting that the behavioural response to the CO₂ control was high compared with *Ae. aegypti* even though their overall activation was lower. This could be due to differences in their perception of CO₂ (as may be expected due to their feeding habits) and may have masked any fine attractant effects caused by the 1-octen-3-ol treatments tested. Further field evaluation of the enantiomers is undoubtedly required. However, the (*R*)-1-octen-3-ol enantiomer could be used for selective trapping for research purposes in areas where *Ae. aegypti* and *Cx. quinquefasciatus* are prevalent. With some further investigation, it may even be possible to use (*R*)-1-octen-3-ol to interfere with host seeking behaviour of *Cx. quinquefasciatus* mosquitoes.

Acknowledgements

This project was supported by a DTI-LINK grant (J0018 K). We thank Stephen Powers for advice on the statistical analysis and Göran Birgersson for technical help. Rothamsted receives grant-aided support from the Biotechnology and Biological Sciences Research Council (BBSRC) of the United Kingdom. This study was also supported by the Linnaeus-program Insect Chemical Ecology, Ethology and Evolution IC-E³.

References

- Becker, N., Zgomba, M., Petric, D. & Ludwig, M. (1995) Comparison of carbon dioxide, octenol and a host-odour as mosquito attractants in the Upper Rhine Valley, Germany. *Medical and Veterinary Entomology* **9**, 377–380.
- Bohbot, J.D. & Dickens, J.C. (2009) Characterization of an enantioselective odourant receptor in the yellow fever mosquito *Aedes aegypti*. *PLoS One* **4**, e7032.
- Burkett, D.A., Lee, W.J., Lee, K.W., Kim, H.C., Lee, H.I., Lee, J.S., Shin, E.H., Wirtz, R.A., Cho, H.W., Claborn, D.M., Coleman, R.E. & Klein, T.A. (2001) Light, carbon dioxide, and octenol-baited mosquito trap and host seeking activity evaluations for mosquitoes in a malarious area of the Republic of Korea. *Journal of the American Mosquito Control Association* **17**, 196–205.
- Bursell, E. (1984) Effects of host odour on the behaviour of tsetse. *Insect Science and its Application* **5**, 345–349.
- Canyon, D.V. & Hii, J.L.K. (1997) Efficacy of carbon dioxide, 1-octen-3-ol, and lactic acid in modified Fay-Prince traps as compared to man-landing catch of *Aedes aegypti*. *Journal of the American Mosquito Control Association* **13**, 66–70.
- Eiras, A.E. & Jepson, P.C. (1994) Responses of female *Aedes aegypti* (Diptera: Culicidae) to host odours and convection currents using an olfactometer bioassay. *Bulletin of Entomological Research* **84**, 207–211.

- Gibson, G. & Torr, S.J. (1999) Visual and olfactory responses of haematophagous Diptera to host stimuli. *Medical and Veterinary Entomology* **13**, 2–23.
- Hall, D.R., Beevor, P.S., Cork, A., Nesbitt, B.F. & Vale, G.A. (1984) 1-Octen-3-ol a potent olfactory stimulant and attractant for tsetse isolated from cattle odours. *Insect Science and its Application* **5**, 335–339.
- Hill, S.R., Hansson, B.S. & Ignell, R. (2009) Characterization of antennal trichoid sensilla from female southern house mosquito, *Culex quinquefasciatus* Say. *Chemical Senses* **34**(3), 231–252.
- Kemme, J.A., Van Essen, P.H.A., Ritchie, S.A. & Kay, B.H. (1993) Response of mosquitoes to carbon dioxide and 1-octen-3-ol in southeast Queensland, Australia. *Journal of the American Mosquito Control Association* **9**, 431–435.
- Khan, A.A., Maibach, H.I., Strauss, W.G. & Fenley, W.R. (1966) Quantitation of effect of several stimuli on the approach of *Aedes aegypti*. *Journal of Economic Entomology* **59**, 690–694.
- Kline, D.L. & Lemire, G.F. (1995) Field evaluation of heat as an added attractant to traps baited with carbon dioxide and octenol for *Aedes taeniorhynchus*. *Journal of the American Mosquito Control Association* **11**, 454–456.
- Kline, D.L., Allan, S.A., Bernier, U.R. & Welch, C.H. (2007) Evaluation of the enantiomers of 1-octen-3-ol and 1-octyn-3-ol as attractants for mosquitoes associated with a freshwater swamp in Florida, U.S.A. *Medical and Veterinary Entomology* **21**, 323–331.
- Knudsen, J.T., Tollsten, L. & Bergstrom, G. (1993) Floral scents – A checklist of volatile compounds isolated by head-space techniques. *Phytochemistry* **33**, 253–280.
- Logan, J.G., Birkett, M.A., Clark, S.J., Powers, S., Seal, N.J., Wadhams, L.J., Mordue, A.J. & Pickett, J.A. (2008) Identification of human-derived volatile chemicals that interfere with attraction of *Aedes aegypti* mosquitoes. *Journal of Chemical Ecology* **34**, 308–322.
- Lu, T., Qiu, Y.T., Wang, G., Kwon, J.Y., Rutzler, M., Kwon, H.W., Pitts, R.J., Van Loon, J.J.A., Takken, W., Carson, J.R. & Zwiebel, L.J. (2007) Odor coding in the maxillary palp of the malaria vector mosquito *Anopheles gambiae*. *Current Biology* **17**, 1533–1544.
- Mboera, L.E.G., Takken, W. & Sambu, E.Z. (2000) The response of *Culex quinquefasciatus* (Diptera: Culicidae) to traps baited with carbon dioxide, 1-octen-3-ol, acetone, butyric acid and human foot odour in Tanzania. *Bulletin of Entomological Research* **90**, 155–159.
- McMahon, C., Guerin, P.M. & Syed, Z. (2001) 1-Octen-3-ol isolated from bont ticks attracts *Amblyomma variegatum*. *Journal of Chemical Ecology* **27**, 471–486.
- Miller, R.J., Wing, J., Cope, S., Davey, R.B. & Kline, D.L. (2005) Comparison of carbon dioxide- and octenol-baited encephalitis virus surveillance mosquito traps at the Shoalwater Bay Training area, Queensland, Australia. *Journal of the American Mosquito Control Association* **21**, 497–500.
- Pates, H.V., Takken, W. & Curtis, C.F. (2005) Laboratory studies on the olfactory behaviour of *Anopheles quadriannulatus*. *Entomologia Experimentalis et Applicata* **114**, 153–159.
- Pierce, A.M., Pierce, H.D. Jr, Borden, J.H. & Oehlschlager, A.C. (1989) Production dynamics of cucujolide pheromones and identification of 1-octen-3-ol as a new aggregation pheromone for *Oryzaephilus surinamensis* and *O. mercator* (Coleoptera: Cucujidae). *Environmental Entomology* **18**, 747–755.
- Rueda, L.M., Harrison, B.A., Brown, J.S., Whitt, P.B., Harrison, R.L. & Gardner, R.C. (2001) Evaluation of 1-octen-3-ol, carbon dioxide, and light as attractants for mosquitoes associated with two distinct habitats in North Carolina. *Journal of the American Mosquito Control Association* **17**, 61–66.
- Russell, R.C. (2004) The relative attractiveness of carbon dioxide and octenol in CDC- and EVS-type light traps for sampling the mosquitoes *Aedes aegypti* (L.), *Aedes polynesiensis* Marks, and *Culex quinquefasciatus* Say in Moorea, French Polynesia. *Journal of Vector Ecology* **29**, 309–314.
- Shone, S.M., Ferrao, P.N., Lesser, C.R., Glass, G.E. & Norris, D.E. (2003) Evaluation of carbon dioxide- and 1-octen-3-ol-baited Centers for Disease Control Fay-Prince traps to collect *Aedes albopictus*. *Journal of the American Mosquito Control Association* **19**, 445–447.
- Syed, Z. & Leal, W.S. (2007) Maxillary palps are broad spectrum odourant detectors in *Culex quinquefasciatus*. *Chemical Senses* **32**, 727–738.
- Takken, W. (1991) The role of olfaction in host-seeking of mosquitoes: a review. *Insect Science and its Application* **12**, 287–295.
- Takken, W. & Kline, D.L. (1989) Carbon dioxide and 1-octen-3-ol as mosquito attractants. *Journal of the American Mosquito Control Association* **5**, 311–316.
- Vale, G.A. & Hall, D.R. (1985a) The role of 1-octen-3-ol, acetone and carbon dioxide in the attraction of tsetse flies, *Glossina* spp. (Diptera: Glossinidae), to ox odour. *Bulletin of Entomological Research* **75**, 209–217.
- Vale, G.A. & Hall, D.R. (1985b) The use of 1-octen-3-ol, acetone and carbon dioxide to improve baits for tsetse flies, *Glossina* spp. (Diptera: Glossinidae). *Bulletin of Entomological Research* **75**, 219–231.
- Van Essen, P.H.A., Kemme, J.A., Ritchie, S.A. & Kay, B.H. (1994) Differential responses of *Aedes* and *Culex* mosquitoes to octenol or light in combination with carbon dioxide in Queensland, Australia. *Medical and Veterinary Entomology* **8**, 63–67.
- Vythilingam, I., Lian, C.G. & Thim, S. (1992) Evaluation of carbon dioxide and 1-octen-3-ol as mosquito attractants. *Southeast Asian Journal of Tropical Medicine and Public Health* **23**, 328–332.
- Zinser, M., Ramberg, F. & Willott, E. (2004) *Culex quinquefasciatus* (Diptera: Culicidae) as a potential West Nile virus vector in Tuscon, Arizona: blood meal analysis indicates feeding on both humans and birds. *Journal of Insect Science* **4**, 20.