

Regulation of the cell cycle and centrosome biology by deubiquitylases.

Sarah Darling, Andrew B Fielding, Dorota Sabat-Pospiech, Ian A Prior, Judy M Coulson*

Cellular and Molecular Physiology, Institute of Translational Medicine, University of Liverpool, Liverpool, L69 3BX, UK

*Corresponding author: Cellular & Molecular Physiology, Institute of Translational Medicine, University of Liverpool, Crown St, Liverpool, L69 3BX, UK. Phone: 44-151-794-5850, Fax: 44-151-794-4434, E-mail: jcoulson@liverpool.ac.uk.

Running title: DUBs in the cell and centrosome cycles

Key words: ubiquitin, post-translational modification, centrosomes, cell cycle, cancer

Subject area: Cell Cycle, Growth & Proliferation, Cancer, Molecular Bases of Health & Disease, Organelles & Localization, Post-Translational Modifications

Abstract

Post-translational modification of proteins by ubiquitylation is increasingly recognised as a highly complex code that contributes to the regulation of diverse cellular processes. In humans, a family of almost one hundred deubiquitylase enzymes (DUBs) are assigned to six sub-families and many of these DUBs can remove ubiquitin from proteins to reverse signals. Roles for individual DUBs have been delineated within specific cellular processes, including many that are dysregulated in diseases, particularly cancer. As potentially druggable enzymes, disease-associated DUBs are of increasing interest as pharmaceutical targets. The biology, structure and regulation of DUBs have been extensively reviewed elsewhere, so here we focus specifically on roles of DUBs in regulating cell cycle processes in mammalian cells. Over a quarter of all DUBs, representing four different families, have been shown to play roles either in the unidirectional progression of the cell cycle through specific checkpoints, or in the DNA damage response and repair pathways. We catalogue these roles and discuss specific examples. Centrosomes are the major microtubule nucleating centres within a cell, and play a key role in forming the bipolar mitotic spindle required to accurately divide genetic material between daughter cells during cell division. To enable this mitotic role, centrosomes undergo a complex replication cycle that is intimately linked to the cell division cycle. Here we also catalogue and discuss DUBs that have been linked to centrosome replication or function, including centrosome clustering, a mitotic survival strategy unique to cancer cells with supernumerary centrosomes.

Reversible ubiquitylation

The post-translational attachment of ubiquitin moieties to substrate proteins, termed ubiquitylation, involves the covalent conjugation of ubiquitin, most commonly to lysine (K) residues. In the simplest form, ubiquitylation is the addition of an ubiquitin monomer, termed monoubiquitylation. However, the ubiquitin signal can be highly complex and is linked to a plethora of cellular processes (1, 2). Polyubiquitin chains linked through K48 target proteins for proteasomal degradation. However, ubiquitin possesses seven lysine residues (K6, K11, K27, K29, K33, K48 and K63) enabling the formation of diverse polyubiquitin chains that may be homotypic or heterotypic in nature, and can have alternative functions, as comprehensively reviewed in (3). The world of ubiquitylation is multifaceted and each layer relies upon families of proteins to write, read or erase this ubiquitin code. The steps to write the ubiquitin code are highly conserved, relying on an E1 ubiquitin activating enzyme, an E2 ubiquitin conjugating enzyme and an E3 ubiquitin protein ligase with substrate specificity. As reviewed in (1), the human genome encodes two ubiquitin E1 enzymes, approximately forty E2 enzymes and more than six hundred E3 ligases, a clear depiction of the complexity involved in functional ubiquitylation.

Ubiquitylation is a reversible post-translational modification, with removal of the ubiquitin signal catalysed by deubiquitylase enzymes (DUBs). The human genome encodes approximately one hundred DUBs, that we refer to here as the DUBome. These DUBs belong to six families: the ubiquitin specific protease (USP), ubiquitin C-terminal hydrolase (UCH), ovarian tumour protease (OTU), Josephin (JOS), JAB1/MPN/MOV34 (JAMM) families (4), or the newly discovered motif-interacting with Ub (MIU)-containing novel DUB (MINDY) family (5). As reviewed in (4), whilst most DUB families are thiol proteases harbouring a catalytic triad, the JAMM metalloproteases require a zinc ion to facilitate ubiquitin chain removal. As editors of ubiquitin signalling, DUBs are regulators of varied essential cellular processes, notably many have been assigned roles in DNA damage repair and cell cycle progression. As these processes are often dysregulated in cancer, DUBs, as potentially druggable enzymes, have quickly become the focus of several pharmaceutical companies vying to develop new cancer therapies.

The cell division cycle

The cell cycle coordinates cellular events to duplicate the genetic material and divide the cellular contents to create two identical daughter cells. The cycle comprises four stages. After division, cells undergo an initial growth phase (G_1), followed by the

replication of the genome (S-phase). A second growth phase (G_2) prepares the cell for division and assembles cytoskeletal structures, before the genetic material divides between the daughter cells during mitosis (M). The unidirectional progression through these cell cycle phases is dependent upon the periodic activation and inactivation of substrate proteins by kinases (including CDKs and PLKs) and ubiquitin-mediated degradation of key effectors by E3 ligases (including the APC/C and SCF complexes). Accordingly, cell cycle effectors are regulated through protein-protein interactions, phosphorylation-dependent activation and ubiquitylation-dependent degradation, all working in concert to achieve an exquisite level of control throughout the cycle (6). Once initiated, the cell cycle can be viewed as a series of autonomic cellular events that cascade until the eventual division into two daughter cells. However, checkpoints are inherent in the system, to temporarily halt the cell cycle if conditions are unfavourable. Many DUBs have direct or indirect roles during the cell cycle (7-9). We discuss here selected examples of those regulating cell cycle progression and checkpoint maintenance, as summarised in Figure 1.

DUBs and the G_1 restriction point

Upon entering G_1 , cells are not committed to a subsequent round of cell division, as entry into the cell cycle requires sufficient mitogenic signalling to overcome a restriction point in late G_1 . Rb, the mediator of this restriction point, inhibits the E2F transcription factor during G_1 (10). Upon CDK4/6 activation by the G_1 -cyclin CCND and then by CCNE, Rb is increasingly phosphorylated. This results in progression through the restriction point as hyperphosphorylated Rb dissociates from E2F, causing the transcription of S-phase genes (importantly CCNE and CCNA) (11). In addition to phosphorylation, Rb is regulated by ubiquitylation, being a target of the E3 ligase MDM2 (12, 13) which promotes its proteasomal degradation (14). The DUB USP7 directly antagonises MDM2-mediated polyubiquitylation of Rb, stalling the cell cycle in G_1 (15). USP7 is not the only DUB that may govern the restriction point, the tumour suppressor BAP1 also indirectly regulates the activity of E2F, via the deubiquitylation of HCF-1, an important transcriptional co-regulator at E2F promoter sites (16, 17). CYLD, another well-established tumour suppressor, plays a protective role during G_1 via the transcription factor BCL-3. CYLD deubiquitylates BCL-3 inhibiting its nuclear translocation and so decreases the transcription of BCL-3 target genes including CCND (18). CYLD therefore indirectly decreases CCND levels preventing cells from passing through the restriction point. During G_1 , the APC/C polyubiquitylates the S-phase cyclin CCNA, targeting it for degradation in order to prevent the cell from entering S-phase. The DUB USP37 directly regulates S-phase

entry through antagonising activity of the APC/C^{CDH1} in G₁ by removing polyubiquitin chains to stabilise CCNA (19).

DUBs and DNA damage checkpoints

Key to successful cell division is maintaining the integrity of the genome during DNA replication in S-phase, and this is monitored by a number of quality control mechanisms. If DNA becomes damaged, checkpoints stall the cell cycle and activate DNA damage repair (DDR) pathways. This response revolves around p53, which is stabilised and activated by DNA damage checkpoint signalling following a range of genotoxic insults. Under normal conditions, p53 is continuously synthesized but maintained at a low level by MDM2 polyubiquitylation targeting p53 for proteasomal degradation (20). Under genotoxic stress, these regulatory mechanisms are reversed, to allow p53 to stall the cell cycle to enable repair or trigger apoptosis. At sites of DNA damage, sensors (e.g. 53BP1) facilitate the activation of DNA damage kinases (notably ATM and CHK2) resulting in p53 phosphorylation. This abolishes the interaction between p53 and MDM2, increasing p53 levels and inducing transcription of p53 target genes (21), as well as activating transcription-independent roles of p53 in many of the major DDR pathways (22).

Given the integral role of p53 in cell cycle fate, it is perhaps unsurprising that many DUBs have been highlighted as direct or indirect p53 regulators, including USP2a, USP5, USP7 (HAUSP), USP10, USP11, USP28, OTUB1 and OTUD5. USP7, a predominantly nuclear DUB, was the first to be associated with the p53-dependent DDR via directly antagonising MDM2 polyubiquitylation of p53 (23). However, USP7 also directly deubiquitylates the auto-polyubiquitylated MDM2, stabilising the E3 ligase as well as its substrate (24). Although this may seem counterintuitive, USP7 exhibits a preference for MDM2 over p53 in unstressed cells ensuring p53 levels are maintained at a low level. Upon DNA damage, USP7 is dephosphorylated by PPM1G reducing activity towards MDM2, leading to increased auto-polyubiquitylation and degradation of MDM2, and the subsequent accumulation of p53 (25).

Other DUBs, including USP10, USP11 and OTUD5, also directly interact with, deubiquitylate and stabilise p53. Interestingly USP10, a predominantly cytoplasmic DUB, is involved in homeostasis of cytoplasmic p53 in unstressed cells, but following DNA damage a fraction of USP10 can translocate into the nucleus where it contributes to p53 activation (26). As described for USP7, other DUBs indirectly control p53 levels via MDM2. For example, USP2a negatively regulates p53 levels

through the stabilisation of MDM2, whilst exhibiting no deubiquitylating activity towards p53 directly (27). OTUB1, another cytoplasmic DUB, can directly interact with p53, but predominantly stabilises p53 indirectly in the cytoplasm, through a non-catalytic mechanism. OTUB1 does this by binding and suppressing polyubiquitylation through the MDM2 associated E2 enzyme UbcH5 (28). In contrast, USP28 was shown to interact with and stabilise both the damage sensor 53BP1 and the checkpoint kinase CHK2, that activate p53 under genotoxic conditions (29). USP5 uses perhaps the most indirect mechanism to stabilise p53 without physically interacting with components of the p53-MDM2 axis. It primarily disassembles unanchored polyubiquitin chains, and loss of USP5 results in accumulation of these chains that compete with ubiquitylated p53, but not MDM2, for proteasome recognition and degradation so that p53 is selectively stabilised (30).

In addition, many DUBs have also been associated with executing specific DDR pathways (8). For example, USP1 can support repair through both the Fanconi anaemia and translesion repair pathways (31). An RNAi-based study has linked USP3 with double-strand DNA break repair; USP3 directly interacts with and removes monoubiquitylation from histones H2A and H2B, and possibly other DDR effectors, to coordinate DNA repair (32). Some DUBs exhibit a more global effect on DDR pathways, for example one screen revealed that UCHL5 was recruited to sites of DNA damage in addition to being involved in double-strand break resection (33).

DUBs with roles in mitotic progression and cytokinesis

Following replication of the genome, and assuming checkpoints are satisfied in G₂, the cell enters mitosis, where the newly replicated sister chromatids must be divided into each daughter cell. To achieve this, the cell passes through a sequence of distinct mitotic phases: prophase, metaphase, anaphase, telophase and cytokinesis (Figure 1). Prior to mitosis, the mitotic kinase CDK1 is held in an inactivate state by WEE1 phosphorylation, until SCF^{βTrCP}-mediated ubiquitylation and degradation of WEE1 triggers mitotic entry; USP50 can repress mitotic entry through stabilising WEE1 (34). Subsequently, USP7 can indirectly regulate the levels of Aurora A, a kinase required for correct maturation of the bipolar mitotic spindle, by stabilising CHFR, an E3 ligase that targets Aurora A for degradation (35).

USP44 was one of the first DUBs to be linked to mitotic progression, with a role in metaphase-anaphase transition (36). Anaphase entry is stimulated by the APC/C and results in the separation of sister chromatids. To ensure the correct chromosome

complement is distributed to each daughter cell, the spindle assembly checkpoint (SAC) monitors attachment of each chromosome pair to opposite poles of the mitotic spindle. Anaphase is arrested until the SAC is satisfied, preventing premature and inaccurate division of genomic content. Three key proteins, MAD2, BUBR1 and BUB3, comprise the mitotic checkpoint complex (MCC) (37). The MCC sequesters the APC/C activator CDC20 at unattached chromosomes, thus inhibiting the APC/C until chromosomes are correctly attached (37). Once the SAC is satisfied, CDC20 is ubiquitylated and subsequently dissociates from the MCC to activate the APC/C (38). USP44 plays a protective role at the SAC, directly antagonising CDC20 ubiquitylation, and so promoting MCC inhibition of the APC/C (36). Once the SAC is satisfied, USP44 dephosphorylation decreases its activity towards CDC20, initiating mitotic exit through APC/C activation (39).

USP44 is not the only DUB that contributes to regulation of the SAC, for example USP39 and USP9X are also essential for the correct alignment of chromosomes at the mitotic spindle and their accurate division during anaphase. The mitotic kinase Aurora B is a key regulator of the attachment of sister chromatids to microtubules in the mitotic spindle. It exists in a complex with Survivin, the ubiquitylation status of which mediates interaction of the complex with chromosomes (40). Depletion of USP39 results in decreased transcription and consequently lower levels of Aurora B kinase in cycling cells (41), whilst USP9X-mediated deubiquitylation of Survivin is required for dissociation from the chromosomes once correctly aligned (42). Another DUB, USP4, plays an indirect role in the SAC through regulating correct splicing of mRNA transcripts, including for the mitotic checkpoint kinase BUB1 (43).

The DUB CYLD plays roles during both metaphase and cytokinesis. CYLD directly interacts with the catalytic domain of HDAC6, inhibiting alpha-tubulin deacetylation and therefore indirectly increasing the stability of microtubules. This governance of microtubule stability by CYLD plays a role in spindle orientation during metaphase (44) and regulates the rate of cytokinesis (45). Lastly, USP8 and AMSH, two DUBs that are usually recruited to endosomes, have an important role in cytokinesis. The scission of the two daughter cells requires components of the ESCRT machinery including VAMP8, which co-localises with, and is deubiquitylated by, both USP8 and AMSH during cytokinesis (46).

The centrosome cycle

Centrosomes are cytoplasmic organelles which act as the dominant microtubule-organizing centres (MTOCs) in animal cells. During the cell cycle, centrosomes determine spatial arrangement of the microtubule arrays to influence cell shape, polarity, motility and organization of the mitotic spindle (47, 48). The core components are two centrioles, small barrel-shaped organelles that are embedded in pericentriolar material (PCM). Each centriole consists of nine microtubule triplets arranged in a highly conserved rotational symmetry, imparted by SAS-6 during centriole assembly (49, 50). The PCM is a dense protein matrix composed of various proteins and exhibiting a high level of spatial organization, its major function is recruitment of gamma-tubulin complexes which are essential for microtubule nucleation (51, 52).

Centrosome replication is strictly coordinated with cell cycle progression (Figure 2). Duplication of the single G_1 centrosome begins at the G_1/S transition and is completed during S-phase so that two centrosomes are present in G_2 . These facilitate bipolar spindle formation at metaphase and are then segregated, one into each daughter cell, during cytokinesis (53, 54). Key to centrosome replication is centriole duplication, as the pre-existing mother centriole duplicates itself to form a daughter centriole. The kinase PLK4 and two SCF ubiquitin E3 ligases ensure that only a single replication event normally occurs. SCF^{FBXW5} ubiquitylates SAS-6 to target it for proteasomal degradation, preventing centriole over-duplication. SCF^{FBXW5} activity is limited by PLK4 to prevent premature SAS-6 degradation. Following G_1/S transition, PLK4 homodimerises and *trans*-autophosphorylates, signalling recruitment of SCF^{βTrCP} which ubiquitylates and degrades PLK4. Decreased PLK4 levels restore SCF^{FBXW5} activity and block re-duplication (49, 55-58). Once duplicated, the daughter centriole elongates during S-phase and G_2 . This process is controlled by several genes including the multifunctional centriolar protein CP110, which becomes ubiquitylated by SCF^{cyclinF} during G_2 and mitosis. Centrosomes then undergo a maturation process which requires recruitment of PCM. Finally, the centrosomes separate during G_2 , through KIF11 kinesin activity, which also facilitates bipolar spindle formation during mitosis (49, 59, 60).

Many human cells also display cilia in a cell cycle-dependent manner. During G_1 (or G_0 in terminally-differentiated cells) centrosomes migrate to the cell cortex, where the mother centriole matures into a basal body which acts as a template for cilia elongation. During S-phase, both mother and daughter centrioles undergo

duplication as normal. Then, prior to mitosis, cilia disassemble and the centrioles migrate back to the cell interior, ready to act as spindle poles during mitosis (61).

Various cell division errors, such as centrosome over-duplication, cytokinesis failure, or cell fusion can cause centrosome amplification, which is observed in many human cancers. The notion that, in addition to acting as MTOCs, centrosomes may function as signalling hubs (62) suggests one way in which amplification of centrosomes may benefit cancer cells. However, supernumerary centrosomes may cause multipolar spindle formation, impaired cell division, aneuploidy and genomic instability (63, 64). If uncorrected, multipolar spindles can lead to multipolar cell division and massive aneuploidy, which is usually lethal for the cell. Some cancer cells use mechanisms such as centrosome inactivation or centrosome loss to avoid multipolar divisions (65). However, centrosome clustering is probably the most common response in cancer cells; this enables aggregation of additional centrosomes into two groups to form a pseudo-bipolar spindle and allow the cell to undergo bipolar cell division (66, 67). Ubiquitylation is increasingly recognised as a key regulator of centrosome biology (68), and our current knowledge of the role for DUBs in specific aspects of the centrosome cycle is summarised in Figure 2.

DUBs regulating centrosome duplication and elongation during S/G₂

During S-phase centrosomes must be duplicated exactly once. A number of the key proteins involved in centrosome duplication are ubiquitylated and therefore also open to regulation by deubiquitylation; the balance between these processes is imperative for precise duplication. For example, CP110 levels are normally tightly controlled during G₂ and mitosis through ubiquitylation by SCF^{cyclinF}, leading to CP110 degradation (69). Countering this, USP33 localises to centrioles during S-phase and G₂/M where it can deubiquitylate and stabilise CP110. Overexpression of either CP110 (70) or USP33 (71) leads to centrosome amplification. Similarly, appropriate expression of CEP131, a centriolar satellite protein, is required for accurate centrosome duplication (72). Affinity purification and mass spectrometry identified USP9X as a CEP131 interactor (73); USP9X localizes to centrosomes in a cell cycle-dependent manner, most strikingly during S-phase and G₂. USP9X gain-of-function leads to CEP131 deubiquitylation, stabilization and centrosome amplification (73). In addition, overexpression of a third DUB, USP1, is also linked with centrosome amplification. Although the mechanism remains unclear, USP1 may act in part through increasing expression of ID1 (74), a fraction of which localises to the centrosome, as ID1 overexpression can induce centrosome amplification (75).

DUBs affecting centrosome maturation, separation and mitotic spindle organisation during G₂ and mitosis

BRCA1/BARD1-dependent ubiquitylation of gamma-tubulin plays a key role in the regulation of centrosome duplication and microtubule nucleation, with BRCA1 loss resulting in centrosome amplification (76, 77). An siRNA screen for DUBs that affect levels of ubiquitylated gamma-tubulin identified BAP1 and UCHL1 as candidates (78). Whilst UCHL1 interacts with gamma-tubulin in G₁, the BAP1 interaction is largely confined to mitosis, suggesting these two DUBs regulate gamma-tubulin in a cell cycle-dependent manner (78). BAP1 removes ubiquitin from gamma-tubulin, and mitotic defects in cells with low BAP1 levels are rescued by expression of BAP1 but not a catalytically inactive mutant. Whilst the mechanism remains to be fully elucidated, it seems deubiquitylation of gamma-tubulin by BAP1 during mitosis allows proper spindle organisation and function (78). CEP192 is a centrosomal protein with roles in maturation of centrosomes at the onset of mitosis and organization of the mitotic microtubule landscape. Mass spectrometry identified the deubiquitylase CYLD as a CEP192 interactor and CYLD co-depletion restores spindle assembly defects in CEP192-depleted cells (79).

In addition to its well described role in the spindle-assembly checkpoint (36), USP44 also independently affects mitotic geometry by regulating centrosome separation and positioning (80). USP44 interacts with CETN2 and, although the targets of USP44 at the centrosome remain to be elucidated, catalytically-inactive or CETN2-binding mutants of USP44 fail to rescue centrosome positioning defects. USP7 also plays a role in maintaining the correct number of centrosomes in a cell. It interacts with and stabilises centrosomal PLK1-phosphorylated 53BP1 at mitosis (81). Depletion of 53BP1 results in lower levels of p53 and CENPF which is required for proper centrosome separation and spindle formation. Cells lacking 53BP1 accumulate supernumerary centrosomes, not through *de novo* amplification but rather due to failure of cytokinesis in cells with incorrect centrosome and spindle positioning and chromosomal missegregation. In contrast, USP37 depletion indirectly results in centrosome fragmentation, and hence multipolar spindle formation, through ubiquitylation and degradation of WAPL, a regulator of sister chromatid resolution and spindle tension (82). Notably, three recent papers have revealed a novel checkpoint, the mitotic surveillance pathway, that can detect centrosome loss or prolonged mitosis and results in cell cycle arrest (83-85). The signalling pathway

involves 53BP1 and the deubiquitylase USP28 acting in a complex to deubiquitylate and stabilise p53, which in turn controls cell fate.

DUBs involved in centrosome clustering during cancer cell mitosis

Centrosome clustering is a mechanism that cancer cells containing supernumerary centrosomes commonly use to gather amplified centrosomes into two poles during mitosis, allowing for bipolar division and cancer cell proliferation (86). Inhibition of centrosome clustering is an attractive, cancer specific, therapeutic intervention. Two genome-wide screens have identified proteins required for centrosome clustering in *Drosophila* or human cells (87, 88). Analysis of the *Drosophila* dataset reveals prominence of proteins involved in ubiquitylation and the proteasomal pathway, including two DUBs, the *Drosophila* orthologues of human USP8 and USP31 (87). The screen in human cells also identified USP54 (88), a DUB that is predicted to be catalytically inactive (89). However, neither the ubiquitylation process nor these DUBs were investigated further in either study. In relation to its role in stabilising CP110 described above, USP33 may also indirectly affect centrosome clustering. Inhibition of CDK2 prevents CP110 phosphorylation that is required for centrosome clustering activity (90, 91) and combining CDK2 inhibition with USP33 depletion has a co-operative effect on CP110, driving anaphase catastrophe via multipolar spindle formation (91). In addition, the functional overlap of other DUBs with centrosome regulation makes it likely there are further DUBs involved in this process. For example, a functional SAC is required for effective centrosome clustering (87) and, as discussed above, several DUBs including USP4, USP9X, USP39 and USP44 are required for SAC activity (36, 41-43).

DUBs involved in ciliogenesis during G₀/G₁

A number of DUBs have been found to be required for the formation of primary cilia during G₀/G₁ phase of the cell cycle, a process termed ciliogenesis. Firstly, the DUB CYLD is recruited to centrosomes and the basal body of cilia via its interaction with CAP350, where it has to be present and catalytically active to promote docking of basal bodies at the plasma membrane and hence ciliogenesis (92). A concurrent study also demonstrated that CYLD is required for docking of basal bodies at the plasma membrane and identified that this can, at least in part, be explained by its ability to deubiquitylate CEP70. Deubiquitylation of CEP70 allows it to interact with gamma-tubulin at the centrosome to mediate ciliogenesis (93). In addition, CYLD inactivates HDAC6, which modulates cilia length (93). Secondly, via an independent mechanism to its roles in centrosome duplication, USP9X also regulates ciliogenesis

(94). During G_0/G_1 , USP9X is recruited to the centrosome where it deubiquitylates and stabilises NPHP5, a positive regulator of ciliogenesis, so favouring cilia formation. However, at G_2/M USP9X becomes cytoplasmic, allowing degradation of NPHP5 and loss of cilia. Lastly, a survey of DUB subcellular localisation found that USP21 localised to centrosomes and microtubules (95). USP21 is required for effective microtubule regrowth from centrosomes, neurite outgrowth, generation of the primary cilium (95), and hedgehog signalling (96).

Conclusions, future challenges and outlook

Here we highlight specific roles for many different DUBs in controlling critical aspects of cell cycle progression, p53 homeostasis and DNA damage repair, as well as centrosome biology. To date, at least 30% of the DUBome has been associated with these processes, with predominant representation from the USP and UCH families. In addition to these roles, it is evident that many other DUBs regulate cellular processes during specific cell cycle phases. One example is the role of USP15 in regulating the transcriptional repressor REST. Like many transcription factors, REST is rapidly degraded at G_2/M prior to cell division, however as it represses cellular differentiation genes it must be reconstituted in G_1 . REST degradation is triggered by phosphorylation-dependent SCF^{BTCP} ubiquitylation (97, 98), and whilst this is reported to be antagonised by USP7 in neural progenitors (99), in cycling cells mitotic REST degradation appears to be unopposed. However, as cells exit mitosis, USP15 acts to deubiquitylate newly synthesized REST and rapidly rescue its expression levels (100). Considering phase-specific roles such as this greatly expands the involvement of the DUBome in cell cycle biology.

This review aims to capture the current state of the field of DUB cell cycle research, but many outstanding questions remain. Whilst certain DUBs have very distinct roles, others like CYLD, USP9X and USP7, play multiple roles at various phases of both the cell and centrosome cycles. Often we do not yet know how the function of a particular DUB is restricted to a cell cycle phase, or directed towards a specific target, to achieve precise temporal regulation of cell cycle effectors. Indeed, although transcriptomics suggest USP1 is the only DUB that is periodically transcribed during the cell cycle (101), proteomics reveals periodic phosphorylation of several DUBs (102), but we at present lack a clear profile of regulated protein expression and activity for the DUBome during the cell cycle.

Although certain DUBs, OTUB1 being a notable example (28), play important roles through scaffolding interactions independent of their catalytic activity, most DUBs have catalytic functions. As highlighted in a recent review (103), unrestricted enzymatic activity of the DUBs would be hazardous for cells, and we are now beginning to appreciate the multi-layered mechanisms by which their activity can be controlled and directed. These include internal regulatory domains within some DUBs, interaction with allosteric regulators, incorporation into macromolecular complexes, and post-translational modifications. Relevant examples for stabilisation of p53 in response to genotoxic stress include phosphorylation-dependent nuclear localization of USP10 (26) and modulation of USP7 activity towards MDM2 (25). Intriguingly, allosteric activation of USP7 by GMPS that stabilizes alignment of the catalytic site can also direct USP7 activity towards p53 under genotoxic stress (104, 105). These findings begin to rationalise the physiological roles of a DUB that is capable of stabilising both p53 and the E3 ligase MDM2, which targets p53 for degradation.

Another open question is why for certain processes, most notably in p53 regulation, there appears to be huge redundancy with multiple DUBs playing similar roles. One potential explanation is the ability of different DUBs to regulate p53 by different mechanisms and in different cellular compartments, as described above for USP7, USP10 and OTUB1. This may help ensure fine control of p53 activation in response to genotoxic stress. Critical roles for many DUBs have also been described in regulating the sharp and irreversible signalling decisions that are made at the G₁ restriction point and the SAC. In both cases, a picture is emerging where DUBs contribute to a regulatory network, and each key component of the cascade is controlled by a specific DUB.

There is emerging interest in the role of the DUBome in centrosome biology, which is less well studied than in the cell cycle. As a distinct organelle, it is easier to visualise how temporal roles for DUBs may be regulated, with some DUBs such as USP33 (71), USP9X (73) and BAP1 (78) already known to be recruited to the centrosome in a cell cycle-dependent manner. Where DUBs have been associated with the centrosome cycle, their mechanistic roles are often not yet well elucidated. For example, screens suggest that USP8 and USP31 may be linked to the centrosome clustering in *Drosophila* (87) and USP54 in human cells (88). However, their centrosome-associated targets remain unknown, and no mechanism of action for these DUBs in regulating centrosome clustering has yet been suggested. It will

be interesting to see whether DUBs predicted from screens and in model organisms do play significant roles in centrosome clustering in human cancer cells. Lastly, in addition to acting as MTOCs, centrosomes have recently also been established as signalling hubs (62). Many of the studies on DUBs at centrosomes we have discussed focus on their roles in duplicating and regulating the centrosome structure, and on their functions in nucleating microtubules. In future it is likely that new roles for DUBs will be discovered in centrosome-based signalling pathways.

Funding information

The authors are supported by Wellcome Trust PhD studentships 096567/Z/11/Z (SD) and 109307/Z/15/Z (DSP), a North West Cancer Research project grant CR1050 (ABF) and a North West Cancer Research endowment (IAP).

Competing interests

The Authors declare no competing interests associated with the manuscript.

Abbreviations

53BP1, p53-binding protein 1; AMSH, Associated molecule with the SH3 domain of STAM; APC/C, anaphase promoting complex/cyclosome; BAP1, BRCA1-associated protein 1; ATM, Ataxia telangiectasia mutated; BARD1, BRCA1-associated RING domain protein 1; BCL-3, B-cell lymphoma 3 protein; BRCA1, Breast cancer type 1 susceptibility protein; β TrCP, beta-transducin repeat containing protein; BUB3, Mitotic checkpoint protein BUB3; BUBR1, Mitotic checkpoint serine/threonine-protein kinase BUB1 beta; CAP350, Centrosome-associated protein of 350 kDa; CCNA, cyclin-A; CCND, cyclin-D; CCNE, cyclin-E; CDC20, Cell division cycle protein 20 homolog; CDH1, CDC20 homolog 1; CDKs, cyclin-dependent kinases; CENPF, Centromere protein F; CEP70, Centrosomal protein of 70 kDa; CEP131, Centrosomal protein of 131 kDa; CEP192, Centrosomal protein of 192 kDa; CETN2, Centrin2; CHFR, Checkpoint with Forkhead and Ring Finger; CHK2, checkpoint kinase 2; CP110, Centriolar coiled-coil protein of 110 kDa; CYLD, Ubiquitin carboxyl-terminal hydrolase Cylindromatosis; DDR, DNA damage repair; DUBs, deubiquitylases; DUBome, the cellular complement of DUBs; ESCRT, Endosomal sorting complexes required for transport; FBXW5, F-box/WD repeat-containing protein 5; H2A, Histone 2A; H2B, Histone 2B; HCF-1, Host cell factor 1; HDAC6, histone deacetylase 6; ID1, Inhibitor of DNA binding 1; JAMM, JAB1/MPN/MOV34; JOS, Josephin; KIF11, Kinesin-related motor protein Eg5; MAD2, Mitotic arrest deficient 2-like protein 1; MCC, mitotic checkpoint complex; MDM2, E3 ubiquitin-protein ligase Mdm2 (Double minute 2 protein); MINDY, motif-interacting with Ub (MIU)-containing novel DUB; NPHP5, Nephrocystin-5/IQ calmodulin-binding motif-containing protein 1; MTOCs, microtubule organizing centres; OTU, ovarian tumour protease; p53, Cellular tumour antigen p53; PCM, pericentriolar material; PLKs, polo-like kinases; PPM1G, Protein phosphatase 1G; Rb, retinoblastoma protein; SAC, spindle assembly checkpoint; SAS-6, Spindle assembly abnormal protein 6 homolog; SCF, skp/cullin/F-box; UbcH5, Ubiquitin-Conjugating Enzyme E2D 1 (UBC4/5 Homolog, Yeast); UCH, ubiquitin C-terminal hydrolase; USP, ubiquitin specific protease; VAMP8, Vesicle-associated membrane protein 8; WAPL, Wings apart-like protein homolog; WEE1, WEE1 G2 Checkpoint Kinase.

References

1. Clague MJ, Heride C, Urbe S. The demographics of the ubiquitin system. *Trends Cell Biol.* 2015;25(7):417-26.
2. Clague MJ, Coulson JM, Urbé S. Cellular functions of the DUBs. *J Cell Sci.* 2012;125(Pt 2):277-86.
3. Swatek KN, Komander D. Ubiquitin modifications. *Cell Res.* 2016;26(4):399-422.
4. Clague MJ, Barsukov I, Coulson JM, Liu H, Rigden DJ, Urbé S. Deubiquitylases from genes to organism. *Physiological reviews.* 2013;93(3):1289-315.
5. Abdul Rehman SA, Kristariyanto YA, Choi SY, Nkosi PJ, Weidlich S, Labib K, et al. MINDY-1 Is a Member of an Evolutionarily Conserved and Structurally Distinct New Family of Deubiquitinating Enzymes. *Mol Cell.* 2016;63(1):146-55.
6. Rhind N, Russell P. Signaling pathways that regulate cell division. *Cold Spring Harb Perspect Biol.* 2012;4(10).
7. Fournane S, Krupina K, Kleiss C, Sumara I. Decoding ubiquitin for mitosis. *Genes Cancer.* 2012;3(11-12):697-711.
8. Pinto-Fernandez A, Kessler BM. DUBbing Cancer: Deubiquitylating Enzymes Involved in Epigenetics, DNA Damage and the Cell Cycle As Therapeutic Targets. *Front Genet.* 2016;7:133.
9. Lim KH, Song MH, Baek KH. Decision for cell fate: deubiquitinating enzymes in cell cycle checkpoint. *Cell Mol Life Sci.* 2016;73(7):1439-55.
10. Helin K, Harlow E, Fattaey A. Inhibition of E2F-1 transactivation by direct binding of the retinoblastoma protein. *Mol Cell Biol.* 1993;13(10):6501-8.
11. Blagosklonny MV, Pardee AB. The restriction point of the cell cycle. *Cell cycle.* 2002;1(2):103-10.
12. Delston RB, Matattal KA, Sun Y, Onken MD, Harbour JW. p38 phosphorylates Rb on Ser567 by a novel, cell cycle-independent mechanism that triggers Rb-Hdm2 interaction and apoptosis. *Oncogene.* 2011;30(5):588-99.
13. Xiao ZX, Chen J, Levine AJ, Modjtahedi N, Xing J, Sellers WR, et al. Interaction between the retinoblastoma protein and the oncoprotein MDM2. *Nature.* 1995;375(6533):694-8.
14. Uchida C, Miwa S, Kitagawa K, Hattori T, Isobe T, Otani S, et al. Enhanced Mdm2 activity inhibits pRB function via ubiquitin-dependent degradation. *EMBO J.* 2005;24(1):160-9.
15. Bhattacharya S, Ghosh MK. HAUSP, a novel deubiquitinase for Rb - MDM2 the critical regulator. *FEBS J.* 2014;281(13):3061-78.
16. Machida YJ, Machida Y, Vashisht AA, Wohlschlegel JA, Dutta A. The deubiquitinating enzyme BAP1 regulates cell growth via interaction with HCF-1. *J Biol Chem.* 2009;284(49):34179-88.
17. Eletr ZM, Wilkinson KD. An emerging model for BAP1's role in regulating cell cycle progression. *Cell Biochem Biophys.* 2011;60(1-2):3-11.
18. Massoumi R, Chmielarska K, Hennecke K, Pfeifer A, Fassler R. Cyld inhibits tumor cell proliferation by blocking Bcl-3-dependent NF-kappaB signaling. *Cell.* 2006;125(4):665-77.

19. Huang X, Summers MK, Pham V, Lill JR, Liu J, Lee G, et al. Deubiquitinase USP37 is activated by CDK2 to antagonize APC(CDH1) and promote S phase entry. *Mol Cell*. 2011;42(4):511-23.
20. Fuchs SY, Adler V, Buschmann T, Wu X, Ronai Z. Mdm2 association with p53 targets its ubiquitination. *Oncogene*. 1998;17(19):2543-7.
21. Lakin ND, Jackson SP. Regulation of p53 in response to DNA damage. *Oncogene*. 1999;18(53):7644-55.
22. Sengupta S, Harris CC. p53: traffic cop at the crossroads of DNA repair and recombination. *Nat Rev Mol Cell Biol*. 2005;6(1):44-55.
23. Li M, Chen D, Shiloh A, Luo J, Nikolaev AY, Qin J, et al. Deubiquitination of p53 by HAUSP is an important pathway for p53 stabilization. *Nature*. 2002;416(6881):648-53.
24. Li M, Brooks CL, Kon N, Gu W. A dynamic role of HAUSP in the p53-Mdm2 pathway. *Mol Cell*. 2004;13(6):879-86.
25. Khoronenkova SV, Dianova II, Ternette N, Kessler BM, Parsons JL, Dianov GL. ATM-dependent downregulation of USP7/HAUSP by PPM1G activates p53 response to DNA damage. *Mol Cell*. 2012;45(6):801-13.
26. Yuan J, Luo K, Zhang L, Cheville JC, Lou Z. USP10 regulates p53 localization and stability by deubiquitinating p53. *Cell*. 2010;140(3):384-96.
27. Stevenson LF, Sparks A, Allende-Vega N, Xirodimas DP, Lane DP, Saville MK. The deubiquitinating enzyme USP2a regulates the p53 pathway by targeting Mdm2. *EMBO J*. 2007;26(4):976-86.
28. Sun XX, Challagundla KB, Dai MS. Positive regulation of p53 stability and activity by the deubiquitinating enzyme Otubain 1. *EMBO J*. 2012;31(3):576-92.
29. Zhang D, Zaugg K, Mak TW, Elledge SJ. A role for the deubiquitinating enzyme USP28 in control of the DNA-damage response. *Cell*. 2006;126(3):529-42.
30. Dayal S, Sparks A, Jacob J, Allende-Vega N, Lane DP, Saville MK. Suppression of the deubiquitinating enzyme USP5 causes the accumulation of unanchored polyubiquitin and the activation of p53. *J Biol Chem*. 2009;284(8):5030-41.
31. Kee Y, Huang TT. Role of Deubiquitinating Enzymes in DNA Repair. *Mol Cell Biol*. 2015;36(4):524-44.
32. Nicassio F, Corrado N, Vissers JH, Areces LB, Bergink S, Marteijn JA, et al. Human USP3 is a chromatin modifier required for S phase progression and genome stability. *Curr Biol*. 2007;17(22):1972-7.
33. Nishi R, Wijnhoven P, le Sage C, Tjeertes J, Galanty Y, Forment JV, et al. Systematic characterization of deubiquitylating enzymes for roles in maintaining genome integrity. *Nat Cell Biol*. 2014;16(10):1016-26, 1-8.
34. Aressy B, Jullien D, Cazales M, Marcellin M, Bugler B, Burlet-Schiltz O, et al. A screen for deubiquitinating enzymes involved in the G₂/M checkpoint identifies USP50 as a regulator of HSP90-dependent Wee1 stability. *Cell cycle*. 2010;9(18):3815-22.
35. Giovinazzi S, Morozov VM, Summers MK, Reinhold WC, Ishov AM. USP7 and Daxx regulate mitosis progression and taxane sensitivity by affecting stability of Aurora-A kinase. *Cell Death Differ*. 2013;20(5):721-31.

36. Stegmeier F, Rape M, Draviam VM, Nalepa G, Sowa ME, Ang XL, et al. Anaphase initiation is regulated by antagonistic ubiquitination and deubiquitination activities. *Nature*. 2007;446(7138):876-81.
37. Sudakin V, Chan GK, Yen TJ. Checkpoint inhibition of the APC/C in HeLa cells is mediated by a complex of BUBR1, BUB3, CDC20, and MAD2. *J Cell Biol*. 2001;154(5):925-36.
38. Reddy SK, Rape M, Margansky WA, Kirschner MW. Ubiquitination by the anaphase-promoting complex drives spindle checkpoint inactivation. *Nature*. 2007;446(7138):921-5.
39. Visconti R, Palazzo L, Della Monica R, Grieco D. Fcp1-dependent dephosphorylation is required for M-phase-promoting factor inactivation at mitosis exit. *Nat Commun*. 2012;3:894.
40. Vader G, Kauw JJ, Medema RH, Lens SM. Survivin mediates targeting of the chromosomal passenger complex to the centromere and midbody. *EMBO Rep*. 2006;7(1):85-92.
41. van Leuken RJ, Luna-Vargas MP, Sixma TK, Wolthuis RM, Medema RH. Usp39 is essential for mitotic spindle checkpoint integrity and controls mRNA-levels of aurora B. *Cell cycle*. 2008;7(17):2710-9.
42. Vong QP, Cao K, Li HY, Iglesias PA, Zheng Y. Chromosome alignment and segregation regulated by ubiquitination of survivin. *Science*. 2005;310(5753):1499-504.
43. Song EJ, Werner SL, Neubauer J, Stegmeier F, Aspden J, Rio D, et al. The Prp19 complex and the Usp4Sart3 deubiquitinating enzyme control reversible ubiquitination at the spliceosome. *Genes Dev*. 2010;24(13):1434-47.
44. Yang Y, Liu M, Li D, Ran J, Gao J, Suo S, et al. CYLD regulates spindle orientation by stabilizing astral microtubules and promoting dishevelled-NuMA-dynein/dynactin complex formation. *Proc Natl Acad Sci U S A*. 2014;111(6):2158-63.
45. Wickstrom SA, Masoumi KC, Khochbin S, Fassler R, Massoumi R. CYLD negatively regulates cell-cycle progression by inactivating HDAC6 and increasing the levels of acetylated tubulin. *EMBO J*. 2010;29(1):131-44.
46. Mukai A, Mizuno E, Kobayashi K, Matsumoto M, Nakayama KI, Kitamura N, et al. Dynamic regulation of ubiquitylation and deubiquitylation at the central spindle during cytokinesis. *J Cell Sci*. 2008;121(Pt 8):1325-33.
47. Bettencourt-Dias M, Glover DM. Centrosome biogenesis and function: centrosomes brings new understanding. *Nature reviews Molecular cell biology*. 2007;8(6):451-63.
48. Bornens M. The centrosome in cells and organisms. *Science*. 2012;335(6067):422-6.
49. Nigg EA, Stearns T. The centrosome cycle: Centriole biogenesis, duplication and inherent asymmetries. *Nat Cell Biol*. 2011;13(10):1154-60.
50. Nakazawa Y, Hiraki M, Kamiya R, Hirono M. SAS-6 is a cartwheel protein that establishes the 9-fold symmetry of the centriole. *Current Biology*. 2007;17(24):2169-74.
51. Moritz M, Agard DA. gamma-Tubulin complexes and microtubule nucleation. *Current Opinion in Structural Biology*. 2001;11(2):174-81.

52. Lawo S, Hasegan M, Gupta GD, Pelletier L. Subdiffraction imaging of centrosomes reveals higher-order organizational features of pericentriolar material. *Nature Cell Biology*. 2012;14(11):1148-+.
53. Nigg EA. Centrosome duplication: of rules and licenses. *Trends in cell biology*. 2007;17(5):215-21.
54. Rivera-Rivera Y, Saavedra HI. Centrosome - a promising anti-cancer target. *Biol-Targets Ther*. 2016;10:167-76.
55. Bettencourt-Dias M, Rodrigues-Martins A, Carpenter L, Riparbelli M, Lehmann L, Gatt MK, et al. SAK/PLK4 is required for centriole duplication and flagella development. *Current Biology*. 2005;15(24):2199-207.
56. Habedanck R, Stierhof YD, Wilkinson CJ, Nigg EA. The Polo kinase Plk4 functions in centriole duplication. *Nature Cell Biology*. 2005;7(11):1140-6.
57. Puklowski A, Homsy Y, Keller D, May M, Chauhan S, Kossatz U, et al. The SCF-FBXW5 E3-ubiquitin ligase is regulated by PLK4 and targets HsSAS-6 to control centrosome duplication. *Nature Cell Biology*. 2011;13(8):1004-U291.
58. Cunha-Ferreira I, Rodrigues-Martins A, Bento I, Riparbelli M, Zhang W, Laue E, et al. The SCF/Slimb ubiquitin ligase limits centrosome amplification through degradation of SAK/PLK4. *Current biology : CB*. 2009;19(1):43-9.
59. Jana SC, Marteil G, Bettencourt-Dias M. Mapping molecules to structure: unveiling secrets of centriole and cilia assembly with near-atomic resolution. *Current Opinion in Cell Biology*. 2014;26:96-106.
60. Mardin BR, Schiebel E. Breaking the ties that bind: New advances in centrosome biology. *Journal of Cell Biology*. 2012;197(1):11-8.
61. Ishikawa H, Marshall WF. Ciliogenesis: building the cell's antenna. *Nature Reviews Molecular Cell Biology*. 2011;12(4):222-34.
62. Arquint C, Gabryjonczyk AM, Nigg EA. Centrosomes as signalling centres. *Philosophical transactions of the Royal Society of London Series B, Biological sciences*. 2014;369(1650).
63. Nigg EA. Centrosome aberrations: Cause or consequence of cancer progression? *Nature Reviews Cancer*. 2002;2(11):815-25.
64. Nigg EA. Origins and consequences of centrosome aberrations in human cancers. *International Journal of Cancer*. 2006;119(12):2717-23.
65. Godinho SA, Pellman D. Causes and consequences of centrosome abnormalities in cancer. *Philos T R Soc B*. 2014;369(1650).
66. Quintyne NJ, Reing JE, Hoffelder DR, Gollin SM, Saunders WS. Spindle multipolarity is prevented by centrosomal clustering. *Science*. 2005;307(5706):127-9.
67. Kwon M, Godinho SA, Chandhok NS, Ganem NJ, Azioune A, They M, et al. Mechanisms to suppress multipolar divisions in cancer cells with extra centrosomes. *Genes & development*. 2008;22(16):2189-203.
68. Zhang Y, Galardy PJ. Ubiquitin, the centrosome, and chromosome segregation. *Chromosome Res*. 2016;24(1):77-91.
69. D'Angiolella V, Donato V, Vijayakumar S, Saraf A, Florens L, Washburn MP, et al. SCF(Cyclin F) controls centrosome homeostasis and mitotic fidelity through CP110 degradation. *Nature*. 2010;466(7302):138-42.

70. Chen Z, Indjeian VB, McManus M, Wang L, Dynlacht BD. CP110, a cell cycle-dependent CDK substrate, regulates centrosome duplication in human cells. *Dev Cell*. 2002;3(3):339-50.
71. Li J, D'Angiolella V, Seeley ES, Kim S, Kobayashi T, Fu W, et al. USP33 regulates centrosome biogenesis via deubiquitination of the centriolar protein CP110. *Nature*. 2013;495(7440):255-9.
72. Staples CJ, Myers KN, Beveridge RD, Patil AA, Lee AJ, Swanton C, et al. The centriolar satellite protein Cep131 is important for genome stability. *Journal of cell science*. 2012;125(Pt 20):4770-9.
73. Li X, Song N, Liu L, Liu X, Ding X, Song X, et al. USP9X regulates centrosome duplication and promotes breast carcinogenesis. *Nat Commun*. 2017;8:14866.
74. Jung JK, Jang SW, Kim JM. A novel role for the deubiquitinase USP1 in the control of centrosome duplication. *Cell Cycle*. 2016;15(4):584-92.
75. Hasskarl J, Duensing S, Manuel E, Munger K. The helix-loop-helix protein ID1 localizes to centrosomes and rapidly induces abnormal centrosome numbers. *Oncogene*. 2004;23(10):1930-8.
76. Starita LM, Machida Y, Sankaran S, Elias JE, Griffin K, Schlegel BP, et al. BRCA1-dependent ubiquitination of gamma-tubulin regulates centrosome number. *Mol Cell Biol*. 2004;24(19):8457-66.
77. Sankaran S, Crone DE, Palazzo RE, Parvin JD. BRCA1 regulates gamma-tubulin binding to centrosomes. *Cancer Biol Ther*. 2007;6(12):1853-7.
78. Zarrizi R, Menard JA, Belting M, Massoumi R. Deubiquitination of gamma-tubulin by BAP1 prevents chromosome instability in breast cancer cells. *Cancer Res*. 2014;74(22):6499-508.
79. Gomez-Ferreria MA, Bashkurov M, Mullin M, Gingras AC, Pelletier L. CEP192 interacts physically and functionally with the K63-deubiquitinase CYLD to promote mitotic spindle assembly. *Cell Cycle*. 2012;11(19):3555-8.
80. Zhang Y, Foreman O, Wigle DA, Kosari F, Vasmatzis G, Salisbury JL, et al. USP44 regulates centrosome positioning to prevent aneuploidy and suppress tumorigenesis. *J Clin Invest*. 2012;122(12):4362-74.
81. Yim H, Shin SB, Woo SU, Lee PC, Erikson RL. Plk1-mediated stabilization of 53BP1 through USP7 regulates centrosome positioning to maintain bipolarity. *Oncogene*. 2017;36(7):966-78.
82. Yeh C, Coyaud E, Bashkurov M, van der Lelij P, Cheung SW, Peters JM, et al. The Deubiquitinase USP37 Regulates Chromosome Cohesion and Mitotic Progression. *Curr Biol*. 2015;25(17):2290-9.
83. Lambrus BG, Daggubati V, Uetake Y, Scott PM, Clutario KM, Sluder G, et al. A USP28-53BP1-p53-p21 signaling axis arrests growth after centrosome loss or prolonged mitosis. *J Cell Biol*. 2016;214(2):143-53.
84. Fong CS, Mazo G, Das T, Goodman J, Kim M, O'Rourke BP, et al. 53BP1 and USP28 mediate p53-dependent cell cycle arrest in response to centrosome loss and prolonged mitosis. *Elife*. 2016;5.
85. Meitinger F, Anzola JV, Kaulich M, Richardson A, Stender JD, Benner C, et al. 53BP1 and USP28 mediate p53 activation and G1 arrest after centrosome loss or extended mitotic duration. *J Cell Biol*. 2016;214(2):155-66.

86. Marthiens V, Piel M, Basto R. Never tear us apart--the importance of centrosome clustering. *J Cell Sci.* 2012;125(Pt 14):3281-92.
87. Kwon M, Godinho SA, Chandhok NS, Ganem NJ, Azioune A, Thery M, et al. Mechanisms to suppress multipolar divisions in cancer cells with extra centrosomes. *Genes Dev.* 2008;22(16):2189-203.
88. Leber B, Maier B, Fuchs F, Chi J, Riffel P, Anderhub S, et al. Proteins required for centrosome clustering in cancer cells. *Sci Transl Med.* 2010;2(33):33ra8.
89. Quesada V, Diaz-Perales A, Gutierrez-Fernandez A, Garabaya C, Cal S, Lopez-Otin C. Cloning and enzymatic analysis of 22 novel human ubiquitin-specific proteases. *Biochem Biophys Res Commun.* 2004;314(1):54-62.
90. Hu S, Danilov AV, Godek K, Orr B, Tafe LJ, Rodriguez-Canales J, et al. CDK2 Inhibition Causes Anaphase Catastrophe in Lung Cancer through the Centrosomal Protein CP110. *Cancer Res.* 2015;75(10):2029-38.
91. Hu S, Lu Y, Orr B, Godek K, Mustachio LM, Kawakami M, et al. Specific CP110 Phosphorylation Sites Mediate Anaphase Catastrophe after CDK2 Inhibition: Evidence for Cooperation with USP33 Knockdown. *Mol Cancer Ther.* 2015;14(11):2576-85.
92. Eguether T, Ermolaeva MA, Zhao Y, Bonnet MC, Jain A, Pasparakis M, et al. The deubiquitinating enzyme CYLD controls apical docking of basal bodies in ciliated epithelial cells. *Nat Commun.* 2014;5:4585.
93. Yang Y, Ran J, Liu M, Li D, Li Y, Shi X, et al. CYLD mediates ciliogenesis in multiple organs by deubiquitinating Cep70 and inactivating HDAC6. *Cell Res.* 2014;24(11):1342-53.
94. Das A, Qian J, Tsang WY. USP9X counteracts differential ubiquitination of NPHP5 by MARCH7 and BBS11 to regulate ciliogenesis. *PLoS Genet.* 2017;13(5):e1006791.
95. Urbe S, Liu H, Hayes SD, Heride C, Rigden DJ, Clague MJ. Systematic survey of deubiquitinase localization identifies USP21 as a regulator of centrosome- and microtubule-associated functions. *Mol Biol Cell.* 2012;23(6):1095-103.
96. Heride C, Rigden DJ, Bertsoulaki E, Cucchi D, De Smaele E, Clague MJ, et al. The centrosomal deubiquitylase USP21 regulates Gli1 transcriptional activity and stability. *J Cell Sci.* 2016;129(21):4001-13.
97. Guardavaccaro D, Frescas D, Dorrello NV, Peschiaroli A, Multani AS, Cardozo T, et al. Control of chromosome stability by the beta-TrCP-REST-Mad2 axis. *Nature.* 2008;452(7185):365-9.
98. Westbrook TF, Hu G, Ang XL, Mulligan P, Pavlova NN, Liang A, et al. SCFbeta-TRCP controls oncogenic transformation and neural differentiation through REST degradation. *Nature.* 2008;452(7185):370-4.
99. Huang Z, Wu Q, Guryanova OA, Cheng L, Shou W, Rich JN, et al. Deubiquitylase HAUSP stabilizes REST and promotes maintenance of neural progenitor cells. *Nat Cell Biol.* 2011;13(2):142-52.
100. Faronato M, Patel V, Darling S, Dearden L, Clague MJ, Urbe S, et al. The deubiquitylase USP15 stabilizes newly synthesized REST and rescues its expression at mitotic exit. *Cell cycle.* 2013;12(12):1964-77.
101. Gauthier NP, Larsen ME, Wernersson R, de Lichtenberg U, Jensen LJ, Brunak S, et al. Cyclebase.org--a comprehensive multi-organism online database of cell-cycle experiments. *Nucleic Acids Res.* 2008;36(Database issue):D854-9.

102. Olsen JV, Vermeulen M, Santamaria A, Kumar C, Miller ML, Jensen LJ, et al. Quantitative phosphoproteomics reveals widespread full phosphorylation site occupancy during mitosis. *Sci Signal*. 2010;3(104):ra3.
103. Sahtoe DD, Sixma TK. Layers of DUB regulation. *Trends Biochem Sci*. 2015;40(8):456-67.
104. Faesen AC, Dirac AM, Shanmugham A, Ovaa H, Perrakis A, Sixma TK. Mechanism of USP7/HAUSP activation by its C-terminal ubiquitin-like domain and allosteric regulation by GMP-synthetase. *Mol Cell*. 2011;44(1):147-59.
105. Reddy BA, van der Knaap JA, Bot AG, Mohd-Sarip A, Dekkers DH, Timmermans MA, et al. Nucleotide biosynthetic enzyme GMP synthase is a TRIM21-controlled relay of p53 stabilization. *Mol Cell*. 2014;53(3):458-70.

Figure Legends

Figure 1. DUBs associated with the cell cycle.

The cell cycle is schematically represented, highlighting key checkpoints and the individual stages of mitosis. DUBs with specific roles are indicated in the appropriate phases: solid colouring shows membership of the DUB families, and coloured edges illustrate the major cell cycle function. SAC: spindle assembly checkpoint.

Figure 2. DUBs associated with the centrosome cycle.

The cell cycle is schematically represented, highlighting the key stages of centrosome replication and function. DUBs with specific roles are indicated in the appropriate phases: solid colouring shows membership of the DUB families, and coloured edges illustrate the major function in the centrosome cycle.




Figure 1. DUBs with roles in the cell cycle.




Figure 2. DUBs with roles in the centrosome cycle.