

LONDON
SCHOOL of
HYGIENE
& TROPICAL
MEDICINE



Gedela, K; Edwards, SG; Benn, P; Grant, AD (2014) Prevalence of vitamin D deficiency in HIV-positive, antiretroviral treatment-naive patients in a single centre study. *International journal of STD & AIDS*, 25 (7). pp. 488-492. ISSN 0956-4624 DOI: 10.1177/0956462413515194

Downloaded from: <http://researchonline.lshtm.ac.uk/1440273/>

DOI: [10.1177/0956462413515194](http://dx.doi.org/10.1177/0956462413515194)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

Full title: Prevalence of vitamin D deficiency in HIV positive, antiretroviral treatment naïve patients in a single centre study.

Short title: Vitamin D deficiency in ART naïve patients

Key words: HIV; Vitamin D; Antiretroviral treatment.

Word Count: 1296 (excluding table, abstract and references)

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Abstract

Objective: To describe the prevalence of vitamin D deficiency among antiretroviral treatment (ART)-naïve, HIV positive individuals.

Methods: We reviewed records of consecutive ART-naïve patients, registering for care for the first time at a London clinic from 01/01/2008 to 31/12/2009. During this period, serum 25-hydroxycholecalciferol (25-OHD) was measured routinely for all new patients. 25-OHD deficiency and severe deficiency were defined as ≤ 50 and ≤ 25 nmol/L, respectively.

Results: Among 253 patients (82% male, median age 36 years, 64% white ethnicity), 148 (58.5%) were 25-OHD deficient, including 32 (12.6%) who were severely deficient. 73.5% (61/83) patients of non-white ethnicity were 25-OHD deficient compared with 50.7% (76/150) of those reporting white ethnicity ($p < 0.001$). Seven of eight (87.5%) patients with hypocalcaemia (< 2.12 nmol/L) were 25-OHD deficient. The prevalence of 25-OHD deficiency was higher in winter and spring vs. summer and autumn (89/129 [69.0%] vs. 59/124 [47.6%], $p < 0.001$). Serum 25-OHD deficiency was not associated with gender, CD4 count, HIV viral load or clinical stage.

Conclusion: Serum 25-OHD deficiency was common among ART-naïve patients, with those of non-white ethnicity at highest risk. CD4 count, HIV viral load and HIV clinical staging do not help to identify those at risk, but low serum calcium should prompt investigation of 25-OHD levels.

Introduction

In addition to vitamin D's established role in bone metabolism, vitamin D deficiency is reported to be associated with a variety of medical conditions including diabetes, cardiovascular disease [1–5] and infectious diseases particularly tuberculosis and HIV [6–11].

Studies investigating vitamin D's role in HIV infection have been conflicting but associations with an increased risk of HIV acquisition and progression to advanced HIV have been shown [9,12,13]. Vitamin D deficiency is also speculated to be associated with an increased risk of cardiovascular disease in non-HIV infected populations [2–5]. One study demonstrated an association between lower vitamin D levels and greater common carotid artery intimal thickness values in HIV positive individuals [14]. Severe vitamin D deficiency (serum 25-hydroxycholecalciferol (25-OHD) concentration $<10\mu\text{g/ml}$ or $<25\text{nmol/L}$) has also been associated with increased levels of serum inflammatory markers including IL-6 and hsCRP [15] indicating vitamin D deficiency as a risk factor for cardiovascular disease specifically in the HIV population.

Specifically, HIV positive individuals are speculated to be at increased risk of vitamin D deficiency [10,11,13,16–18]. Dark-skinned individuals in the UK have a higher risk of vitamin D deficiency; one study of over eight hundred patients from Birmingham revealed a prevalence of vitamin D deficiency (25-OHD $<10\mu\text{g/L}$) of 26% among Afro-Caribbeans and 31% among Asians compared to 12% of Caucasians [19]. As many HIV positive individuals in the UK are migrants from African countries, this may in part explain the high prevalence observed.

Recent studies in the UK investigating the prevalence of vitamin D deficiency in HIV positive individuals have involved populations where the majority of individuals were of black ethnicity [16,17]. Importantly, previous studies have also investigated the prevalence of vitamin D deficiency predominantly in individuals established on antiretroviral treatment (ART) [10,16–18]. ART itself is associated with vitamin D deficiency, non-nucleoside reverse transcriptase inhibitors (NNRTI's) and protease inhibitors (PI's) appear to interfere with vitamin D metabolism [17,18,20–22] whereas tenofovir may be a culprit by decreasing bone mineralisation and causing secondary hyperparathyroidism through effects on renal phosphate regulation [16,23]. The impact of immune reconstitution on vitamin D levels is less clear.

A recent systematic review of the literature on vitamin D deficiency in HIV infection reported that vitamin D deficiency, secondary hyperparathyroidism and low bone mineral density were common in HIV positive patients but that HIV infection may not be specifically associated with vitamin D deficiency (24). Importantly all the studies reviewed investigating vitamin D deficiency were observational studies that included HIV positive patients on ART or who were ART experienced. Although some adjusted for ART use and individual ART/drug classes, they are still subject to confounding.

The objective of this study was to describe the prevalence of vitamin D deficiency specifically among ART-naïve individuals newly diagnosed with HIV in a central London clinic.

Methods

Patients registering for HIV care between 01/01/2008 and 31/12/2009 for the first time at the Mortimer Market centre, a large HIV clinic in central London were included in this retrospective record review. During this period, serum 25-OHD levels were measured routinely for new patients registering for HIV care. We restricted the analysis to ART-naïve patients with an available serum 25-OHD result, all patients included were not taking vitamin D supplements. Vitamin D status was defined as “normal” when 25-OHD was $>50\text{nmol/L}$, “deficient” when $25\text{-OHD} \leq 50\text{nmol/L}$ and “severely deficient” when $25\text{-OHD} \leq 25\text{nmol/L}$.

We reviewed data fields including demographics, CD4 cell count, HIV RNA viral load; self-reported ethnicity; season when the 25-OHD sample was taken (winter [December – February]; spring [March – May]; summer [June – August]; autumn [September - November]); serum glucose, calcium, phosphate, alkaline phosphatase (ALP) and HIV clinical stage (asymptomatic; symptomatic; AIDS, as recorded by the clinician).

Continuous data were grouped into categories and Fisher’s exact test was used to assess associations. Associations with vitamin D deficiency were explored by calculating unadjusted odds ratios.

This was a retrospective audit of anonymised data and specific ethical approval was not requested.

Results

Among 253 patients fulfilling inclusion criteria, 207 (81.8%) were male and the median age was 36 years (range 16 - 75 years). Among 233 with ethnicity data available, 150 (64.4%) were white, and 83 (35.6%) black or other ethnicity. The median CD4 count and HIV viral load were 450 cells/mm³ (range 10 – 1100 cells/mm³) and 8,600 copies/mL (range; less than 50copies/ml – 11,000,000 copies/mL) respectively. 236 (93.3%) presented with asymptomatic HIV infection.

Among 253, 148 (58.5%) patients were 25-OHD deficient (25-OHD \leq 50nmol/L), including 32 (12.6%) who were severely deficient (25-OHD \leq 25 nmol/L) (Table 1).

Among 83 patients of non-white ethnicity 61, (73.5%) were 25-OHD deficient compared with 50.7% (76/150) of those reporting white ethnicity ($p < 0.001$).

Seven of eight (87.5%) patients with hypocalcaemia (< 2.12 mmol/L) were 25-OHD deficient, as were seven of 15 (46.7%) with hypophosphataemia (< 0.8 mmol/L) and seven of 11 (63.6%) with high alkaline phosphatase (> 130 IU/L). There was no association between 25-OHD deficiency and hypophosphataemia. The prevalence of 25-OHD deficiency was higher in winter and spring vs. summer and autumn (89/129 [69.0%] vs. 59/124 [47.6%], $p < 0.001$). Serum 25-OHD deficiency was not associated with age, gender, CD4 count, HIV viral load, serum glucose or HIV clinical stage.

On univariable analysis, there was trend towards vitamin D deficiency and hypocalcaemia [$p = 0.14$, odds ratio 5.44 (95% confidence interval 0.7 – 45.0)] larger numbers may have produced a significant association and a smaller confidence interval with these results reflecting small numbers. Season and ethnicity were the only factors

significantly associated with vitamin D deficiency on univariable analysis (where $p < 0.05$), therefore unadjusted odds ratios were calculated to report measures of association for season and ethnicity (see table).

Discussion

This study found a high prevalence of vitamin D deficiency among newly diagnosed, ART-naïve patients; which, therefore cannot be attributed to the effect of antiretroviral agents. Our study used a vitamin D level of ≤ 50 nmol/L to define deficiency, optimal vitamin D status is reflected by serum concentrations of 25-OHD of 75 nmol/l (30 μ g/l) and above [5]. We found that 58.5% were deficient including 75% reporting black ethnicity and 51% of those reporting white ethnicity. This compares with a study in South London which found that 57% (131 of 227 HIV positive subjects) were vitamin D deficient (< 50 nmol/L), however in their study population only 15% (33/227) were ART-naïve and over two thirds of the sample were of black ethnicity [16]. Another study in South London of 1077 HIV positive patients found that 73.5% were vitamin D deficient (< 50 nmol/L) but only 12.4% of their study population were ART-naïve and 60% were of black ethnicity [17].

A recent study investigating vitamin D levels in Swiss HIV-positive patients found a positive association between vitamin D and time since HIV diagnosis on multivariable analysis. They speculated that this may be related to the potential poorer health of recently diagnosed patients, however the relationship to ART use was unclear [18].

As expected, we found a higher prevalence of vitamin D deficiency in non-white ethnic groups and an association in the winter and spring months [1,8,16–19]. There were no other demographic factors, which identified individuals at higher risk. Among biochemical variables, most of our patients with hypocalcaemia were vitamin D deficient; but a significant association was not observed probably due to relatively small numbers. Other markers of bone metabolism (phosphate, alkaline phosphatase) were not associated with vitamin D deficiency.

Our study population was predominantly white males with asymptomatic HIV infection, however it has demonstrated a high prevalence of vitamin D deficiency in HIV positive, ART-naïve individuals, which thus cannot be attributed to ART. The only clinical factor associated with vitamin D deficiency was non-white ethnicity. This suggests that all HIV positive individuals should be considered for routine screening, particularly individuals of non-white ethnicity. CD4 count, HIV viral load and HIV clinical staging do not help to identify those at risk, but low serum calcium should prompt investigation of 25-OHD levels.

References

1. Tahrani AA, Ball A, Shepherd L, Rahim A, Jones AF, Bates A. The prevalence of vitamin D abnormalities in South Asians with type 2 diabetes mellitus in the UK. *Int J Clin Pract*. 2010 Feb;64(3):351–5.
2. Cheng S, Massaro JM, Fox CS, Larson MG, Keyes MJ, McCabe EL, et al. Adiposity, cardiometabolic risk, and vitamin D status: the Framingham Heart Study. *Diabetes*. 2010 Jan;59(1):242–8.
3. Judd SE, Tangpricha V. Vitamin D deficiency and risk for cardiovascular disease. *Am J Med Sci*. 2009 Jul;338(1):40–4.
4. Kilkkinen A, Knekt P, Aro A, Rissanen H, Marniemi J, Heliövaara M, et al. Vitamin D status and the risk of cardiovascular disease death. *Am J Epidemiol*. 2009 Oct 15;170(8):1032–9.
5. Pearce SHS, Cheetham TD. Diagnosis and management of vitamin D deficiency. *BMJ*. 2010;340:b5664.
6. Yamshchikov AV, Desai NS, Blumberg HM, Ziegler TR, Tangpricha V. Vitamin D for Treatment and Prevention of Infectious Diseases: A Systematic Review of Randomized Controlled Trials. *Endocrine Practice*. 2009 Jul 1;15(5):438–49.
7. Martineau AR, Honecker FU, Wilkinson RJ, Griffiths CJ. Vitamin D in the treatment of pulmonary tuberculosis. *J Steroid Biochem Mol Biol*. 2007 Mar;103(3-5):793–8.
8. Ustianowski A, Shaffer R, Collin S, Wilkinson RJ, Davidson RN. Prevalence and associations of vitamin D deficiency in foreign-born persons with tuberculosis in London. *J Infect*. 2005 Jun;50(5):432–7.
9. Villamor E. A potential role for vitamin D on HIV infection? *Nutr Rev*. 2006 May;64(5 Pt 1):226–33.
10. Fox J, Peters B, Prakash M, Arribas J, Hill A, Moecklinghoff C. Improvement in vitamin D deficiency following antiretroviral regime change: Results from the MONET trial. *AIDS Res Hum Retroviruses*. 2011 Jan;27(1):29–34.
11. Yin M. Vitamin D, bone, and HIV infection. *Top Antivir Med*. 2012 Dec;20(5):168–72.
12. Mehta S, Hunter DJ, Mugusi FM, Spiegelman D, Manji KP, Giovannucci EL, et al. Perinatal outcomes, including mother-to-child transmission of HIV, and child mortality and their association with maternal vitamin D status in Tanzania. *J Infect Dis*. 2009 Oct 1;200(7):1022–30.

13. Mehta S, Giovannucci E, Mugusi FM, Spiegelman D, Aboud S, Hertzmark E, et al. Vitamin D status of HIV-infected women and its association with HIV disease progression, anemia, and mortality. *PLoS ONE*. 2010;5(1):e8770.
14. Ross AC, Judd S, Kumari M, Hileman C, Storer N, Labbato D, et al. Vitamin D is linked to carotid intima-media thickness and immune reconstitution in HIV-positive individuals. *Antivir Ther (Lond)*. 2011;16(4):555–63.
15. Ansemant T, Mahy S, Piroth C, Ornetti P, Ewing S, Guiland J-C, et al. Severe hypovitaminosis D correlates with increased inflammatory markers in HIV infected patients. *BMC Infect Dis*. 2013;13:7.
16. Rosenvinge MM, Gedela K, Copas AJ, Wilkinson A, Sheehy CA, Bano G, et al. Tenofovir-linked hyperparathyroidism is independently associated with the presence of vitamin D deficiency. *J Acquir Immune Defic Syndr*. 2010 Aug;54(5):496–9.
17. Welz T, Childs K, Ibrahim F, Poulton M, Taylor CB, Moniz CF, et al. Efavirenz is associated with severe vitamin D deficiency and increased alkaline phosphatase. *AIDS*. 2010 Jul 31;24(12):1923–8.
18. Mueller NJ, Fux CA, Ledergerber B, Elzi L, Schmid P, Dang T, et al. High prevalence of severe vitamin D deficiency in combined antiretroviral therapy-naïve and successfully treated Swiss HIV patients. *AIDS*. 2010 May 15;24(8):1127–34.
19. Ford L, Graham V, Wall A, Berg J. Vitamin D concentrations in an UK inner-city multicultural outpatient population. *Ann Clin Biochem*. 2006 Nov;43(Pt 6):468–73.
20. Brown TT, McComsey GA. Association between initiation of antiretroviral therapy with efavirenz and decreases in 25-hydroxyvitamin D. *Antivir Ther (Lond)*. 2010;15(3):425–9.
21. Cozzolino M, Vidal M, Arcidiacono MV, Tebas P, Yarasheski KE, Dusso AS. HIV-protease inhibitors impair vitamin D bioactivation to 1,25-dihydroxyvitamin D. *AIDS*. 2003 Mar 7;17(4):513–20.
22. Bouvier G. Protease inhibitors can interfere with vitamin D metabolism. *HIV Clin*. 2009;21(3):9–10.
23. Klassen K, Martineau AR, Wilkinson RJ, Cooke G, Courtney AP, Hickson M. The Effect of Tenofovir on Vitamin D Metabolism in HIV-Infected Adults Is Dependent on Sex and Ethnicity. Slominski AT, editor. *PLoS ONE*. 2012 Sep 12;7(9):e44845.
24. Childs K, Welz T, Samarawickrama A, Post FA. Effects of vitamin D deficiency and combination antiretroviral therapy on bone in HIV-positive patients. *AIDS*. 2012 Jan 28;26(3):253–62.

Table: Prevalence of vitamin D deficiency by demographic characteristics and relevant laboratory variables

	Total Sample (n=253)	Normal Vitamin D (>50nmol/L)	Deficient Vitamin D (≤50nmol/L)	p-value (Fishers Exact)	Odds ratio (95% confidence interval)
Sex, n (%)					
Male	207 (81.8%)	90 (43.5%)	117 (56.5%)	p=0.19	
Female	46 (18.2%)	15 (32.6%)	31 (67.4%)		
Age Range					
<30	80 (31.6%)	48 (60%)	32 (40%)	p=0.90	
31 – 40	88 (34.8%)	52 (59.1%)	36 (40.9%)		
>40	85 (33.6%)	48 (56.5%)	37 (43.5%)		
Ethnicity, n (%) (excl unknown, n=233)					
White	150 (59.3%)	74 (49.3%)	76 (50.7%)	p=0.01	
Black	63 (24.9%)	16 (25.4%)	47 (74.6%)		
Other/non-white (Asian, Chinese, mixed race)	20 (8.6%)	6 (30%)	14 (70%)		
Ethnicity, n (%) (excl unknown, n=233)					
White	150 (64.4%)	74 (49.3%)	76 (50.7%)	p<0.001	2.7 (1.5 – 4.8)
Non white	83 (35.6%)	22 (55.4%)	61 (73.5%)		
Season, n (%)					
Summer/Autumn	124 (49%)	65 (52.4%)	59 (47.6%)	p=0.001	2.5 (1.5 – 4.1)
Winter/Spring	129 (51%)	40 (31%)	89 (69%)		
KC60* HIV Stage, n (%)					
E1 (asymptomatic)	236 (93.3%)	135 (57.2%)	101 (42.8%)	p=0.241	
E2 (symptomatic)	7 (2.8%)	6 (85.7% %)	1 (14.3%)		
E3 (AIDS)	10 (3.95%)	7 (70%)	3 (30%)		

CD4 count (cells/mm³ x10) (missing values) Median IQR	(12/253) 0.45 0.60 – 0.30	(9/148) 0.43 0.59 – 0.25	(3/105) 0.47 0.61 – 0.36	p=0.651	
HIV viral load (copies/mL) (missing values) Median IQR	(9/253) 8,600 66,500 – 335	(6/148) 8,450 70,000 – 110	(3/105) 11,500 62,000 – 730	p=0.374	
Calcium (mmol/L) Normal (2.12 -2.65) Hypocalcaemia (<2.12)	224(96.5%) 8 (3.5%)	98 (43.7%) 1 (12.5%)	126 (56.3%) 7 (87.5%)	p=0.14	5.4 (0.7 – 45.0)
Phosphate (mmol/L) Normal (0.8-1.4) Hypophosphataemia (<0.8)	210 (90.5%) 15 (6.5%)	89 (42.4%) 8 (53.3%)	121 (57.6%) 7 (46.7%)	p=0.43	
ALP (IU/L) Normal (30-130) High ALP (>130)	233 (95.1%) 11 (4.5%)	98 (42.1%) 4 (36.4%)	135 (57.9%) 7 (63.6%)	p=0.77	

* the KC60 code, is a Health Protection Agency (HPA) required ‘diagnoses made’ and ‘services provided’ code provided by all genitourinary clinic services in the UK for each patient presentation; for the purpose of internal management, audit and reporting data to the HPA.