

MPRA

Munich Personal RePEc Archive

Continuity and completeness of strongly independent preorders

David McCarthy and Kalle Mikkola

Dept. of Philosophy, University of Hong Kong, Hong Kong, Dept. of Mathematics and Systems Analysis, Aalto University, Finland

17 June 2017

Online at <https://mpa.ub.uni-muenchen.de/79757/>

MPRA Paper No. 79757, posted 19 June 2017 14:29 UTC

Continuity and completeness of strongly independent preorders*

David McCarthy[†] Kalle Mikkola[‡]

Abstract

A strongly independent preorder on a possibly infinite dimensional convex set that satisfies two of the following conditions must satisfy the third: (i) the Archimedean continuity condition; (ii) mixture continuity; and (iii) comparability under the preorder is an equivalence relation. In addition, if the preorder is nontrivial (has nonempty asymmetric part) and satisfies two of the following conditions, it must satisfy the third: (i') a modest strengthening of the Archimedean condition; (ii) mixture continuity; and (iii') completeness. Applications to decision making under conditions of risk and uncertainty are provided.

1 Introduction and main results

The completeness axiom of expected utility has long been regarded as dubious, while the usual continuity axioms are typically seen as innocuous. However, given a strongly independent preorder on a convex set, we show that the standard Archimedean and mixture continuity axioms together imply that the possibilities for incompleteness are highly restricted, in a sense made precise below. In particular, they rule out the most natural preference structures for agents who find they cannot exactly compare two alternatives. If the Archimedean axiom is slightly strengthened in a natural direction, the room for incompleteness vanishes entirely: the preorder must be complete. These claims generalize results of Aumann (1962) and Dubra (2011).

*David McCarthy thanks the Research Grants Council of the Hong Kong Special Administrative Region, China (HKU 750012H) for support. We are grateful to Teruji Thomas for comments.

[†]Corresponding author, Dept. of Philosophy, University of Hong Kong, Hong Kong, davidmccarthy1@gmail.com.

[‡]Dept. of Mathematics and Systems Analysis, Aalto University, Finland, kalle.mikkola@iki.fi.

In more detail, let X be a nonempty convex set, and \succsim_X a preorder (a reflexive, transitive binary relation) on X . Consider the following axioms. The first is the standard strong independence axiom.

(SI) For $x, y, z \in X$ and $\alpha \in (0, 1)$,

$$x \succsim_X y \iff \alpha x + (1 - \alpha)z \succsim_X \alpha y + (1 - \alpha)z.$$

Thus \succsim_X is an ‘SI preorder’. We will be considering the following three Archimedean or continuity axioms.

(Ar) For $x, y, z \in X$, if $x \succ_X y \succ_X z$, then $(1 - \epsilon)x + \epsilon z \succ_X y$ for some $\epsilon \in (0, 1)$.

(Ar⁺) For $x, y, z \in X$, if $x \succ_X y$, then $(1 - \epsilon)x + \epsilon z \succ_X y$ for some $\epsilon \in (0, 1)$.

(MC) For $x, y, z \in X$, if $\epsilon x + (1 - \epsilon)y \succ_X z$ for all $\epsilon \in (0, 1]$, then $y \succsim_X z$.

The axiom Ar is weaker than, but for SI preorders equivalent to, the standard Archimedean axiom introduced by Blackwell and Girshick (1954).¹ It is weaker than the axiom Ar⁺, essentially introduced by Aumann (1962).² But both Ar and Ar⁺ express a similar heuristic. Suppose x is strictly preferred to y , and z is some third alternative. Then Ar says that z cannot be so radically worse than y that a sufficiently small chance of z would disturb the original preference. The axiom Ar⁺ extends this by replacing ‘worse than’ with ‘worse than or incomparable with’. For SI preorders, MC is equivalent to the ‘mixture-continuity’ axiom of Herstein and Milnor (1953), that $\{\alpha \in [0, 1] : \alpha x + (1 - \alpha)y \succsim_X z\}$ is closed in $[0, 1]$. The displayed formulation is especially normatively natural, and suggests that MC should be seen as just as much an Archimedean condition as Ar and Ar⁺.

The final two axioms restrict the possibilities for incomparability. Say that two members x and y of X are *comparable* if $x \succsim_X y$ or $y \succsim_X x$. They are *incomparable* if they are not comparable. They have a common upper bound if $z \succsim_X x$ and $z \succsim_X y$ for some $z \in X$, and similarly for common lower bound. The next axiom is nonstandard, while the last is the standard completeness axiom.

(Eq) Comparability is an equivalence relation.

¹See Proposition 6 below.

²The axiom Aumann actually discusses is $\epsilon_0 x + (1 - \epsilon_0)z \succ_X y \Rightarrow \epsilon x + (1 - \epsilon)z \succ_X y$ for all ϵ close enough to ϵ_0 , but for SI preorders, this is equivalent to Ar⁺.

(C) All members of X are comparable.

Note that Eq is equivalent to the claim that pairs of incomparable elements have neither a common upper bound nor a common lower bound. This is a demanding requirement. In realistic cases where an agent finds it hard to compare two alternatives, she will find it easy to imagine an alternative that she finds superior to both, and also one she finds inferior.

To state our main result, say that \succsim_X is *nontrivial* if it has a nonempty strict part; that is, for some $x, y \in X$, $x \succ_X y$.

Theorem 1. *For any SI preorder \succsim_X on a convex set X :*

- (1) *Any two of the following imply the third: MC, Ar and Eq.*
- (2) *For nontrivial \succsim_X , any two of the following imply the third: MC, Ar⁺, and C.*

The conditions in (1) do not suffice for C for trivial or nontrivial SI preorders, and nontriviality is essential to (2):

- Example 2.** (a) Let X contain at least two elements. Set $x \succsim_X y \Leftrightarrow x = y$. Then \succsim_X is trivial, Ar, Ar⁺ and MC hold, but C fails.
 (b) Let $X = \mathbb{R}^2$. Set $x \succsim_X y \Leftrightarrow x_1 \geq y_1 \wedge x_2 = y_2$. Then \succsim_X is nontrivial, Ar, MC and Eq hold, but Ar⁺ and C fail.

Theorem 1 has several precedents. In a seemingly overlooked observation, Aumann (1962) claimed without proof that MC and Ar⁺ imply Eq. Thus both parts of the theorem strengthen his claim. Aumann also claimed that when $X = V$ is a finite dimensional vector space, MC and Ar⁺ imply that X may be written as the direct sum of two subspaces such that two elements are comparable if and only if their second coordinates are identical. The following strengthens this claim, by dropping finite dimensionality and weakening Ar⁺ to Ar.

Corollary 3. *Suppose \succsim_X is an SI preorder satisfying Ar and MC, and that $X = V$ is a vector space. Then $V = V_1 \oplus V_2$, and $v, w \in V$ are comparable if and only if $v_2 = w_2$.*

In the case where X is the set of probability functions on a given finite set, and thus can be identified with the standard simplex of a finite dimensional vector space, the second part of Theorem 1 was proved by Dubra (2011), building on Schmeidler (1971). Dubra's proof makes essential use of finite dimensionality. But placing no restrictions on the dimension of X allows

for considerably broader applications,³ including general sets of probability measures.

Schmeidler’s result was that if a nontrivial preorder on a connected topological set has closed weak upper and lower contour sets, and open strict upper and lower contour sets, it must be complete. The axioms we discuss are purely algebraic, making them applicable to cases in which X is not naturally equipped with a topology. Our proof of Theorem 1 is purely algebraic. Indeed the main technical tool, stated in Theorem 7 below, states equivalences between the three continuity conditions and conditions involving algebraic openness or closedness in ambient vector spaces.

1.1 Discussion

The abstract structure of incomplete SI preorders on convex sets has been discussed, but the relevance of Theorem 1 perhaps has more to do with its compatibility with the typical concrete settings that are used to represent objective risk and subjective uncertainty. To illustrate, let Y be an arbitrary set, Y_c be a compact metric space, and Y_m an arbitrary measurable space; these are typical consequence spaces. Let $P(Y)$ be the set of finitely supported probability measures on Y , $P(Y_c)$ be the set of Borel probability measures on Y_c , and $P(Y_m)$ be an arbitrary convex set of probability measures on Y_m . These are obviously all convex sets, and cover typical cases involving objective risk. Let S_0 and S be a finite and arbitrary sets of states of nature respectively. Then $P(Y)^{S_0}$ is the set of Anscombe-Aumann ‘horse lotteries’. Here, members of S_0 are bearers of subjective uncertainty, while the outcomes of horse lotteries are ‘roulette lotteries’ involving objective risk. For any $x, y \in P(Y)^{S_0}$ and $\alpha \in [0, 1]$, $\alpha x + (1 - \alpha)y \in P(Y)^{S_0}$ is defined by setting $(\alpha x + (1 - \alpha)y)(s) = \alpha x(s) + (1 - \alpha)y(s)$ for any $s \in S_0$, making $P(Y)^{S_0}$ into a convex set. Finally, the set Y^S is the set of Savage-acts associating states of nature with consequences; states of nature continue to be the bearers of subjective uncertainty, but no objective risk is modelled. The space Y^S is not naturally a convex set, but given a preorder on Y^S that satisfies reasonably modest axioms, Y^S can be endowed with convex structure; see for example Ghirardato, Maccheroni, Marinacci and Siniscalchi (2003). The importance of allowing X to be infinite dimensional can be seen from the fact that none of these typical domains can be identified with a finite dimensional X . There are many works discussing incomplete SI preorders in the settings just mentioned. Some focus only on SI strict partial orders,

³Recall that the dimension of a convex set X is the dimension of $\text{Span}(X - X)$, or, equivalently, the dimension of the smallest affine space containing X .

but these are consistent with our model as they can be seen as studies of the class of SI preorders that are compatible with the partial orders.⁴

Given Theorem 7, it is natural to think of Ar and Ar⁺ as ‘open’ conditions, and MC as a ‘closed’ condition. Both styles of condition have been used extensively in discussions of incomplete SI preorders on the kinds of convex sets just described. In almost every case we know of,⁵ the open conditions are at least as strong as Ar in the given model, and the closed conditions are at least as strong as, and typically much stronger than, MC.⁶ Thus Theorem 1 has considerable relevance.

The continuity conditions in the literature just mentioned are typically presented without discussion, and are sometimes said just to be ‘technical assumptions’. But it is not so clear what this means. Each of our three continuity conditions expresses a normatively natural idea that might nevertheless be questioned, but so too are conditions like completeness and strong independence. Both Ar⁺ and MC express different ways of ruling out infinitesimal value differences, and since this basic idea is so widely accepted in discussions of complete SI preorders (when all three continuity conditions are equivalent), it is rather remarkable that ruling out infinitesimal value differences across the board forces one to accept the heavily criticized completeness axiom. Thus we suggest that Theorem 1 deserves to be seen as an impossibility result. We end by canvassing some possible responses.

First, one could try to argue that of the two styles of continuity condition, open and closed, one is more normatively or descriptively plausible than the other. But we side with Aumann (1962) and Manzini and Mariotti (2008) in thinking that Ar⁺ and MC are comparably plausible in the abstract. Thus such an argument would have to pay attention to the specific interpretation of the preorder in question. Second, for applications, one could try to develop mirror theories for open and closed conditions, and analyze

⁴This class is always nonempty, as the reflexive closure of an SI strict partial order is an SI preorder whose asymmetric part is identical to the partial order.

⁵The exceptions are Aumann (1962), who imposes a continuity condition that is strictly weaker than both Ar and MC, and Seidenfeld, Schervish and Kadane (1995) who impose a similar condition in the Anscombe-Aumann setting.

⁶For open conditions, see Bewley (2002); Manzini and Mariotti (2008); Galaabaatar and Karni (2012, 2013); Evren (2014); McCarthy, Mikkola, and Thomas (2017b). For closed conditions, see Shapley and Baucells (1998); Ghirardato *et al* (2003); Dubra, Maccheroni and Ok (2004); Nau (2006); Baucells and Shapley (2008); Evren (2008); Kopylov (2009); Gilboa, Maccheroni, Marinacci and Schmeidler (2010); Danan, Guerdjikova and Zimper (2012); Ok, Ortoleva and Riella (2012); McCarthy, Mikkola, and Thomas (2017a). Without any continuity condition, one faces incomplete analogues of the situation first discussed by Hausner and Wendel (1952), discussed further in McCarthy, Mikkola, and Thomas (2017c).

the sensitivity of their implications to these conditions. Third, one could choose between the two styles of conditions on the basis of the convenience of the representation theorems they support (compare Evren, 2014). Fourth, to try to bypass the impossibility, one could adopt a nonstandard model of the relationship between strict partial orders and associated preorders (see further Karni, 2007; Galaabaatar and Karni, 2012). Fifth, one could argue that in some settings, the case for both styles of condition is strong enough that they provide a novel normative argument for completeness (in a different context, compare Broome, 1999), or even a new argument against strong independence.

2 Proofs

2.1 Preliminaries

When X is a nonempty convex set of a vector space V , the following provides a useful representation of the subspace $\text{Span}(X - X)$.

Lemma 4. *Let X be a nonempty convex subset of a vector space V . Then*

$$\text{Span}(X - X) = \{\lambda(x - x') : x, x' \in X, \lambda > 0\}.$$

Proof. The right-hand side is clearly included in the left. For the converse, let $v \in \text{Span}(X - X)$. The case $v = 0$ is trivial, so let $v = \sum_{i=1}^n \lambda_i(x_i - x'_i)$ with $x_i, x'_i \in X$, $\lambda_i \neq 0$ for all i , and $n \in \mathbb{N}$. Exchange x_i with x'_i if necessary to have each $\lambda_i > 0$. Set $\lambda = \sum_{i=1}^n \lambda_i$, $x = \frac{1}{\lambda} \sum_{i=1}^n \lambda_i x_i$, and $x' = \frac{1}{\lambda} \sum_{i=1}^n \lambda_i x'_i$. Then $x, x' \in X$ by convexity, and $v = \lambda(x - x')$ as needed. \square

Recall that a vector preorder \succsim_V is a preorder on a vector space V such that for any $v, w, u \in V$ and $\alpha > 0$, $v \succsim_V w$ implies $\alpha v + u \succsim_V \alpha w + u$. We define $\{\succsim_V 0\} := \{v \in V : v \succsim_V 0\}$, and similarly $\{\succ_V 0\}$, $\{\sim_V 0\}$. We also define \succsim_X , \succsim_V and sets such as $\{\succsim_V 0\}$ in the obvious way; for example, $x \succsim_X y \Leftrightarrow y \succsim_X x$.

Proposition 5. *Let \succsim_X be a SI preorder on a nonempty convex subset X of a vector space V . For any $v, w \in V$, define \succsim_V by*

$$v \succsim_V w \iff v - w = \lambda(x - y) \text{ for some } x, y \in X, \lambda > 0 \text{ with } x \succsim_X y.$$

Then \succsim_V is a vector preorder on V , and \succsim_X is its restriction to $X \times X$. Moreover, \succsim_X is complete if and only if \succsim_V is complete on $\text{Span}(X - X) = \text{Span}\{\succsim_V 0\}$.

Proof. Clearly \succsim_V is reflexive. Suppose $u \succsim_V v$ and $v \succsim_V w$. Then for some $\lambda, \mu > 0$ and $x_1, x_2, y_1, y_2 \in X$, we have $u - v = \lambda(x_1 - x_2)$, $v - w = \mu(y_1 - y_2)$, $x_1 \succsim_X x_2$ and $y_1 \succsim_X y_2$. The former imply $u - w = (\lambda + \mu)\left(\frac{\lambda x_1 + \mu y_1}{\lambda + \mu} - \frac{\lambda x_2 + \mu y_2}{\lambda + \mu}\right)$; the latter and applications of SI imply $\frac{\lambda x_1 + \mu y_1}{\lambda + \mu} \succsim_X \frac{\lambda x_2 + \mu y_2}{\lambda + \mu}$. It follows that $u \succsim_V w$, establishing transitivity of \succsim_V , so \succsim_V is a preorder on V .

Suppose $v \succsim_V w$, $u \in V$, $\alpha > 0$. For some $\lambda > 0$, $x, y \in X$, we have $v - w = \lambda(x - y)$ with $x \succsim_X y$. Then $(\alpha v + u) - (\alpha w + u) = \alpha\lambda(x - y)$. This implies $\alpha v + u \succsim_V \alpha w + u$, so \succsim_V is a vector preorder.

Clearly $x \succsim_X y$ implies $x \succsim_V y$. Conversely, suppose $x \succsim_V y$ for some $x, y \in X$. Then for some $x', y' \in X$, $\lambda > 0$, $x - y = \lambda(x' - y')$ with $x' \succsim_X y'$. The former implies $\alpha x + (1 - \alpha)y' = \alpha y + (1 - \alpha)x'$ where $\alpha := \frac{1}{1 + \lambda}$; the latter and SI imply $\alpha x + (1 - \alpha)x' \succsim_X \alpha y + (1 - \alpha)y'$. Substituting, then using SI again, yields $x \succsim_X y$, hence \succsim_X is the restriction of \succsim_V .

The completeness claim follows from Lemma 4. \square

A convex set can always be embedded in a vector space, so without loss of generality we henceforth assume that (X, \succsim_X) and (V, \succsim_V) are as in Proposition 5. It clearly follows that $\text{Span}\{\succsim_V 0\} \subset \text{Span}(X - X)$.

2.2 Algebraic conditions

The following assembles facts about the Archimedean conditions.

Proposition 6.

(a) *Ar is equivalent to each of the following two conditions.*

$$\begin{aligned} \text{For all } x, y, z \in X: x \succ_X y \text{ and } y \succ_X z &\implies (1 - \epsilon)x + \epsilon z \succ_X y & (1) \\ \text{and } y \succ_X \epsilon x + (1 - \epsilon)z &\text{ for some } \epsilon \in (0, 1). \end{aligned}$$

$$\text{For all } v, w \in V: v, w \succ_V 0 \implies v \succ_V \epsilon w \text{ for some } \epsilon > 0. \quad (2)$$

(b) *If one of Ar, (1), (2) and Ar^+ holds for some ϵ_0 , it holds for all $\epsilon \in [0, \epsilon_0]$.*

(c) *Ar^+ implies Ar (but not conversely).*

Proof. (a) To show (2) \implies (1), assume (2) and suppose $x \succ_X y \succ_X z$. Set $x_\alpha := (1 - \alpha)(x - y) + \alpha(z - y) \forall \alpha \in [0, 1]$. Then $x_0 \succ_V 0 \succ_V x_1$, and from (2) one deduces that $(1 - \epsilon)x_0 \succ_V \epsilon(-x_1)$ for small $\epsilon > 0$, i.e. $(1 - \epsilon)(x - y) \succ_V \epsilon(y - z)$, implying $(1 - \epsilon)x + \epsilon z \succ_X y$ for some $\epsilon \in (0, 1)$. From (2) one also deduces $(1 - \epsilon)(-x_1) \succ_V \epsilon x_0$ for small $\epsilon > 0$, hence $y \succ_X \epsilon x + (1 - \epsilon)z$ for some $\epsilon \in (0, 1)$. This establishes (1), and clearly (1) \implies Ar.

To show Ar \implies (2), assume Ar and suppose $v, w \succ_V 0$. By Proposition 5, $v = \lambda(x - y)$, $w = \mu(s - t)$ for some $x \succ_X y$, $s \succ_X t$, $\lambda, \mu > 0$. By SI, $\frac{1}{2}(x + s) \succ_X \frac{1}{2}(y + s) \succ_X \frac{1}{2}(y + t)$. By Ar, for some $\epsilon \in (0, 1)$, $(1 - \epsilon)\frac{1}{2}(x +$

$s) + \epsilon \frac{1}{2}(y+t) \succ_X \frac{1}{2}(y+s)$. This implies $(1-\epsilon)(x-y) \succ_V \epsilon(s-t)$, hence $v \succ_V \frac{\epsilon}{1-\epsilon} \cdot \frac{\lambda}{\mu} w$, establishing (2).

(b) Assume Ar^+ , and suppose $x \succ_X y$ and $(1-\epsilon_0)x + \epsilon_0 z \succ_X y$ for some $\epsilon_0 > 0$. These and SI imply $(1-\epsilon)x + \epsilon z \succ_X y$ for all $\epsilon \in [0, \epsilon_0]$. The claims about Ar and (1) are proved similarly, and the claim about (2) is clear.

(c) That Ar^+ implies Ar is obvious. The failure of the converse is shown by Example 2(b). \square

Recall that a subset S of a vector space W is *algebraically open in W* if for all $v \in S$, $w \in W$, $v + \epsilon w \in S$ for all sufficiently small $\epsilon > 0$. S is *algebraically closed* if for all $v, w \in W$: $(1-\alpha)v + \alpha w \in S$ for all $\alpha \in (0, 1] \Rightarrow v \in S$. Given $v, w \in W$, we sometimes write $[v, w] \subset W$ for the line segment $\{(1-\alpha)v + \alpha w : \alpha \in [0, 1]\}$. Then S is algebraically closed if $w \in S$ whenever $[v, w] \subset S$. The following connects these algebraic notions with our continuity axioms.

Theorem 7.

- (a) Ar holds if and only if $\{\succ_V 0\}$ is algebraically open in $\text{Span}\{\succ_V 0\}$.⁷
- (b) Ar^+ holds if and only if $\{\succ_V 0\}$ is algebraically open in $\text{Span}(X - X)$.
- (c) MC holds if and only if $\{\succ_V 0\}$ is algebraically closed.

Proof. (a) Suppose \succ_X satisfies Ar . Let $v \in \{\succ_V 0\}$, $w \in \text{Span}\{\succ_V 0\}$. Clearly we can write $w = a - b$ where each of a and b is either 0 or in $\{\succ_V 0\}$. Since \succ_V is a vector preorder, $v + \epsilon_1 a \succ_V 0$ for all $\epsilon_1 > 0$. By Proposition 6(a)(2), we have $v \succ_V \epsilon_2 b$ for all sufficiently small $\epsilon_2 > 0$. These imply $v + \epsilon w \succ_V 0$ for all small enough $\epsilon > 0$. This shows that $\{\succ_V 0\}$ is algebraically open in $\text{Span}\{\succ_V 0\}$.

Conversely, suppose $\{\succ_V 0\}$ is algebraically open in $\text{Span}\{\succ_V 0\}$, and that $v, w \succ_V 0$. Since $-w \in \text{Span}\{\succ_V 0\}$, $v + \epsilon(-w) \in \{\succ_V 0\}$ for all sufficiently small $\epsilon > 0$. By Proposition 6, \succ_X satisfies Ar .

(b) Assume Ar^+ . Let $c \in \{\succ_V 0\}$, $v \in \text{Span}(X - X)$. By Lemma 4 and Proposition 5, $c = \alpha(x-y)$, $v = \beta(p-q)$ for some $x, y, p, q \in X$ with $x \succ_X y$, $\alpha, \beta > 0$. Then SI, Ar^+ and Proposition 6(b) imply $(1-\epsilon)\frac{1}{2}(x+q) + \epsilon\frac{1}{2}(x+p) \succ_X \frac{1}{2}(y+q)$, hence $(x-y) + \epsilon(p-q) \succ_V 0$, for all for all sufficiently small $\epsilon > 0$. Consequently $c + \epsilon\frac{\alpha}{\beta}v \succ_V 0$ for small enough $\epsilon > 0$.

Conversely, suppose $\{\succ_V 0\} \cap \text{Span}(X - X)$ is algebraically open in $\text{Span}(X - X)$. Suppose $x \succ_X y$ and $z \in X$. Then $x - y \in \{\succ_V 0\}$ and $z - y \in \text{Span}(X - X)$, hence for some $\epsilon \in (0, 1)$, $(1-\epsilon)(x-y) + \epsilon(z-y) \in \{\succ_V 0\}$, implying $(1-\epsilon)x + \epsilon z \succ_X y$.

(c) This is proved in McCarthy *et al* (2017a, Thm. 2.2). \square

⁷The latter condition is often phrased as ‘ $\{\succ_V 0\}$ is relatively algebraically open.’

Corollary 8. \succsim_X satisfies MC if and only if \succsim_X satisfies MC.

Proof. The set $\{\succsim_V 0\} = -\{\succsim_V 0\}$ is algebraically closed if and only if $\{\succsim_V 0\}$ is. \square

2.3 Proof of Theorem 1 and Corollary 3.

Define $x \bowtie_X y \Leftrightarrow x \succsim_X y \vee y \succsim_X x$; that is, x and y are comparable.

Lemma 9. Let $x_i \bowtie_X y_i$, $\alpha_i \in \mathbb{R}$ ($i = 1, \dots, n$). Then $\sum_{i=1}^n \alpha_i(x_i - y_i) = \alpha(p - q)$, where $\alpha > 0$ and $p \bowtie_X z \bowtie_X q$ for some $z \in X$.

Proof. 1° Case $n = 2$. Without loss of generality, assume $\alpha_2, \alpha_1 > 0$; $\alpha_2 \geq \alpha_1$; and $\alpha_2 = 1$. Set $p_k = x_k$, $q_2 = y_2$, $q_1 = (1 - \alpha_1)x_1 + \alpha_1 y_1$ to have $\sum_{i=1}^2 \alpha_i(x_i - y_i) = p_1 - q_1 + p_2 - q_2$. Clearly $x_1 \bowtie_X q_1$. Set $p := \frac{1}{2}(p_1 + p_2)$, $q := \frac{1}{2}(q_1 + q_2)$, $z := \frac{1}{2}(q_1 + p_2)$, to have $\sum_{i=1}^2 \alpha_i(x_i - y_i) = 2(p - q)$ and $p \bowtie_X z \bowtie_X q$.

2° General case. Without loss of generality, assume $x_i \succsim_X y_i$ for all i . If $\alpha_1, \alpha_2 > 0$, then

$$\alpha_1(x_1 - y_1) + \alpha_2(x_2 - y_2) = \alpha'(x'_1 - y'_1)$$

where $\alpha' = \alpha_1 + \alpha_2 > 0$, $x'_1 = (1 - \alpha'')x_1 + \alpha''x_2$, $y'_1 = (1 - \alpha'')y_1 + \alpha''y_2$, $\alpha'' = \alpha_2/(\alpha_1 + \alpha_2)$, and hence $x'_1 \succsim_X y'_1$, by SI. This way, by induction, we combine all terms having $\alpha_i > 0$. If all the α_i are strictly positive (or similarly, strictly negative), the result is immediate. Otherwise, similarly combine the terms with $\alpha_i < 0$, then apply 1° to the two. \square

Lemma 10. Let W be any vector space, and $S \subset W$. If S is algebraically open, then $W \setminus S$ is algebraically closed. The converse holds if S is convex.

Proof. The claim is Ok (2007, Exercise G.1.5.30). The first claim is clear. For the converse, suppose S is not algebraically open. Then for some $v \in S$, $w \in W$, $\{(1 - \alpha)v + \alpha w : \alpha \in [0, \epsilon]\} \not\subset S$ for all $\epsilon > 0$. If S is convex, this implies $\{(1 - \alpha)v + \alpha w : \alpha \in (0, \epsilon_0]\} \subset W \setminus S$ for some $\epsilon_0 > 0$. If $W \setminus S$ is algebraically closed, we have $v \in W \setminus S$, a contradiction. \square

Recall that \succsim_X is nontrivial if $\succ_X \neq \emptyset$.

Proposition 11.

(a) The following conditions are equivalent.

(i) Both Ar and MC hold.

(ii) $\{\succsim_V 0\} = \{\sim_V 0\} + [0, \infty)c$ for some $c \succsim_V 0$.

(iii) $V = V_1 \oplus V_2$ where $V_1 = \{\sim_V 0\} + \mathbb{R}c$ for some $c \succ_V 0$, and $v \bowtie_V w \Leftrightarrow v_2 = w_2$.

(iv) Ar holds and \succ_V is complete on $\text{Span}\{\succ_V 0\}$.

(b) \succ_V is complete on $\text{Span}\{\succ_V 0\}$ if and only if \bowtie_X is an equivalence relation.

(c) If \succ_X is nontrivial, then both Ar⁺ and MC hold if and only if \succ_X is complete and Ar holds.

(d) If \succ_X is nontrivial, then $\text{Span}\{\succ_V 0\} = \text{Span}\{\succ_V 0\}$.

Proof. (a) We show (iii) \Leftrightarrow (ii) \Leftrightarrow (i) \Leftrightarrow (iv). It is clear that (iii) \Leftrightarrow (ii). We obtain (ii) \Rightarrow (i) by using Proposition 6(a)(2) for Ar and by Theorem 7(c) for MC. Conversely, assume (i). If $\succ_V = \emptyset$, (ii) is immediate, so pick $c \succ_V 0$. Suppose for a contradiction $\{\succ_V 0\} \neq \{\sim_V 0\} + [0, \infty)c =: Q$. Clearly Q is contained in $\{\succ_V 0\}$, so there exists $d \succ_V 0$ such that $d \notin Q$. By Ar and Theorem 7(a), since $\{\succ_V 0\}$ is algebraically open in $\text{Span}\{\succ_V 0\}$, $\alpha(-c) + (1 - \alpha)d \succ_V 0$ for sufficiently small $\alpha > 0$. Since \succ_V is a vector preorder, the set of such α is an interval and is bounded above by 1. Let α_0 be its supremum, and set $e := \alpha_0(-c) + (1 - \alpha_0)d$. By MC and Theorem 7(c), $e \succ_V 0$, and hence $\alpha_0 \in (0, 1)$. By Ar and Theorem 7(a), $e \not\sim_V 0$. Hence $e \sim_V 0$, implying $(1 - \alpha_0)d \sim_V e + \alpha_0 c \in Q$, a contradiction. Thus (ii) \Leftrightarrow (i).

Assume (i). Since (i) implies (ii), we have $\{\succ_V 0\} = \{\sim_V 0\} + [0, \infty)c$ and $\text{Span}\{\succ_V 0\} = \{\sim_V 0\} + \mathbb{R}c$ for some $c \succ_V 0$, so \succ_V is complete on $\text{Span}\{\succ_V 0\}$, establishing (iv). Conversely, assume (iv), and for a contradiction suppose MC does not hold. By Theorem 7(c), there is some $[a, b) \subset \{\succ_V 0\}$ with $b \not\sim_V 0$. Clearly $b \in \text{Span}\{\succ_V 0\}$, so by completeness of \succ_V on $\text{Span}\{\succ_V 0\}$, $-b \succ_V 0$. This is a contradiction, since Ar and Theorem 7(a) imply $\{\succ_V 0\}$ is algebraically open in $\text{Span}\{\succ_V 0\}$, but $[-a, -b) \subset \{\succ_V 0\}$. Hence (i) \Leftrightarrow (iv).

(b) Suppose \succ_V is complete on $\text{Span}\{\succ_V 0\}$. Then $x \bowtie_X y$ and $y \bowtie_X z \Rightarrow x - y, y - z \in \pm\{\succ_V 0\} \subset \text{Span}\{\succ_V 0\}$, implying $x - z \in \text{Span}\{\succ_V 0\}$, so $x \bowtie_X z$, implying \bowtie_X is transitive, and hence an equivalence relation as it is clearly reflexive and symmetric.

Conversely, suppose \bowtie_X is an equivalence relation. Let $v \in \text{Span}\{\succ_V 0\} = \text{Span}\{x - y : x \succ_X y\}$, by Proposition 5. Then $v = \sum_{i=1}^n \alpha_i(x_i - y_i)$ for some $\alpha_i \in \mathbb{R}$, $x_i \succ_X y_i$. By Lemma 9, $v = \alpha(p - q)$, where $\alpha > 0$, $p, q \in X$ with $p \bowtie_X z$ and $z \bowtie_X q$ for some $z \in X$. Transitivity of \bowtie_X implies $p \bowtie_X q$, hence $v \bowtie_V 0$, implying that \succ_V is complete on $\text{Span}\{\succ_V 0\}$.

(c) Assume \succ_X is nontrivial. Suppose \succ_X satisfies Ar⁺ and MC. By Proposition 6(c), \succ_X satisfies Ar. By Theorem 7(b), $\{\succ_V 0\}$ is algebraically open in $\text{Span}(X - X)$. Let $w \in X - X$, and by nontriviality, pick $v \in \{\succ_V 0\}$.

Then $v + \epsilon w \in \{\succ_V 0\}$ for sufficiently small $\epsilon > 0$, so $w \in \text{Span}\{\succ_V 0\}$; that is, $X - X \subset \text{Span}\{\succ_V 0\}$. By (a), \succsim_V is complete on $X - X$, hence \succsim_X is complete.

Conversely, when \succsim_X is complete and satisfies Ar, it must satisfy Ar⁺ by SI. By Proposition 5, \succsim_V is complete on $\text{Span}\{\succsim_V 0\}$, hence by (a), \succsim_X satisfies MC.

(d) The left-hand side is clearly contained in the right. But for nontrivial \succsim_V , $\{\sim_V 0\} \subset \text{Span}\{\succ_V 0\}$, hence $\text{Span}\{\succsim_V 0\} \subset \text{Span}\{\succ_V 0\}$. \square

Proof of Theorem 1. (1) If \succsim_X is trivial, clearly all three of Ar, MC, and Eq hold, so suppose \succsim_X is nontrivial. Now if \succsim_X satisfies Ar and MC, Eq must hold by Proposition 11(a, b).

Assume Eq. Then \succsim_V is complete on $\text{Span}\{\succsim_V 0\}$ by Proposition 11(b). If Ar holds, then so does MC, by Proposition 11(a)(iv, i). Assume MC. Then $\{\succsim_V 0\}$ is algebraically closed, by Corollary 8 and Theorem 7(c), hence $\{\succ_V 0\}$ is algebraically open in $\text{Span}\{\succsim_V 0\}$, by Lemma 10 and the completeness on $\text{Span}\{\succsim_V 0\}$. But $\text{Span}\{\succsim_V 0\} = \text{Span}\{\succ_V 0\}$, by Proposition 11(d), hence Ar holds, by Theorem 7(a).

(2) By Proposition 11(c), if Ar⁺ and MC hold, then so does C. By Proposition 6(c) and Proposition 11(c), if Ar⁺ and C hold, then so does MC. Finally, MC and C imply Ar by Theorem 1(1), and C and Ar imply Ar⁺ by SI. \square

Proof of Corollary 3. This is immediate from Proposition 11(a). \square

References

- Aumann, R., 1962. ‘Utility theory without the completeness axiom.’ *Econometrica* 30, 455–462.
- Baucells, M. and Shapley, L. 2008. ‘Multiperson utility.’ *Games and Economic Behavior* 62: 329–347.
- Bewley, T., 2002. ‘Knightian decision theory: Part I.’ *Decisions in Economics and Finance* 25(2): 79–110.
- Blackwell, D. and Girshick, M., 1954. *Theory of Games and Statistical Decisions*. New York, John Wiley.
- Broome, J., 1999. ‘Is incommensurability vagueness?’ In *Ethics out of Economics* Cambridge University Press, Cambridge, 123–161.
- Danan, E., Guerdjikova, A. and Zimper, A., 2012. ‘Indecisiveness aversion and preference for commitment.’ *Theory and Decision* 72: 1–13.

- Dubra, J., 2011. ‘Continuity and completeness under risk.’ *Mathematical Social Sciences* 61, 80–81.
- Dubra, J., Maccheroni, F., and Ok, E., 2004. ‘Expected utility theory without the completeness axiom.’ *Journal of Economic Theory* 115: 118–133.
- Evren, Ö., 2008. ‘On the existence of expected multi-utility representations.’ *Economic Theory* 35: 575–592.
- Evren, Ö., 2014. ‘Scalarization methods and expected multi-utility representations.’ *Journal of Economic Theory* 151: 30–63.
- Galaabaatar, T. and Karni, E., 2012. ‘Expected multi-utility representations.’ *Mathematical Social Sciences* 64: 242–246.
- Galaabaatar, T. and Karni, E., 2013. ‘Subjective expected utility with incomplete preferences.’ *Econometrica* 81: 255–284.
- Ghirardato, P., Maccheroni, F., Marinacci, M., and Siniscalchi, M., 2003. ‘A subjective spin on roulette wheels.’ *Econometrica* 71: 1897–1908.
- Gilboa, I., Maccheroni, F., Marinacci, M. and Schmeidler, D., 2010. ‘Objective and subjective rationality in a multiple prior model.’ *Econometrica* 78: 755–770.
- Hausner, M., Wendel, J., 1952 ‘Ordered vector spaces.’ *Proceedings of the American Mathematical Society* 3, 977–982.
- Herstein, I. and Milnor, J., 1953. ‘An axiomatic approach to measurable utility.’ *Econometrica* 21: 291–297.
- Karni, E., 2007. ‘Archimedean and continuity.’ *Mathematical Social Sciences* 53: 332–334.
- Kopylov, I., 2009. ‘Choice deferral and ambiguity aversion.’ *Theoretical Economics* 4: 199–225.
- Manzini, P. and Mariotti, M., 2008. ‘On the representation of incomplete preferences over risky alternatives.’ *Theory and Decision* 65: 303–323.
- McCarthy, D., Mikkola, K., Thomas, T., 2017a. ‘Representation of strongly independent preorders by sets of scalar-valued functions.’ MPRA Paper No. 79284 <https://mpra.ub.uni-muenchen.de/79284/>
- McCarthy, D., Mikkola, K., Thomas, T., 2017b. ‘Representation of strongly independent strict partial orders by scalar-valued utility functions.’ Mimeo.

- McCarthy, D., Mikkola, K., Thomas, T., 2017c. ‘Representation of strongly independent preorders by vector-valued functions.’ Mimeo
- Nau, R., 2006. ‘The shape of incomplete preferences’ *Annals of Statistics* 34; 2430–2448.
- Ok, E., 2007. *Real Analysis with Economic Applications*. Princeton, Princeton University Press.
- Ok, E., Ortoleva, P. and Riella, G., 2012. ‘Incomplete preferences under uncertainty: Indecisiveness in beliefs vs. tastes.’ *Econometrica* 80: 1791–1808.
- Schmeidler, D., 1971. ‘A condition for the completeness of partial preference relations.’ *Econometrica* 39: 403–404.
- Seidenfeld, T., Schervish, M. and Kadane, J., 1995. ‘A representation of partially ordered preferences.’ *Annals of Statistics* 23: 2168–2217.
- Shapley, L. and Baucells, M., 1998. ‘Multiperson utility.’ UCLA Working Paper 779.