

# Continuity for the Maximal Bochner-Riesz operators on the weighted Weak Hardy spaces

Zhu Shihong

College of Mathematics and Computer ,Tongling University ,Tongling 24400,China

**Abstract:** In this papers ,we generalize some results of other authors to weighted spaces and gain the boundedness of maximal Bochner-Riesz operator on weighted Herz-Hardy spaces,weighted Hardy spaces and weighted weak Hardy spaces ,where  $\omega \in A_1$ .

**Key words:** Weighted Herz-Hardy spaces, Maximal Bochner-Riesz operator,Weighted weak Hardy space

**CLC number:**O175.14     **Document code:** A

## 1 Introduction

Jiang Yinsheng and other authors <sup>[1,2]</sup> discuss the boundednes of maximal Bochner-Riesz and it's commutator .Recently ,Tongseng Quek <sup>[3]</sup>proved some C-Z type operators are bounded on weighted Hardy spaces and weighted weak Hary spaces . Inspiring by the above and the paper<sup>[4]</sup> ,we will prove the boundedness of the maximal Bochner-Riesz operator on these spaces.

Let maximal Bochner-Riesz operator  $B_\delta^*$  be

$$T_\delta^* f(x) = \sup_{t>0} |B_\delta^t(f)(x)|,$$

where  $B_\delta^t(x) = t^{-n} B_\delta(x/t)$  is the kernel of  $T_\delta^t$ , and satisfies  $|\frac{\partial^\beta}{\partial x^\beta} B_\delta(x)| \leq C(1 + |x|)^{(\delta+(n+1)/2)}$  for any  $x \in \mathbb{R}^n$  and multi-index  $\beta \in \mathbb{Z}_+^n$ .

Suppose  $0 < p < \infty$ , we denote the weak  $L_\omega^p(\mathbb{R}^n)$  by  $WL_\omega^p(\mathbb{R}^n)$  and set

$$\|f\|_{WL_\omega^p(\mathbb{R}^n)} = \sup_{\lambda>0} \lambda[\omega(\{x \in \mathbb{R}^n : |f(x)| > \lambda\})]^{1/p},$$

where, and in what follows,  $\omega(E) = \int_E \omega(x)dx$ .

Receive date: .

Foundation item: The Natural Science Foundation of Anhui Province(kj2010b460), 《Continuity of the multilinear operators》 , Peroject director: Zhu Shihong

Biography: Zhu Shihong(1972-),male,Lecturer, engaged in harmonic analysis. email: [zhu\\_shihong@126.com](mailto:zhu_shihong@126.com)

## 2 $[\dot{K}_q^{\alpha,p}(\omega_1, \omega_2), H\dot{K}_q^{\alpha,p}(\omega_1, \omega_2)]$ -Type Continuity.

Let us first recall the definition of Herz spaces .For  $k \in \mathbb{Z}$ , let  $B_k = \{x \in \mathbb{R}^n : |x| \leq 2^k\}$ ,  $C_k = B_k \setminus B_{k-1}$  and  $\chi_k = \chi_{C_k}$ .

**Definition 1**<sup>[4]</sup> Let  $0 \leq \alpha < \infty$ ,  $1 < q < \infty$ ,  $0 < p < \infty$ , and  $\omega_1, \omega_2$  be non-negative weight function. The homogeneous weighted Herz spaces  $\dot{K}_q^{\alpha,p}(\omega_1, \omega_2)$  are defined by

$$\dot{K}_q^{\alpha,p}(\omega_1, \omega_2) = \{f \in L_{loc}^q(\mathbb{R}^n \setminus \{0\}, \omega_2) : \|f\|_{\dot{K}_q^{\alpha,p}(\omega_1, \omega_2)} < \infty\},$$

where

$$\|f\|_{\dot{K}_q^{\alpha,p}(\omega_1, \omega_2)} = \left\{ \sum_{k=-\infty}^{\infty} \omega_1(B_k)^{\frac{\alpha p}{n}} \|f \chi_k\|_{L_{\omega_2}^q}^p \right\}^{1/p}.$$

**Definition 2**<sup>[4]</sup> Let  $\alpha \in \mathbb{R}^n$ ,  $1 < q < \infty$ ,  $0 < p < \infty$  and  $\omega_1, \omega_2$  be non-negative weight function. The homogeneous weighted Herz-Hardy spaces are defined by  $H\dot{K}_q^{\alpha,p}(\omega_1, \omega_2) = \{f \in S' : G(f) \in \dot{K}_q^{\alpha,p}(\omega_1, \omega_2)\}$  and

$$\|f\|_{H\dot{K}_q^{\alpha,p}(\omega_1, \omega_2)} = \|G(f)\|_{\dot{K}_q^{\alpha,p}(\omega_1, \omega_2)},$$

where  $G(f)$  is usually called the grand maximal function of  $f$ .

**Definition 3**<sup>[4]</sup> Let  $\omega_1, \omega_2 \in A_1$ ,  $n(1 - \frac{1}{q}) \leq \alpha < \infty$  and  $s \geq [\alpha + n(\frac{1}{q} - 1)]$  be non-negative integral. The function  $a(x)$  is called a center atom of  $(\alpha, q, \omega_1, \omega_2)$  - type, if  $a(x)$  satisfies

- (1)  $\text{supp } a_k \subset B_k = B(0, 2^k r)$ ,
- (2)  $\|a_k\|_{L_{\omega_2}^q} \leq \omega_1(B_k)^{\frac{-\alpha}{n}}$ , (3)  $\int a(x) x^\beta dx = 0$ ,  $|\beta| \leq s$ .

**Theorem 1** Let  $\delta > \frac{n-1}{2}$ ,  $0 < p < \infty$ ,  $1 < q < \infty$ , and  $n(1 - \frac{1}{q}) \leq \alpha < n(1 - \frac{1}{q}) + \varepsilon$ ,  $\varepsilon = \min\{1, (\delta - \frac{n-1}{2})\}$ ,  $\omega_1, \omega_2 \in A_1$ , Then  $T_\delta^*$  is a bounded operator from  $H\dot{K}_q^{\alpha,p}(\omega_1, \omega_2)$  to  $\dot{K}_q^{\alpha,p}(\omega_1, \omega_2)$ .

**Proof of Theorem 1** Since  $\omega_1, \omega_2 \in A_1(\mathbb{R}^n)$ , a temperate distribution of  $f$  can be written as  $f = \sum_{-\infty}^{\infty} \lambda_j a_j$  and  $(\sum_{-\infty}^{\infty} |\lambda_j|^p)^{\frac{1}{p}} \leq C \|f\|_{H\dot{K}_q^{\alpha,p}(\omega_1, \omega_2)}$ , where  $a_j$  is the central atom of  $(\alpha, q, \omega_1, \omega_2)$  - type. Let  $n(1 - \frac{1}{q}) \leq \alpha < n(1 - \frac{1}{q}) + \varepsilon$  and  $f \in H\dot{K}_q^{\alpha,p}(\omega_1, \omega_2)$ . Then we have

$$\begin{aligned} \|T_\delta^t(f)\|_{HK_q^{\alpha,p}(\omega_1, \omega_2)} &= \left\{ \sum_{k=-\infty}^{\infty} \omega_1(B_k)^{\frac{\alpha p}{n}} \|T_\delta^t(f) \chi_k\|_{L_{\omega_2}^q}^p \right\}^{\frac{1}{p}} \\ &\leq C \left\{ \sum_{k=-\infty}^{\infty} \omega_1(B_k)^{\frac{\alpha p}{n}} \left( \sum_{j=-\infty}^{k-3} |\lambda_j| \|T_\delta^t(a_j) \chi_k\|_{L_{\omega_2}^q} \right)^p \right\}^{\frac{1}{p}} \\ &+ C \left\{ \sum_{k=-\infty}^{\infty} \omega_1(B_k)^{\frac{\alpha p}{n}} \left( \sum_{j=k-2}^{\infty} |\lambda_j| \|T_\delta^t(a_j) \chi_k\|_{L_{\omega_2}^q} \right)^p \right\}^{\frac{1}{p}} = D_1 + D_2. \end{aligned}$$

For  $D_2$ , when  $0 < p \leq 1$ . We notice  $\frac{\omega(E)}{\omega(B)} \leq C(\frac{|E|}{|B|})^\xi$  as  $E \subset B$  and  $0 < \xi < 1$ . By the boundedness of operator  $T_\delta^*$  on  $L_{\omega_2}^q(\mathbb{R}^n)^{[5]}$ , we have

$$\begin{aligned} D_2 &\leq \left\{ \sum_{k=-\infty}^{\infty} \omega_1(B_k)^{\frac{\alpha p}{n}} \left( \sum_{j=k-2}^{\infty} |\lambda_j|^p \|T_\delta^t(a_j)\chi_k\|_{L_{\omega_2}^q}^p \right) \right\}^{\frac{1}{p}} \\ &\leq C \left\{ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \left[ \sum_{k=-\infty}^{j-1} \left( \frac{\omega_1(B_k)}{\omega_1(B_j)} \right)^{\frac{\alpha p}{n}} + 1 + \left( \frac{\omega_1(B_{j+1})}{\omega_1(B_j)} \right)^{\frac{\alpha p}{n}} + \left( \frac{\omega_1(B_{j+2})}{\omega_1(B_j)} \right)^{\frac{\alpha p}{n}} \right] \right\}^{\frac{1}{p}} \\ &\leq C \left\{ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \left[ \sum_{k=-\infty}^{j-1} \left( \frac{|B_k|}{|B_j|} \right)^{\frac{\alpha p \xi}{n}} + 1 + 2^{\alpha p} + 2^{2\alpha p} \right] \right\}^{\frac{1}{p}} \leq C \left\{ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \right\}^{\frac{1}{p}}. \end{aligned}$$

When  $p > 1$ , by Hölder's inequality, we have

$$\begin{aligned} D_2 &\leq \left\{ \sum_{k=-\infty}^{\infty} \omega_1(B_k)^{\frac{\alpha p}{n}} \left[ \sum_{j=k-2}^{\infty} |\lambda_j| \omega_1(B_j)^{-\frac{\alpha}{n}} \right]^p \right\}^{1/p} \\ &\leq \left\{ \sum_{k=-\infty}^{\infty} \left( \sum_{j=k-2}^{\infty} |\lambda_j|^p \omega_1(B_j)^{-\frac{\alpha p}{2n}} \omega_1(B_k)^{\frac{\alpha p}{2n}} \right) \left( \sum_{j=k-2}^{\infty} \omega_1(B_j)^{-\frac{\alpha p'}{2n}} \omega_1(B_k)^{\frac{\alpha p'}{2n}} \right)^{\frac{p}{p'}} \right\}^{\frac{1}{p}} \\ &\leq \left\{ \sum_{k=-\infty}^{\infty} \left[ \sum_{j=k-2}^{\infty} |\lambda_j|^p \left( \frac{\omega_1(B_k)}{\omega_1(B_j)} \right)^{\frac{\alpha p}{2n}} \left[ \sum_{j=k+1}^{\infty} 2^{\frac{(k-j)\alpha p' \xi}{2}} + 1 + 2^{\frac{\alpha p'}{2}} + 2^{\alpha p'} \right]^{\frac{p}{p'}} \right] \right\}^{\frac{1}{p}} \\ &\leq \left\{ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \left( \sum_{k=-\infty}^{j+2} \left( \frac{\omega_1(B_k)}{\omega_1(B_j)} \right)^{\frac{\alpha p}{2n}} \right) \right\}^{\frac{1}{p}} \leq \left\{ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \sum_{k=-\infty}^{j+2} 2^{\frac{(k-j)\alpha p \xi}{2}} \right\}^{\frac{1}{p}} \leq C \left\{ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \right\}^{\frac{1}{p}} \end{aligned}$$

Now, let us turn to the estimate for  $D_1$ . For any  $j$  respect to a fixed  $k$  satisfied  $j \leq k-3$ , let  $B_j = B(0, 2^j r) = B(0, r_1)$ ,  $B_k = B(0, 2^{k-j} r_1)$ . As  $x \in B_k, y \in B_j$ , we know  $|x-y| \sim |x-0|$ . We will consider two cases for  $D_1$ .

**case 1**  $t < r_1$ . By Hölder's inequality and the condition of core  $B_\delta^t$ , we have

$$\begin{aligned} &\left\{ \int_{B_k} \left| \int_{B_j} B_\delta^t(x-y) a_j(y) dy \right|^q \omega_2(x) dx \right\}^{\frac{1}{q}} \\ &\leq C \left\{ \int_{B_k} |t^{-n} \int_{B_j} \left( 1 + \frac{|x-y|}{t} \right)^{-(\delta + \frac{n+1}{2})} a_j(y) dy|^q \omega_2(x) dx \right\}^{\frac{1}{q}} \\ &\leq C t^{\delta - \frac{n-1}{2}} (2^{(k-j)} r_1)^{-(\delta + \frac{n+1}{2})} |B_j| \left( \frac{\omega_2(B_k)}{\omega_2(B_j)} \right)^{\frac{1}{q}} \left( \int_{B_j} |a_j(y)|^q \omega_2(y) dy \right)^{\frac{1}{q}} \\ &\leq C 2^{(j-k)(\delta + \frac{n+1}{2} - \frac{n}{q})} \|a_j\|_{L_{\omega_2}^q} \leq C 2^{(j-k)(\delta + \frac{n+1}{2} - \frac{n}{q})} \omega_1(B_j)^{-\frac{\alpha}{n}}. \end{aligned}$$

When  $0 < p \leq 1$ , we gain

$$\begin{aligned} D_1 &\leq C \left\{ \sum_{k=-\infty}^{\infty} \left( \sum_{j=-\infty}^{k-3} |\lambda_j|^p 2^{(j-k)p(\delta + \frac{n+1}{2} - \frac{n}{q})} \left( \frac{\omega_1(B_k)}{\omega_1(B_j)} \right)^{\frac{\alpha p}{n}} \right)^{\frac{1}{p}} \right\} \\ &\leq C \left\{ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \left( \sum_{k=j+3}^{\infty} 2^{(j-k)p(\delta + \frac{n+1}{2} - \frac{n}{q} - \alpha)} \right)^{\frac{1}{p}} \right\} \leq C \left\{ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \right\}^{\frac{1}{p}}. \end{aligned}$$

When  $p > 1$ . By Hölder's inequality, we have

$$\begin{aligned} D_1 &\leq C \left\{ \sum_{j=-\infty}^{\infty} \left( \sum_{k=j+3}^{\infty} |\lambda_j|^p 2^{(j-k)\frac{p}{2}(\delta + \frac{n+1}{2} - \frac{n}{q} - \alpha)} \right) \left( \sum_{k=j+3}^{\infty} 2^{(j-k)\frac{p'}{2}(\delta + \frac{n+1}{2} - \frac{n}{q} - \alpha)} \right)^{\frac{p}{p'}} \right\}^{\frac{1}{p}} \\ &\leq C \left\{ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \left( \sum_{k=j+3}^{\infty} 2^{(j-k)\frac{p}{2}(\delta + \frac{n+1}{2} - \frac{n}{q} - \alpha)} \right)^{\frac{1}{p}} \right\} \leq C \left\{ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \right\}^{\frac{1}{p}}. \end{aligned}$$

**case 2**  $t \geq r_1$ . By the mean value theorem and the vanishing moment condition of atom  $a_j$ , we have

$$\begin{aligned} &\left( \int_{B_k} \left| \int_{B_j} B_{\delta}^t(x-y) a_j(y) dy \right|^q \omega_2(x) dx \right)^{\frac{1}{q}} \\ &= \left( \int_{B_k} \left| \int_{B_j} [B_{\delta}^t(x-y) - B_{\delta}^t(x-0)] a_j(y) dy \right|^q \omega_2(x) dx \right)^{\frac{1}{q}} \\ &\leq C \left\{ \int_{B_k} \left| t^{-(n+1)} \int_{B_j} \left( 1 + \frac{|x-0|}{t} \right)^{-(\delta + \frac{n+1}{2})} y a_j(y) dy \right|^q \omega_2(x) dx \right\}^{\frac{1}{q}} \end{aligned}$$

(1) When  $\delta < \frac{n+1}{2}$ .

$$\begin{aligned} &\left( \int_{B_k} \left| \int_{B_j} B_{\delta}^t(x-y) a_j(y) dy \right|^q \omega_2(x) dx \right)^{\frac{1}{q}} \\ &\leq C \left( \frac{r_1}{t} \right)^{\frac{n+1}{2} - \delta} 2^{(j-k)(\delta + \frac{n+1}{2})} \left[ \frac{\omega_2(B_k)}{\omega_2(B_j)} \right]^{\frac{1}{q}} \|a_j\|_{L_{\omega_2}^q} \\ &\leq C 2^{(j-k)(\delta + \frac{n+1}{2} - \frac{n}{q})} \omega_1(B_j)^{-\frac{\alpha}{n}}. \end{aligned}$$

When  $0 < p \leq 1$ , we obtain

$$D_1 \leq C \left\{ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \left( \sum_{k=j+3}^{\infty} 2^{(j-k)p(\delta + \frac{n+1}{2} - \frac{n}{q} - \alpha)} \right)^{\frac{1}{p}} \right\} \leq C \left\{ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \right\}^{\frac{1}{p}}.$$

When  $p > 1$ . By Hölder's inequality, we also have

$$D_1 \leq C \left\{ \sum_{k=-\infty}^{\infty} \left( \sum_{j=-\infty}^{k-3} |\lambda_j|^p 2^{(j-k)\frac{p}{2}(\delta + \frac{n+1}{2} - \frac{n}{q})} \left( \frac{\omega_1(B_k)}{\omega_1(B_j)} \right)^{\frac{\alpha p}{2n}} \right)^{\frac{1}{p}} \right\} \leq C \left\{ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \right\}^{\frac{1}{p}}.$$

(2) When  $\delta \geq \frac{n+1}{2}$ . By the mean value theorem and the vanishing moment condition of atom  $a_j$ , we have

$$\left( \int_{B_k} \left| \int_{B_j} B_\delta^t(x-y) a_j(y) dy \right|^q \omega_2(x) dx \right)^{\frac{1}{q}} \leq C 2^{(j-k)(n+1-\frac{n}{q})} \omega_1(B_j)^{-\frac{\alpha}{n}}$$

As the above, we consider  $D_1$  under the condition of  $0 < p \leq 1$  at first. We obtain

$$D_1 \leq C \left\{ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \left( \sum_{k=j+3}^{\infty} 2^{(j-k)p(n+1-\frac{n}{q}-\alpha)} \right)^{\frac{1}{p}} \right\} \leq C \left\{ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \right\}^{\frac{1}{p}}.$$

Also, we secondly consider  $D_1$  under the condition of  $p > 1$ . By Hölder's inequality, we have

$$D_1 \leq C \left\{ \sum_{k=-\infty}^{\infty} \left( \sum_{j=-\infty}^{k-3} |\lambda_j|^p 2^{(j-k)\frac{p}{2}(n+1-\frac{n}{q}-\alpha)} \right)^{\frac{1}{p}} \right\} \leq C \left\{ \sum_{j=-\infty}^{\infty} |\lambda_j|^p \right\}^{\frac{1}{p}}.$$

Thus, we have

$$\|T_\delta^t(f)\|_{\dot{K}_q^{\alpha,p}(\omega_1,\omega_2)} \leq C \|T_\delta^t(f)\|_{H\dot{K}_q^{\alpha,p}(\omega_1,\omega_2)}.$$

We take the supreme of the left side respect to the inequality above for any  $t > 0$ , and obtain desirable result.

### 3 $[L_\omega^p(\mathbb{R}^n), H_\omega^p(\mathbb{R}^n)]$ -Type Continuity

**Definition 4** Let  $\omega_\infty(\mathbb{R}^n)$  and  $p \in (0, 1]$ . The weighted Hardy spaces  $H_\omega^p(\mathbb{R}^n)$  is defined by

$$H_\omega^p(\mathbb{R}^n) = \{f \in S'(\mathbb{R}^n) : \phi_t * (f)(x) = \sup_{t>0} |\phi_t * f(x)| \in L_\omega^p(\mathbb{R}^n)\},$$

where  $\phi \in S(\mathbb{R}^n)$  is a fixed function with  $\int_{\mathbb{R}^n} \phi(x) dx \neq 0$  and  $\phi_t = t^{-n} \phi(y/t)$  for any  $t > 0$ . Moreover, we define  $\|f\|_{H_\omega^p(\mathbb{R}^n)} = \|\phi * (f)\|_{L_\omega^p(\mathbb{R}^n)}$ .

**Definition 5** Let  $\omega \in A_\infty$ ,  $p \in (0, 1]$ . A  $p$ -atom with respect to  $\omega$  is a function supported in a ball  $B$  such that

$$\|a\|_{L^\infty(\mathbb{R}^n)} \leq \omega(B)^{-\frac{1}{p}}$$

$\int_{\mathbb{R}^n} a(x) x^\alpha dx = 0$  for every multi-index  $\alpha$  with  $|\alpha| \leq [n(q_\omega/p - 1)]$ , where, and in what follows,  $[s]$  denotes the greatest integer less than or equal to  $s$ .

**Theorem 2** Let  $\omega \in A_1$ ,  $\delta > \frac{n-1}{2}$ ,  $\min\{\frac{n}{\delta+(n+1)/2}, \frac{n}{n+1}\} < p \leq 1$ , Then  $T_\delta^*$  is a bounded map from  $H_\omega^p(\mathbb{R}^n)$  into  $L_\omega^p(\mathbb{R}^n)$ .

**Proof of Theorem 2** We only need to show that for any  $p$ -atom  $a$  with respect to  $\omega$ ,  $\|T_\delta^*(a)\|_{L_\omega^p(\mathbb{R}^n)} \leq C$  with  $C$  independent of  $a$ . Suppose  $\text{supp } a \subset B(x_0, r)$ . Let  $\omega \in A_{q_0}(\mathbb{R}^n)$  with  $q_0 > 1$ . We choose  $p_0 > 1$ , and write

$$\|T_\delta^t(a)\|_{L_\omega^p(\mathbb{R}^n)}^p \leq \int_{B(x_0, 4r)} |T_\delta^t(a)(x)|^p \omega(x) dx + \int_{\mathbb{R}^n \setminus B(x_0, 4r)} |T_\delta^t(a)(x)|^p \omega(x) dx = L_1 + L_2.$$

By the boundedness of operator  $T_\delta^*$  on  $L_\omega^q$ <sup>[5]</sup>, we then have

$$\begin{aligned} L_1 &\leq C \left( \int_{B(x_0, 4r)} |T_\delta^t(a)(x)|^{p_0} \omega(x) dx \right)^{\frac{p}{p_0}} \left( \int_{B(x_0, 4r)} \omega(x) dx \right)^{1 - \frac{p}{p_0}} \\ &\leq C \left( \int_{B(x_0, r)} |a(x)|^{p_0} \omega(x) dx \right)^{\frac{p}{p_0}} \omega(B(x_0, r))^{1 - \frac{p}{p_0}} \leq C, \end{aligned}$$

where  $C$  is independent of  $a$ .

Noticing  $y \in B(x_0, r)$  and  $x \in \mathbb{R}^n \setminus B(x_0, 4r)$ , we gain  $|x - x_0| \sim |x - y|$ .

**case 1**  $t < r$  By the vanishing moments of  $a$ -atom, we gain

$$\begin{aligned} T_\delta^t(a)(x) &\leq Ct^{-n} \int_{B(x_0, r)} \left(1 + \frac{|x - y|}{t}\right)^{-(\delta + \frac{n+1}{2})} |a(y)| dy \\ &\leq Ct^{\delta - \frac{n-1}{2}} |x - x_0|^{-(\delta + \frac{n+1}{2})} \|a\|_\infty |B| \leq Ct^{\delta - \frac{n-1}{2}} |x - x_0|^{-(\delta + \frac{n+1}{2})} \omega(B)^{-\frac{1}{p}} |B| \\ L_2 &\leq Ct^{p(\delta - \frac{n-1}{2})} \omega(B)^{-1} |B|^p \int_{\mathbb{R}^n \setminus B(x_0, 4r)} |x - x_0|^{-p(\delta + \frac{n+1}{2})} \omega(x) dx \\ &\leq Ct^{p(\delta - \frac{n-1}{2})} \omega(B)^{-1} |B|^p \sum_{k=2}^{\infty} \int_{B_{k+1} \setminus B_k} |x - x_0|^{-p(\delta + \frac{n+1}{2})} \omega(x) dx \\ &\leq C \left(\frac{t}{r}\right)^{p(\delta - \frac{n-1}{2})} \sum_{k=2}^{\infty} \left(\frac{\omega(B_{k+1})}{\omega(B)}\right) 2^{-kp(\delta + \frac{n+1}{2})} \leq C \sum_{k=2}^{\infty} 2^{-k[p(\delta + \frac{n+1}{2}) - n]} \leq C \end{aligned}$$

**case 2**  $t \geq r$ . By the vanishing moments of  $a$ -atom and the mean value theorem, we have

$$T_\delta^t(a)(x) \leq Ct^{-(n+1)} \int_{B(x_0, r)} \left(1 + \frac{|x - x_0|}{t}\right)^{-(\delta + \frac{n+1}{2})} |y - x_0| |a(y)| dy$$

When  $\delta < \frac{n+1}{2}$ , we obtain

$$L_2 \leq C \left(\frac{r}{t}\right)^{p(\frac{n+1}{2} - \delta)} \sum_{k=2}^{\infty} 2^{-kp(\delta + \frac{n+1}{2})} \frac{\omega(B_{k+1})}{\omega(B)} \leq C \sum_{k=2}^{\infty} 2^{-k[p(\delta + \frac{n+1}{2}) - n]} \leq C.$$

When  $\delta \geq \frac{n+1}{2}$ , we obtain

$$\begin{aligned} T_\delta^t(a)(x) &\leq |x - x_0|^{-(n+1)} r^{n+1} \|a\|_\infty \\ L_2 &\leq r^{p(n+1)} \omega(B)^{-1} \sum_{k=2}^{\infty} \int_{B_{k+1} \setminus B_k} |x - x_0|^{-p(n+1)} \omega(x) dx \leq C \sum_{k=2}^{\infty} 2^{-k[p(n+1) - n]} \leq C. \end{aligned}$$

Thus, we have

$$\|T_\delta^t(a)\|_{L_\omega^p}^p \leq C$$

For any  $t > 0$ , we take the supreme of the left side and can gain the desire result.

#### 4 $[WL_\omega^p(\mathbb{R}^n), WH_\omega^p(\mathbb{R}^n)]$ -Type Continuity .

**Lemma 1** Let  $p \in (0, 1]$  and  $\omega \in A_\infty(\mathbb{R}^n)$ . For  $f \in WH_\infty^p(\mathbb{R}^n)$ , there exists a sequence  $\{f_k\}_{k=-\infty}^\infty$  of bounded measurable functions such that

$$f = \sum_{k=-\infty}^{\infty} f_k \quad \text{in } S'(\mathbb{R}^n). \quad (1)$$

Each  $f_k$  can be further decomposed into  $f_k = \sum_i b_{ki}$ , where the sequence  $\{b_{ki}\}_i$  satisfies

$$\text{supp } b_{ki} \subset Q_{ki} \text{ and } Q_{ki} \text{ is a cube}, \quad (2)$$

$$\sum_i \omega(Q_{ki}) \leq c_1 2^{-kp}; \quad \sum_i \chi_{Q_{ki}}(x) \leq c_1,$$

$\chi_E$  being the characteristic function of the set  $E$ ,  $c_1$  a constant and  $c_1 \leq C \|f\|_{WH_\omega^p(\mathbb{R}^n)}^p$ ;

$$\|b_{ki}\|_{L^\infty(\mathbb{R}^n)} \leq C 2^k \quad \text{and} \quad \int_{\mathbb{R}^n} b_{ki}(x) x^\alpha dx = 0, \text{ for } |\alpha| \leq [n(q_\omega/p - 1)].$$

Conversely, if  $f \in S'(\mathbb{R}^n)$  has a decomposition satisfying (1) and (2), then  $f \in WH_\omega^p(\mathbb{R}^n)$  and  $\|f\|_{WH_\omega^p(\mathbb{R}^n)}^p \leq C c_1$ , where  $C$  is a constant.

**Theorem 3** Let  $\delta > \frac{n-1}{2}$ ,  $\max\{\frac{n}{\delta+(n+1)/2}, \frac{n}{n+1}\} < p \leq 1$ ,  $\omega \in A_q$ ,  $q \geq 1$ , Then  $T_\delta^*$  is a bounded operator from  $WH_\omega^p(\mathbb{R}^n)$  to  $WL_\omega^p(\mathbb{R}^n)$ .

**Proof of Theorem 3** For any  $\lambda > 0$ , let  $k_0 \in \mathbb{Z}$  such that  $2^{k_0} \leq \lambda < 2^{k_0+1}$ . Let  $\omega \in A_q(\mathbb{R}^n)$  and  $f \in WH_\omega^p(\mathbb{R}^n)$ . Then by Lemma 1, we write

$$f = \sum_{k=-\infty}^{\infty} f_k = \sum_{k=-\infty}^{\infty} \sum_i b_{ki} = \sum_{k=-\infty}^{k_0} \sum_i b_{ki} + \sum_{k=k_0+1}^{\infty} \sum_i b_{ki} = F_1 + F_2,$$

where  $b_{k,i}$ s are as in Lemma 1. Suppose  $A_k = \text{supp } f_k$ , then  $A_k = \cup_i Q_{ki}$  and  $\omega(A_k) \leq \sum_i \omega(Q_{ki}) \leq C 2^{-kp} \|f\|_{WH_\omega^p}^p$ . Since  $B_\delta^*$  is bounded on  $L_\omega^q$  spaces, we have

$$\begin{aligned} \omega(\{x \in \mathbb{R}^n : |T_\delta^t(F_1)(x)| > \lambda\}) &\leq C \|T_\delta^t(F_1)\|_{L_\omega^2}^2 / \lambda^2 \\ &\leq C \|F_1\|_{L_\omega^2}^2 / \lambda^2 \leq C \left[ \sum_{k=-\infty}^{k_0} \|f_k\|_{L_\omega^2} \right]^2 / \lambda^2 \leq C \left[ \sum_{k=-\infty}^{k_0} \left( \sum_i \int_{B_{ki}} |b_{ki}|^2 \omega(x) dx \right)^{\frac{1}{2}} \right]^2 / \lambda^2 \\ &\leq C \left\{ \sum_{k=-\infty}^{k_0} 2^k \left[ \sum_i \omega(B_{ki}) \right]^{\frac{1}{2}} \right\}^2 / \lambda^2 \leq C \left[ \sum_{k=-\infty}^{k_0} 2^{k(1-p/2)} \right]^2 \|f\|_{WH_\omega^p}^p / \lambda^2 \leq C \|f\|_{WH_\omega^p}^p / \lambda^p. \end{aligned}$$

Now set  $Q_{ki}^* = Q(x_{ki}, (5\sqrt{n})r_{ki})$  and  $A_{k_0} = \cup_{k=k_0+1}^\infty \cup_i Q_{ki}^*$ , where  $Q_{ki}^*$  is the cube with the same  $x_{ki}$ -center as  $Q_{ki}$  and side length  $5\sqrt{n}$  times the side length of  $Q_{ki}$ . We get

$$\omega(Q_{k_0}) \leq \sum_{k=k_0+1}^{\infty} \sum_i \omega(Q_{ki}^*) \leq \sum_{k=k_0+1}^{\infty} \sum_i \omega(Q_{ki}) \leq \sum_{k=k_0+1}^{\infty} 2^{-kp} \|f\|_{WH_\omega^p}^p \leq \|f\|_{WH_\omega^p}^p / \lambda^p.$$

To finish the proof, we still need to estimate

$$\omega(\{x \in A_{k_0}^c : |T_\delta^t(F_2)(x)| > \lambda\}) \leq C \|T_\delta^t(F_1)\|_{L_\omega^p}^p / \lambda^p.$$

For  $x \in (A_{k_0}^*)^c, y \in Q_{ki}$ , we know  $|x - y| \sim |x - x_{ki}|$ .

**case 1**  $t < r_{ki}$ , We then obtain

$$\begin{aligned} \int_{A_{k_0}^c} |T_\delta^t(b_{ki})(x)|^p \omega(x) dx &= \int_{A_{k_0}^c} \left\{ \int_{Q_{ki}} t^{-n} \left(1 + \frac{|x-y|}{t}\right)^{-(\delta + \frac{n+1}{2})} |b_{ki}(y)| dy \right\}^p \omega(x) dx \\ &\leq C \sum_{j=2}^{\infty} \int_{Q_{ki}^{j+1} \setminus Q_{ki}^j} t^{p(\delta - \frac{n-1}{2})} |x - x_{ki}|^{-p(\delta + \frac{n+1}{2})} 2^{kp} |Q_{ki}|^p \omega(x) dx \\ &\leq C \sum_{j=2}^{\infty} 2^{-jp(\delta + \frac{n+1}{2})} \left(\frac{t}{r_{ki}}\right)^{p(\delta - \frac{n-1}{2})} 2^{kp} \omega(Q_{ki}^{j+1}) \leq C \sum_{j=2}^{\infty} 2^{-j[p(\delta + \frac{n+1}{2}) - n]} 2^{kp} \omega(Q_{ki}) \leq C 2^{kp} \omega(Q_{ki}), \end{aligned}$$

where  $Q_{ki}^j$  is the cube with the same  $x_{ki}$ -center as  $Q_{ki}$  and side length  $2^j$  times the side length- $r_{ki}$  of  $Q_{ki}$ .

**case 2**  $t \geq r_{ki}$ .

we consider  $\delta < \frac{n+1}{2}$  at first. By the vanishing moments of  $b_{ki}$  and the mean value theorem, we have

$$\begin{aligned} \int_{A_{k_0}^c} |T_\delta^t(b_{ki})(x)|^p \omega(x) dx &= \int_{A_{k_0}^c} \left| \int_{Q_{ki}} [B_\delta^t(x-y) - B_\delta^t(x-x_{ki})] b_{ki}(y) dy \right|^p \omega(x) dx \\ &\leq C \sum_{j=2}^{\infty} \int_{Q_{ki}^{j+1} \setminus Q_{ki}^j} t^{p(\delta - \frac{n+1}{2})} (2^j r_{ki})^{-p(\delta + \frac{n+1}{2})} r_{ki}^p 2^{kp} |Q_{ki}|^p \omega(x) dx \\ &\leq C \sum_{j=2}^{\infty} \left(\frac{r_{ki}}{t}\right)^{p(\frac{n+1}{2} - \delta)} 2^{-j[p(\delta + \frac{n+1}{2}) - n]} 2^{kp} \omega(Q_{ki}) \leq C 2^{kp} \omega(Q_{ki}) \end{aligned}$$

Secondly, we consider  $\delta \geq \frac{n+1}{2}$ ,

$$\begin{aligned} \int_{A_{k_0}^c} |T_\delta^t(b_{ki})(x)|^p \omega(x) dx &= \int_{Q_{k_0}^c} \left| \int_{Q_{ki}} [B_\delta^t(x-y) - B_\delta^t(x-x_{ki})] b_{ki}(y) dy \right|^p \omega(x) dx \\ &\leq C \sum_{j=2}^{\infty} \int_{Q_{ki}^{j+1} \setminus Q_{ki}^j} \left( \int_{B_{ki}} |x - x_{ki}|^{-(n+1)} r_{ki} |b_{ki}(y)| dy \right)^p \omega(x) dx \\ &\leq C \sum_{j=2}^{\infty} 2^{-j[p(n+1) - n]} 2^{kp} \omega(Q_{ki}) \leq C 2^{kp} \omega(Q_{ki}) \end{aligned}$$

These yields the desired results .

**Acknowledgements:**

**References**



- [1] Jiang Yinsheng ,Tang Lin , Continuity for maximal commutators of Bochner-Riesz operators with BMO functions[J],Acta Mathematica. Scientia.2002,22B(3):339-349.
- [2] Zhu Shihong, Continuity of of maximal Bochner-Riesz operators and its commutators[D],Anhui normal University,2006.
- [3] Tongseng Quek, Yang Dachun, Calderón-Zygmund-Type operators on weighted weak Hardy spaces over  $\mathbb{R}^n$ [J] , Acta Mathematica Sinica ,English Series,2000,16(1):141-160.
- [4] Lu Shanzhen , Yang Dachun, Weighted Herz-Hardy spaces and its Applications[J], Science in China (Series A),1995,25(3):235-245.
- [5] Wang Hua ,Some estimates of Bocher-Riesz operators on weighted Morrey spaces[J],2012,55(3):551-560.