

# **The effect of expertise on diagnosis accuracy, non-technical skills and thought processes during simulated high-fidelity anaesthetist scenarios**

Allistair P McRobert

Research Institute for Sport and Exercise Sciences, Liverpool John Moores University,  
Liverpool, Merseyside, UK

Simon J Mercer, David Raw & Jeff Goulding

Centre for Simulation and Patient Safety, Aintree University Hospitals NHS Foundation  
Trust, Merseyside, UK

A. Mark Williams

Department of Health, Kinesiology, and Recreation, The University of Utah, Salt Lake  
City, Utah, USA

Please address correspondence to:

Allistair McRobert

Liverpool John Moores University

Tom Reilly Building

Byrom Street

Liverpool, L3 3AF

United Kingdom

Tel - +44 151 231 4320

Email: [a.p.mcrobot@ljmu.ac.uk](mailto:a.p.mcrobot@ljmu.ac.uk)

Keywords: Decision-making, Anaesthesia, Verbal reports, Medical simulation.

Word count (excluding title page, abstract, references, figures and tables): 3472

**The effect of expertise on diagnosis accuracy, non-technical skills and thought processes during simulated high-fidelity anaesthetist scenarios**

**RUNNING HEAD: Decision-making in anaesthetists**

1        **ABSTRACT**

2        We examined the thought processes and performance of anaesthetists during  
3 simulated environments. Verbal reports of thinking and the Anaesthetists Non-  
4 Technical Skills (ANTS) were recorded to examine cognitive processes, non-technical  
5 behaviours and diagnosis accuracy during fully immersive, high-fidelity medical  
6 scenarios. Skilled (n = 6) and less skilled (n = 9) anaesthetists were instructed to  
7 respond to medical scenarios experienced in theatre. Skilled participants demonstrated  
8 higher diagnosis accuracy and ANTS scores compared to less skilled participants.  
9 Furthermore, skilled participants engaged in deeper thinking and verbalised more  
10 evaluation, prediction and deep planning statements. The ability to employ an  
11 effective cognitive processing strategy, more efficient non-technical behaviours and  
12 superior diagnosis is associated with superior performance in skilled participants.

13

14

15

16

17

18

19

20

21

1       **WHAT THIS PAPER ADDS**

- 2       • High-fidelity medical simulation is an ideal tool to examine differences in  
3       knowledge and performance of anaesthetists because scenarios can be  
4       standardised and replicated.
- 5       • Diagnosis accuracy, ANTS and concurrent verbal reports can be used as  
6       performance measures during simulation to compare anaesthetists of different  
7       levels of expertise.
- 8       • Concurrent verbal reports are a technique that offers important inferences  
9       about thought processes.
- 10      • Skilled anaesthetists are characterised by their ability to make more evaluation  
11      statements about the current situation, predict future events and outcomes, and  
12      forward plan actions.
- 13      • Simulation training may present an ideal context for developing higher levels  
14      of expertise and the associated cognitive and technical skills

15  
16  
17  
18  
19  
20  
21

# 1 INTRODUCTION

2 Anaesthesia in the modern operating environment requires a range of attributes  
3 and skills to ensure patient care and safety. Furthermore, each patient poses a unique  
4 set of challenges and constraints that make it difficult to directly compare  
5 performance levels between and across individuals. For example, some patients  
6 present with complex and difficult problems, compared to more straightforward  
7 diagnosis and treatment pathways. The challenge for educators and researchers is to  
8 capture and examine technical skills, non-technical skills, thought processes and  
9 performance using standardised (i.e., valid, objective and reliable), representative  
10 tasks.(1) Recent developments in high-fidelity medical simulators provide more  
11 realistic, fully immersive environments for examining cognitive (decision-making),  
12 behavioural (technical and non-technical skills) and performance (diagnosis accuracy)  
13 differences between participants.(2-4) Furthermore, simulators allow control and  
14 replication of scenarios while maintaining ecological validity. In this study, the expert  
15 performance approach was used to develop anaesthetist, high-fidelity medical  
16 scenarios to examine differences in performance, cognition and behaviour.(5)

17 The expert performance approach has been used in sport, music, education and  
18 medicine to examine expertise.(6-11) First, experts are observed *in situ* so that the  
19 essential skills and competencies underpinning expertise can be captured. Based on  
20 these observations, simulated representative laboratory or field-based task(s) are  
21 created so that skill based differences in performance can be reliably and objectively  
22 measured under more controlled conditions. Second, to promote understanding of  
23 expertise, process-tracing measures such as think-aloud verbal reports and /or eye-  
24 movement recordings to examine perceptual and cognitive mechanisms that support

1 performance. Finally, development of expert skills and mechanisms are traced to  
2 identify how and when they were acquired, while considering the practical  
3 implications for guiding practice and instruction.

4 In medicine, high-fidelity simulators are used as a valid and reliable tool for  
5 assessment of competency and training because scenarios can be standardised,  
6 repeated and require the participant to interact with the patient to provide care.(3,12-  
7 14) Performance is assessed using diagnosis accuracy, response time, rating scales,  
8 non-technical behaviours (e.g., Anaesthesia Non-Technical Skill, ANTS) and  
9 technical skills with little focus on examining the thought processes and decision-  
10 making (i.e., cognition) simultaneously.(3,4,15,16)

11 Cognitions are recorded and analysed using think-aloud verbal reports and  
12 protocol analysis, providing a quantifiable measure that represents the thought  
13 processes underpinning participant actions.(17) Fox et al. (18) conducted a meta-  
14 analysis of 94 studies and found no difference in task performance when comparing  
15 think-aloud verbal reports to a matched condition without verbal reports. Therefore,  
16 they suggested that the cognitive processes mediating task performance do not alter,  
17 allowing researchers to make inferences about performance based on verbalised  
18 cognitions (see Ericsson and Simon (17) for a more detailed theoretical rationale).  
19 Furthermore, Whyte et al. (19) reported that concurrent think-aloud verbal reports  
20 provide a more complete representations of nurse's cognitions when admitting a  
21 patient suffering from an acute exacerbation of congestive heart failure to a simulated  
22 intensive care unit.

23 Think-aloud protocols have been used by researchers to examine cognitions and  
24 make inferences on performance in law enforcement, sport and medicine (20-22). For

1 example, Cormier et al. (2) examined cognitions and performance of low and high  
2 performing student nurses during high-fidelity medical simulations. They measured  
3 verbal statements coded as ‘observations’ (e.g., context, patient and monitor) or  
4 ‘actions’ (e.g., based on orders or patient condition) and physiological data of the  
5 patient’s respiratory status. High performing student nurses were better at observing  
6 relevant cues suggesting patient deterioration and forward plan responses that  
7 positively altered the patient’s physiological trajectory. In comparison, low  
8 performing student nurses verbalised more irrelevant cues and failed to perform  
9 action directly related to the patient’s condition.

10 In the current study, we use high-fidelity scenarios so that the cognitive processes  
11 associated with expertise can be fully understood. Diagnosis accuracy and non-  
12 technical behaviours of skilled and less skilled anaesthetists were examined while  
13 they provided think-aloud concurrent verbal reports during the scenarios. It was  
14 hypothesised that skilled anaesthetists would demonstrate superior diagnosis accuracy  
15 compared to less skilled anaesthetists. Second, the Anaesthetist Non-Technical Skills  
16 (ANTS) was used as a valid and reliable tool for assessing non-technical  
17 behaviours.(23) It was hypothesised that skilled participants would score a higher  
18 global rating compared to less skilled anaesthetists. Finally, skilled anaesthetists’  
19 superior performance would be supported by more evaluation, prediction and deep  
20 planning statements when compared to less skilled counterparts.

## 21 **METHOD**

### 22 **Participants**

23 Nine less skilled (mean age = 29.4 years, SD = 2.3) and 6 skilled (mean age =  
24 37.2, SD = 4.5) anaesthetists were recruited. Less skilled participants were pre-FRCA

1 trainees in their first 2 years of anaesthesia with a mean of 4.2 years (SD = 1.1)  
2 experience as a doctor. Skilled participants were at consultant or experience staff  
3 grade level with a mean of 13.3 years (SD = 3.1) experience. The study was approved  
4 by the lead institute's Research Ethics Committee and carried out under its ethical  
5 guidelines.

## 6 **Simulated Test Environment (STE)**

7 A simulated task environment (STE) was developed. The STE consisted of a high-  
8 fidelity adult METI human patient simulator (CAE Healthcare, Florida, USA) and  
9 fully equipped operating theatre, that included an anaesthetic machine, monitoring  
10 familiar to the anaesthetists, realistic functioning medical gases, intravenous pumps  
11 (IV), a real crash cart with defibrillator, medication and medical supplies. The STE  
12 allowed real-life scenarios to be dynamically presented to participants with the onset  
13 and choice of patient response under full experimenter control. METI human patient  
14 simulator responded to treatment in a realistic manner exhibiting applicable changes  
15 in cardiovascular, pulmonary, physiological, pharmacological, metabolic,  
16 genitourinary and neurological states.

## 17 **Scenarios**

18 Three tightly scripted and highly standardised simulator scenarios were  
19 developed. Each scenario lasted approximately 15 minutes, involved an emergency  
20 developing during the course of an operation and required the anaesthetist to intervene  
21 and administer treatment.

22 In scenario one (anaphylaxis), the patient is a 68-year-old man, non-smoker, no  
23 medication, allergies or airway problems, drinks alcohol occasionally, no previous



1 history of general anaesthetic (GA) and suffers from active reflux so requires an  
2 endotracheal tube (ETT). In addition, he has requested a GA, as he 'didn't like the  
3 sound of a spinal'. A confederate Surgeon, Scrub nurse and Operating Department  
4 Practitioner (ODP) are on hand to provide assistance when requested. The participant  
5 enters the room and is guided over to the anaesthetic machine where they receive a  
6 handover from a colleague who is being relieved for lunch. "The patient has just been  
7 anaesthetised for a left knee replacement under GA. Skin is prepared for surgery and  
8 the surgeon and scrub nurse are ready. Patient-controlled analgesia (PCA) is planned  
9 for post-operative pain relief". The Surgeon requests 1.5g of Cefuroxime prior to  
10 tourniquet inflation. Administration is rapidly followed by cardiovascular system  
11 (CVS) collapse and bronchospasm. The patient will require 100% oxygen,  
12 termination of the suspected agent, administration of adrenaline and referral to the  
13 intensive treatment unit (ITU).

14 In the second scenario (Malignant Hyperpyrexia), the patient is a 69-year-old  
15 man, non-smoker, no medication, allergies or airway problems, drinks alcohol  
16 occasionally, no previous history of GA or anaesthetic problems and has active reflux  
17 so was intubated for the procedure. A confederate Surgeon and Scrub nurse are busy  
18 operating with an ODP on hand to provide assistance if requested. The participant  
19 receives a handover in the theatre from a colleague who is being relieved for a  
20 meeting. "The patient was anaesthetised 20 minutes ago for removal of metalwork  
21 from the left ankle. PCA is planned for post-operative pain relief". Malignant  
22 hyperpyrexia gradually starts once the participant is settled. The participant will be  
23 required to stop the volatile, hyperventilate the patient to reduce end tidal CO<sub>2</sub>, give  
24 patient 100% oxygen, change breathing circuit/anaesthetic machine, administer  
25 Dantrolene and actively cool patient.

1        In the third scenario (Supraventricular Tachycardia post induction), the patient is a  
2 59-year-old man, who is fit and well, no previous GA problems, no allergies or  
3 airway problems, ex-smoker and drinks alcohol occasionally. The patient takes an oral  
4 dose of Lansporazola (30mg) and has well-controlled reflux. A confederate ODP is on  
5 hand to provide assistance when requested. The participant receives drugs ready  
6 drawn up in a kidney bowl and a handover, before they enter the theatre. “The patient  
7 is due to have an elective laparoscopic cholecystectomy for gallstones. PCA for post-  
8 operative pain relief is planned if an open procedure is required”. The participant will  
9 induce anaesthesia. Once intubated, the patient develops narrow complex tachycardia  
10 and his cardiovascular function becomes compromised. The participant will be  
11 required to give DC shock cardio version to the unstable patient.

## 12        **Procedure**

13        Participants completed a biographical (i.e., age, number of years training and  
14 details of training activity) information sheet and provided informed consent.  
15 Participants were instructed in a think-aloud verbal reporting method by Ericsson and  
16 Simon (17) that has been used previously in sport, law enforcement and medicine  
17 .(19,20,24) Participants practised giving verbal reports while solving non-clinical  
18 alphanumerical problems until the criterion for providing concurrent verbal reports  
19 were met. For example, if the researcher asked, “What the third letter after A is?”, you  
20 might respond “D, I counted three letters forward” but your actual thoughts might  
21 have been “A, B, C, D”. The former is an example of a summary, whereas the latter  
22 are your actual thoughts. During the training, participants accuracy on alphanumerical  
23 tasks were not recorded, as the objective was to provide feedback so that they could  
24 understand the difference between verbalising a summary of their thoughts compared

1 to think-aloud statements about their thoughts. Following their verbal report training,  
2 participants were familiarised with the STE using a structured protocol and then  
3 completed a simple practice trial while providing a concurrent verbal report. The  
4 practice trail required them to check the equipment and conduct an objective  
5 assessment of the patient. Concurrent verbal reports were selected as they provide a  
6 more complete cognitive representation of performance within the STE, compared to  
7 retrospective reports.(19) Following their practice trial, participants were given an  
8 opportunity to ask further questions prior to the experimental scenarios and further  
9 feedback were provided.

10 Participants were fitted with a lapel microphone and radio transmitter and in  
11 conjunction with 2-angle video cameras, visual and audio observations were captured.  
12 The orders of scenarios presented to each participant were counterbalanced across the  
13 participants to reduce order and learning effects. Participants interacted with the  
14 patient and confederate staff involved. In order to standardise the scenarios,  
15 confederates were equipped with a two-way radio transceiver so that the  
16 experimenters could control the information provided to the participant. If the  
17 participant remained silent for an extended period of time, the confederate ODP was  
18 instructed to remind them to concurrently verbalise their thoughts using the prompt  
19 “think-aloud”. Each scenario terminated when the 15 minutes elapsed.

## 20 **Data analysis**

### 21 **Diagnosis data**

22 A diagnosis response was marked as correct and awarded a score of ‘1’ if the  
23 participants verbalised the correct scenario diagnosis. If participants failed to provide  
24 or provided an incorrect diagnosis response, a score of ‘0’ was recorded for that

1 scenario. Diagnosis accuracy was defined as the total number of correct responses  
2 (i.e., frequency score) divided by the total number of scenarios ( $N = 3$ ) and multiplied  
3 by 100 to create a percentage score. An independent samples *t*-test was conducted to  
4 compare diagnosis accuracy between skilled and less skilled participants.

#### 5 Anaesthetists' Non-Technical Skills (ANTS) data

6 The Anaesthetists' Non-Technical Skills (ANTS) system was used to assess  
7 behavioural markers during the medical scenarios. The ANTS system's validity and  
8 reliability has previously been established when assessing non-technical  
9 behaviours.(23) A consultant anaesthetist trained and qualified in the use of the ANTS  
10 system reviewed videotapes of the participants' non-technical skills during the  
11 scenarios. ANTS describes the main observable non-technical skills associated with  
12 good anaesthetic practice.

13 The ANTS system comprises of four skill categories and beneath these are fifteen  
14 skill elements (see Table 1). Each element has a definition and examples of good and  
15 poor behaviours. The reviewer rated the 15 skill elements based on a four-point scale  
16 (i.e., 4 = good; 3 = acceptable, 2 = marginal; 1- poor; 0 – not observed). The rating  
17 score for each element was added together to calculate the participant's global  
18 performance score for each scenario. A mean ANTS score was calculated based on  
19 the participant's global scores from the three scenarios. A higher score on the ANTS  
20 represented better non-technical behaviours during the scenarios. An independent  
21 samples *t*-test was conducted to compare differences in ANTS scores between skilled  
22 and less skilled participants.

23

1 Table 1: ANTS four skill categories and fifteen skill elements

<b>Category</b>	<b>Elements</b>
Task Management	Planning and preparing Prioritising Providing and maintaining standards Identifying and utilising resources
Team Working	Co-ordinating activities with team members Exchanging information Using authority and assertiveness Assessing capabilities Supporting others
Situation Awareness	Gathering information Recognising and understanding Anticipating
Decision Making	Identifying options Balancing risks and selecting options Re-evaluating

2

3 Verbal Report data

4 Scenario videos were captured and analysed using Studiocode, version 3.5  
 5 (Sportstec Ltd, Australia), an analytical tool that allows the researcher to mark and  
 6 code segments of video and audio into categories. Following the procedures outlined  
 7 in Ericsson and Simon (17) verbal reports were initially segmented using natural  
 8 speech and syntactical markers.

9 Verbal statements were coded based on categories developed and previously used  
 10 by McRobert et al. (22) to monitor thought processes of participants during  
 11 emergency medicine simulated scenarios. To ensure logical validity and remove the  
 12 individual bias of the researcher a panel of anaesthetists (N = 3), medical educators (N

1 = 2) and researchers (N = 2) reviewed the sample of statements from the simulated  
2 scenarios that had been assigned to the coding categories. (25,26) The panel provided  
3 written feedback and agreed that the categories and operational definitions were  
4 appropriate for analysing concurrent verbal statements.

5 Table 2: Verbal report statement categories and definitions

<b>Category</b>	<b>Definition</b>
Monitor	Statements representing information that was present or previously present in the current environment.
Evaluation	Some form of positive or negative assessment of a prior action, event or statement.
Prediction	Statements about what could, would and should occur next in the environment.
Deep Planning	Statements are about future actions and options in a future situation.

6

7 The primary experimenter analysed all participant data and assigned each  
8 concurrent verbal statement to the monitor, evaluation, prediction or deep planning  
9 category (Table 2). Fifty verbal statements were selected at random to establish  
10 objectivity and reliability using the inter- and intra-observer percentage agreement  
11 formulas.(27) The primary experimenter coded the 50 statements on four separate  
12 occasions (i.e., prior to coding, after 6, 12 and 18 participants) with an intra-observer  
13 agreement range from 95% to 98%. Inter-observer agreement was assessed prior to  
14 coding (92%) the statements, and after 12 participants (93%) by an independent  
15 experimenter.

1 Two-way analysis of variance (ANOVA) was conducted with group (less  
2 skilled/skilled) as the between participant factor and verbal statement type  
3 (monitor/evaluation/prediction/deep planning) as the within-participant factors. Partial  
4 eta squared ( $\eta^2$ ) values are provided as a measure of effect size. When making  
5 comparisons between two means, Cohen's *d* measures are reported. Posthoc  
6 Bonferroni corrected pairwise comparisons are reported as follow-ups where  
7 appropriate.

## 8 **RESULTS**

### 9 **Diagnosis data**

10 An independent samples *t*-test was conducted to compare differences in diagnosis  
11 accuracy between skilled and less skilled participants. Levene's test for homogeneity  
12 of variance was violated ( $p < .05$ ); therefore the test statistic for equal variance not  
13 assumed was reported. There was a significant difference in diagnosis accuracy scores  
14 for less skilled and skilled participants,  $t_{13} = -2.688$ ,  $p = .02$ . Skilled participants ( $M$   
15  $= 94\%$ ,  $SD = 14$ ) diagnosis accuracy scores were significantly higher than less skilled  
16 participants ( $M = 63\%$ ,  $SD = 31$ ).

### 17 **Anaesthetists' Non-Technical Skills (ANTS) data**

18 The independent samples *t*-test was conducted to compare ANTS score  
19 differences between skilled and less skilled participants. There was a significant  
20 difference in ANTS scores between skilled and less skilled participants,  $t_{13} = -3.215$ ,  
21  $p = .02$ . Skilled participants ( $M = 22$ ,  $SD = 4$ ) ANTS scores were significantly higher  
22 than less skilled participants ( $M = 13$ ,  $SD = 6$ ).

### 23 **Verbal Report data**

1 ANOVA showed a significant Group  $\times$  Verbal Statement Type interaction ( $F_{3,39}$   
2 = 20.33,  $p < .001$ ,  $\eta p^2 = .52$ ). As predicted, less skilled participants made more  
3 monitoring statements ( $d = 1.19$ ) compared to their skilled counterparts (see Figure  
4 1). In contrast, skilled participants made more evaluation ( $d = 2.2$ ) and deep planning  
5 ( $d = 1.76$ ) statements compared to less skilled participants. Skilled participants did  
6 make more prediction statements compared to less skilled participants, however this  
7 was only a medium effect size ( $d = 0.52$ ). The Cohen's  $d$  effect size score quantifies  
8 the magnitude of the difference between the skilled and less skilled groups. There is a  
9 larger difference between the skilled and less skilled groups for evaluation and deep  
10 planning, compared to prediction. It is worth noting that skilled and less skilled  
11 standard deviations for the monitor, evaluation and deep planning are relative to the  
12 mean. In contrast, the standard deviation for prediction is large relative to the mean  
13 score. The large variance could be due to the low number of prediction statements  
14 compared to the other categories.

15 Significant main effects for group ( $F_{3,39} = 10.83$ ,  $p < .05$ ,  $\eta p^2 = .45$ ) and verbal  
16 statement type ( $F_{3,39} = 67.12$ ,  $p < .001$ ,  $\eta p^2 = .83$ ) were observed. Skilled  
17 participants ( $M = 12$ ,  $SD = 8$ ) made more verbal statements, compared to less skilled  
18 participants ( $M = 10$ ,  $SD = 5$ ). Pairwise comparisons demonstrated that all  
19 participants made more evaluation and deep planning than monitor and prediction  
20 statements ( $p < .05$ ). Second, a higher number of monitor statements were coded  
21 compared to prediction statements ( $p < .05$ ).

22

23

24



1 INSERT FIGURE 1 HERE.

2

3 **DISCUSSION**

4 We examined the cognitive processes, diagnosis accuracy and non-technical  
5 behaviours in anaesthetists during simulated medical scenarios. Skilled and less  
6 skilled participants were required to diagnose the patient's condition and engage in  
7 effective treatment during fully immersive, high-fidelity, medical scenarios. Verbal  
8 reports were collected concurrently during the simulation as an index of thought  
9 processes. In addition, the Anaesthetists' Non-Technical Skills (ANTS) were assessed  
10 retrospectively to rate the effectiveness of the participant's behaviours. It was  
11 hypothesised that skilled participants would demonstrate superior diagnosis accuracy  
12 and a higher global rating score on the ANTS, compared to less skilled participants.  
13 Moreover, skilled participants' would be characterised by a higher total number of  
14 statements, specifically more evaluation, prediction and deep planning statements.

15 As predicted, skilled participants outperformed less skilled participants on  
16 diagnosis accuracy. Findings support differences in skilled and less skilled  
17 performance reported previously by researchers in medicine and other domains such  
18 as aviation, law enforcement and sport using dynamically evolving simulated  
19 tasks.(3,20,28,29) In addition, skilled participants had higher ANTS scores suggesting  
20 that as a group their task management, team working, situation awareness and  
21 decision-making skills when managing the scenarios are better.

22 Think-aloud concurrent verbal reports and protocol analysis was conducted to  
23 examine if there were any differences between less skilled and skilled anaesthetists

1 thought processes. Skilled participants made more verbal statements in total,  
2 compared to less skilled participants. In addition, skilled participants' made more  
3 evaluation, prediction and deep planning and fewer monitoring statements, compared  
4 to less skilled participants'. Moreover, these differences are larger for evaluation and  
5 deep planning statements between the groups compared to prediction. Skilled  
6 participants appear to be characterised by their ability to make more evaluation  
7 statements about the current situation, predict future events and outcomes, and  
8 forward plan actions. In addition, the verbal report data in this study support previous  
9 work demonstrating skilled participants thought processing during dynamically  
10 evolving tasks in sport and law enforcement.(20,21)

11 To provide further context, monitoring statements often refer to an observation of  
12 oxygen saturation, core temperature or actions such as administering oxygen,  
13 adrenaline and reducing end tidal CO<sub>2</sub> that have taken place during the scenario.  
14 Monitoring statements are just observations, whereas evaluation statements involve an  
15 assessment of the observed information or action. Evaluation statements would  
16 involve the observation of oxygen saturation or core temperature followed by an  
17 assessment of the value compared to the norm value expected. Prediction statements  
18 are assumptions or forecasts about what might or should happen next based on  
19 evaluation of observed information (e.g., change in patient vitals) or most recent  
20 action (e.g., patient intervention). Deep planning statements provide information on  
21 future actions that the participant will to undertake in order to manage potential future  
22 situations that might occur. This action plan profile is often based on a primary  
23 diagnosis and differential diagnosis for so that they can plan a series of future actions  
24 to deal with anaphylaxis (e.g., provide 100% oxygen, terminate suspected agent,  
25 administer adrenalin and referral to eth intensive care unit), malignant hyperpyrexia

1 (e.g., stop the volatile, hyperventilate the patient to reduce end tidal CO<sub>2</sub>, give 100%  
2 oxygen, change breathing circuit/anaesthetic machine, administer Dantrolene and  
3 actively cool) and supraventricular tachycardia (e.g., give DC shock cardio version).

4 The current findings have implications for those in medicine that have used the  
5 recent developments in high-fidelity simulation for the training and assessment of  
6 competency.(2-4) Our data provides construct validity for the use of high-fidelity  
7 medical simulation as a vehicle for assessment of skill competency. Moreover, such  
8 simulations may provide an ideal vehicle for training because it presents a safe  
9 environment, where scenarios can be replicated so that the examiner is confident that  
10 each individual can experience and be evaluated on the same factors. In sum, it  
11 presents a valid, reliable and objective environment for testing and training expertise  
12 in medicine. Furthermore, simulators can conduct scenarios that occur on a rare basis.  
13 This ability to present novel or rare scenarios will allow participants to develop  
14 domain-specific knowledge that can facilitate diagnosis and patient care if the  
15 scenario was to occur during their normal working hours. However, it is worth noting  
16 that the size, expense and staff support required to manage a simulation environment  
17 are often a barrier to widespread use, particular outside of large hospitals and major  
18 cities. Finally, the recording of verbal reports provides an opportunity to design  
19 educational programmes that specifically target clinical reasoning based on an expert-  
20 model approach (30,31). Immersive simulation techniques often involve uninterrupted  
21 real-time cases, followed by an individual or group debriefing. Introducing think-  
22 aloud verbal reporting during a scenario, allows participants to retrospectively reflect  
23 on their thought processes in conjunction with the educator to identify knowledge  
24 gaps or reframe clinical reasoning (i.e., cognitive processing).

1 In conclusion, we examined the cognitive processes and non-technical skills used  
2 by skilled and less skilled anaesthetists during high-fidelity medical scenarios. Skilled  
3 participants demonstrated superior diagnosis accuracy, better non-technical  
4 behaviours and verbalised more evaluation, prediction and deep planning statements,  
5 and, compared to less skilled participants. In comparison to less skilled participants,  
6 skilled participants were characterised by the ability to evaluate the current situation  
7 and engage in prediction of future events and to forward plan actions rather than  
8 thinking about immediately available information. Our findings suggest that  
9 simulations present a valid method for quantifying and exploring expertise in medical  
10 contexts, whereas at a practical level the significant potential to use such  
11 environments for training health care practitioners is highlighted.

## 12 **Contributions**

13 MAW, DR and APM were responsible for the conception and design of the  
14 current study. Second, DR, SM, JG and APM collected the data and engaged in many  
15 hours of subsequent analysis and interpretation of this data. Third, ARM was  
16 responsible for the initial draft of the article, revisions and approval of the final  
17 version submitted. Finally, MAW, DR and SM were responsible for critically revising  
18 drafts for intellectual content and quality of submission.

## 19 **Funding**

20 There are no funders to report for this submission.

## 21 **Competing interests**

22 There are no competing interests.

1           **REFERENCES**

- 2    1.    Ericsson KA, Ward P. Capturing the Naturally Occurring Superior  
3           Performance of Experts in the Laboratory: Toward a Science of Expert and  
4           Exceptional Performance. *Curr Dir Psychol Sci.* 2007;16(6):346–50.
- 5    2.    Cormier EM, Pickett-Hauber R, Whyte J. Cognitions and clinical performance:  
6           a comparison of high and low performing baccalaureate nursing students. *Int J*  
7           *Nurs Educ Scholarsh.* 2010;7(1):Article27.
- 8    3.    Weller J, Bloch M, Young S, Maze M, Oyesola S, Wyner J, et al. Evaluation of  
9           high fidelity patient simulator in assessment of performance of anaesthetists. *Br*  
10           *J Anaesth.* 2003;90(1):43–7.
- 11   4.    Zheng B, Tien G, Atkins SM, Swindells C, Tanin H, Meneghetti A, et al.  
12           Surgeon's vigilance in the operating room. *Am J Surg.* Elsevier Inc. 2011  
13           May;201(5):673–7.
- 14   5.    Ericsson KA, Smith J. *Toward a general theory of expertise: Prospects and*  
15           *limits.* Cambridge: Cambridge University Press; 1991.
- 16   6.    Ericsson KA, Krampe RT, Tesch-Römer C. The role of deliberate practice in  
17           the acquisition of expert performance. *Psychol Rev.* 1993;100(3):363.
- 18   7.    Ford PR, Low J, McRobert AP, Williams AM. Developmental activities that  
19           contribute to high or low performance by elite cricket batters when recognizing  
20           type of delivery from bowlers' advanced postural cues. *J Sport Exer Psychol.*  
21           2010;32(5):638–54.
- 22   8.    Plant EA, Ericsson KA, Hill L, Asberg K. Why study time does not predict  
23           grade point average across college students: Implications of deliberate practice  
24           for academic performance. *Contemp Educ Psychol.* 2005;30(1):96–116.
- 25   9.    Whyte J, Ward P, Eccles DW. The relationship between knowledge and  
26           clinical performance in novice and experienced critical care nurses. *Heart*  
27           *Lung.* 2009 Nov;38(6):517–25.
- 28   10.   Williams A, WARD P. Perceptual Expertise: Development in Sport. In: Starke  
29           J, Ericsson K, editors. *Expert Performance in Sport: Recent Advances in*  
30           *Research on Sport Expertise.* Human Kinetics, Champaign, IL; 2003. pp. 220–  
31           49.
- 32   11.   Causer J, Barach P, Williams AM. Expertise in medicine: using the expert  
33           performance approach to improve simulation training. *Med Educ.*  
34           2014;48(2):115–23.
- 35   12.   Lorroway PG, Savoldelli GL, Joo HS, Chandra DB, Chow R, Naik VN.  
36           Management of simulated oxygen supply failure: is there a gap in the  
37           curriculum? *Anesth Analg.* 2006;102(3):865–7.
- 38   13.   Nehring WM. U.S. boards of nursing and the use of high-fidelity patient

- 1           simulators in nursing education. *J Prof Nurs*. 2008;24(2):109–17.
- 2   14.   Weller J, Robinson B, Jolly B. Psychometric characteristics of simulation-  
3           based assessment in anaesthesia and accuracy of self- assessed scores\*.  
4           *Anaesth*. 2005;60:245–50.
- 5   15.   Fletcher GCL, McGeorge P, Flin RH, Glavin RJ, Maran NJ. The role of non-  
6           technical skills in anaesthesia: a review of current literature. *Br J Anaesth*.  
7           2002;88(3):418–29.
- 8   16.   Schwid HA, Rooke GA, Carline J, Steadman RH, Murray WB, Olympio M, et  
9           al. Evaluation of anesthesia residents using mannequin-based simulation: a  
10          multiinstitutional study. *Anesthesiol*. 2002;97(6):1434.
- 11  17.   Ericsson KA, Simon HA. Protocol Analysis: Verbal Reports as Data, revised  
12          edition. revised edition. A Bradford Book; 1993.
- 13  18.   Fox MC, Ericsson KA, Best R. Do procedures for verbal reporting of thinking  
14          have to be reactive? A meta-analysis and recommendations for best reporting  
15          methods. *Psychol Bull*. 2011;137(2):316–44.
- 16  19.   Whyte J IV, Cormier E, Pickett-Hauber R. Cognitions associated with nurse  
17          performance: A comparison of concurrent and retrospective verbal reports of  
18          nurse performance in a simulated task environment. *Int J Nurs Stud*.  
19          2010;47(4):446–51.
- 20  20.   Ward P, Suss J, Eccles DW, Williams AM, Harris KR. Skill-based differences  
21          in option generation in a complex task: a verbal protocol analysis. *Cog Process*.  
22          2011;12(3):289–300.
- 23  21.   Roca A, Ford PR, McRobert AP, Mark Williams A. Identifying the processes  
24          underpinning anticipation and decision-making in a dynamic time-constrained  
25          task. *Cog Process*. 2011;12(3):301–10.
- 26  22.   McRobert AP, Causer J, Vassiliadis J, Watterson L, Kwan J, Williams MA.  
27          Contextual information influences diagnosis accuracy and decision making in  
28          simulated emergency medicine emergencies. *BMJ Qual Saf*. 2013; 22(6):478–  
29          84.
- 30  23.   Fletcher G, Flin R, McGeorge P, Glavin R, Maran N, Patey R. Rating non-  
31          technical skills: developing a behavioural marker system for use in anaesthesia.  
32          *Cog Tech Work*. 2004;6(3):1–8.
- 33  24.   North JS, Ward P, Ericsson A, Williams AM. Mechanisms underlying skilled  
34          anticipation and recognition in a dynamic and temporally constrained domain.  
35          *Memory*. 2011;19(2):155–68.
- 36  25.   George K. Validity in clinical research: a review of basic concepts and  
37          definitions. *Phys Ther Sport*. 2003;4(3):115–21.
- 38  26.   Cushion C, Harvey S, Muir B, Nelson L. Developing the Coach Analysis and  
39          Intervention System (CAIS): Establishing validity and reliability of a

1 computerised systematic observation instrument. *J Sports Sci.* 2012;30(2):201–  
2 16.

3 27. Thomas J, Nelson J, Silverman S. *Research Methods in Physical Activity.*  
4 Champaign, IL: Human Kinetics; 2010.

5 28. Doane SM, Sohn YW, Jodlowski MT. Pilot ability to anticipate the  
6 consequences of flight actions as a function of expertise. *Hum Factors.*  
7 2004;46(1):92–103.

8 29. Mann DTY, Williams AM, Ward P, Janelle CM. Perceptual-cognitive expertise  
9 in sport: a meta-analysis. *J Sport Exerc Psychol.* 2007;29(4):457–78.

10 30. Ericsson KA, Whyte J, Ward P. Expert performance in nursing: reviewing  
11 research on expertise in nursing within the framework of the expert-  
12 performance approach. *Adv Nurs Sci.* 2007;30(1):58–71.

13 31. Kneebone RL. Simulation reframed. *Adv Simul.* 2016;1(27):1–8.

14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

**Figure legend**

Figure 1: Number of verbal statement types (with *SD* bars) for skilled and less skilled participants.



