



LJMU Research Online

Ballard, A, Ahmad, HO, Narduolo, S, Rosa, L, Chand, N, Cosgrove, DA, Varkonyi, P, Asaad, N, Tomasi, S, Buurma, NJ and Leach, AG

Quantitative Prediction of Rate Constants for Aqueous Racemization To Avoid Pointless Stereoselective Syntheses

<http://researchonline.ljmu.ac.uk/7450/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Ballard, A, Ahmad, HO, Narduolo, S, Rosa, L, Chand, N, Cosgrove, DA, Varkonyi, P, Asaad, N, Tomasi, S, Buurma, NJ and Leach, AG (2017) Quantitative Prediction of Rate Constants for Aqueous Racemization To Avoid Pointless Stereoselective Syntheses. *Angewandte Chemie*

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

A quantitative approach to predicting rate constants for aqueous racemization allows pointless stereoselective syntheses to be avoided.

Andrew Ballard, Hiwa O. Ahmad, Stefania Narduolo, Lucy Rosa, Nikki Chand, David A. Cosgrove, Peter Varkonyi, Nabil Asaad, Simone Tomasi, Niklaas J. Buurma,* Andrew G. Leach*

Abstract: Racemization has a large impact upon the biological properties of molecules but the chemical scope of compounds with known rate constants for racemization in aqueous conditions was hitherto limited. To address this remarkable blind spot, we have measured the kinetics for racemization of 28 compounds using circular dichroism and ^1H NMR spectroscopy. We show that rate constants for racemization (measured by ourselves and others) correlate well with deprotonation energies from quantum mechanical (QM) and group contribution calculations. Such calculations thus provide predictions of the second-order rate constants for general-base-catalyzed racemization that are usefully accurate. When applied to recent publications describing the stereoselective synthesis of compounds of purported biological value, the calculations reveal that racemization would be sufficiently fast to render these expensive syntheses pointless.

Thalidomide racemizes in a matter of hours and yet it remains a poster child for enantioselective synthesis which would not have saved its victims.^[1] The status quo in enantioselective synthesis thus ignores the cruel blind spot that we address in this paper: racemization.

Although necessary in dynamic kinetic resolution protocols,^[2,3] racemization and epimerization can cause safe compounds to become toxic or lose efficacy,^[4–11] lead to misidentification of chiral compounds extracted from natural sources,^[12] etc. Ignoring racemization leads to wasted material and human resources.

Racemization is a particular problem because its detection requires chiral analytical methods.^[13,14] Hence, few reports disclose rate constants for racemization under aqueous conditions.^[1,15–20] Chiral centers with certain combinations of substituents have been posited to be prone to general-base-catalyzed racemization although with little supporting data.^[21–23]

We therefore classified stereogenic carbon atoms according to their attached substituents. Each substituent is identified as one of sixty types,^[24] which encompass more than 99.95% of all such substituents in the GOSTAR database.^[25] The ten most frequently occurring substituents are listed in Figure 1; the H required for general-base-catalyzed racemization is prominent.^[24] Groups labelled * were selected for experimental study.

Based on prevalence, earlier work,^[21–23] and chemical intuition, several compounds were selected for detailed kinetic studies.^[26–28] The rate constants for general-base-catalyzed racemization were derived for a range of 11 arylglycine derivatives (**1**, **2** and **3**), 12 hydantoins (**4**, **5** and **6**) and 5 thiohydantoins (**7** and **8**). Briefly, at several buffer concentrations, circular dichroism spectroscopy (CD) followed the decrease in ellipticity and/or ^1H NMR spectroscopy the incorporation of D from deuterated buffers. The pseudo-first-order rate constants for these processes were corrected for hydrolysis side reactions if required.^[24] Plotting the first-order rate constants for racemization or H/D exchange against the concentration of the basic component of the buffer yielded the second-order rate constants for general-base-catalyzed racemization. These were corrected for reaction temperature and substrate protonation state.^[24]




Figure 1. The ten substituents appearing most frequently adjacent to carbon stereogenic centers in the GOSTAR database. Alk = alkyl, C = carbon-linked alkyl or aromatic and X = any group. * = selected for experimental study.

For predictive modeling, a mechanistic understanding is beneficial. Racemization of the stereogenic centers studied here could occur by either an $\text{S}_{\text{E}}1$ or an $\text{S}_{\text{E}}2$ mechanism. For hydantoins (e.g. **4–6**) both the $\text{S}_{\text{E}}1$ and $\text{S}_{\text{E}}2$ mechanisms have been proposed previously,^[18,29] but we have shown that these reactions occur via the $\text{S}_{\text{E}}1$ mechanism.^[30] Further, Hammett plots show a better correlation with σ^- than σ suggesting that a negative charge is formed on the stereogenic center in the rate-determining step of the racemization reaction, in line with an $\text{S}_{\text{E}}1$ mechanism.^[24] The experimental data were correlated with deprotonation energies ($\Delta\Delta\text{G}(\text{R}_1, \text{R}_2, \text{R}_3)$, Scheme 1) from B3LYP/6-31+G** calculations incorporating aqueous solvation via the PCM protocol.^[24] Second-order rate constants for

Dr A. Ballard, Dr H.O. Ahmad, Dr S. Narduolo and Dr N.J. Buurma
Physical Organic Chemistry Centre, School of Chemistry, Cardiff
University, Main Building, Park Place, Cardiff, CF10 3AT, UK.
E-mail: Buurma@Cardiff.ac.uk

Dr H.O. Ahmad
Pharmaceutical Chemistry Department, College of Pharmacy,
Hawler Medical University, Erbil, Kurdistan Region, Iraq


L. Rosa, N. Chand, Dr. D. A. Cosgrove, Dr N. Asaad
AstraZeneca Pharmaceuticals, Mereside, Alderley Park,
Macclesfield, SK10 4TG, UK.

Dr P. Varkonyi
AstraZeneca R+D, Pepparedsleden 1, S-431 83 Mölndal, Sweden.


Dr S. Tomasi
AstraZeneca, Charter Way, Silk Road Business Park, Macclesfield,
SK10 2NA, UK

Dr A. G. Leach
School of Pharmacy and Biomolecular Sciences, Liverpool John
Moore's University, Liverpool, L3 3AF, UK.
E-mail: a.g.leach@ljmu.ac.uk


general-base-catalyzed racemization, k_{gb} , correlate well with $\Delta\Delta G(R_1, R_2, R_3)$ for **1-3** and **4-8**.^[24]



The set of compounds was supplemented with literature data,^[24] leading to the relationship with $\Delta\Delta G(R_1, R_2, R_3)$ shown at the top of Figure 2. The line of best fit has equation $\log(k_{gb}) = -0.20 \times \Delta\Delta G(R_1, R_2, R_3) - 14.28$, with an R^2 value of 0.68 and root mean square error of 0.61, i.e. reproducing rate constants to within a factor of ~ 4 . Clopidogrel is excluded from this analysis due to large experimental uncertainties.^[24]



Scheme 1.



The computational procedure was extended to include a group-contribution approach that is amenable to rapid analysis of chiral compounds and is described in the supporting information. A simple representative (R) of each substituent type was selected and $\Delta\Delta G(R_1, R_2, R_3)$ computed with $R_1=R$ and $R_2=R_3=H$. These $\Delta\Delta G(R, H, H)$ values indicate how much R stabilizes an adjacent anion. The sum $\Delta\Delta G(R_1, H, H) + \Delta\Delta G(R_2, H, H) + \Delta\Delta G(R_3, H, H)$, for the three non-H substituents around a chiral carbon atom is referred to here as $\Sigma\Delta\Delta G$. When two or three of the groups provide stabilization through charge delocalization (as indicated in Section S7), a cross-conjugation correction is applied to

reflect the reduced ability of the second group to stabilize the anion caused by the presence of the first.^[24]

The 35 compounds studied fall into fourteen chiral carbon atom types (A – N, Table 1) that have one hydrogen attached.^[1,15–19,31] The second-order rate constants for general-base-catalyzed racemization are plotted against $\Sigma\Delta\Delta G$ at the bottom of Figure 2. When a chiral center type is represented by more than one compound, the mean value of $\log(k_{gb})$ for all representatives was used and the full range of values is shown as a vertical line. This prevents any center type from dominating the linear fit.




Figure 2. Second-order rate constants for racemization under aqueous general-base-catalyzed conditions plotted against: computed $\Delta\Delta G(R_1, R_2, R_3)$ values (top) and $\Sigma\Delta\Delta G$ values (bottom). Points are shown as Top: **1a** (■), **1b** (◆), **1c** (▲), **1d** (●), **1e** (□), **1f** (◇), **1g** (△), **1h** (○), **2** (■), **3a** (■), **3b** (◆), **4a** (■), **4b** (◆), **4c** (▲), **4d** (●), **4e** (□), **4f** (◇), **4g** (△), **5a** (■), **5b** (◆), **5c** (▲), **5f** (◇), **6** (■), **7a** (■), **7b** (◆), **7c** (▲), **7d** (●), **8** (■), **9** (■), **10** (◆), **11** (▲), **12** (●), **13** (□), **14** (◇), **15** (△), **16** (○). Bottom: **A** (■), **B** (◆), **C** (▲), **D** (■), **D'** (◇), **E** (◆), **F** (▲), **G** (●), **H** (●), **I** (□), **J** (◇), **K** (□), **L** (△), **M** (○) and **N** (○). Clopidogrel (**16**) is excluded from the line fitting.

Table 1. Stereogenic center types.

Center Type	Non-H Substituents	Representative compounds [Number]
A	Phenyl, Reversed 2° amide, Ester	1a-h [8]
B	Phenyl, Reversed 2° amide, 1° amide	2 [1]
C	5-Membered aromatic, Reversed 2°	3 [2]

	amide, Ester	
D/D'	Alkyl, Reversed 2° amide, Acidic 2° amide	D = 4a-g, 5a-c+f [11] D' = 14 [1]
E	Phenyl, Reversed 2° amide, Acidic 2° amide	6 [1]
F	Alkyl, Aminothiooxo imide, Acidic 2° amide	7a-d [4]
G	Alkyl, Reversed 2° thioamide, Acidic 2° amide	8 [1]
H	Ketone, Dialkyl 3° amine, Alkyl	9 [1]
I	Ketone, 1° amine, Alkyl	10 [1]
J	Carboxylic acid, 5-Membered aromatic, Alkyl	11 [1]
K	Thioether, Alkyl, Acidic 2° amide	12 [1]
L	Imide, Alkyl, Acidic 2° amide	13 [1]
M	Phenyl, Phenyl, 5-Membered aromatic	15 [1]
N	Ester, Dialkyl tertiary amine, Phenyl	16 [1]

For the phenylglycine esters (A), substituent effects can cause up to a log unit variation from the line of best fit. This is likely to be representative of general substituent effects.^[24] Grouping five-membered aromatic rings together (C) masks variation of 1.6 log units, likely reflecting the more direct influence of heteroatoms in aromatic rings. A relatively diverse set of alkyl substituents in the 5-position on a hydantoin ring (group D) or thiohydantoin (F) causes little variation in rate constants for racemization. In general, variation caused by substitution or structural variation within classes is less than two orders of magnitude and typically less than one order of magnitude.

At the bottom of Figure 2, two subgroups are apparent: those involving a cyclic anion with the potential to be aromatic and those that do not. For the non-aromatic set (shown in red) a line of best fit with equation $\log(k_{\text{gb}}) = -0.11 \times \Sigma\Delta\Delta G - 9.81$ was found ($R^2 = 0.78$ and $\text{RMSE} = 0.40$ log units) and for the aromatic anion set (shown in blue) the line of best fit has equation $\log(k_{\text{gb}}) = -0.26 \times \Sigma\Delta\Delta G - 16.95$ ($R^2 = 0.92$ and $\text{RMSE} = 0.39$ log units). The RMSE for all compounds computed individually is 0.64 for non-aromatic anions (excluding clopidogrel) and 0.37 for aromatic anions i.e. predictions are typically within 5-fold. Although not a perfect guide, the group contribution approach provides an easily applied, useful and rapid filtering that can even be used for very large databases.

For the particular example of chiral pharmaceuticals, our analysis can be applied to predict half lives of racemization in physiological conditions. The rate constants for racemization of thalidomide at different phosphate buffer concentrations compared to that in blood suggest that, in terms of availability of catalytically active general bases, blood is approximately equivalent to a 0.15 M phosphate buffer at pH 7.2. Therefore, with k_{gb} predicted by the QM or group contribution method, the required half lives can be predicted.^[1]

A comprehensive workflow has now arisen: rapid analysis with a group contribution based method can trigger quantum mechanical calculations, which in turn can trigger an experimental protocol (Figure 3). Compounds at high risk of racemization can be avoided and racemization risk can be suppressed by design.




Figure 3. Workflow for identifying compounds at risk of racemization, in parentheses is the typical time taken to process one compound at each step.

To illustrate the degree to which racemization in aqueous conditions is an overlooked problem, we have surveyed recent editions of leading chemistry journals, using our knowledge of the group contributions, to identify several articles envisaging biological applications. This was not an exhaustive search. Compounds described were subject to group contribution calculations. It is disappointing to reveal (Table 2) that liability to racemize under physiological conditions is more commonplace than would be possible if it were properly understood and controlled, as many chemists seem to believe.

Table 2. Examples of potentially pointless stereoselective syntheses from recent literature.

Reference	Representative compound	$\Sigma\Delta\Delta G$ (kcal/mol)	Predicted % racemized within 24 hours
[32]		-48.8	28 %
[33]		-50.7	40 %
[34]		-54.1	70 %
[35]		-46.9	19 %

In summary, we describe an approach to quantitative predictions of racemization risk that is generally applicable and allows synthetic chemists to avoid racemization-prone targets or

understand erosion of e.e. It allows users of chiral compounds, as pharmaceuticals or otherwise, to assess the risk of them becoming mixtures of compounds.

Acknowledgements

The EPSRC and AstraZeneca are thanked for a CASE for New Academics award, the Kurdistan Regional Government is thanked for a PhD studentship (HOA), Advanced Research Computing @ Cardiff (ARCCA) are thanked for computational resources.

Keywords: racemization • kinetics • computational • stereogenic center • stereoselective synthesis

References

- [1] M. Reist, P. A. Carrupt, E. Francotte, B. Testa, *Chem. Res. Toxicol.* **1998**, *11*, 1521–1528.
- [2] K. Nakano, M. Kitamura, *Sep. Enantiomers* **2014**.
- [3] M. T. El Gihani, J. M. Williams, *Curr. Opin. Chem. Biol.* **1999**, *3*, 11–15.
- [4] E. J. Ariens, W. Soudijn, P. B. M. W. M. Timmermans, *Stereochemistry and Biological Activity of Drugs*, Blackwell Scientific Publications, Oxford, England, **1983**.
- [5] E. J. Ariens, in *Stereochem Biol Act Drugs Rev Int Symp*, **1983**, pp. 11–32.
- [6] E. J. Ariens, *Trends Pharmacol. Sci.* **1993**, *14*, 68–75.
- [7] S. W. Smith, *Toxicol. Sci.* **2009**, *110*, 4–30.
- [8] A. K. Peepliwal, S. B. Bagade, C. G. Bonde, *J. Biomed. Sci. Res.* **2010**, *2*, 29–45.
- [9] A. G. Leach, E. A. Pilling, A. A. Rabow, S. Tomasi, N. Asaad, N. J. Buurma, A. Ballard, S. Narduolo, *MedChemComm* **2012**, *3*, 528–540.
- [10] F. Lovering, J. Bikker, C. Humblet, *J. Med. Chem.* **2009**, *52*, 6752–6756.
- [11] J. T. Bagdanoff, Y. Xu, G. Dollinger, E. Martin, *J. Med. Chem.* **2015**, *58*, 5781–5788.
- [12] F. Pellati, G. Cannazza, S. Benvenuti, *J. Chromatogr. A n.d.*, *1217*, 3503–3510.
- [13] L. Calabrese, A. Fleischer, *Am J Med* **2000**, *108*, 487–495.
- [14] I. Ali, V. K. Gupta, H. Y. Aboul-Enein, P. Singh, B. Sharma, *Chirality* **2007**, *19*, 453–463.
- [15] M. Reist, L. H. Christiansen, P. Christoffersen, P.-A. Carrupt, B. Testa, *Chirality* **1995**, *7*, 469–473.
- [16] M. Reist, M. Roy-de Vos, J. P. Montseny, J. M. Mayer, P. A. Carrupt, Y. Berger, B. Testa, *Drug Metab. Dispos. Biol. Fate Chem.* **2000**, *28*, 1405–1410.
- [17] C. Danel, C. Foulon, J.-F. Goossens, J.-P. Bonte, C. Vaccher, *Tetrahedron Asymmetry* **2006**, *17*, 2317–2321.
- [18] M. Reist, P.-A. Carrupt, B. Testa, S. Lehmann, J. J. Hansen, *Helv. Chim. Acta* **1996**, *79*, 767–778.
- [19] M. Brandl, D. Conley, D. Johnson, D. Johnson, *J. Pharm. Sci.* **1995**, *84*, 1045–1048.
- [20] B. Jamali, I. Bjornsdottir, O. Nordfang, S. H. Hansen, *J. Pharm. Biomed. Anal.* **2008**, *46*, 82–87.
- [21] B. Testa, P.-A. Carrupt, J. Gal, *Chirality* **1993**, *5*, 105–111.
- [22] P. Jenner, B. Testa, *Drug Metab. Rev.* **1973**, *2*, 117–184.
- [23] E. J. Ebbers, G. J. A. Ariaans, J. P. M. Houbiers, A. Bruggink, B. Zwanenburg, *Tetrahedron* **1997**, *53*, 9417–9476.
- [24] For further details, see supporting information.
- [25] <http://www.gostardb.com/gostar/loginEntry.do>
- [26] A. Ballard, Kinetics and Mechanism of H/D Exchange Reactions and Racemization in Aqueous Solutions: Configurational Stability of Ester and Amide Arylglycine Derivatives, Kinetics and mechanism of H/D exchange reactions and racemization in aqueous solutions: configurational stability of ester and amide arylglycine derivatives, Cardiff University, **2011**.
- [27] S. Narduolo, The Mechanism of Racemization of 5-Substituted Hydantoins in Aqueous Solution, The mechanism of racemization of 5-substituted hydantoins in aqueous solution, Cardiff University, **2011**.
- [28] H. O. Ahmad, Kinetics and Mechanism of Racemization Reactions of Configurationally Labile Stereogenic Centres in Drug-like Molecules in Aqueous Solutions; Thiohydantoins and Related Compounds., Kinetics and mechanism of racemization reactions of configurationally labile stereogenic centres in drug-like molecules in aqueous solutions; thiohydantoins and related compounds., Cardiff University, **2015**.
- [29] R. A. Lazarus, *J. Org. Chem.* **1990**, *55*, 4755–4757.
- [30] Manuscript in preparation.
- [31] A. K. Mills, A. E. W. Smith, *Helv. Chim. Acta* **1960**, *43*, 1915–1927.
- [32] J. P. Lutz, S. T. Chau, A. G. Doyle, *Chem. Sci.* **2016**, *7*, 4105–4109.
- [33] F. Wang, D. Wang, X. Wan, L. Wu, P. Chen, G. Liu, *J. Am. Chem. Soc.* **2016**, *138*, 15547–15550.
- [34] R. Y. Liu, Y. Yang, S. L. Buchwald, *Angew. Chem.* **2016**, *128*, 14283–14286.
- [35] I. V. Pavlidis, M. S. Weiß, M. Genz, P. Spurr, S. P. Hanlon, B. Wirz, H. Ilding, U. T. Bornscheuer, *Nat. Chem.* **2016**, *8*, 1076–1082.