



The University of  
**Nottingham**

UNITED KINGDOM · CHINA · MALAYSIA

Brand, Oliver J. and Pasini, Alice and Habgood, Antony and Knox, Alan J. and Jenkins, Gisli and Pang, Linhua (2016) S52 Suberanilohydroxamic acid (SAHA) inhibits collagen deposition in a transforming growth factor  $\beta$ 1-driven precision cut lung slice (PCLS) model of pulmonary fibrosis. *Thorax*, 71 (Supp 3). A31. ISSN 1468-3296

**Access from the University of Nottingham repository:**

<http://eprints.nottingham.ac.uk/47500/1/BTS%202016%20-Abstract-Oliver%20Brand%20Accepted.pdf>

**Copyright and reuse:**

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see: [http://eprints.nottingham.ac.uk/end\\_user\\_agreement.pdf](http://eprints.nottingham.ac.uk/end_user_agreement.pdf)

**A note on versions:**

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact [eprints@nottingham.ac.uk](mailto:eprints@nottingham.ac.uk)

**Suberanilohydroxamic Acid (SAHA) Inhibits Collagen deposition in a Transforming Growth Factor  $\beta_1$ -driven Precision Cut Lung Slice (PCLS) model of Pulmonary Fibrosis.**

Oliver J Brand, Alice Pasini, Anthony Habgood, Alan J Knox, Gisli Jenkins, Linhua Pang

**Introduction and Objectives:** Idiopathic Pulmonary Fibrosis (IPF) is a chronic, progressive interstitial lung disease that is refractory to current treatment options. Transforming growth factor (TGF)- $\beta_1$  is a key pro-fibrotic cytokine that plays a crucial role in IPF pathogenesis. Our group previously demonstrated distinct epigenetic modifications involved in repression of the antifibrotic gene cyclooxygenase-2 (COX-2) in fibroblasts from IPF (F-IPF) lungs compared with fibroblasts from non-fibrotic lungs (F-NL). Epigenetic drugs capable of inhibiting DNA and histone modifications may, therefore, represent a putative novel therapy. The aim of this study was to investigate the ability of 4 epigenetic inhibitors to regulate TGF- $\beta$ -driven fibrosis in *ex vivo* mouse lung.

**Methods:** A precision-cut lung slice (PCLS) model of fibrosis was established using the previously described [1] CC10-tTS-rtTA-TGF $\beta_1$  transgenic (tgTGF- $\beta_1$ ) mouse. The model was first assessed by investigating PCLS overexpression of TGF- $\beta_1$  in response to stimulation of the transgene by doxycycline treatment. Gene expression of COX-2 and fibrotic markers including collagen were assessed after 4 days of treatment. The anti-fibrotic potential of 4 epigenetic inhibitors; BIX01294 (BIX, inhibitor of G9a histone methyltransferase), 3-deazaneplanocin A (DZNep, inhibitor of EZH2 histone methyltransferase), SAHA (inhibitor of histone deacetylases, HDACs) and Decitabine (DAC, DNA demethylating agent) was investigated. Viability of PCLS was assessed by MTT and Prestoblu<sup>®</sup> viability assay.

**Results:** Treatment of PCLS from tgTGF- $\beta_1$  mice with doxycycline induced a concentration-dependent increase in global TGF- $\beta_1$ , pro-fibrotic markers including collagen and pro-inflammatory COX-2, which was comparable to recombinant TGF- $\beta_1$  treatment. Treatment with three of the epigenetic inhibitors BIX01294, DZNep and DAC did not reduce the pro-fibrotic response following doxycycline treatment. However SAHA demonstrated a significant suppressive effect on COX-2 and collagen expression, while not directly affecting TGF- $\beta_1$  transgene expression.

**Conclusions:** The data suggests that SAHA has the potential to reduce fibrosis in a TGF- $\beta_1$  driven model of pulmonary fibrosis. Further work is currently underway to assess the anti-fibrotic potential of this drug in tgTGF- $\beta_1$  animals.

## Reference

[1] Lee CG, Cho SJ, Kang MJ et al. Early Growth Response Gene 1-mediated Apoptosis Is Essential for Transforming Growth Factor  $\beta_1$ -induced Pulmonary Fibrosis. *J Exp Med.* 2004; 200(3):377-389.