



The University of
Nottingham

UNITED KINGDOM · CHINA · MALAYSIA

Fernandes, Gwen Sascha and Parekh, Sanjay M. and Moses, Jonathan and Fuller, Colin and Scammell, Brigitte E. and Batt, Mark E. and Zhang, Weiya and Doherty, Michael (2017) Prevalence of knee pain, radiographic osteoarthritis and arthroplasty in retired professional footballers compared to men in the general population: a cross-sectional study. *British Journal of Sports Medicine* . ISSN 1473-0480 (In Press)

Access from the University of Nottingham repository:

http://eprints.nottingham.ac.uk/47791/1/BJSM_FootballPaper_manuscript.September_AAM.pdf

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see: http://eprints.nottingham.ac.uk/end_user_agreement.pdf

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

Prevalence of Knee Pain, Radiographic Osteoarthritis and Arthroplasty in Retired Professional Footballers compared to men in the General Population: A Cross-Sectional Study

Gwen Sascha Fernandes, research fellow^{1,2,3}, Sanjay Parekh, doctoral student ^{1,2}, Jonathan Moses, foundation doctor^{1,2,5}, Colin Fuller, risk management consultant^{2,4}, Brigitte Scammell, professor of orthopaedics^{1,2,3,5}, Mark Edward Batt, professor of sports and exercise medicine^{1,2,5}, Weiya Zhang, professor of epidemiology^{1,2,3}, Michael Doherty, professor of rheumatology^{1,2,3}

1: Division of Rheumatology, Orthopaedics and Dermatology (School of Medicine, University of Nottingham), Academic Rheumatology, Nottingham City Hospital, NG5 1PB

2: Arthritis Research UK Centre for Sports, Exercise and Osteoarthritis, Queens Medical Centre, Derby Road, NG7 2UH

3: Arthritis Research UK Pain Centre, Nottingham City Hospital, NG5 1PB

4: Colin Fuller Consultancy Ltd, Sutton Bonnington, LE12 6PF

5: Nottingham University Hospitals NHS Trust, Queens Medical Centre, Derby Road, NG7 2UH

Correspondence to: Professor Weiya Zhang

Address: Academic Rheumatology, Clinical Sciences Building, Nottingham City Hospital, University of Nottingham, NG5 1PB.

Tel: 01158231756

Email: weiya.zhang@nottingham.ac.uk

Manuscript Word Count: 3625

What are the new findings?

- This is the first large scale cross-sectional study to compare outcomes of knee pain, radiographic osteoarthritis and requirement for knee arthroplasty in ex-footballers to men in the general population. The prevalence of all knee outcomes was almost 2-3 times higher in ex-footballers even after adjustment for known risk factors including significant knee injury.
- The prevalence of knee pain by age reached its peak 10-15 years earlier in ex-footballers compared to men in the general population.
- Knee injury, a high body mass index and a high risk occupation (post-retirement from football) are the main attributable risk factors for the increases in knee pain, radiographic osteoarthritis and knee replacement in ex-professional footballers.

Abstract

Objectives: To determine the prevalence of knee pain, radiographic knee osteoarthritis (RKO), total knee replacement (TKR) and associated risk factors in male ex-professional footballers compared to men in the general population (comparison group).

Methods: 1207 male ex-footballers and 4085 men in the general population in the UK were assessed by postal questionnaire. Current knee pain was defined as pain in or around the knees on most days of the previous month. Presence and severity of RKO were assessed on standardised radiographs using the Nottingham Line Drawing Atlas (NLDA) in a sub-sample of 470 ex-footballers and 491 men in the comparison group. The adjusted risk ratio (aRR) and risk difference (aRD) with 95% confidence interval (CI) in ex-footballers compared to the general population were calculated using the marginal model in Stata.

Results: Ex-footballers were more likely than the comparison group to have current knee pain [aRR 1.91, 95%CI 1.77-2.06], RKO [aRR 2.21, 95%CI 1.92-2.54] and TKR (aRR 3.61, 95%CI 2.90–4.50). Ex-footballers were also more likely to present with chondrocalcinosis [aRR 3.41, 95%CI 2.44-4.77]. Prevalence of knee pain and RKO were higher in ex-footballers at all ages. However, even after adjustment for significant knee injury and other risk factors, there was more than a doubling of risk of these outcomes in footballers.

Conclusions: The prevalence of all knee osteoarthritis outcomes (knee pain, RKO and TKR) were 2-3 times higher in male ex-footballers compared to men in the general population group. Knee injury is the main attributable risk factor. Even after adjustment for recognised risk factors, knee osteoarthritis appear to be an occupational hazard of professional football.

Abstract Word Count: 265 words

Keywords:

Knee osteoarthritis, professional football, epidemiology, knee radiographs, knee pain, risk factors, and chondrocalcinosis

Introduction

Football is the world's most popular team sport. Worldwide over 265 million people are estimated to play football and of these 110,000 are professional male footballers¹. The average career of a professional footballer lasts 13.5 years and, despite typically being extremely fit, engaging in high intensity match-play and training can result in sport-related health risks². Professional football has a high injury rate and 17% of all injuries involve the knee³. Apart from overt acute injury, the cumulative effect of repetitive micro-trauma and joint overloading could also prove deleterious to the knee joint, as is recognised in other physically demanding occupations such as coal mining⁴.

Knee osteoarthritis (KOA) is a common complex disorder with multiple risk factors including injury². Although there have been long-standing concerns over the risk of KOA in ex-footballers, only six studies have examined this association⁵⁻¹⁰. However, these studies used different definitions of KOA and often focused on just one or two outcomes (e.g., knee symptoms, structural KOA on imaging, self-reported diagnosis of KOA). Most studies were small, four had no comparison groups and adjustment for other known risk factors for KOA was absent or limited². Indeed, a recent systematic review cited the same limitations and concluded that only 'very low quality' evidence suggests football increases the risk of KOA¹¹. No studies have used a general population group for comparison, so whether ex-footballers have more KOA than the general population remains unknown.

We hypothesise that, when recognised risk factors are adjusted for, several outcomes relevant to KOA, including knee pain, radiographic KOA (RKOA), radiographic chondrocalcinosis (CC) (this co-associates with KOA and has been associated with

prior joint trauma¹²), physician-diagnosed KOA, and knee arthroplasty (a surrogate for clinically severe KOA) are more prevalent in ex-footballers compared to the general population. Therefore we undertook the following cross-sectional study comparing male ex-footballers to men in the general population.

The objectives of this study were:

1. To determine the prevalence of KOA outcomes (specifically knee pain, RKOA, CC, requirement for knee replacement surgery (TKR)) and risk factors for knee OA in a sample of retired male professional footballers in the UK and in a random sample of men in the general population.
2. To compare the prevalence of these outcomes in ex-footballers and the general population with adjustment for other known risk factors.
3. To determine the main attributable risks for any increased prevalence in knee OA outcomes in ex-footballers.

Methods

The study was approved by the Nottingham Research Ethics Committee 1 (Refs 14/EM/0045 & 14/EM/0015). A cross-sectional design was used, involving postal questionnaire surveys to ex-footballers and to a sample of men in the general population to gain information on knee pain, knee surgery and KOA risk factors as well as simple demographics, occupational history, general health and current medications. Participants who indicated interest in attending for knee radiographs, irrespective of knee pain status and who provided written informed consent, were invited to undergo bilateral knee radiographs to determine RKOA and CC.

Participants

Ex-footballers were recruited via the Professional Footballers' Association (PFA) and former players' associations (n=21 professional clubs). Inclusion criteria for ex-footballers were men over age 40 who had played professionally (in the top four tiers of the English Football League). The comparison group were recruited from the Knee Pain and Related Health in the Community Study (KPIC), involving recruitment via 12 general practitioner/family medicine (GP) practices in the UK Midlands region. All men on these UK National Health Service GP registers aged 40 years and older who were not terminally ill, were able to give written informed consent and had no other reason judged by the GPs to exclude them from the study were sent the questionnaire.

Questionnaire Survey

The postal questionnaire was developed based on previously published questionnaires^{13,14}. Through public and patient involvement, two pilot versions were evaluated to identify any problems with content, language and layout.

The 44,775 questionnaires were similarly constructed to capture detailed information about the participant, their medical history and putative risk factors for KOA⁵. A validated screening question was used to determine presence of current knee pain: "Have you ever had knee pain for most days of the past one month?"¹⁴. Additionally, a body pain mannequin¹⁵ was used to locate pain in other body regions. There were specific enquiries about comorbidities, current medications, and any past knee surgery including TKR. Constitutional knee alignment (in early 20's), current knee alignment, and the index to ring finger length ratio (2D4D) were assessed using validated line-drawings^{16, 17}. These drawings, which illustrated the direction and severity of each alignment grade, allowed participants to choose their knee alignments grades separately for early adult life and for current alignment. The grades being: A = severe varus, B=mild varus, C=straight legs, D = mild valgus, and E= severe valgus. The 2D4D ratio can be visually classified as pattern 1 (index longer than ring finger), pattern 2 (index length equal to ring finger), or pattern 3 (index shorter than ring finger) and it is pattern 3 that has been associated with RKOA¹⁷. Nodal OA was determined using a validated diagram¹⁸ and classified as present in those reporting nodes on at least two rays of both hands¹⁹. Significant knee injury was defined as '*one which caused pain for most days for at least a three month period and resulted in an absence from all training and matches during this time*'. Occupations were classed as 'high risk for KOA' based on published evidence²⁰. Each listed occupation per individual was

analysed and the data dichotomised into high- or low-risk groups (excluding professional football careers).

Radiographic Knee Osteoarthritis and Chondrocalcinosis

All ex-footballers who returned a questionnaire indicating willingness to have knee radiographs, and who lived within 40 miles of a Spire Healthcare Hospital were invited to attend. Similarly, all general population men in KPIC who had indicated interest were invited to attend the Nottingham University Hospitals Radiology Department for bilateral knee radiographs. Participants who had undergone bilateral TKR were excluded from radiographic assessment. All radiographs (weight-bearing semi-flexed posterior-anterior and 30° flexion skyline views) were performed using published protocols^{16,21}. Radiographs were scored by one observer (GSF), blinded to knee pain status, as a single mixed batch using HIPAX Dicom software. The Nottingham Line Drawing Atlas (NLDA) is a logically derived, interval (not ordinal) based atlas and was used to score individual compartment and composite joint space narrowing (JSN), individual compartment and subsequent composite osteophyte, and combined (global JSN and osteophyte) scores for each knee²¹. The NLDA uniquely provides separate illustrations for JSN for men and women to account for the normally wider joint space width in men and gives 0-5 interval scores for osteophyte at 8 sites in the three compartments with consideration for natural variations in osteophyte shape. Separate line drawings for joint space width and osteophyte removes the distraction of combined features and replaces the ordinal grading inherent in photographic atlases. The Kellgren and Lawrence (KL) composite grade [0-4] was also scored for the combined tibiofemoral compartments (medial and lateral) and separately for combined

patellofemoral compartments (medial and lateral) using verbal descriptors for features of OA (JSN and/or osteophyte presence) as opposed to a photographic atlas..

Definite dichotomised RKOA using the NLDA was defined as definite JSN (score ≥ 2) and definite osteophyte (score ≥ 2) in any compartment. CC in either hyaline or fibrocartilage was defined as present or absent. RKOA using KL was defined as grade ≥ 3 (definite osteophyte and definite narrowing) and additionally, grade ≥ 2 (definite osteophyte and possible narrowing) in any compartment. Intra-observer agreement (GSF) and inter-observer agreement (GSF and AS) were assessed for NLDA and KL grades using kappa coefficients.

Statistical Analyses

Based on a 15% prevalence of RKOA in the general population²², the sample size needed was calculated using the z test and a logistic regression model with an a priori type power analysis. We assumed our data would include multiple confounding factors with a multiple correlation coefficient of 0.3 between them. With a power of 80% at 0.05, one tail (assuming footballers would have greater risk of knee OA than the general population) and a relative risk (RR) of 2.0, the sample size required was 424 participants per group. The power calculations were done using GPower, version 3.1.9.2.

Categorical variables were reported as frequencies and percentages and continuous variables as means and standard deviations (SD). To determine whether distributions of the variables were statistically significantly different between ex-footballers and the general population, a t-test (continuous variables) or chi-square test (categorical variables) was used. Statistical significance was defined as $p < 0.05$.

The relative risk or rate ratio (RR), i.e., the ratio of the prevalence of knee OA outcomes between the ex-footballers versus that in the general population was calculated. A multivariable logistic regression model was used to adjust for other confounding factors such as age, body mass index (BMI) and injury. This was followed by the `adjrr` STATA command²³. RR was determined instead of odds ratios (ORs) as the outcomes of interest, such as KP and RKOA are not rare and therefore the use of ORs would inflate the estimate. The primary purpose of this analysis was to confirm whether playing professional football is a risk factor independent from other potential knee OA risk factors. We therefore ran four analyses to follow this up. Firstly, we

calculated crude RR for playing professional football without any adjustment. Secondly, we estimated the RR adjusted for two common confounders for OA, age and BMI (model 1). Thirdly, we brought injury into the model as it is a major risk factor of OA for anyone in the general population to examine whether footballing per se without a major injury is a risk factor for knee OA (model 2). Finally we included all putative risk factors/confounders previously established in the literature and collected in the study (age, body mass index, nodal OA, 2D4D ratio, alignment, injury, high risk occupation and comorbidities)²⁴. We also calculated the risk difference (RD) as an absolute measure of association according to the STROBE²⁵. Whilst the RR is the ratio of knee OA outcomes between the ex-footballers and controls, the RD is the difference between the prevalence of knee OA outcomes between these groups. The RD was calculated using a logistic regression model for each knee OA outcome (dichotomous) between the ex-footballers and general population followed by the same `adjrr` STATA command. The RR and RD in cross sectional studies have also been termed as prevalence proportion ratio (PPR) and prevalence proportion difference (PPD) respectively to differ from those measured in cohort studies where the incidence rather than prevalence is used²⁶. However, despite the different terminologies, the calculations are identical, i.e., $RR \text{ or } PPR = r1/r2$, whereas $RD \text{ or } PPD = r1 - r2$, where $r1$ is the risk (prevalence in cross sectional study or incidence in cohort study) in the exposure group (e.g., ex-footballers) and $r2$ is the risk in the non-exposure group (e.g., general population control). We therefore opted to use the RR and RD for this paper as they are more commonly used terms for relative risk and attributable risk and understood by the majority of clinicians. We had very few missing

data at random (for example, where BMI was not reported by a participant). Imputation or modelling was therefore not undertaken for the occasional missing values.

All analysis was conducted using Stata IC Version 14 on Windows 7 Operating System and power calculations undertaken using Power and Precision Version 2.1.

Results

Of 4775 questionnaires sent to ex-footballers, 1207 responses (25%) were received. Of 40,000 questionnaires sent to men and women in the community (KPIC), 9517 completed questionnaires (24%) were returned, including 4085 men (42.9%) (Fig.1).

Figure 1. Selection of the Ex-Footballer and the general population groups.

The mean age of ex-footballers was 59 years (± 11.7), 3.9 years younger than the comparison group (62.9 years, ± 10.4), but mean BMI were comparable (Table 1). Ex-footballers had a higher prevalence of nodal OA, 2D:4D pattern 3 and body pain (all $p < 0.01$) and were more likely to report constitutionally malaligned knees (varus or valgus) compared to men in the general population. Ex-footballers had fewer comorbidities such as diabetes and cancer ($p < 0.01$), (Table 1). Details on missing data for each variable has been provided in Appendix 1.

Table 1. Demographics of the Footballer and General Population groups.

	Footballers	General Population	p-value
Questionnaires, n	1207	4085	
Age (years), mean (SD)	59.0 (11.7)	62.9 (10.4)	<0.001**
BMI (kg/m²), mean (SD)	27.3 (3.2)	27.5 (4.7)	0.139
Right Handed, n (%)	1057 (87.6)	3474 (85.0)	0.529
Right Lower Limb Dominance, n (%)	1000 (82.85)	N/A	N/A
Pattern 3 Digit Ratio, n (%)	733 (60.7)	2237 (54.8)	0.003**
Nodal OA, n (%)	86 (7.1)	218 (5.6)	<0.001**
Knee Injury, n (%)	778 (64.5)	953 (23.3)	<0.001**
High Risk Occupation, n (%)	742 (61.5)	2185 (53.5)	<0.001**
Malalignment, n (%) †			
Constitutional	193 (16.0)	278 (6.8)	<0.001**
Current	289 (24.6)	434 (11.2)	<0.001**
Proportion with change in alignment since 20's, (%)			
• Varus	88 (8.9)	132(3.7)	<0.001**
• Valgus	32 (3.2)	42(1.2)	
Body Pain, n (%) ‡	901 (74.7)	2574 (69.8)	0.001**
Painkillers, n (%) #	423 (35.0)	1289 (31.6)	0.02*
Comorbidities, n (%) ^	355 (29.4)	1868 (45.7)	<0.001**

† Malalignment: right knee varus or valgus deformities (symmetry assumed);

‡ Pain reported in any region of the body for most days of the past month;

Painkillers include non-steroidal anti-inflammatories, opioids, over the counter medications and auxiliary medications that have a pain relieving effect
^Co-morbidities: diabetes / hypertension / myocardial infarction / cancer / fibromyalgia
P<0.05 denoted as * and p<0.01 denoted as **

Self-reported Knee Osteoarthritis outcomes

Overall prevalence of knee pain in ex-footballers was 52.2% compared to 26.9% in the general population ($p < 0.01$). Across all age groups, ex-footballers had more current knee pain than the general population (Fig.2a), especially in younger age groups (45-54 years). There was no effect of laterality in knee pain prevalence in the right, left and both knees (Appendix 2).

Figure 2a. Prevalence of current knee pain by age categories in the footballer and general population

After adjusting for age and BMI, the aRR of current knee pain in ex-footballers was 1.91 (95% CI 1.77 – 2.06) compared to the general population. A higher proportion of ex-footballers (28.3%) than men in the general population (12.2%) had received a diagnosis of KOA from a physician (aRR 3.73 (95% CI 3.33 -4.17). Furthermore, 11.1% of ex-footballers reported TKR's compared with 3.8% of the general population ($p < 0.01$), giving an aRR of 3.61 (95% CI: 2.90-4.50). The prevalence of TKR by age categories is presented in Fig.2b.

Radiographic findings:

Agreement between observers (AS and GSF) examining 21 participants (40 knees) on two occasions using the NLDA was substantial (kappa 0.78), and intra-observer agreement kappas were 1.00 for each observer.

RKOA in any knee using NLDA scoring was present in 64% of footballers and 35.2% of the general population. Ex-footballers had significantly more RKOA in their right, left and both knees (27.2%, 34.6% and 14.4%) compared to the general population (12.2, 10.1%, and 5.6%) (Appendix 3). 15.7% of ex-footballers had RKOA in their right knee compared to 4.6% in the general population. The prevalence of RKOA by ages is presented in Fig.2c.

After adjusting for age and BMI, the aRR of RKOA increased from 1.82 to over 2 times more likely (RR 2.21, 95% CI: 1.92-2.54) than in the general population. The prevalence of RKOA using KL grading (≥ 3) in any compartment of either knee, was also higher in ex-footballers with an aRR of 2.46 (95%CI: 1.89-3.22) (Table 2).

1 **Table 2. Prevalence of Knee Osteoarthritis and its related outcomes in ex-footballers compared to the general population**

	Prevalence, n (%)		Relative Risk (95% Confidence Interval)			
	Footballers	General Population	Crude	Model 1	Model 2	Model 3
<u>Self-Reported Outcomes</u>						
Current Knee Pain	630 (52.2)	1100 (26.9)	1.94 (1.80 – 2.09)	1.91 (1.77 – 2.06)	1.48 (1.38 – 1.63)	1.50 (1.28 – 1.76)
Physician-Diagnosed Knee OA	341 (28.3)	500 (12.2)	3.53 (3.15 – 3.96)	3.73 (3.33-4.17)	2.69 (2.36 – 3.07)	2.18 (1.73 – 2.77)
Total Knee Replacement	134 (11.1)	157 (3.8)	2.88 (2.31 – 3.60)	3.61 (2.90 – 4.50)	2.33 (1.84 – 2.95)	2.10 (1.42 – 3.14)
<u>Radiographic Outcomes</u>						
Nottingham Line Drawing Atlas (≥ 2 osteophyte and ≥ 2 JSN)	301 (64.0)	173 (35.2)	1.82 (1.58 – 2.08)	2.21 (1.92-2.54)	1.91 (1.65 – 2.22)	1.92 (1.66 – 2.23)
Kellgren Lawrence (Grade ≥ 3)	134 (28.5)	69 (14.1)	2.02 (1.56 – 2.63)	2.46 (1.89 – 3.22)	2.10 (1.58 – 2.80)	2.08 (1.56-2.79)
Kellgren Lawrence (Grade ≥ 2)	257 (54.7)	154 (31.4)	2.06 (1.69-2.53)	2.46 (2.11 – 3.02)	1.99 (1.60 – 2.49)	1.97 (1.58 – 2.46)
Chondrocalcinosis (CC)	114 (24.3)	43 (8.8)	2.77 (1.99 – 3.84)	3.41 (2.44 – 4.77)	2.63 (1.84 – 3.75)	2.57 (1.80 – 3.66)

2 1 RR adjusted for football-status, age, BMI
3 2 RR adjusted for football-status, age, BMI and injury
4 3 RR adjusted for football-status, age, BMI, nodal OA, injury, constitutional alignment, high-risk occupation, 2D:4D ratio, nodal OA, comorbidities

5 Ex-footballers had more RKOA in their left compared to their right knee. They also
6 showed most JSN in the PF compartment whereas the general population had most
7 JSN in the TF compartment (Appendix 3). Ex-footballers also had more RKOA in their
8 right and left TF compartment (15.7% and 11.6%) compared to the general population
9 (4.6% and 5%) (Appendix 4).

10

11 Ex-footballers were more likely than the general population to have CC (24.3% versus
12 8.8%) with an aRR of 3.41 (95%CI: 2.44 – 4.77) (Table 2), and particularly had more
13 CC in the left knee (Appendix 5).

14

15 Table 3 presents the crude and adjusted risk difference (RD) for each knee OA
16 outcome in the ex-footballers compared to the controls. The absolute measure of
17 association for knee pain suggests that even after adjustment for age and bmi (Model
18 1), 24.46% (21.30% - 27.62%) more ex-footballers would present with knee pain
19 compared to the general population. Similarly, even after adjustment for age and bmi,
20 37.49% (31.82% -43.16%) more ex-footballers would have RKOA using the NLDA
21 compared to the general population.

22

23

24

25

26

27

28

29

30

31 **Table 3. Risk Differences in Knee Osteoarthritis and its related outcomes in ex-footballers compared to the general**
 32 **population**

33
 34
 35
 36

Knee OA Outcomes	Risk Differences % (95 % CI)			
	Crude	Model 1	Model 2	Model 3
Current Knee Pain	25.27 (22.14 – 28.40)	24.46 (21.30 – 27.62)	14.59 (11.32 – 17.87)	14.50 (7.97 – 21.03)
Total Knee Replacement	7.26 (5.39 – 9.13)	9.53 (7.44 – 11.63)	5.40 (3.63 – 7.17)	4.52 (1.39 – 7.65)
Nottingham Line Drawing Atlas (≥ 2 osteophyte and ≥ 2 JSN)	28.81 (22.75 – 34.86)	37.49 (31.82 – 43.16)	31.11 (24.61 – 37.60)	31.45 (24.87 – 38.03)
Kellgren Lawrence (Grade ≥ 3)	14.46 (9.35 – 19.57)	18.77 (13.46 – 24.08)	15.33 (9.56 – 21.09)	15.02 (9.17 – 20.88)
Kellgren Lawrence (Grade ≥ 2)	21.98 (16.26 – 27.71)	27.61 (21.82 – 33.39)	21.21 (14.79 – 27.63)	20.75 (14.22 – 27.28)
Chondrocalcinosis	15.50 (10.89 – 20.11)	18.91 (14.04 – 23.77)	14.69 (9.56 – 19.82)	14.30 (9.18 – 19.41)

37 1 RD adjusted for football-status, age, BMI
 38 2 RD adjusted for football-status, age, BMI and injury
 39 3 RD adjusted for football-status, age, BMI, nodal OA, injury, constitutional alignment, high-risk occupation, 2D:4D ratio, nodal OA, comorbidities

40

Risk Factors

Apart from being professional footballers, knee injury, BMI, other high risk occupation, constitutional knee malalignment and 2D4D finger ratio were also significantly associated with knee pain, RKOA and TKR (Appendix 6).

Discussion

This is the first study comparing the risk of KOA in ex-footballers and the general population. The main findings in footballers are: firstly, a near two-fold increased risk of current knee pain (aRR 1.91, 95% CI 1.77-2.06) which was most marked in younger age groups; secondly, a two-fold increased prevalence of RKOA (aRR 2.21, 95%CI 1.92-2.54) at all age groups and a three-fold increased prevalence of radiographic CC (aRR 3.41, 95%CI 2.44-4.77; and thirdly, almost a three times higher requirement for TKR (aRR 3.61, 95%CI 2.90 – 4.50). The major attributable risk factors are knee injury (aRR 1.89 for knee pain, 1.44 for RKOA and 3.32 for TKR), body mass index (aRR 1.49 for knee pain and 2.24 for TKR) and nodal OA (aRR 1.46 for knee pain and 1.93 for TKR). However, even after adjusting for injury and other risk factors there is still over a two-fold increased risk of KOA outcomes which supports an important role for repetitive micro-trauma associated with playing football.

The knee pain prevalence in the general population group accords with previous general population surveys^{14,27} and that in ex-footballers also is similar to previous reports². Ex-footballers were three times more likely to report a physician-diagnosis of KOA with a prevalence (28.3%) in line with previous questionnaire surveys^{5,6}. In this study, ex-footballers had three times more RKOA and specifically, JSN, in their right TF articulation (15.7%) than the general population (4.6%), which is similar to one study⁹, but lower than others^{7,8}. This discrepancy may arise from smaller population samples in previous studies, selection bias from recruiting from a single club, and radiographic cut-offs using 'possible' JSN. When using this cut-off (KL ≥ 2), our results remained unchanged with a two-fold increased risk of RKOA. Our preferred RKOA definition requiring definite osteophyte and definite narrowing (KL ≥ 3) accords with

pathological definitions of KOA that require both focal loss of hyaline cartilage and bone hypertrophy²⁸. Furthermore we used the interval scale NLDA to score narrowing and osteophyte separately, which has several advantages over more commonly used ordinal scale photographic atlases and the composite KL grading system, particularly accounting for wider joint space widths in men and assessing the entire joint (all three compartments) with a global score²¹. In addition to this, the NLDA was constructed to score changes in the TF and PF compartments of the knee whereas KL grading has only subsequently been adapted for use in the PF compartment (using the same verbal descriptors as for the combined TF compartments) and was not specifically designed for this purpose originally. Nevertheless, despite these caveats we included the KL grading to permit comparison with other studies that used KL grading to assess radiographic OA severity. The prevalence of CC was significantly higher in ex-footballers (24.3%) compared to the general population (8.8%) and previous population estimates (7-10%)²⁹. Both KOA and prior joint insult are recognized risk factors for CC¹¹, so this increased prevalence is consistent with more biomechanical knee trauma in footballers. Furthermore concurrence of CC with KOA may associate with greater clinical severity and worse outcomes¹². Interestingly, whilst our general population had more right knee RKOA, ex-footballers had significantly more left knee RKOA, despite the majority indicating right limb dominance. As most football injuries are non-body contact in nature³, this finding might be attributed to playing technique where players cannot respond quickly enough to rapid, unpredictable movements and where rotational strain on the weight-bearing, non-dominant/kicking limb may cause damage⁸. Additionally, in ex-footballers JSN particularly targeted the PF compartment which is an important component of the knee extensor mechanism that could be stressed by kicking with a partially flexed knee, by constitutional malalignment (which

was increased in footballers (Table 1)) or by torsional movements that influence patellar tracking, all of which may initiate PF degeneration³⁰.

Ex-footballers reported more knee pain at all ages, and this was particularly marked in younger age groups (Figure 2a). Having adjusted for all significant risk factors, including injury, ex-footballers still showed increased risks of knee pain, RKOA and TKR suggesting lasting damage from repetitive microtrauma sustained over the course of their footballing career. This is the first nationwide study to examine structural and person-centred outcomes relating to KOA in ex-footballers. The reported degree of increased adjusted risk (at least doubled) is in the order required by many national bodies to recognise KOA as an industrial disease for professional football. Importantly, the study also identifies risk factors for KOA in footballers and in particular, key modifiable risk factors, such as knee injury, being overweight/obese and undertaking high risk occupations after retiring from football (Appendix 6).

Caveats

There are several limitations to this study. Firstly, the use of self-reported questionnaires might involve recall bias and possible misclassification of self-reported outcomes³¹. A structured clinical enquiry and assessment of every participant would have been preferable but was impractical due to logistics (UK nationwide survey) and limited resources. Nevertheless, we used validated instruments designed for questionnaire use and involved ex-footballers, patients and general public volunteers to help optimize clarity and ease of use of the questionnaire. Secondly, the suboptimal response rate to the questionnaire and subsequently the ex-footballers and general population sub-samples indicating willingness to have knee x-rays, could have

resulted in selection bias. Unfortunately we could not compare simple demographics or health records of responders and non-responders in either group due to data protection of externally held databases and patient confidentiality in the general population group. However, although knee pain positive individuals might be more likely to respond, this bias should be present in both groups. Furthermore, this is a large cross-sectional study and given that the prevalence of knee pain, RKOA, CC and risk factors in the general population generally aligned with results from other population-based studies, it seems unlikely that the estimates are unduly biased. Thirdly, although we surveyed ex-footballers over a wide area of the UK, for logistical reasons the general population was recruited from just one region. Nevertheless, the East Midlands was a major coal-mining region, a male occupation with an established risk of RKOA⁴. Thus, any regional bias in the general population would have been expected to inflate rather than lower the prevalence of KOA, making the demonstrated increased risk in ex-footballers even more confident. Fourthly, although radiographs are widely used to assess structural KOA in population-based studies, they are insensitive to soft tissue and other changes indicative of early KOA that can be identified by alternative imaging techniques such as MRI³². Therefore, we are likely to have underestimated the true prevalence of structural OA in both ex-footballers and the general population. However, use of MRIs in this study was not financially or logistically viable. Another caveat is that whilst the authors were able to include known risk factors of KOA such as age, BMI and injury, there are still unmeasured confounders such as physical activity or unknown confounders that might contribute to the increased KOA outcomes in ex-footballers compared to the general population. In summary, the prevalence of knee pain and RKOA in ex-footballers is twice as high as in men in the general population and ex-professional players develop knee pain

earlier and require three times more knee joint replacements. This study took into account other risk factors for knee OA and demonstrates that professional football is a significant risk factor for the development of the symptoms (knee pain), structural change (RKOA, CC) and requirement for TKR. Knee injury is a major attributable risk factor for each of these outcomes. These findings have important consequences for football associations/unions and stakeholders for whom the health of retired and current professional footballers is paramount.

Acknowledgements and Affiliations:

Acknowledgements

The study was funded by the Arthritis Research UK Centre for Sport, Exercise and Osteoarthritis (Grant 20194). The study was also funded by FIFA's Medical Assessment and Research Centre as well as the Professional Footballers' Association.

We are grateful to the Arthritis Research UK Pain Centre in Nottingham for access to KPIC participants. We would like to acknowledge the invaluable assistance and advice from Dr Charlotte Cowie (Football Association and SPIRE), Richard Jobson, John Bramhall and Ben Simpson in the administration of players' database through the support of the Professional Football Association. We would also like to acknowledge our research assistants, Laura Schimpf, Emily Cooper and Dr Chijioke Nwankor for their work on the occupation data as well as Mrs. Sally Doherty (SD) and Dr Aliya Sarmanova (AS) for their help with x-ray scoring and validation work.

Affiliations

None of the study sponsors/funders were involved in the study design, in the collection, analysis and interpretation of data, in the writing of the report and in the decision to submit the article for publication.

All researchers were independent from funders.

All authors had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Acronyms:

2D4D: 2nd Digit 4th Digit (finger ratio)

AS: Aliya Sarmanova

BMI: Body Mass Index

BS: Brigitte Scammell

CC: Chondrocalcinosis

CF: Colin Fuller

CI: Confidence Interval

GLM: Generalised Linear Modelling

GSF: Gwen Sascha Fernandes

IQR: Inter Quartile Range

JPM: Jonathan Philip Moses

KL: Kellgren Lawrence

KOA: Knee Osteoarthritis

MD: Michael Doherty

MEB: Mark Edward Batt

NLDA: Nottingham Line Drawing Atlas

Pe: Percentage of Exposure

PF: Patellofemoral

RKOA: Radiographic Knee Osteoarthritis

RD: Risk Difference

RR: Relative Risk

SD: Sally Doherty

SMP: Sanjay Mukundh Parekh

TF: Tibiofemoral

TKR: Total Knee Replacement

WZ: Weiya Zhang

References:

1. FIFA Big Count 2006. FIFA Facts, 2006. http://www.fifa.com/mm/document/fifafacts/bcoffsurv/bigcount.statspackage_7024.pdf. Accessed 10th October, 2016.
2. Kuijt MT, Inklaar H, Gouttebauge V, Frings-Dresen MH. Knee and ankle osteoarthritis in former elite soccer players: a systematic review of the recent literature. *J Sci Med Sport* 2012;15:480-7.
3. Ekstrand J, Hägglund M, Waldén M. Injury incidence and injury patterns in professional football: the UEFA injury study. *British Journal of Sports Medicine* 2011;45:553-8.
4. McMillan G, Nichols L. Osteoarthritis and meniscus disorders of the knee as occupational diseases of miners. *Occup Environ Med* 2005;62:567-75.
5. Drawer S, Fuller CW. Propensity for Osteoarthritis and Lower Limb Joint Pain in Retired Professional Soccer Players. *British Journal of Sports Medicine* 2001;402–408.
6. Turner AP, Barlow JH, Heathcote-Elliott C. Long term health impact of playing professional football in the United Kingdom. *British Journal of Sports Medicine* 2000;34:332-6.
7. Elleuch MH, Guermazi M, Mezghanni M, Ghroubi S, Fji H, Mefteh S, Baklouti S, et al. Knee osteoarthritis in 50 former top-level soccer players: a comparative study. *Ann Readapt Med Phys* 2008;51:174-8.
8. Krajnc Z, Vogrin M, Recnik G, Crnjac A, Drobic M, Antolic V. Increased risk of knee injuries and osteoarthritis in the non-dominant leg of former professional football players. *Wien Klin Wochenschr* 2010;2:40-3.
9. Roos H, Lindberg H, Gardsell P, Lohmander LS, Wingstrand H. The prevalence of gonarthrosis and its relation to meniscectomy in former soccer players. *Am J Sports Med* 1994;22:219-22.
10. Arliani, G. G., Astur, D. C., Yamada, R. K. F., Yamada, A. F., Miyashita, G. K., Mandelbaum, B., & Cohen, M. Early osteoarthritis and reduced quality of life after retirement in former professional soccer players. *Clinics* 2014;69:589–594.
11. Tran G, Smith TO, Grice A, Kingsbury SR, McCrory P, Conaghan PG. Does sports participation (including level of performance and previous injury) increase risk of osteoarthritis? A systematic review and meta-analysis. *British Journal of Sports Medicine* 2016;50(23):1459-66.
12. Ledingham J, Regan M, Jones A, Doherty M. Radiographic patterns and associations of osteoarthritis of the knee in patients referred to hospital. *Annals of the Rheumatic Diseases* 1993;52(7):520-26.
13. O'Reilly SC, Muir K, and Doherty M. Screening for Pain in Knee Osteoarthritis: Which Question? *Annals of the Rheumatic Diseases* 1996;55 12:931–933.
14. Ingham SL, Zhang W, Doherty SA, McWilliams DF, Muir KR, Doherty M. Incident knee pain in the Nottingham community: a 12-year retrospective cohort study. *Osteoarthritis and Cartilage* 2011;19:847-52.
15. Hunt IM, Silman AJ, Benjamin S, McBeth J, Macfarlane GJ. The prevalence and associated features of chronic widespread pain in the community using the 'Manchester' definition of chronic widespread pain. *Rheumatology* 1999;38:275-9.
16. Ingham SL, Moody A, Abhishek A, Doherty SA, Zhang W, Doherty M. Development and validation of self-reported line drawings for assessment of

- knee malalignment and foot rotation: a cross-sectional comparative study. *BMC Medical Research Methodology* 2010;10:1-6.
17. Zhang W, Robertson J, Doherty S, Liu JJ, Maciewicz RA, Muir KR, et al. Index to ring finger length ratio and the risk of osteoarthritis. *Arthritis Rheum* 2008;58:137-44.
 18. Rees F, Doherty S, Hui M, Maciewicz R, Muir K, Zhang W, et al. Distribution of finger nodes and their association with underlying radiographic features of osteoarthritis. *Arthritis Care & Research* 2012;64:533-8.
 19. Zhang W, Doherty M, Leeb BF, Alekseeva L, Arden N, Bijlsma JW, et al. EULAR evidence-based recommendations for the diagnosis of hand osteoarthritis: report of a task force of ESCISIT. *Ann Rheum Dis* 2009;68:8-17.
 20. Palmer, Keith T. Occupational Activities and Osteoarthritis of the Knee. *British Medical Bulletin* 2012;102:147–170.
 21. Nagaosa, Y. Mateus M, Hassan B, Lanyon P, Doherty M, Development of a Logically Devised Line Drawing Atlas for Grading of Knee Osteoarthritis. *Annals of the Rheumatic Diseases* 2000;59:587–595.
 22. Lawrence, JS, Bremner, JM, Bier, F. Osteoarthrosis. Prevalence in the population and relationship between symptoms and x-ray changes. *Annals of Rheumatic Diseases* 1966;25(1):1-24.
 23. Norton EC, Miller MM, Kleinman, LC. Computing risk ratios from data with complex survey design. *The Stata Journal*. 2013: 13(3):492-509.
 24. Silverwood, V.; Blagojevic-Bucknall, M.; Jinks, C.; Jordan, J.L.; Protheroe, J.; Jordan, K.P. Current evidence on risk factors for knee osteoarthritis in older adults: A systematic review and meta-analysis. *Osteoarthr. Cartil.* 2014;23(4):507-515.
 25. Moher D, Schulz KF, Simera I, Altman DG .Guidance for Developers of Health Research Reporting Guidelines. 2010:PLOS Medicine 7(2): e1000217
 26. Nielsen RO, Bertelsen ML, Verhagen E, et al. When is a study result important for athletes, clinicians and team coaches/staff? *British Journal of Sports Medicine* 2017 doi: 10.1136/bjsports-2017-097759
 27. Arthritis Research UK Primary Care Centre, Keele. Communication to Arthritis Research UK. Based on annual consultation prevalence figures for 2010. 2013. Available from: www.arthritisresearchuk.org [Accessed 2nd September 2016].
 28. Pritzker KPH, Gay S, Jimenez SA, Ostergaard K, Pelletier JP, Revell PA, et al. Osteoarthritis cartilage histopathology: grading and staging. *Osteoarthritis and Cartilage* 2006;14:13-29.
 29. Richette P, Bardin T. Chondrocalcinosis. *Rev Prat* 2010;60:14-9.
 30. Kim, Young-Mo, and Yong-Bum Joo. Patellofemoral Osteoarthritis. *Knee Surgery & Related Research* 2012;24(4):193-200.
 31. E Hassan. Recall Bias can be a Threat to Retrospective and Prospective Research Designs. *The Internet Journal of Epidemiology* 2005;3:2.
 32. Braun HJ, Gold GE. Diagnosis of Osteoarthritis: Imaging. *Bone* 2012;51:278-88.

Declarations:

Consent:

The study was approved by Nottingham University Hospitals NHS Trust and the Nottingham Research Ethics Committee 1 (Refs 14/EM/0045 & 14/EM/0015) and registered (clinicaltrials.gov portal: NCT02098044 & NCT02098070).

All participants offered consent by responding to the postal questionnaire survey and written informed consent prior to radiographic assessment at the SPIRE Hospitals and the Nottingham City Hospital.

Contributor's statement:

GSF designed the data collection tools, wrote the statistical plan, implemented the study design (conducted the study), conducted cleaning and analysis of the data, wrote the initial drafts of the paper and subsequently revised the paper after feedback from the team. SMP designed the data collection tools, wrote the statistical plan, implemented the study design (conducted the study), conducted cleaning and analysis of the data, and drafted and revised the paper. JPM conducted cleaning and analysis of the data, and drafted and revised the paper. CWF conceptualised the study and drafted and revised the paper. BS conceptualised the study and drafted and revised the paper. MEB monitored data collection and drafted and revised the paper. WZ conceptualised and designed the study and data collection tools, wrote the statistical plan, monitored data collection and drafted and revised the paper. MD conceptualised and designed the study and data collection tools, monitored data collection and drafted and revised the paper. He is guarantor.

Conflict of Interest Disclosures:

All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare: financial support (research grant) for the submitted work from FIFA Medical Assessment and Research Centre, other from Arthritis Research UK Centre for Sports, Exercise and Osteoarthritis (Grant Reference 20194); research grant from the Professional Footballers' Association and financial support from the SPIRE Healthcare Group at the Football Association (St George's Park); CF did paid consultancy for FIFA Medical Assessment and Research Centre, the Football Association and the Premier League in the past three years and has received personal fees from these bodies outside the remit of the submitted work; MD received research funding by AstraZeneca, Nordic Biosciences, Roche, outside the submitted work; Dr. Zhang reports grants from Arthritis Research UK, grants from Arthritis Research UK, during the conduct of the study; other from AstraZeneca, other from Daiichi Sankyo, other from Biobarica, other from Hisun, outside the submitted work; no financial relationships with any organisation that might have an interest in the submitted work in the previous three years for any other authors; no other relationships or activities that could appear to have influenced the submitted work.

Copyright / license for publication

"The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, a worldwide license to the Publishers and its licensees in perpetuity, in all forms, formats and media (whether known now or created in the future), to i) publish, reproduce, distribute, display and store the Contribution, ii) translate the Contribution into other languages, create adaptations, reprints, include within collections and create summaries, extracts and/or, abstracts of the Contribution,

iii) create any other derivative work(s) based on the Contribution, iv) to exploit all subsidiary rights in the Contribution, v) the inclusion of electronic links from the Contribution to third party material where-ever it may be located; and, vi) license any third party to do any or all of the above.”

Data Sharing: No additional data available.

Transparency Declaration: Professor Michael Doherty (MD) affirms that the manuscript is an honest, accurate, and transparent account of the study being reported and that no important aspects of the study have been omitted.

Figure 1.




Figure 2.a Prevalence of knee pain by age categories




Figure 2.b Prevalence of RKOA by age categories




Figure 2.c Prevalence of TKR by age categories

