

Elsevier Editorial System(tm) for Virology  
Manuscript Draft

Manuscript Number: VIRO-17-534R1

Title: Development and characterization of a human monoclonal antibody targeting the N-terminal region of hepatitis C virus envelope glycoprotein E1

Article Type: Regular Article

Section/Category: 8 Immunity

Keywords: Hepatitis C virus; envelope protein; antibody; entry; vaccine

Corresponding Author: Professor Philip Meuleman, Ph.D

Corresponding Author's Institution: Ghent University

First Author: Ahmed Atef Mesalam

Order of Authors: Ahmed Atef Mesalam; Isabelle Desombere; Ali Farhoudi; Freya Van Houtte; Lieven Verhoye; Jonathan Ball; Jean Dubuisson; Steven Fount; Arvind H Patel; Mats A Persson; Geert Leroux-Roels; Philip Meuleman

Abstract: Monoclonal antibodies (mAbs) targeting the hepatitis C virus (HCV) envelope have been raised mainly against envelope protein 2 (E2), while the antigenic epitopes of envelope protein 1 (E1) are not fully identified. Here we describe the detailed characterization of a human mAb, designated A6, generated from an HCV genotype 1b infected patient. ELISA results showed reactivity of mAb A6 to full-length HCV E1E2 of genotypes 1a, 1b and 2a. Epitope mapping identified a region spanning amino acids 230-239 within the N-terminal region of E1 as critical for binding. Antibody binding to this epitope was not conformation dependent. Neutralization assays showed that mAb A6 lacks neutralizing capacity and does not interfere with the activity of known neutralizing antibodies. In summary, mAb A6 is an important tool to study the structure and function of E1 within the viral envelope, a crucial step in the development of an effective prophylactic HCV vaccine.

1           **Development and characterization of a human monoclonal antibody targeting the N-**  
2           **terminal region of hepatitis C virus envelope glycoprotein E1**

3  
4 3  
5  
6 4  
7  
8  
9 5 Ahmed Atef Mesalam<sup>1,2</sup>, Isabelle Desombere<sup>1,#</sup>, Ali Farhoudi<sup>1</sup>, Freya Van Houtte<sup>1</sup>, Lieven Verhoye<sup>1</sup>,  
10  
11 6 Jonathan Ball<sup>3,4</sup>, Jean Dubuisson<sup>5</sup>, Steven K. H. Fong<sup>6</sup>, Arvind H. Patel<sup>7</sup>, Mats A. A. Persson<sup>8</sup>, Geert  
12  
13 7 Leroux-Roels<sup>1</sup>, Philip Meuleman<sup>1,\*</sup>  
14  
15  
16 8

17  
18 9 <sup>1</sup> Department of Clinical Chemistry, Microbiology and Immunology, Ghent University, Ghent, Belgium.  
19

20  
21 10 <sup>2</sup> Therapeutic Chemistry Department, National Research Centre (NRC), Dokki, Cairo, Egypt.  
22

23 11 <sup>3</sup> School of Life Sciences, The University of Nottingham, Nottingham, NG7 2RD, UK.  
24

25  
26 12 <sup>4</sup> Nottingham Digestive Diseases Centre, National Institute for Health Research (NIHR) Nottingham Biomedical  
27  
28 13 Research Centre, Nottingham University Hospitals NHS Trust and The University of Nottingham, Nottingham, NG7  
29  
30 14 2UH, UK.  
31

32 15 <sup>5</sup> Univ. Lille, CNRS, Inserm, CHU Lille, Institut Pasteur de Lille, U1019 - UMR 8204 - CIIL – Center for Infection  
33  
34 16 & Immunity of Lille, F-59000, Lille, France.  
35  
36

37 17 <sup>6</sup> Department of Pathology, Stanford University School of Medicine, Stanford, California, USA.  
38

39 18 <sup>7</sup> MRC-University of Glasgow Centre for Virus Research, University of Glasgow, Glasgow, UK.  
40

41 19 <sup>8</sup> Karolinska Institutet, Department of Clinical Neurosciences, Center for Molecular Medicine, Karolinska University  
42  
43 20 Hospital Solna, 171 76 Stockholm, Sweden.  
44  
45

46 21 <sup>#</sup> Present address: Division of Immunology, OD Communicable & Infectious Diseases, Scientific Institute of Public  
47  
48 22 Health (WIV-ISP), Brussels, Belgium.  
49  
50

51 23  
52  
53 24 <sup>\*</sup> Corresponding author: Prof. Dr. Philip Meuleman, Laboratory of Liver Infectious Diseases (LLID) - Ghent  
54  
55 25 University, UZ Gent, MRB2 (Entrance 38), De Pintelaan 185, B-9000 Gent, Belgium. Phone: +32 9 332 36 58.  
56  
57 26

58  
59 27 **E-mail addresses:**  
60  
61  
62  
63  
64  
65

- 28 Ahmed.Mesalam@ugent.be
- 1  
2 29 Isabelle.Desombere@wiv-isp.be
- 3  
4 30 Ali.Farhoudi@ugent.be
- 5  
6 31 Freya.VanHoutte@ugent.be
- 7  
8 32 Lieven.Verhoye@ugent.be
- 9  
10 33 Jonathan.Ball@nottingham.ac.uk
- 11  
12 34 Jean.Dubuisson@ibl.cnrs.fr
- 13  
14 35 sfoung@stanford.edu
- 15  
16 36 Arvind.Patel@glasgow.ac.uk
- 17  
18 37 Mats.Persson@ki.se
- 19  
20 38 Geert.LerouxRoels@ugent.be
- 21  
22 39 Philip.Meuleman@ugent.be
- 23  
24 40
- 25  
26 41 **Conflict of interest:** The authors declare no conflict of interest.

27  
28 42

29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

43 **Abstract**

1  
2 44 Monoclonal antibodies (mAbs) targeting the hepatitis C virus (HCV) envelope have been raised mainly  
3  
4 45 against envelope protein 2 (E2), while the antigenic epitopes of envelope protein 1 (E1) are not fully  
5  
6 46 identified. Here we describe the detailed characterization of a human mAb, designated A6, generated  
7  
8 47 from an HCV genotype 1b infected patient. ELISA results showed reactivity of mAb A6 to full-length  
9  
10 48 HCV E1E2 of genotypes 1a, 1b and 2a. Epitope mapping identified a region spanning amino acids 230-  
11  
12 49 239 within the N-terminal region of E1 as critical for binding. Antibody binding to this epitope was not  
13  
14 50 conformation dependent. Neutralization assays showed that mAb A6 lacks neutralizing capacity and does  
15  
16 51 not interfere with the activity of known neutralizing antibodies. In summary, mAb A6 is an important tool  
17  
18 52 to study the structure and function of E1 within the viral envelope, a crucial step in the development of an  
19  
20 53 effective prophylactic HCV vaccine.  
21  
22  
23

24 54

25  
26 55 **Keywords:** hepatitis C virus, envelope protein, antibody, entry, vaccine.  
27

28 56

29  
30 57

31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 58 1. Introduction

1  
2 59 Hepatitis C virus (HCV) is a member of the *Hepacivirus* genus within the *Flaviviridae* family. Based  
3  
4 60 on the 9.6kb-long RNA genome sequence, HCV is classified into 7 genotypes (1-7) and multiple subtypes  
5  
6 61 (a, b, c, etc.) (Bukh, 2016). More than 170 million people worldwide are estimated to be infected with  
7  
8 62 HCV (Petruzzello et al., 2016). More than 70% of individuals with acute HCV infection will become  
9  
10 63 chronically infected, which after several decades may lead to liver cirrhosis and hepatocellular carcinoma.  
11  
12 64 Despite the remarkable improvement of HCV treatment regimens using direct-acting antivirals (DAAs),  
13  
14 65 mutations occur that bring about resistance and negatively impact treatment outcome (Li and De Clercq,  
15  
16 66 2017; Pawlotsky, 2016). The risk of acquiring HCV infection is high among injection drug users due to  
17  
18 67 the high prevalence of HCV in this population. The high mutation rate of HCV, leading to continuous  
19  
20 68 release of closely related viral variants that escape the host's adaptive immune response, represents a  
21  
22 69 major hurdle for the development of a vaccine. Identification of conserved epitopes that induce protective  
23  
24 70 immunity would strongly promote vaccine design.

25  
26  
27  
28 71 Inside the host cell, viral RNA is translated into a single polyprotein precursor that is cleaved into  
29  
30 72 structural (core, envelope E1 and E2) and non-structural (P7, NS2, NS3, NS4A, NS4B, NS5A, NS5B)  
31  
32 73 proteins (Bartenschlager et al., 2011). Structural proteins are responsible for the formation of the HCV  
33  
34 74 virus particle, while the non-structural proteins are essential for viral replication, translation and assembly.  
35  
36 75 Our insight in the HCV life cycle was hampered for a long time but the development of the HCV  
37  
38 76 pseudoparticle (HCVpp) and HCV cell culture (HCVcc) systems paved the way for a more  
39  
40 77 comprehensive exploration of virus infectivity (Hsu et al., 2003; Wakita et al., 2005). Both systems are  
41  
42 78 now widely used for studying HCV entry and screening for viral inhibitors (Catanese and Dorner, 2015).  
43  
44 79 The envelope proteins E1 and E2 are highly glycosylated membrane-associated proteins consisting of an  
45  
46 80 N-terminal ectodomain and a C-terminal transmembrane domain. The E1 protein is 192 amino acids (AA)  
47  
48 81 long (from AA 192 to 383) while the E2 glycoprotein is 363 residues (AA 384 to 746) in length according  
49  
50 82 to the reference strain H77 (accession no. AF011751). E2 functions as mediator for viral entry through  
51  
52 83 interaction with host attachment factors and receptors (Dubuisson and Cosset, 2014). It is also the main  
53  
54 84 target for the host's adaptive immune system. Recently the E1 protein was also shown to be involved in  
55  
56 85 viral entry and virion assembly (Haddad et al., 2017).  
57  
58  
59  
60  
61  
62  
63  
64  
65

86 The essential role of cell-mediated immunity in HCV clearance has been extensively reported  
1  
2 87 (Abdelwahab, 2016). The early generation of neutralizing antibodies has been associated with resistance  
3  
4 88 to infection in individuals at high-risk of exposure, spontaneous clearance during acute infection and  
5  
6 89 sustained virologic response after therapy (Ndongo et al., 2010; Osburn et al., 2014; Swann et al., 2016).  
7  
8 90 Although the envelope of HCV contains multiple immunogenic epitopes, the majority of monoclonal  
9  
10 91 antibodies (mAbs) isolated from infected patients or vaccinated animals have been identified as E2-  
11  
12 92 specific (Tabll et al., 2015). The biological activity of these antibodies is diverse and varies from  
13  
14 93 neutralizing to non-neutralizing or even interfering (Wang et al., 2011a). Polyclonal antibodies from  
15  
16 94 HCV-infected patients were able to protect animal models like humanized mice and chimpanzees from  
17  
18 95 HCV challenge (Bukh et al., 2015; Meuleman et al., 2011; Vanwolleghem et al., 2008). We and others  
19  
20 96 previously reported that neutralizing antibodies targeting the E2 protein could protect from HCV infection  
21  
22 97 *in vitro* and *in vivo* (Desombere et al., 2016; Keck et al., 2016; Mesalam et al., 2016). In addition,  
23  
24 98 administration of mAb MBL-HCV1, targeting E2, delayed viral rebound following liver transplantation,  
25  
26 99 while complete protection was reported when combined with the polymerase inhibitor sofosbuvir (Chung  
27  
28 100 et al., 2013; Smith et al., 2017). These observations point towards the importance of humoral immunity  
29  
30  
31 101 and neutralizing antibodies in HCV clearance.  
32  
33

34  
35 102 Despite the availability of multiple *in vitro* systems as well as experimental animal models, little is  
36  
37 103 still known about the structure and actual function of the E1 glycoprotein (Douam et al., 2014; Haddad et  
38  
39 104 al., 2017; Lavillette et al., 2007; Wahid et al., 2013). Evidence for the immunogenicity and the induction  
40  
41 105 of neutralizing antibodies by E1 has been reported. However, only few mAbs have been raised against  
42  
43 106 this protein compared to the numerous anti-E2 mAbs described in the literature. This may be related to  
44  
45 107 the difficulty to express the E1 protein as a correctly folded monomer (Op De Beeck et al., 2001).  
46  
47 108 Polyclonal antibodies from mice vaccinated with E1-HCVpp or recombinant E1 protein were able to  
48  
49 109 neutralize HCVcc (Dreux et al., 2006; Pietschmann et al., 2006). Also, synthetic peptides covering the C  
50  
51 110 terminal region of E1 were able to react with 32% of sera from infected patients in one study (Siemoneit  
52  
53 111 et al., 1995) while 92% reactivity was reported in another study (Ray et al., 1994). In addition, immune  
54  
55 112 sera of rabbits vaccinated with a synthetic peptide encompassing AA 315-323 prevented the binding and  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

113 entry of HCV particles into HepG2 cells (El-Awady et al., 2006). However, this finding has not been  
114 confirmed using the HCVpp or HCVcc systems.

115 Two main regions have been identified as reactive domains for anti-E1 antibodies. The first region is  
116 the N- terminal region identified by the human mAb H-111 (AA 192-202) (Keck et al., 2004b) and the  
117 murine mAb A4 (AA 197-207) (Dubuisson et al., 1994). While these antibodies recognize E1 presented  
118 on the viral envelope, mAb H-111 is only weakly neutralizing while no neutralization has been reported  
119 for mAb A4. The second region encompasses AA 313-327, which is located at the C terminus and is  
120 identified mainly by mAbs IGH505 and IGH526 (Kong et al., 2015; Meunier et al., 2008). We isolated a  
121 monoclonal antibody, designated A6, from a HCV genotype 1b infected patient. Epitope mapping  
122 identified AA 230-239 within the N-terminal region of E1 as the critical site for binding. This mAb  
123 showed affinity towards a panel of HCV envelope proteins of genotypes 1a, 1b and 2a, but lacked any  
124 neutralizing or interfering activity.

125

## 126 2. Materials and Methods

### 127 2.1. Human sera, cell lines and antibodies

128 Blood samples were collected from HCV infected patients who were followed at the Ghent  
129 University Hospital. The study was approved by the local ethical committee and all patients involved  
130 gave informed consent. Human embryonic kidney cells (293T) and human hepatoma cell lines (Huh-  
131 7.5RFP-NLS-IPS and Hep3B) were grown in Dulbecco's Modified Eagle's Medium supplemented with  
132 10% fetal calf serum, 1% non-essential amino acids, 1% L-glutamine and antibiotics. The human anti-E1  
133 mAb IGH526, the murine anti-E1 mAb A4, the murine anti-E2 mAb AP33 and the human anti-E2 mAbs  
134 MRCT10, 1:7, HC84.26, CBH-7 and HC-1AM were previously described (Allander et al., 2000;  
135 Dubuisson et al., 1994; Keck et al., 2004a; Keck et al., 2012; Kong et al., 2015; Owsianka et al., 2001;  
136 Pantua et al., 2013; Wang et al., 2011b).

### 138 2.2. Amplification and expression of HCV envelope

139 RNA was extracted from the plasma of infected patients using ZR Viral RNA kit (Zymo Research)  
140 followed by cDNA synthesis using superscript III reverse transcriptase (Invitrogen) and random primers.  
141 The full-length E1E2 sequence was amplified by nested PCR. The first PCR was performed using  
142 LongAmp DNA polymerase (NEB) and the following primers: (F) 5'-CGT AGG TCG CGT AAC TTG  
143 GGT AA-3' and (R) 5'-GTG CGC CTC GGC CCT GGT GAT AAA-3'. The second-round PCR was  
144 performed using Pfu DNA polymerase (Promega) and primers: (F) 5'-TAT AGA TAT CAT GGG GTA  
145 CAT TCC GCT CGT C-3' and (R) 5'-ATA TGA TAT CTT ACT CAG CCT GAG CTA TCA G-3'. PCR  
146 products were analyzed using 1% agarose gel electrophoresis and the bands corresponding to E1E2 were  
147 eluted and cloned into the pCDNA3.1/Hygro expression vector (Invitrogen). The inserts were sequenced  
148 (Center for Medical Genetics, Ghent University, Belgium), multiple aligned and analyzed using BioEdit  
149 version 7.2.0, Clone manager 9 professional and CLC main workbench version 7.6.4 (QIAGEN). HCV  
150 E1E2 sequences covering all 7 genotypes (1a, 1b, 2a, 2b, 3a, 4a, 5a, 6a and 7a) were retrieved from the  
151 European HCV database (<https://euhcvdb.ibcp.fr/euHCVdb/>) and the NCBI  
152 (<https://www.ncbi.nlm.nih.gov/nucleotide/>) and used for alignment with Belgian isolates. Based on amino  
153 acid sequences of the whole E1E2, neighbor joining phylogenetic tree was constructed using 1000



154 replicates bootstrapping analysis of CLC main workbench version 7.6.4. For expression of E1E2 in  
1  
2 155 mammalian cells, constructs encoding the E1E2 region of genotypes 1-6 were used for transfection of  
3  
4 156 293T cells using ProFection mammalian transfection kit (Promega). After 48 hours, lysis buffer (Promega)  
5  
6 157 and protease inhibitors (Roche) were added and the cell lysate was centrifuged at 13,000 rpm and 4°C.  
7  
8 158 The supernatant was collected and stored at -80°C until use. Constructs containing the following viral  
9  
10  
11 159 strains were used: H77, UKN1A20.8, UKN1A14.38, J4, UKN1B5.23, UKN1B12.16, P5VD, P5VE,  
12  
13 160 P5VF, JFH1, UKN2A1.2, UKN2B2.8, S52, UKN4.11.1, UKN5.14.4 and UKN6.5.8 in addition to 28  
14  
15 161 Belgian isolates (Desombere et al., 2016; Fafi-Kremer et al., 2010; Lavillette et al., 2005; Owsianka et al.,  
16  
17 162 2005).

### 19 163 20 21 22 164 *2.3. Generation of anti-HCV mAb*

23  
24 165 The hybridoma cells were generated from the peripheral blood mononuclear cells (PBMCs) of a  
25  
26 166 genotype 1b HCV infected patient as described previously (Depraetere et al., 2001). Clones that secreted  
27  
28 167 E1-specific antibodies were identified using a prototype version of the INNO-LIA® HCV antibody test  
29  
30  
31 168 (Innogenetics, Belgium). After subcloning, one E1-specific hybridoma cell line was retained and  
32  
33 169 designated A6. Cells were propagated in a specific antibody production bioreactor (Integra) and the  
34  
35 170 culture supernatant was harvested. mAb A6 was purified using a protein G column (GE Healthcare Life  
36  
37 171 Sciences), concentrated using Amicon centrifugal filters with 50 kDa cut-off (Merck Millipore) and the  
38  
39  
40 172 concentration was estimated using a commercial human IgG ELISA quantification kit according to  
41  
42 173 manufacturer instructions (Bethyl Laboratories).

### 44 174 45 46 175 *2.4. GNA binding, competition and denaturation ELISA*

47  
48  
49 176 The binding affinity of mAb A6 to cell lysate containing recombinant HCV envelope glycoproteins  
50  
51 177 was tested using GNA binding ELISA as previously described with some modifications (Owsianka et al.,  
52  
53 178 2005). Briefly, ELISA plates were coated with Galanthus nivalis lectin (GNA) and incubated overnight at  
54  
55 179 4°C. The next day, plates were washed (PBS-0.05% Tween 20) and blocked (PBS-5% BSA) for 1 hour at  
56  
57  
58 180 room temperature (RT), after which cell lysate was added and incubated for 2 hours. After washing,  
59  
60 181 diluted mAbs (PBS-5% BSA, 20% goat serum, 0.05% Tween 20) were added and plates were incubated

182 1.5 hours followed by addition of HRP-conjugated goat anti-host IgG (anti-human for A6 and 1:7 or anti-  
183 mouse for A4). In case of peptide competition assay, serial dilutions of peptides were mixed with a fixed  
184 concentration of mAb A6 (20 µg/mL) before addition to wells. TMB substrate was added and the optical  
185 density (OD) was read at 450nm. Competition ELISA was performed using a previously described  
186 protocol with some modifications (Potter et al., 2012). Briefly, the GNA-coated plates were loaded with  
187 E1E2 cell lysate diluted in blocking buffer (2.5% BSA, 2.5% goat serum, 0.1% Tween 20) and incubated  
188 for 2 hours. Plates were washed and serial dilutions of competing mAbs (A6 or MRCT10) were added.  
189 After 1 hour, biotinylated AP33 was added at a concentration corresponding to 60-75% of the maximum  
190 OD value and incubated for 1 hour. Plates were washed and HRP-conjugated streptavidin was added for  
191 30 min. In denaturation ELISA, cell lysates were incubated with 0.5% sodium dodecyl sulfate (SDS) and  
192 5 mM dithiothreitol (DTT) for 15 min at 56°C (Keck et al., 2012).

193

#### 194 2.5. Protein electrophoresis and Immunoblotting

195 To test the specificity of mAb A6 binding, lysate of 293T cells expressing H77 E1E2 was used in  
196 sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). Samples were mixed with 4X  
197 Bolt LDS sample and reducing buffers (Life Technologies) followed by heating at 70°C for 10 minutes.  
198 In case of deglycosylation, the cell lysate was first treated with PNGase F or Endo H (NEB) according to  
199 manufacturer instructions. Samples were then resolved on 12% acrylamide gel followed by transfer to  
200 nitrocellulose membrane and blocked for one hour at RT using 5% non-fat milk in TBS-T (TBS-0.05%  
201 Tween 20). After washing (TBS-T), mAb A6 was added (1 µg/mL) followed by 2 hours incubation at RT.  
202 HRP-conjugated goat anti-human IgG (Fc-specific) was added followed by one hour incubation. Bands  
203 were visualized using Pierce ECL Plus western blotting substrate (Thermo Scientific) and imaged on an  
204 ImageQuant LAS 4000 (GE Healthcare Life Sciences). Molecular weights were estimated based on the  
205 MagicMark XP western protein standard (Invitrogen).

#### 207 2.6. Epitope mapping

208 To identify the region critical for mAb binding, a 15-mer, 14-AA overlapping peptide microarray  
209 (PEPperCHIP) was used according to the manufacturer's instructions (PEPperPRINT GmbH, Germany).

210 The peptides printed on the chip were designed to cover the entire E1E2 sequence of a HCV genotype 1b  
1  
2211 isolate. The sequences and the layout of peptides spotted on the chip are presented in Supplementary File  
3  
4212 1. For peptide ELISA, plates were loaded with 2.5 µg/well of dissolved peptides (GenScript, USA) and  
5  
6213 incubated overnight at 4°C. After washing and blocking for 1 hour, serial dilutions of mAb A6 were  
7  
8214 added followed by incubation for 1 hour. Plates were washed and HRP-conjugated goat-anti-human IgG  
9  
10215 was added. After 1 hour of incubation, plates were washed, TMB substrate was added and the OD was  
11  
12216 measured as described previously.  
13  
14

15217

16

17218

18

19219

20

21

22220

23

24221

25

26222

27

28223

29

30

31224

32

33225

34

35226

36

37227

38

39

40228

41

42229

43

44230

45

46231

47

48

49232

50

51233

52

53234

54

55235

56

57236

58

59

60237

61

62

63

64

65

## 2.7. Generation and neutralization of HCVpp

HCV pseudoparticles were produced as previously described (Hsu et al., 2003). Briefly, using calcium phosphate method (Promega), 293T cells were co-transfected with two constructs: pcDNA3.1/Hygro-E1E2 and pNL4-3.Luc.R<sup>-</sup>E<sup>-</sup> (envelope-deficient HIV-1 proviral genome with a luciferase reporter). Isolates used for HCVpp production included genotype 1a (H77), genotype 1b (UG-P09-779, UG-P10-252 and UG-P12-763) and genotype 2a (JFH1) strains. Supernatant containing HCVpp was collected 72 hours post transfection, sterile filtered through 0.45 µm filter and stored at -80°C. For HCVpp neutralization, 1x10<sup>4</sup> Hep3B cells were seeded in 96-well tissue culture plate and incubated at 37°C and 5% CO<sub>2</sub>. The following day, HCVpp were mixed with mAb A6 (100 µg/mL) or anti-E1 mAb IGH526 (10 µg/mL) or anti-E2 mAbs (MRCT10, 1:7, HC84.26, CBH-7 and HC-1AM; 10 µg/mL) or combinations. The mixture was incubated for 1 hour at 37°C before it was added to the Hep3B culture. After 72 hours, intracellular luciferase activity was measured to estimate HCVpp infection (Promega).

## 2.8. HCVcc production and neutralization

Cell culture-derived HCV (HCVcc) of isolates H77c/JFH1 (genotype 1a), J4/JFH1 (genotype 1b) and J6/JFH1 (Jc1; genotype 2a) was produced as described previously with some modifications (Wakita et al., 2005). Briefly, XbaI linearized HCV plasmids were *in vitro* transcribed (Promega) and the RNA was used for transfection of Huh-7.5RFP-NLS-IPS cells using Lipofectamine 2000 reagent (Invitrogen). Culture supernatant was harvested, sterile filtered and stored at -80°C. For HCVcc neutralization, 1.3x10<sup>4</sup> Huh-7.5RFP cells were seeded in a 96-well plate. The next day, mAb A6 (100 µg/mL) or MRCT10 (10 µg/mL)

238 was pre-incubated with 50-100 focus forming units (FFUs) of HCVcc at 37°C for 1 hour. The mixture  
1  
239 was then added to Huh-7.5RFP cells and incubated for 4 hours followed by washing and incubation for  
3  
4240 another 48 hours. Infected cells were immunostained with mouse anti-NS5A mAb 9E10 (kindly provided  
5  
6241 by Dr. Charles Rice, Rockefeller University, USA) and Alexa 647 conjugated goat-anti-mouse IgG  
7  
8242 (Invitrogen). The number of FFUs were determined using the BD Pathway 435 High Content Bioimager  
9  
10243 (BD Biosciences). To test the suitability of mAb A6 for immunofluorescence imaging, H77c/JFH1  
11  
12244 infected cells were incubated overnight at RT with mAb A6 (1 µg/mL), after which Alexa 647-conjugated  
13  
14245 goat-anti-human IgG was added for visualization. As a positive control, the same cells were stained with  
15  
16246 the anti-NS5A mAb 9E10, in combination with Alexa 488-conjugated goat-anti-mouse IgG.  
17  
18

19  
20247

21  
22248

23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

### 249 3. Results

#### 1 250 3.1. Amplification of E1E2 from Belgian isolates

3  
4251 Following E1E2 amplification and agarose gel electrophoresis, bands of approximately 1.7kb,  
5  
6252 corresponding to full length E1E2 coding genes were cloned into mammalian expression vector and  
7  
8253 sequenced (data not shown). The names and accession numbers of previously published sequences are  
9  
10  
11254 listed in Table 1. Multiple sequence alignment and phylogenetic analysis showed that our sequences  
12  
13255 correspond to 28 new viral isolates (Supplementary Fig. 1) and are of genotype 1b, except for one isolate  
14  
15256 that is classified as genotype 1a (Fig. 1).

#### 17 18 19257 20258 3.2. Human mAb A6 efficiently binds to recombinant E1E2 proteins

21  
22259 Hybridoma cell lines were prepared from B lymphocytes of a chronic HCV carrier as described  
23  
24260 before (Depraetere et al., 2001). Screening of hybridoma culture supernatants identified a clone secreting  
25  
26261 an HCV E1-specific antibody, designated A6. To evaluate the binding affinity of this antibody to  
27  
28262 recombinant HCV envelope (E1E2), expression vectors encoding the HCV envelope proteins were used  
29  
30  
31263 to transfect 293T cells and cell lysates were used in GNA binding ELISA. As shown in Fig. 2A and B,  
32  
33264 sigmoidal dose response curves were observed when cell lysate containing E1E2 of three genotype 1a  
34  
35265 (H77, UKN1A20.8 and UKN1A14.38), one genotype 1b (UKN1B12.16) and two genotype 2a  
36  
37266 (UKN2A1.2 and JFH1) isolates were used. Similar binding experiments were performed using cell lysates  
38  
39  
40267 containing E1E2 of multiple isolates. These included previously published genotype 1b isolates (J4,  
41  
42268 UKN1B5.23, P5VD, P5VE and P5VF), the 28 Belgian isolates of genotype 1a and 1b mentioned above as  
43  
44269 well as isolates of genotype 2b (UKN2B2.8), genotype 3a (S52), genotype 4 (UKN4.11.1), genotype 5  
45  
46270 (UKN5.14.4) and genotype 6 (UKN6.5.8). Overall, GNA ELISA binding data showed that mAb A6  
47  
48  
49271 antibody efficiently reacts with full length E1E2 envelope glycoproteins of genotypes 1a, 1b and 2a, but  
50  
51272 not with E1E2 of genotypes 2b, 3a, 4, 5 and 6 (Fig. 2C).

#### 53 54 55274 3.3. Antibody A6 recognizes the region AA 230-239 of E1

56  
57  
58275 To identify the region critical for mAb A6 binding, we used a custom-made peptide microarray  
59  
60276 spanning the complete E1E2 sequence. Upon incubation of array chip with mAb A6, reactive spots were

277 observed with peptides spanning amino acids 225-243 of the HCV E1 protein, while no reactivity was  
1  
278 seen in any other regions within E1 or E2 (Fig. 3A). The sequences of peptides spanning the full length  
3  
4279 E1E2 are presented in Supplementary File 1 and peptides with positive reactivity to A6 are highlighted  
5  
6280 with grey background. To confirm this result, 11 individual peptides spanning the region 222-246 were  
7  
8281 produced for use in traditional peptide ELISA. As shown in Figure 3B, antibody A6 reacted with peptides  
9  
10282 spanning the region AA 225-244, thereby confirming the results obtained with the PEPperPRINT  
11  
12283 microarray assay. Since the peptides used in these assays are not glycosylated, we can assume that the  
13  
14284 glycosylation of the HCV envelope protein in this region is not essential for mAb binding. To test this  
15  
16285 hypothesis, we performed peptide-based competition assay. In this assay, ELISA plates pre-coated with  
17  
18286 recombinant E1E2 protein of H77 isolate were loaded with serial dilutions of individual peptides together  
19  
20287 with a fixed concentration of mAb A6. This showed the ability of the peptides spanning AA 225-244 to  
21  
22288 efficiently compete with recombinant E1E2 for binding to mAb A6, which confirms that A6-binding is  
23  
24289 glycosylation independent (Fig. 3C). The glycosylation-independent reactivity of mAb A6 was confirmed  
25  
26290 using Western blotting analysis of PNGase F-treated, Endo H-treated and non-treated E1E2 cell lysate,  
27  
28291 where mAb A6 binds to both glycosylated and deglycosylated E1 (Fig. 4). Overall, our data indicates that  
29  
30292 the minimal epitope of antibody A6 encompasses AA 230-239 of HCV E1.  
31  
32  
33293

### 37294 *3.4. The epitope identified by mAb A6 has a linear conformation*

38  
39

40295 In order to assess whether the A6 epitope has a linear or conformational structure, we performed a  
41  
42296 denaturation binding assay. Binding of A6 to native and denatured E1E2-containing cell lysate was  
43  
44297 assessed and compared to the binding activity of antibodies A4 and 1:7, targeting respectively a linear E1-  
45  
46298 epitope and a conformational epitope within E2. While denaturation of H77 E1E2 only minimally  
47  
48299 affected binding of antibodies A4 and A6, that of mAb 1:7 was completely abolished (Fig. 5A). Likewise,  
49  
50300 denaturation of JFH1 E1E2 did not negatively impact A6 binding, while the binding of mAb 1:7 was  
51  
52301 completely lost (Fig. 5B). The absence of A4 binding to both native and denatured JFH1 E1E2 protein is  
53  
54302 explained by its non-reactivity towards genotype 2 isolates. Altogether, these data show that the epitope  
55  
56303 targeted by mAb A6 has a linear, denaturation-resistant structure.  
57  
58  
59  
60  
61  
62  
63  
64  
65

304 To investigate potential conservation of the A6 epitope among different HCV genotypes, amino acid  
1  
2305 sequences of E1 for isolates used in the binding assay were aligned using BioEdit and CLC main  
3  
4306 workbench. As seen from the sequence alignment and the percentage of conservation, certain amino acids  
5  
6307 in region 230-239, which is critical for mAb A6 binding, are highly conserved especially among  
7  
8308 genotypes 1a and 1b (Fig. 6). Amino acids cysteine and tryptophan at sites 238 and 239 respectively are  
9  
10309 conserved in all 7 genotypes. The residues 234N and 236S are also conserved, except in case of genotype  
11  
12310 2b where they are replaced by glycine and leucine, respectively. Likewise for 230V, which is only  
14  
15311 different for genotype 2. Other positions such as 231, 232 and 237 show high conservation among  
16  
17312 genotypes 1a, 1b and 2a compared to genotypes 2b, 3a, 4, 5 and 6.  
18

### 19313 20 21 22314 3.5. Antibody A6 does not neutralize HCVpp and HCVcc

23  
24315 Since mAb A6 efficiently binds to E1E2, we next wanted to investigate its neutralizing activity.  
25  
26316 Therefore, HCVpp from genotypes 1a (H77 isolate), 1b (UG-P09-779, UG-P10-252 and UG-P12-763)  
27  
28317 and 2a (JFH1 isolate) were produced and used in neutralization assays. Incubation of mAb A6 with  
29  
30318 HCVpp before addition to Hep3B cell cultures did not protect the target cells from infection, even at high  
31  
32319 mAb concentration (100 µg/mL) (Fig. 7A-E). In parallel, 10 µg/mL of the AP33-derived humanized mAb  
34  
35320 MRCT10 was included as positive control and completely prevented HCVpp infection of all tested  
36  
37321 genotypes. These results indicate that mAb A6 does not exhibit any neutralizing activity in the HCVpp  
38  
39  
40322 system, despite its capacity to efficiently bind the viral envelope.

41  
42323 Subsequently we also tested the neutralizing capacity of mAb A6 in the context of the HCV cell  
43  
44324 culture (HCVcc) system. Three universally used chimeric viruses H77c/JFH1, J4/JFH1 and Jc1 (J6/JFH1),  
45  
46325 incorporating the envelope proteins of isolates H77c (genotype 1a), J4 (genotype 1b) and J6 (genotype 2a)  
47  
48326 respectively, were pre-incubated with A6 or MRCT10, after which the mixture was transferred to Huh7.5  
49  
50  
51327 hepatoma cells. Enumeration of the number of infected foci 2 days later showed that mAb A6 was unable  
52  
53328 to protect Huh7.5 cells from HCVcc challenge, while MRCT10 did (Fig. 7F-H). Altogether, our HCVpp  
54  
55329 and HCVcc data indicate that mAb A6 has no neutralizing activity. However, staining of H77c/JFH1  
56  
57330 infected Huh7.5RFP cells with mAb A6 enabled the visualization of infected cells which indicates the  
58  
59  
60331 applicability of mAb A6 for use in immunofluorescence (Fig. 8).

### 3.6. mAb A6 does not interfere with neutralizing antibodies

A broadly neutralizing determinant encompassing residues 412-423 of E2 has previously been shown to be susceptible to antibody interference (Kachko et al., 2015; Zhang et al., 2009), although this is controversial (Tarr et al., 2012). In this regard, we examined if antibody A6 could interfere with the binding and/or protective activity of other known neutralizing antibodies. Murine mAb AP33 is one of the most extensively studied antibodies that target a linear epitope within HCV E2 protein and exhibits broad neutralizing activity both *in vitro* and *in vivo* (Desombere et al., 2016; Owsianka et al., 2005). In a first setup, full length E1E2 protein of isolate H77 was incubated with serial dilutions of mAb A6 before addition of biotinylated AP33. Subsequent addition of HRP-conjugated streptavidin enabled us to assess the amount of biotinylated antibody that still was able to bind to the envelope proteins. As can be seen in Figure 9A, mAb A6 did not interfere with the binding the biotinylated AP33 whereas the humanized version of AP33, mAb MRCT10, inhibited the subsequent binding of biotinylated AP33 in a dose-dependent manner. To confirm the non-interfering nature of A6, HCVpp neutralization was performed using H77 and JFH1 isolates in the presence of mAbs A6, MRCT10, IGH526, 1:7, HC84.26, CBH-7 and HC-1AM alone or in combination. Measurement of luciferase activity showed that MRCT10 efficiently inhibited HCVpp infection of both genotypes, while the co-incubation of mAb A6 did not interfere with this activity (Fig. 9B and C). Similarly, mAb A6 did not negatively impact the neutralizing activity of mAbs IGH526, 1:7, HC84.26, CBH-7 and HC-1AM. These mAbs have been previously shown to have cross-neutralizing activity against HCVpp and HCVcc (Johansson et al., 2007; Keck et al., 2004a; Keck et al., 2012; Wang et al., 2011b). Based on our ELISA and HCVpp competition experiments we can conclude that our anti-E1 mAb A6 does not interfere with the neutralizing activity of known anti-E2 mAbs.



#### 357 4. Discussion

1  
2358 In the current study, we describe the development and characterization of a novel human monoclonal  
3  
4359 antibody (designated A6) isolated by hybridoma technology from the PBMCs of a patient chronically  
5  
6360 infected with HCV genotype 1b. Epitope mapping identified the region encompassing AA 230-239,  
7  
8361 located at the N-terminal region of E1, as the critical region for binding. Previously, two other regions  
9  
10362 within E1 have been described that elicit E1-specific antibodies. The first is the N-terminal region  
11  
12363 recognized by human mAb H-111 (AA 192-202) (Keck et al., 2004b) and murine mAb A4 (AA 197-207)  
13  
14364 (Dubuisson et al., 1994), which lies upstream of the epitope recognized by A6. The second antigenic  
15  
16365 region within E1 encompasses AA 313-327, which is located at the C-terminal region and is recognized  
17  
18366 by cross-neutralizing mAbs such as IGH505 and IGH526 (Meunier et al., 2008). mAb H-111 was isolated  
19  
20367 from a patient chronically infected with HCV of genotype 1b and showed high binding activity to E1  
21  
22368 protein from genotypes 1a, 1b, 2b. It bound only moderately to HCV of genotype 3a, and no binding was  
23  
24369 observed to genotype 2a and 4a. It also exhibited some neutralizing activity towards HCV-like particles  
25  
26370 and HCVpp, but complete prevention of infection was not achieved (Dreux et al., 2006; Keck et al.,  
27  
28371 2004b). To evaluate the binding activity of mAb A6, we used full length E1E2 protein that was expressed  
29  
30372 in mammalian cells. New isolates from Belgian patients, genotyped as 1a and 1b, have been used together  
31  
32373 with well-known isolates covering genotypes 1 to 6. Active binding of mAb A6 to all isolates of  
33  
34374 genotypes 1a (4 isolates), 1b (33 isolates) and 2a (2 isolates) was observed, while no binding activity was  
35  
36375 seen with the other genotypes (2b, 3a, 4, 5 and 6; one isolate for each genotype). This resembles the mAb  
37  
38376 H-111 in binding to genotypes 1a and 1b (Keck et al., 2004b). The major difference between the mAb H-  
39  
40377 111 study and A6 is that we used full length E1E2 proteins instead of E1. Altogether these data confirm  
41  
42378 that the activity of antibodies raised against the N-terminal region of E1 is genotype dependent.

43  
44379 Although the sequence of the HCV envelope exhibits great variability across genotypes, conserved  
45  
46380 glycosylation sites in E1, including the N-terminal region, and E2 have been identified and shown to play  
47  
48381 a role in evasion from host immunity (Dubuisson et al., 1994; Goffard and Dubuisson, 2003; Helle et al.,  
49  
50382 2011; Helle et al., 2010). Since the recombinant E1E2 proteins used in our binding ELISA were  
51  
52383 expressed in mammalian cells, their proper glycosylation and post-translational modification may be  
53  
54384 expected. In contrast, synthetic peptides used in epitope mapping are non-glycosylated. For this reason,  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

385 we have also examined the effect of these peptides in a competition assay together with recombinant  
1  
2386 E1E2 protein. The strong inhibition of binding of mAb A6 to E1E2 glycoprotein upon addition of  
3  
4387 synthetic peptides indicates that this binding is not relying on the glycosylation of E1. This was  
5  
6388 corroborated by Western blot analysis, which showed positive reactivity of mAb A6 to both glycosylated  
7  
8389 and (PNGase/Endo H) deglycosylated E1 (Dubuisson et al., 2000). In addition, the envelope of HCV  
9  
10390 contains multiple intramolecular disulfide bonds which are essential for proper folding and stabilizing  
11  
12391 (Krey et al., 2010). Incubation of recombinant envelope glycoprotein with SDS and DTT at high  
13  
14392 temperature destroys these disulfide bonds and disrupts the secondary structure of the protein. This  
15  
16393 approach has been used to differentiate between antibodies targeting linear versus conformation-  
17  
18394 dependent epitopes (Keck et al., 2012; Tarr et al., 2006). mAb A6 efficiently recognized the envelope of  
19  
20395 both native and denatured E1E2 proteins and a similar result was obtained for mAb A4 that previously  
21  
22396 shown to target a linear epitope within E1. In parallel, mAb 1:7 that targets a conformational epitope  
23  
24397 within E2 completely lost its binding upon denaturation of the envelope proteins, confirming previous  
25  
26398 findings (Allander et al., 2000; Johansson et al., 2007). Altogether, our results indicate that the mAb A6  
27  
28399 targets a linear epitope within the N-terminus of the E1 protein. Multiple sequence alignment of the E1  
29  
30400 protein using all isolates used in the current study in addition to previously published sequences revealed  
31  
32401 the highly conserved nature of residues 230, 234, 236, 238 and 239 within the A6 epitope among different  
33  
34402 genotypes. However, the change of polar amino acids at residues 234 and 236 to more hydrophobic ones  
35  
36403 may be the underlying cause of the absence of reactivity of mAb A6 towards genotype 2b isolates. On the  
37  
38404 other hand, residues at sites 231, 232 and 237 are highly conserved among isolates of genotypes 1a, 1b  
39  
40405 and 2a compared to the high variation observed in other genotypes like 2b, 3a, 4, 5 and 6. Altogether this  
41  
42406 may explain the efficient and specific binding of mAb A6 with the E1 envelope protein of isolates of  
43  
44407 genotypes 1a, 1b and 2a. Similar sequence conservation has been shown upon alignment of the C-  
45  
46408 terminal region, a region that is identified by cross-neutralizing antibodies (Bukh et al., 1993; Meunier et  
47  
48409 al., 2008).

55410 Using the HCVpp system, mAb A6 could not inhibit the entry of pseudoparticles of genotypes 1a, 1b  
56  
57411 and 2a, while complete protection was achieved by mAb MRCT10 - the humanized version of the cross-  
58  
59412 neutralizing mAb AP33 (Owsianka et al., 2001; Pantua et al., 2013). Previous studies showed low  
60  
61  
62  
63  
64  
65

413 neutralizing activity of the human mAb H-111 while the neutralizing potential of mAb A4 has not been  
1  
2 414 reported (Dubuisson et al., 1994; Keck et al., 2004b). Nevertheless, the non-neutralizing function of  
3  
4 415 antibodies raised against the N- terminal region cannot be generalized to all E1-specific mAbs since the  
5  
6 416 two mAbs, IGH505 and IGH526, that target the region AA 313-327 showed cross-neutralizing capacity  
7  
8 417 when tested in the HCVpp and HCVcc systems (Meunier et al., 2008). Since HCVpp are produced in  
9  
10 418 293T cells, a non-hepatic cell line, its lipoprotein profile may differ from the that of genuine HCV virions  
11  
12 419 produced in infected patients and from HCVcc produced in hepatic cell lines. In order to confirm the  
13  
14 420 results obtained with the HCVpp system, we performed similar neutralization experiments in the HCVcc  
15  
16 421 system. Despite the use of high antibody concentration mAb A6 was also unable to protect Huh7.5 cells  
17  
18 422 from infection.  
19  
20

21  
22 423 Some studies have reported that antibodies raised against certain epitopes within E2 interfere with the  
23  
24 424 neutralizing activity of other antibodies. For example, antibodies targeting epitope II (AA 434-446)  
25  
26 425 within E2 protein have been shown to interfere with the neutralizing activity of other mAbs targeting  
27  
28 426 epitope I (AA 412-424) (Kachko et al., 2015; Zhang et al., 2007; Zhang et al., 2009) although others  
29  
30 427 contradicted this finding (Tarr et al., 2012). No data is available about the potential interfering activity of  
31  
32 428 E1 antibodies. We therefore examined whether A6 interfered with the activity of anti-E1 antibody  
33  
34 429 IGH526 and anti-E2 mAbs MRCT10 (epitope I-specific), 1:7 (AA 523-535; domain B-specific), HC84.26  
35  
36 430 (epitope II-specific), CBH-7 (domain C-specific) and HC-1AM (domain B-specific) and found out that it  
37  
38 431 did not. A recent study reported that antibodies against non-neutralizing epitopes within E1 and E2 can be  
39  
40 432 involved in antibody-dependent cytotoxic responses by natural killer cells (Long et al., 2017). This new  
41  
42 433 finding suggests that non-neutralizing, non-interfering antibodies like A6 may still play a role in the  
43  
44 434 immune control of HCV. More in depth exploration of this mechanism is warranted.  
45  
46

47  
48  
49 435 In conclusion, we report that the region spanning AA 230-239 at the N-terminal region of E1 is  
50  
51 436 immunogenic and can elicit non-neutralizing Abs during a chronic HCV infection cross-reactive to  
52  
53 437 genotypes 1a, 1b and 2a. mAb A6 can be considered as one more tool that could help to elucidate the  
54  
55 438 structure of the viral envelope, an important step in the development of an effective prophylactic HCV  
56  
57 439 vaccine.  
58

59  
60 440

61  
62  
63  
64  
65

441 **Acknowledgments**

1  
2 442 AAM is a recipient of a PhD scholarship provided by the Egyptian Government. The study was supported  
3  
4 443 by Ghent University Special Research Fund [CRA grant 01G01712], the Research Foundation Flanders  
5  
6 444 (FWO Vlaanderen), the Belgian state [IUAP P7/47-HEPRO-2] and the European Union [FP7, HepaMab].  
7  
8 445 The work in AHP's laboratory was supported by the Medical Research Council, UK [MC\_UU 12014/2].  
9  
10 446 We would like to thank Dr. Charles Rice (Rockefeller University, USA) for Huh7.5RFP cells and mAb  
11  
12 447 9E10, Dr. Jane McKeating (University of Birmingham, UK) for 293T cells, Dr. Mansun Law (Scripps  
13  
14 448 Research Institute, USA), Dr. Thomas Baumert (University of Strasbourg, France) for P05 E1E2  
15  
16 449 constructs, Dr. Jens Bukh (University of Copenhagen, Denmark) for H77c/JFH1 plasmid, Dr. Nathaniel  
17  
18 450 Landau (NIH AIDS Research and Reference Reagent Program) for the pNL4-3.Luc.R<sup>-</sup>E<sup>-</sup> plasmid, Dr.  
19  
20 451 Ralf Bartenschlager (Heidelberg university, Germany) for Jc1 plasmid, Dr. Takaji Wakita (National  
21  
22 452 Institute of Infectious Diseases, Tokyo, Japan) for JFH1 plasmid and Genentech, USA for the provision  
23  
24 453 of MRCT10.  
25  
26  
27

28 454

29

30 455

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

- 1  
2 457 Abdelwahab, S.F., 2016. Cellular immune response to hepatitis-C-virus in subjects without viremia or  
3 458 seroconversion: is it important? *Infectious agents and cancer* 11, 23.
- 4 459 Allander, T., Drakenberg, K., Beyene, A., Rosa, D., Abrignani, S., Houghton, M., Widell, A., Grillner, L.,  
5 460 Persson, M.A., 2000. Recombinant human monoclonal antibodies against different conformational  
6 461 epitopes of the E2 envelope glycoprotein of hepatitis C virus that inhibit its interaction with CD81. *The*  
7 462 *Journal of general virology* 81, 2451-2459.
- 8 463 Bartenschlager, R., Penin, F., Lohmann, V., Andre, P., 2011. Assembly of infectious hepatitis C virus  
9 464 particles. *Trends in microbiology* 19, 95-103.
- 10 465 Bukh, J., 2016. The history of hepatitis C virus (HCV): Basic research reveals unique features in phylogeny,  
11 466 evolution and the viral life cycle with new perspectives for epidemic control. *J Hepatol* 65, S2-S21.
- 12 467 Bukh, J., Engle, R.E., Faulk, K., Wang, R.Y., Farci, P., Alter, H.J., Purcell, R.H., 2015. Immunoglobulin with  
13 468 High-Titer In Vitro Cross-Neutralizing Hepatitis C Virus Antibodies Passively Protects Chimpanzees from  
14 469 Homologous, but Not Heterologous, Challenge. *Journal of virology* 89, 9128-9132.
- 15 470 Bukh, J., Purcell, R.H., Miller, R.H., 1993. At least 12 genotypes of hepatitis C virus predicted by sequence  
16 471 analysis of the putative E1 gene of isolates collected worldwide. *Proc Natl Acad Sci U S A* 90, 8234-8238.
- 17 472 Catanese, M.T., Dorner, M., 2015. Advances in experimental systems to study hepatitis C virus in vitro  
18 473 and in vivo. *Virology* 479-480, 221-233.
- 19 474 Chung, R.T., Gordon, F.D., Curry, M.P., Schiano, T.D., Emre, S., Corey, K., Markmann, J.F., Hertl, M.,  
20 475 Pomposelli, J.J., Pomfret, E.A., Florman, S., Schilsky, M., Broering, T.J., Finberg, R.W., Szabo, G., Zamore,  
21 476 P.D., Khettry, U., Babcock, G.J., Ambrosino, D.M., Leav, B., Leney, M., Smith, H.L., Molrine, D.C., 2013.  
22 477 Human Monoclonal Antibody MBL-HCV1 Delays HCV Viral Rebound Following Liver Transplantation: A  
23 478 Randomized Controlled Study. *Am J Transplant* 13, 1047-1054.
- 24 479 Depraetere, S., Verhoye, L., Leclercq, G., Leroux-Roels, G., 2001. Human B cell growth and differentiation  
25 480 in the spleen of immunodeficient mice. *Journal of immunology* 166, 2929-2936.
- 26 481 Desombere, I., Fafi-Kremer, S., Van Houtte, F., Pessaux, P., Farhoudi, A., Heydmann, L., Verhoye, L., Cole,  
27 482 S., McKeating, J.A., Leroux-Roels, G., Baumert, T.F., Patel, A.H., Meuleman, P., 2016. Monoclonal anti-  
28 483 envelope antibody AP33 protects humanized mice against a patient-derived hepatitis C virus challenge.  
29 484 *Hepatology* 63, 1120-1134.
- 30 485 Douam, F., Dao Thi, V.L., Maurin, G., Fresquet, J., Mompelat, D., Zeisel, M.B., Baumert, T.F., Cosset, F.L.,  
31 486 Lavillette, D., 2014. Critical interaction between E1 and E2 glycoproteins determines binding and fusion  
32 487 properties of hepatitis C virus during cell entry. *Hepatology* 59, 776-788.
- 33 488 Dreux, M., Pietschmann, T., Granier, C., Voisset, C., Ricard-Blum, S., Mangeot, P.E., Keck, Z., Fong, S.,  
34 489 Vu-Dac, N., Dubuisson, J., Bartenschlager, R., Lavillette, D., Cosset, F.L., 2006. High density lipoprotein  
35 490 inhibits hepatitis C virus-neutralizing antibodies by stimulating cell entry via activation of the scavenger  
36 491 receptor BI. *The Journal of biological chemistry* 281, 18285-18295.
- 37 492 Dubuisson, J., Cosset, F.L., 2014. *Virology and cell biology of the hepatitis C virus life cycle: an update.* *J*  
38 493 *Hepatol* 61, S3-S13.
- 39 494 Dubuisson, J., Duvet, S., Meunier, J.C., Op De Beeck, A., Cacan, R., Wychowski, C., Cocquerel, L., 2000.  
40 495 Glycosylation of the hepatitis C virus envelope protein E1 is dependent on the presence of a  
41 496 downstream sequence on the viral polyprotein. *The Journal of biological chemistry* 275, 30605-30609.
- 42 497 Dubuisson, J., Hsu, H.H., Cheung, R.C., Greenberg, H.B., Russell, D.G., Rice, C.M., 1994. Formation and  
43 498 intracellular localization of hepatitis C virus envelope glycoprotein complexes expressed by recombinant  
44 499 vaccinia and Sindbis viruses. *Journal of virology* 68, 6147-6160.
- 45 500 El-Awady, M.K., Tabll, A.A., Atef, K., Yousef, S.S., Omran, M.H., El-Abd, Y., Bader-Eldin, N.G., Salem, A.M.,  
46 501 Zohny, S.F., El-Garf, W.T., 2006. Antibody to E1 peptide of hepatitis C virus genotype 4 inhibits virus  
47 502 binding and entry to HepG2 cells in vitro. *World journal of gastroenterology* 12, 2530-2535.
- 48 503 Fafi-Kremer, S., Fofana, I., Soulier, E., Carolla, P., Meuleman, P., Leroux-Roels, G., Patel, A.H., Cosset, F.L.,  
49 504 Pessaux, P., Doffoel, M., Wolf, P., Stoll-Keller, F., Baumert, T.F., 2010. Viral entry and escape from  
50 505 antibody-mediated neutralization influence hepatitis C virus reinfection in liver transplantation. *J Exp*  
51 506 *Med* 207, 2019-2031.

507 Goffard, A., Dubuisson, J., 2003. Glycosylation of hepatitis C virus envelope proteins. *Biochimie* 85, 295-  
1508 301.

2509 Haddad, J.G., Rouille, Y., Hanouille, X., Descamps, V., Hamze, M., Dabboussi, F., Baumert, T.F., Duverlie,  
3510 G., Lavie, M., Dubuisson, J., 2017. Identification of Novel Functions for Hepatitis C Virus Envelope  
4511 Glycoprotein E1 in Virus Entry and Assembly. *Journal of virology* 91.

5512 Helle, F., Duverlie, G., Dubuisson, J., 2011. The hepatitis C virus glycan shield and evasion of the humoral  
6513 immune response. *Viruses* 3, 1909-1932.

8514 Helle, F., Vieyres, G., Elkrief, L., Popescu, C.I., Wychowski, C., Descamps, V., Castelain, S., Roingeard, P.,  
9515 Duverlie, G., Dubuisson, J., 2010. Role of N-Linked Glycans in the Functions of Hepatitis C Virus Envelope  
10516 Proteins Incorporated into Infectious Virions. *Journal of virology* 84, 11905-11915.

11517 Hsu, M., Zhang, J., Flint, M., Logvinoff, C., Cheng-Mayer, C., Rice, C.M., McKeating, J.A., 2003. Hepatitis C  
12518 virus glycoproteins mediate pH-dependent cell entry of pseudotyped retroviral particles. *P Natl Acad Sci*  
13519 *USA* 100, 7271-7276.

1520 Johansson, D.X., Voisset, C., Tarr, A.W., Aung, M., Ball, J.K., Dubuisson, J., Persson, M.A.A., 2007. Human  
16521 combinatorial libraries yield rare antibodies that broadly neutralize hepatitis C virus. *P Natl Acad Sci USA*  
17522 104, 16269-16274.

18523 Kachko, A., Frey, S.E., Sirota, L., Ray, R., Wells, F., Zubkova, I., Zhang, P., Major, M.E., 2015. Antibodies to  
19524 an interfering epitope in hepatitis C virus E2 can mask vaccine-induced neutralizing activity. *Hepatology*  
20525 62, 1670-1682.

22526 Keck, Z.Y., Op De Beeck, A., Hadlock, K.G., Xia, J., Li, T.K., Dubuisson, J., Fong, S.K., 2004a. Hepatitis C  
23527 virus E2 has three immunogenic domains containing conformational epitopes with distinct properties  
24528 and biological functions. *Journal of virology* 78, 9224-9232.

25529 Keck, Z.Y., Sung, V.M., Perkins, S., Rowe, J., Paul, S., Liang, T.J., Lai, M.M., Fong, S.K., 2004b. Human  
26530 monoclonal antibody to hepatitis C virus E1 glycoprotein that blocks virus attachment and viral  
27531 infectivity. *Journal of virology* 78, 7257-7263.

29532 Keck, Z.Y., Wang, Y., Lau, P., Lund, G., Rangarajan, S., Fauvelle, C., Liao, G.C., Holtsberg, F.W., Warfield,  
30533 K.L., Aman, M.J., Pierce, B.G., Fuerst, T.R., Bailey, J.R., Baumert, T.F., Mariuzza, R.A., Kneteman, N.M.,  
31534 Fong, S.K., 2016. Affinity maturation of a broadly neutralizing human monoclonal antibody that  
32535 prevents acute hepatitis C virus infection in mice. *Hepatology* 64, 1922-1933.

34536 Keck, Z.Y., Xia, J., Wang, Y., Wang, W., Krey, T., Prentoe, J., Carlsen, T., Li, A.Y., Patel, A.H., Lemon, S.M.,  
35537 Bukh, J., Rey, F.A., Fong, S.K., 2012. Human monoclonal antibodies to a novel cluster of conformational  
36538 epitopes on HCV E2 with resistance to neutralization escape in a genotype 2a isolate. *Plos Pathog* 8,  
37539 e1002653.

38540 Kong, L., Kadam, R.U., Giang, E., Ruwona, T.B., Nieuwma, T., Culhane, J.C., Stanfield, R.L., Dawson, P.E.,  
39541 Wilson, I.A., Law, M., 2015. Structure of Hepatitis C Virus Envelope Glycoprotein E1 Antigenic Site 314-  
40542 324 in Complex with Antibody IGH526. *J Mol Biol* 427, 2617-2628.

42543 Krey, T., d'Alayer, J., Kikuti, C.M., Saulnier, A., Damier-Piolle, L., Petitpas, I., Johansson, D.X., Tawar, R.G.,  
43544 Baron, B., Robert, B., England, P., Persson, M.A., Martin, A., Rey, F.A., 2010. The disulfide bonds in  
44545 glycoprotein E2 of hepatitis C virus reveal the tertiary organization of the molecule. *Plos Pathog* 6,  
45546 e1000762.

47547 Lavillette, D., Pecheur, E.I., Donot, P., Fresquet, J., Molle, J., Corbau, R., Dreux, M., Penin, F., Cosset, F.L.,  
48548 2007. Characterization of fusion determinants points to the involvement of three discrete regions of  
49549 both E1 and E2 glycoproteins in the membrane fusion process of hepatitis C virus. *Journal of virology* 81,  
50550 8752-8765.

51551 Lavillette, D., Tarr, A.W., Voisset, C., Donot, P., Bartosch, B., Bain, C., Patel, A.H., Dubuisson, J., Ball, J.K.,  
52552 Cosset, F.L., 2005. Characterization of host-range and cell entry properties of the major genotypes and  
53553 subtypes of hepatitis C virus. *Hepatology* 41, 265-274.

55554 Li, G., De Clercq, E., 2017. Current therapy for chronic hepatitis C: The role of direct-acting antivirals.  
56555 *Antiviral Res* 142, 83-122.

57556 Long, L., Jia, M., Fan, X., Liang, H., Wang, J., Zhu, L., Xie, Z., Shen, T., 2017. Non-neutralizing epitopes  
58557 induce robust hepatitis C virus (HCV)-specific antibody-dependent CD56+ natural killer cell responses in  
59558 chronic HCV-infected patients. *Clinical and experimental immunology* 189, 92-102.

61  
62  
63  
64  
65

559 Mesalam, A.A., Vercauteren, K., Meuleman, P., 2016. Mouse Systems to Model Hepatitis C Virus  
560 Treatment and Associated Resistance. *Viruses* 8.

561 Meuleman, P., Bukh, J., Verhoye, L., Farhoudi, A., Vanwolleghem, T., Wang, R.Y., Desombere, I., Alter, H.,  
562 Purcell, R.H., Leroux-Roels, G., 2011. In Vivo Evaluation of the Cross-Genotype Neutralizing Activity of  
563 Polyclonal Antibodies Against Hepatitis C Virus. *Hepatology* 53, 755-762.

564 Meunier, J.C., Russell, R.S., Goossens, V., Priem, S., Walter, H., Depla, E., Union, A., Faulk, K.N., Bukh, J.,  
565 Emerson, S.U., Purcell, R.H., 2008. Isolation and characterization of broadly neutralizing human  
566 monoclonal antibodies to the E1 glycoprotein of hepatitis C virus. *Journal of virology* 82, 966-973.

567 Ndongo, N., Berthillon, P., Pradat, P., Vieux, C., Bordes, I., Berby, F., Maynard, M., Zoulim, F., Trepo, C.,  
568 Petit, M.A., 2010. Association of anti-E1E2 antibodies with spontaneous recovery or sustained viral  
569 response to therapy in patients infected with hepatitis C virus. *Hepatology* 52, 1531-1542.

570 Op De Beeck, A., Cocquerel, L., Dubuisson, J., 2001. Biogenesis of hepatitis C virus envelope  
571 glycoproteins. *The Journal of general virology* 82, 2589-2595.

572 Osburn, W.O., Snider, A.E., Wells, B.L., Latanich, R., Bailey, J.R., Thomas, D.L., Cox, A.L., Ray, S.C., 2014.  
573 Clearance of Hepatitis C Infection Is Associated With the Early Appearance of Broad Neutralizing  
574 Antibody Responses. *Hepatology* 59, 2140-2151.

575 Owsianka, A., Clayton, R.F., Loomis-Price, L.D., McKeating, J.A., Patel, A.H., 2001. Functional analysis of  
576 hepatitis C virus E2 glycoproteins and virus-like particles reveals structural dissimilarities between  
577 different forms of E2. *Journal of General Virology* 82, 1877-1883.

578 Owsianka, A., Tarr, A.W., Juttla, V.S., Lavillette, D., Bartosch, B., Cosset, F.L., Ball, J.K., Patel, A.H., 2005.  
579 Monoclonal antibody AP33 defines a broadly neutralizing epitope on the hepatitis C virus E2 envelope  
580 glycoprotein. *Journal of virology* 79, 11095-11104.

581 Pantua, H., Diao, J., Ultsch, M., Hazen, M., Mathieu, M., McCutcheon, K., Takeda, K., Date, S., Cheung,  
582 T.K., Phung, Q., Hass, P., Arnott, D., Hongo, J.A., Matthews, D.J., Brown, A., Patel, A.H., Kelley, R.F.,  
583 Eigenbrot, C., Kapadia, S.B., 2013. Glycan shifting on hepatitis C virus (HCV) E2 glycoprotein is a  
584 mechanism for escape from broadly neutralizing antibodies. *J Mol Biol* 425, 1899-1914.

585 Pawlotsky, J.M., 2016. Hepatitis C Virus Resistance to Direct-Acting Antiviral Drugs in Interferon-Free  
586 Regimens. *Gastroenterology* 151, 70-86.

587 Petruzzello, A., Marigliano, S., Loquercio, G., Cozzolino, A., Cacciapuoti, C., 2016. Global epidemiology of  
588 hepatitis C virus infection: An up-date of the distribution and circulation of hepatitis C virus genotypes.  
589 *World journal of gastroenterology* 22, 7824-7840.

590 Pietschmann, T., Kaul, A., Koutsoudakis, G., Shavinskaya, A., Kallis, S., Steinmann, E., Abid, K., Negro, F.,  
591 Dreux, M., Cosset, F.L., Bartenschlager, R., 2006. Construction and characterization of infectious  
592 intragenotypic and intergenotypic hepatitis C virus chimeras. *Proc Natl Acad Sci U S A* 103, 7408-7413.

593 Potter, J.A., Owsianka, A.M., Jeffery, N., Matthews, D.J., Keck, Z.Y., Lau, P., Fong, S.K.H., Taylor, G.L.,  
594 Patel, A.H., 2012. Toward a Hepatitis C Virus Vaccine: the Structural Basis of Hepatitis C Virus  
595 Neutralization by AP33, a Broadly Neutralizing Antibody. *Journal of virology* 86, 12923-12932.

596 Ray, R., Khanna, A., Lagging, L.M., Meyer, K., Choo, Q.L., Ralston, R., Houghton, M., Becherer, P.R., 1994.  
597 Peptide immunogen mimicry of putative E1 glycoprotein-specific epitopes in hepatitis C virus. *Journal of*  
598 *virology* 68, 4420-4426.

599 Siemoneit, K., Cardoso Mda, S., Koerner, K., Wolpl, A., Kubanek, B., 1995. Human monoclonal antibodies  
600 for the immunological characterization of a highly conserved protein domain of the hepatitis C virus  
601 glycoprotein E1. *Clinical and experimental immunology* 101, 278-283.

602 Smith, H.L., Chung, R.T., Mantry, P., Chapman, W., Curry, M.P., Schiano, T.D., Boucher, E., Cheslock, P.,  
603 Wang, Y., Molrine, D.C., 2017. Prevention of allograft HCV recurrence with peri-transplant human  
604 monoclonal antibody MBL-HCV1 combined with a single oral direct-acting antiviral: A proof-of-concept  
605 study. *J Viral Hepat* 24, 197-206.

606 Swann, R.E., Mandalou, P., Robinson, M.W., Ow, M.M., Fong, S.K., McLauchlan, J., Patel, A.H., Cramp,  
607 M.E., 2016. Anti-envelope antibody responses in individuals at high risk of hepatitis C virus who resist  
608 infection. *J Viral Hepat* 23, 873-880.

59  
60  
61  
62  
63  
64  
65

609 Tabll, A., Abbas, A.T., El-Kafrawy, S., Wahid, A., 2015. Monoclonal antibodies: Principles and applications  
610 of immunodiagnosis and immunotherapy for hepatitis C virus. *World journal of hepatology* 7, 2369-  
611 2383.

612 Tarr, A.W., Owsianka, A.M., Timms, J.M., McClure, C.P., Brown, R.J.P., Hickling, T.P., Pietschmann, T.,  
613 Bartenschlager, R., Patel, A.H., Ball, J.K., 2006. Characterization of the hepatitis C virus E2 epitope  
614 defined by the broadly neutralizing monoclonal antibody AP33. *Hepatology* 43, 592-601.

615 Tarr, A.W., Urbanowicz, R.A., Jayaraj, D., Brown, R.J.P., McKeating, J.A., Irving, W.L., Ball, J.K., 2012.  
616 Naturally Occurring Antibodies That Recognize Linear Epitopes in the Amino Terminus of the Hepatitis C  
617 Virus E2 Protein Confer Noninterfering, Additive Neutralization. *Journal of virology* 86, 2739-2749.

618 Vanwollegem, T., Bukh, J., Meuleman, P., Desombere, I., Meunier, J.C., Alter, H., Purcell, R.H., Leroux-  
619 Roels, G., 2008. Polyclonal immunoglobulins from a chronic hepatitis C virus patient protect human liver-  
620 chimeric mice from infection with a homologous hepatitis C virus strain. *Hepatology* 47, 1846-1855.

621 Wahid, A., Helle, F., Descamps, V., Duverlie, G., Penin, F., Dubuisson, J., 2013. Disulfide bonds in  
622 hepatitis C virus glycoprotein E1 control the assembly and entry functions of E2 glycoprotein. *Journal of*  
623 *virology* 87, 1605-1617.

624 Wakita, T., Pietschmann, T., Kato, T., Date, T., Miyamoto, M., Zhao, Z.J., Murthy, K., Habermann, A.,  
625 Krausslich, H.G., Mizokami, M., Bartenschlager, R., Liang, T.J., 2005. Production of infectious hepatitis C  
626 virus in tissue culture from a cloned viral genome. *Nat Med* 11, 791-796.

627 Wang, Y., Keck, Z.Y., Fong, S.K., 2011a. Neutralizing antibody response to hepatitis C virus. *Viruses* 3,  
628 2127-2145.

629 Wang, Y., Keck, Z.Y., Saha, A., Xia, J., Conrad, F., Lou, J., Eckart, M., Marks, J.D., Fong, S.K., 2011b.  
630 Affinity maturation to improve human monoclonal antibody neutralization potency and breadth against  
631 hepatitis C virus. *The Journal of biological chemistry* 286, 44218-44233.

632 Zhang, P., Wu, C.G., Mihalik, K., Virata-Theimer, M.L., Yu, M.Y., Alter, H.J., Feinstone, S.M., 2007.  
633 Hepatitis C virus epitope-specific neutralizing antibodies in Igs prepared from human plasma. *Proc Natl*  
634 *Acad Sci U S A* 104, 8449-8454.

635 Zhang, P., Zhong, L., Struble, E.B., Watanabe, H., Kachko, A., Mihalik, K., Virata-Theimer, M.L., Alter, H.J.,  
636 Feinstone, S., Major, M., 2009. Depletion of interfering antibodies in chronic hepatitis C patients and  
637 vaccinated chimpanzees reveals broad cross-genotype neutralizing activity. *Proc Natl Acad Sci U S A* 106,  
638 7537-7541.

35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



641 **Figure legends**

1  
2 642 **Fig. 1. Phylogenetic tree for new E1E2 isolates.** Following isolation and sequencing of full length E1E2  
3  
4 643 from Belgian patients infected with HCV, sequences were multiple aligned together with previously  
5  
6 644 published sequences retrieved from the HCV database. Neighbor joining phylogenetic tree was  
7  
8 645 constructed using CLC main workbench based on the whole E1E2 amino acid sequence. Analysis shows  
9  
10 646 that all Belgian isolates belong to genotype 1b, except for one isolate genotyped as 1a. The 28 Belgian  
11  
12 647 genotype 1a and 1b isolates are highlighted in red. Bootstrap values are indicated.

13 648 **Fig. 2. Binding of mAb A6 to recombinant E1E2.** Constructs encoding the full length E1E2 of HCV  
14  
15 649 genotypes 1-6 were used for expression in 293T cells. GNA-coated ELISA plates were loaded with cell  
16  
17 650 lysate containing E1E2. (A and B) Binding of serially diluted mAb A6 was assessed using ELISA plates  
18  
19 651 coated with E1E2 of three genotype 1a isolates (H77, UKN1A20.8, UKN1A14.38), one genotype 1b  
20  
21 652 (UKN1B12.16) and two genotype 2a isolates (UKN2A1.2 and JFH1). (C) mAb A6 binding to the  
22  
23 653 envelope proteins of multiple other isolates of different genotypes was evaluated at fixed antibody  
24  
25 654 concentration (10 µg/mL). Lysate of non-transfected 293T cells was included as a negative control. All  
26  
27 655 conditions were performed in duplicate and error bars represent the standard deviation.

28  
29 656 **Fig. 3. Epitope mapping of mAb A6.** (A) To identify the epitope within E1E2 recognized by mAb A6, a  
30  
31 657 custom-made chip (PEPperPRINT) with printed overlapping peptides spanning the complete E1E2-  
32  
33 658 sequence was used. After incubation with mAb A6 and an Alexa647-conjugated goat anti-human IgG,  
34  
35 659 spots corresponding to 5 different E1-specific peptides appeared (highlighted with black square). The  
36  
37 660 reactive peptides were AA 225-239, AA 226-240, AA 227-241, AA 228-242 and AA 229-243. The  
38  
39 661 positive spots in the outer rim of the chip are controls that are used for localization purposes only. (B)  
40  
41 662 Standard peptide ELISA was performed using synthetic peptides spanning the region AA 222-246 (25  
42  
43 663 µg/mL). mAb A6 was used at 10 µg/mL and binding activity is shown by OD values measured at 450nm.  
44  
45 664 (C) A peptide competition assay was performed on ELISA plates coated with recombinant E1E2 (H77  
46  
47 665 isolate), using a fixed concentration of mAb A6 (20 µg/mL) in the presence of serial dilutions of the  
48  
49 666 indicated synthetic peptides. Binding of mAb A6 is denoted as % of maximal binding in the absence of  
50  
51 667 synthetic peptide. The amino acid sequence of the peptides covering the region AA 222-246 used in (A-C)  
52  
53 668 is indicated.

669 **Fig. 4. Reactivity of mAb A6 to glycosylated and deglycosylated E1E2 protein.** Lysate of 293T cells  
1  
2670 expressing H77 E1E2 was treated with PNGase F or Endo H. Following 12% SDS-PAGE, proteins were  
3  
4671 transferred to a nitrocellulose membrane and revealed using mAb A6 as primary antibody. Positive bands  
5  
6672 were visible at a molecular weight around 20-22 kDa in case of PNGase F and Endo H-treated cell lysate  
7  
8673 (lane 3 and 4), which correspond to deglycosylated E1, while the untreated lysate showed reactive bands  
9  
10674 at around 25-32 kDa (lane 2). The position and molecular weights of protein standard are presented in  
11  
12675 kDa (lane M). The deglycosylated E1 protein is indicated by an asterisk.

13676 **Fig. 5. Antibody A6 targets a linear epitope.** To investigate if the region targeted by mAb A6 is  
14  
15676 conformation sensitive or not, cell lysate containing E1E2 of isolate H77 (genotype 1a) (A) or JFH1  
16  
17677 (genotype 2a) (B) was denatured by incubation with SDS and DTT at 56°C. Native and denatured cell  
18  
19678 lysate were coated on ELISA plates and binding of mAb A6 (5 µg/mL) was assessed. In parallel, murine  
20  
21679 mAb A4 and human mAb 1:7 were included as controls for linear and conformational epitopes  
22  
23680 respectively. All conditions were performed in duplicate and results are shown as the mean +/- standard  
24  
25681 deviation.  
26  
27682

28683 **Fig. 6. Multiple alignment of the region corresponding to AA 192-250 and AA 310-330 of HCV E1.**  
29  
30684 The E1 sequence of all HCV isolates used in the current study was retrieved from GenBank and the HCV  
31  
32685 database and multiple aligned using BioEdit software. The epitopes targeted by antibodies H-111, A4, A6,  
33  
34686 IGH505 and IGH526 are indicated using colored squares. The dots indicate conservation of the residue  
35  
36687 relative to the reference prototype H77 (genotype 1a). The lower part of the alignment shows the  
37  
38688 consensus sequence and the percentage of overall conservation calculated using CLC Main Workbench.  
39  
40689

41  
42689 **Fig. 7. HCVpp and HCVcc neutralization.** In order to verify the neutralizing capacity of mAb A6,  
43  
44690 HCVpp expressing E1E2 of H77 (genotype 1a) (A), UG-P09-779 (genotype 1b) (B), UG-P10-252  
45  
46691 (genotype 1b) (C), UG-P12-763 (genotype 1b) (D) and JFH1 (genotype 2a) (E) were incubated for 1 hour  
47  
48692 at 37°C with mAb A6 (100 µg/mL) or with mAb MRCT10 (10 µg/mL) before addition to Hep3B cells.  
49  
50693 After 72 hours, HCVpp infection was assessed by quantifying the luciferase reporter gene expression and  
51  
52694 the results were normalized to untreated controls (Ctrl). Chimeric HCVcc of genotypes 1a/2a (F), 1b/2a  
53  
54695 (G) and 2a/2a (H) were pre-incubated with mAb A6 at a concentration of 100 µg/mL or with 10 µg/mL of  
55  
56696 mAb MRCT10 for 1 hour at 37 °C before exposure to Huh7.5RFP cells. The number of HCV-infected  
57  
58  
59  
60  
61  
62  
63  
64  
65

697 foci was counted using NS5A-specific antibody staining. The infection was normalized to that in cultures  
1  
2698 that were exposed to virus only (Ctrl). All conditions were performed in triplicate and results are shown  
3  
4699 as the mean +/- standard deviation.  
5

6700 **Fig. 8. mAb A6-based visualization of infected Huh7.5RFP cells using immunofluorescence.**  
7

8701 H77c/JFH1 infected Huh7.5RFP cells (upper panel) and non-infected cells (-ve; lower panel) were  
9  
10702 incubated overnight with mAb A6 (middle column), after which Alexa 647-conjugated goat-anti-human  
11  
12703 IgG was added. As a positive control, the same cells were stained with anti-NS5A mAb 9E10 in  
13  
14704 combination with Alexa 488 conjugated goat-anti-mouse IgG (right column). DAPI was used as  
15  
16705 counterstain (left column).  
17  
18

19706 **Fig. 9. mAb A6 does not interfere with neutralizing antibodies.** (A) Competition ELISA was  
20

21  
22707 performed using biotinylated AP33 and E1E2 of isolate H77 as a coating antigen. Serial dilutions of mAb  
23  
24708 A6 were used starting at 30 µg/ml and the binding of biotinylated AP33 was measured using HRP-  
25  
26709 conjugated streptavidin. As a positive control for inhibition, serial dilutions of MRCT10 were used in  
27  
28710 parallel. To investigate the interference with the neutralizing potential of anti-E2 antibodies, HCVpp from  
29  
30711 genotype 1a (H77) (B) and 2a (JFH1) (C) were used. Co-incubation of mAb A6 (100 µg/mL) and one of  
31  
32712 anti-E1 or anti-E2 mAbs (IGH526, MRCT10<sub>1</sub> 1:7, HC84.26, CBH-7, HC-1AM; 10 µg/mL) and HCVpp  
33  
34  
35713 was done one hour before addition to Hep3B cells. The luciferase activity was determined and the  
36  
37714 percentage of neutralization was calculated relative to infection in the absence of antibody (Ctrl). All  
38  
39715 conditions were performed in duplicate for ELISA and 5 replicates for HCVpp inhibition. Results are  
40  
41  
42716 shown as the mean +/- standard deviation.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Figure 1  
[Click here to download high resolution image](#)

Figure 1




Figure 2






Figure 3


A




B



C




**Figure 4**




# Figure 5

**A**



**B**



 Native E1E2


 Denatured E1E2



Figure 6  
[Click here to download high resolution image](#)

Figure 6







Figure 7



# Figure 8



## Figure 9



**Table 1: Names, genotypes and accession numbers for the E1E2 isolates used in ELISA.**

<b>Isolate ID</b>	<b>Genotype</b>	<b>Accession number</b>
<b>H77</b>	1a	AF011751
<b>UKN1A20.8</b>	1a	EU155192
<b>UKN1A14.38</b>	1a	AY734971
<b>J4</b>	1b	AF054250
<b>UKN1B5.23</b>	1b	AY734976
<b>UKN1B12.16</b>	1b	AY734974
<b>P5VD</b>	1b	(Fafi-Kremer et al., 2010)
<b>P5VE</b>	1b	(Fafi-Kremer et al., 2010)
<b>P5VF</b>	1b	(Fafi-Kremer et al., 2010)
<b>UG-P08-503</b>	1b	(Desombere et al., 2016)
<b>UG-P09-779</b>	1b	(Desombere et al., 2016)
<b>UG-P10-252</b>	1b	(Desombere et al., 2016)
<b>UG-P12-763</b>	1b	(Desombere et al., 2016)
<b>JFH1</b>	2a	AB047639
<b>UKN2A1.2</b>	2a	AY734977
<b>UKN2B2.8</b>	2b	AY734983
<b>S52</b>	3a	GU814263
<b>UKN4.11.1</b>	4	AY734986
<b>UKN5.14.4</b>	5	AY785283
<b>UKN6.5.8</b>	6	EF427671

**Supplementary Figure 1**

[Click here to download Supplementary Material \(To be Published\): Supplementary Figure 1.pdf](#)

**Supplementary File 1**

[Click here to download Supplementary Material \(To be Published\): Supplementary file 1.xlsx](#)

**Agreements additional co-author**

**[Click here to download Supplementary Material \(Not to be Published\): Agreements additional coauthor.pdf](#)**