

SCIENTIFIC REPORTS

OPEN

The world's earliest Aral-Sea type disaster: the decline of the Loulan Kingdom in the Tarim Basin

Steffen Mischke¹, Chenglin Liu², Jiafu Zhang³, Chengjun Zhang⁴, Hua Zhang², Pengcheng Jiao² & Birgit Plessen⁵

Received: 25 August 2016
Accepted: 19 January 2017
Published: 27 February 2017

Remnants of cities and farmlands in China's hyperarid Tarim Basin indicate that environmental conditions were significantly wetter two millennia ago in a region which is barren desert today. Historical documents and age data of organic remains show that the Loulan Kingdom flourished during the Han Dynasty (206 BCE–220 CE) but was abandoned between its end and 645 CE. Previous archaeological, geomorphological and geological studies suggest that deteriorating climate conditions led to the abandonment of the ancient desert cities. Based on analyses of lake sediments from Lop Nur in the eastern Tarim Basin and a review of published records, we show that the Loulan Kingdom decline resulted from a man-made environmental disaster comparable to the recent Aral Sea crisis rather than from changing climate. Lop Nur and other lakes within the Han Dynasty realm experienced rapidly declining water levels or even desiccation whilst lakes in adjacent regions recorded rising levels and relatively wet conditions during the time of the Loulan Kingdom decline. Water withdrawal for irrigation farming in the middle reaches of rivers likely caused water shortage downstream and eventually the widespread deterioration of desert oases a long time before man initiated the Aral Sea disaster in the 1960s.

The Tarim Basin (Xinjiang Province) in northwestern China hosts the Taklamakan Sand Sea, the second largest sand desert of the world. Precipitation in the basin is lower than 20 mm a^{-1} but streams from towering mountain ranges in the north, west and south flow along its margins and charged the closed-basin lake Lop Nur until the 1970s when the lake eventually desiccated^{1–4}. According to historical documents of the Han Dynasty, the lake was once called “Puchang Sea” and estimated to have covered 17,000 to 50,000 km^2 which places the Han-period lake in the range between today's Lake Michigan (the 5th largest lake on Earth) and Lake Balkhash (the 15th largest lake⁵). Ruins of ancient cities, remnants of fields, irrigation channels and roads have been found in the desert region^{1,6,7}. Most of these cities existed in the northwest and west of the lake where the Tarim and Konqi Rivers flowed towards the ancient Lop Nur (Fig. 1). Loulan at the ancient Silk Road is among the most famous ruined cities, reported in historical documents almost 2000 years ago⁸ and re-discovered by the Swedish researcher and traveller Sven Hedin in 1899¹. The flourishing Loulan Civilization was an important trade centre at the ancient Silk Road established at the beginning of the Han Dynasty⁹. Some of the well preserved mummies from the Tarim Basin are significantly older with ages up to 3800 years¹⁰, but most of the cities and farmlands were established during the first expansion of China's mainland towards the west during the Han Dynasty^{1,6,7}. This westward expansion of the Chinese Empire was accompanied by a large population migration and intensification of irrigation farming along the rivers in mountain forelands in the Tarim Basin and the Hexi Corridor (Gansu Province) farther to the east^{1,5,11}. The expansion towards the west was enabled by stationed troops which not only secured the occupation but were used for the establishment of fortified cities and trade posts^{12,13}. Even more important, the soldiers were the ones responsible for the large-scale reclamation of arable land and the establishment of irrigation farming in many regions which had not been intensively cultivated before^{14–16}. Large-scale irrigation farming started in the Tarim Basin in the middle of the first century BCE⁸. According to the historical book by Ban Gu⁸, a total of 300,830 people lived in more than 20 kingdoms upstream of Lop Nur in the Tarim Basin including

¹Faculty of Earth Sciences, University of Iceland, 101 Reykjavík, Iceland. ²Institute of Mineral Resources, Chinese Academy of Geological Sciences, Beijing 100037, China. ³MOE Laboratory for Earth Surface Processes, Department of Geography, College of Urban and Environmental Sciences, Peking University, Beijing 100871, China. ⁴School of Earth Sciences and Key Laboratory of Mineral Resources in Western China, Lanzhou University, Lanzhou 730000, China. ⁵Helmholtz Centre Potsdam, German Research Centre for Geosciences, Potsdam 14473, Germany. Correspondence and requests for materials should be addressed to S.M. (email: smi@hi.is)




Figure 1. The ancient Lop Nur Lake in the Tarim Basin. (a) Tarim Basin with rivers, Lop Nur (black arrow) and ancient sites (S - Sampula, Z - Zaghunluq, R - Ruoqiang, X - Xiaohe, G - Gumugou), area of irrigated farmland during the Han Dynasty^{1,5} indicated by grey shading; inset shows location in China. Frame in the right indicates area shown in Fig. 1b. Background map was generated with data available from the Chinese Geospatial Data Cloud using ESRI's ArcGIS (version 10.2; <http://www.esri.com/>). (b) The dry Lop Nur Basin with position of pit section (black dot) and Loulan and other sites. Ancient lake extent¹ indicated by red dotted line. Background map generated with ESRI's ArcGIS (version 10.2) and Landsat data (NASA Landsat Program, 2006, Landsat ETM+ scenes 218–312 and 218–300, Orthorectified, USGS, Sioux Falls, 03/05/2006 and 04/15/2006, respectively; <http://glcfapp.glc.fumd.edu:8080/esdi/index.jsp>).

Loulan sometime during the Han Dynasty. Among them, the Loulan Kingdom included 14,100 inhabitants⁸. Agriculture was significantly intensified even in remote desert oases such as the lower Keriya and Niya River valleys where land use was restricted to the use of natural oases before^{5,17}. Cultivated areas were significantly larger in the Tarim Basin during the Han Dynasty than during the Qing Dynasty (1644–1911) or even around 1950⁵. The extensive transformation of alluvial fans and plains to irrigated fields in the Tarim Basin and Hexi Corridor not only supported the migration of people from the previous mainland of the Chinese Empire during the Han Dynasty but led also to intensive trade along the Silk Road two millennia ago^{1,2,5–7,12}. Historical documents describe diminishing river discharge and subsequent reductions in soldier's grain rations after ca. 270 CE^{1,18}. The cities had been abandoned when the monk Xuanzang travelled in the region in 645 CE^{1,18}. Correspondingly, radiocarbon dating results for organic remains suggest that the abandonment of the cities started not earlier than 230 ± 66 CE⁶, leaving a period from ca. 200 to 645 CE for the decline of the Loulan Civilization. We investigated




Figure 2. Grain size and stable isotope data for the pit section (upper 4 m) in the Lop Nur depression. Grain size frequency distributions shown for sand- (upper) and silt-dominated (lower) sediments. Red dots indicate position of samples used for grain-shape analysis (Fig. S1). $\delta^{18}\text{O}$ values for bulk carbonate and ostracod shells (green square – *Limnocythere inopinata*, blue triangle – *Eucypris mareotica*, red inverted triangle – *Cyprideis torosa*). Age data of optically stimulated luminescence dating at left¹⁹.

the sediments of Lop Nur and compared the results with palaeoclimate records from other lakes in the region to examine whether climate change triggered the deterioration of the flourishing desert oases suggested as result of previous studies^{1,2,5–7,9}.

Results and Discussion

A 5.7-m deep pit section in the centre of the Lop Nur Basin provides grain-size and stable isotope records for the past ca. 9000 years^{19,20}. Poor to very poorly sorted sediments with a mean grain size in the silt fraction prevail apart from two layers of coarser sediments centered at 0.70 and 0.50 m depth (Fig. 2). The sand layers with grain size and shape similar to aeolian sands in the region were formed ca. 1800 and 1000 years ago (Fig. S1). The two layers of coarser sediments show partly better sorting than sediments beneath or above (Fig. 2) although both include a significant fraction of silt-sized material (39% for the lower and 14% for the upper layer)²⁰. The lower layer of coarser sediments is underlain by a 0.20 m thick sequence of sediments with increasing mean grain sizes. The gradual increase from a mean grain size of fine silt at 0.90 m to fine sand at 0.70 m probably indicates a continuously increasing proportion of aeolian sediment accumulation and steady lake level decrease. In contrast, the upper layer of coarser sediments with a mean grain size of medium sand covers fine-grained sediments with a mean grain size of fine silt without transitional grain size change (Fig. 2). The observed abrupt grain size increase implies that the depositional setting either changed abruptly, or that transitional sediments with inverse grading were possibly accumulated at first during a gradual lake-level decline but were subsequently removed by wave-driven currents before the accumulation of coarser sediments. Soluble salt concentrations in the sediments beneath and in the two coarser layers are increasing and relatively high, respectively, and provide supporting evidence that the coarser sediment layers were accumulated during periods of low lake levels and more saline conditions in Lop Nur²⁰.

The stable oxygen isotope values for ostracod shells are in the same range as the $\delta^{18}\text{O}$ values for bulk carbonate from the same stratigraphic levels in the section (Fig. 2). Thus, the $\delta^{18}\text{O}$ record for bulk carbonate represents a reliable record of the relative $\delta^{18}\text{O}$ changes of past lake waters. The increasing $\delta^{18}\text{O}$ values for bulk carbonate at 0.90 m depth indicate a rapid reduction of inflows to the lake and a salinity rise ca. 2300 years ago further amplified ca. 1000 years ago (Fig. 2). The drop of the $\delta^{18}\text{O}$ values at 0.46 m probably reflects a higher discharge which entered Lop Nur again, and a recovery of the lake system for some time before the final desiccation in the last century. Earlier in the stable isotope record, the lowest $\delta^{18}\text{O}$ values 8500 years ago indicate the wettest conditions in the Lop Nur region. The large increase at 2.00 m depth suggests a significant decrease in inflows and moisture availability ca. 7000 years ago. However, the stable isotope data from Lop Nur suggest that the lake level started




Figure 3. Comparison of climate inferences for the last 12,000 years based on lake records from Central Asia. Data are arranged to show wetter conditions towards the top. Lake numbers refer to information source in reference list.

to decline as early as 2300 years ago whilst the grain-size data indicate very low lake levels and near-desiccation periods starting ca. 1800 years ago. Sand-dominated sediments were not accumulated during the entire preceding period between 9000 and 1800 years which suggests that a fundamental change in the depositional environment occurred ca. 1800 years ago.

Accumulation of aeolian sand in the basin centre almost two millennia ago roughly corresponds to the time of reported grain-ratio reductions in the Loulan Kingdom¹ and the first abandonment of Tarim Basin cities inferred from radiocarbon dating⁶. Thus, Lop Nur apparently experienced lowest lake levels or possibly first periods of desiccation more or less during the time of the Loulan Civilization decline at the end of the Han Dynasty (220 CE) and afterwards.

The intensification of farming was at least locally accompanied by an inappropriate use of water resources which resulted in decreasing natural vegetation, mobilization of aeolian sand, and soil and water salinization^{13,14}. Accordingly, shortage of water and desertification of oases in the Hexi Corridor was reported for a period ca. 2100 years ago¹³. Sanctions against the felling of trees to combat desertification in the Loulan Kingdom were reported for the 3rd century CE²¹; but such evidence for desertification as a result of inappropriate use of water resources during the Han Dynasty is generally rare. However, the systematic assessment of geographical information available in historical documents and literature from the Han Dynasty shows that the farmland area was significantly larger upstream of Lop Nur and around Bosten Lake than it was in the last 400 years⁵. Remnants of extensive irrigated fields and irrigation channels from the Han Dynasty have been discovered near Loulan and in the Ruoqiang region^{12,22,23} (Fig. 1). Thus, water withdrawal for irrigation farming in the middle reaches of the Tarim and Konqi Rivers must have occurred on a large scale and probably caused water shortage in the lower reaches and declining Lop Nur lake levels.

In contrast to changing climate conditions, the decline of the Loulan Civilization and abandonment of the desert cities may have resulted from the intensification of irrigation farming in the middle reaches of the Tarim River and its tributaries and water shortage further downstream. Other regional lake records from within and outside the realm of the Han Empire potentially provide a clue to differentiate between the regional climate conditions and possible human impact on the environment during the period of the Loulan Kingdom decline.

Pollen preserved in lake sediments indicate relatively wet conditions in the 3rd–7th century in the north of Lop Nur at lakes Sayram²⁴, Aibi²⁵, Wulungu²⁶ and Balikun²⁷ (Figs 3 and 4). In contrast, a drop in the *Artemisia/Chenopodiaceae* (A/C) moisture index at Bosten Lake²⁸ around 1800 years ago and the desiccation of the Eastern Juyan Lake at ca. 1700 years²⁹ indicate drier conditions. Drier conditions at Bosten Lake were also inferred from geochemical data suggesting a drop of the lake level at ca. 1900 years ago³⁰. Stable oxygen isotope data of lake carbonates or hydrogen isotopes of alkanolic acids indicate relatively wet conditions for lakes in the west^{31–33} and north³⁴ of Lop Nur (Figs 3 and 4). The level of Issyk-Kul in the Tianshan Mountains was generally low during the late Holocene in comparison to the early Holocene but the slight $\delta^{18}\text{O}$ decrease after 1800 years was assigned to a lake level highstand 16 m above the present level³³. Lake Manas further in the northeast experienced a shift from a fluvial depositional setting to a lake environment ca. 2000 years ago³⁴, providing further support for relatively wet conditions (Figs 3 and 4). The levels of Hurlig³⁵, Qinghai³⁶ and Hala Lake³⁷ in the southeast of Lop Nur




Figure 4. Positions of lake records within the Han Dynasty (206 BCE–220 CE) realm (green shading Han area) and outside. Lake level declines or periods of desiccation during the decline of the Loulan Kingdom were recorded at lakes indicated by red circles. The dotted line marks the realm of the preceding empire of the Qin Dynasty (221–206 BCE). Lake numbers refer to information source in reference list (*information for Bosten Lake also provided by ref. 30; HC – Hexi Corridor, G – Guliya ice core). Background map generated using a freely available map (https://commons.wikimedia.org/wiki/File:China_topography_full_res.jpg) and CorelDRAW version 12 (<http://www.coreldraw.com/>).

were relatively high in the 3rd–7th century whilst those of Zhuyezhe Lake³⁸ declined (Figs 3 and 4). The resulting spatial distribution of lake conditions in the 3rd–7th century shows low levels or even desiccation for lakes Lop Nur, Bosten, Eastern Juyan and Zhuyezhe (Fig. 4). Relatively wet conditions were recorded at all other locations. This pattern corresponds to lowered lake levels or lake desiccation within the Han Empire realm, and relatively wet conditions adjacent to this region. Thus, regional climate conditions were generally relatively wet during the Loulan Kingdom decline. The inference of wetter conditions in western China in the 3rd–7th century is also supported by higher ice-accumulation rates on the Guliya ice cap in the Kunlun Mountains south of the Tarim Basin³⁹ (Fig. 4) and increased winter precipitation in the Kyrgyz Tianshan⁴⁰. Thus, declining levels of the lakes Lop Nur, Bosten, Eastern Juyan and Zhuyezhe most likely resulted from water withdrawal during the Han Empire intensification of irrigation farming along the mountain forelands. The decline of the Loulan Civilization was not a consequence of climate change but probably the result of a major man-made environmental disaster comparable to the recent Aral Sea crisis.

The reviewed lake records^{24–38} (Figs 3 and 4) and the ice core record from the nearby Kunlun Mountains³⁹ indicate that climate conditions were relatively wet at the end of the Han Dynasty. If water withdrawal for irrigation farming in the mountain forelands of the Tarim Basin during the Han Dynasty caused the decline of the Loulan Civilization, a migration of ancient desert oases from more central positions towards the basin margins is expected over the Han period. Age data for sub-recent remains of vegetation and traces of human activities in the Tarim Basin will provide information about periods favourable for the growth of riverine vegetation and human activities. Thus, the lack of such age data only provides indirect evidence for less favourable conditions. Dating results for vegetation remains and remnants of settlements and agriculture in the Tarim Basin are generally rare and insufficient for Loulan and the vicinity of Lop Nur, but dated organic materials from the lower sections of the Keriya and Niya Rivers allow first assessments of available data from two river systems in the central Tarim Basin^{7,41} (Fig. 1).

Radiocarbon age data were presented for organic remains from the lowermost 100 km of the Keriya River⁷. Organic remains (sheep manure, *Populus* stems from house or city wall remnants, chopped *Populus* stems) representing direct or indirect evidence for human activities show a tendency of older ages further downstream (as old as 2600 years) and younger ages (not older than 2000 years) upstream⁷ (Fig. S2). *Tamarix* roots and *Populus* stems recovered from river terraces possibly represent natural gallery forest vegetation of a desert river. The six available age data cluster in a wide range with oldest and youngest age data for the most distal positions of the Keriya River. The data for the *Tamarix* roots and *Populus* stems possibly indicate that natural vegetation was able to survive in favorable positions with near-surface groundwater levels. The small data set of 14 dated organic remains does not include materials with an age between ca. 1600 and 600 years (Fig. S2). The three age data younger than ca. 600 years possibly indicate that climate conditions and/or human impact were apparently not supporting sufficient runoff and vegetation growth in the Keriya River region until ca. 600 years ago.

Exact locations of radiocarbon dating samples from Niya River were not presented by the authors but samples were grouped into northern, middle and southern parts⁴¹. Remains of trees and wooden building materials from the ancient Niya site to the east of the Keriya River similarly provided oldest ages in its northern part (ca. 2400 years) and successively younger ages in its middle (ca. 1900 years) and southern parts (ca. 1500 years⁴¹; Fig. S3).

Ancient sites from both rivers in the central Tarim Basin indicate a migration of human activities upstream which could have resulted also from climate change. However, the implied migration of ancient desert oases towards more marginal positions in the Tarim Basin over the duration of the Han Dynasty is at least also consistent with our assumed shortage of runoff farther downstream as a result of water withdrawal for irrigation farming in the middle reaches of the rivers. Based on the decline of lake levels or even desiccation of lakes within the Han Empire realm in contrast to relatively wet conditions in adjacent regions, we conclude that the Loulan Civilization decline was caused by intensified consumption of water for irrigation farming upstream of the oases and represents an early man-made environmental disaster.

It is puzzling that pre-Han age data for organic remains from ancient sites along the courses of the Keriya and Niya Rivers are confined to the lowermost reaches (Figs S2 and S3). Apparently, settlements and land use prior to the Han Dynasty were restricted to the lowermost river sections. We only may speculate that hazardous flooding, frequent river-channel displacements and/or poorer edaphic conditions (coarser and less nutrient-rich sediments) inhibited the occupation of such locations as long as water was sufficient farther downstream.

It also remains unclear whether significantly older ancient sites further downstream in the Tarim Basin (Gumugou and Xiaohe in its eastern part, 4000–3500 years) and younger but mostly pre-Han sites at its margins (Zaghunluq and Sampula in its south, ca. 2700–2000 years) possibly represent a climate or man-controlled shift of water resources in northwestern China prior to the Han Dynasty, or whether this tendency is not representative due to insufficient data^{42,43} (Fig. 1).

Materials and Methods

Grain size analysis of Lop Nur section sediments. Sediment samples from the pit section in the Lop Nur Basin were used for grain size analysis at Peking University in Beijing. Samples were treated with 30% H₂O₂ at room temperature to remove organic material, with 10% HCl to remove carbonates and deflocculated using a solution of 0.01 M (NaPO₃)₆. Following each step, the sample solution was boiled to promote the removal of remaining organic matter or carbonates. After 3-min sonication in deionized water, the sample solution was analyzed using a Malvern Mastersizer 2000 grain-size analyzer with 0.015 μm resolution and a range of 0.002–2000 μm with a relative error of <1%. The mean grain size was calculated using the Malvern software (Ver. 5.12 C), and sorting was examined with GRADISTAT⁴⁴.

Grain shape examination of Lop Nur section sediments. Grain shape was analyzed for sediments from the pit section collected at two stratigraphic levels. One sample from 0.87–0.86 cm depth was regarded to represent the bulk of the silt-dominated sediments of the section and a second sample from 0.49–0.48 cm depth represents the coarser, sand-dominated sediments from the upper part of the section. Subsamples were treated with 3% H₂O₂ for 48 hours and washed through 100 and 250 μm sieves. Sand grains remaining on the 250 μm sieve were dried, evenly spread on a stub and examined using a Zeiss Supra 40 VP Scanning Electron Microscope (SEM) at Freie Universität Berlin. SEM images of the first 25 separately placed sand grains (excluding overlaying grains) were obtained.

Stable isotope analyses of bulk carbonate and ostracod shells. Stable isotope analysis of bulk carbonate was carried out using a Finnigan MAT 252 isotopic ratio mass spectrometer (IRMS) at the Lanzhou Institute of Geology, Chinese Academy of Sciences. Bulk samples were powdered and baked at 300 °C for 1 hour in a vacuum system, before the samples were reacted with 100% H₃PO₄ at 90 °C. The produced CO₂ was trapped in a cold finger with liquid nitrogen and transferred into the mass spectrometer for the stable isotope analysis. The reliability of bulk carbonate for stable isotope analysis was tested at stratigraphic sections containing ostracod shells. Ostracod shells form within a few hours or days during moulting and reflect the stable isotope composition of host waters biased by a vital effect of commonly 1–2‰^{45,46}. For stable isotope analysis of ostracod shells, sub-samples were treated with 3% H₂O₂ for 48 hours and washed through 100 and 250 μm sieves. Ostracod shells were picked from the >250 μm sieve residues. On average, 50 μg of shell material (11 shells of *Limnocythere inopinata*, six shells of *Eucypris mareotica*, two shells of *Cyprideis torosa*) was reacted with 103% H₃PO₄ at 70 °C in an automated carbonate preparation device (KIEL IV) coupled to a Finnigan MAT 253 IRMS at the German Research Centre for Geosciences in Potsdam. Isotopic ratios are given in delta notation relative to VPDB (Vienna Pee Dee Belemnite) calibrated with NBS-19 (δ¹⁸O = −2.20‰). The analytical precision is better than ±0.06‰ for the ostracod shell and ±0.3‰ for the bulk carbonate samples (1σ).

Review of lake record data. Available lake records from northwestern China and adjacent regions were examined to distinguish between climate change presumably affecting a larger region and human impact expected to be higher in the realm of the Han Dynasty in comparison to the regions outside the Chinese Empire. Inferences from the lake records are based on four different types of evidence: (1) pollen data (six records), (2) stable isotope data (four records), (3) directly dated shorelines above the present lake level (three records), and (4) the sedimentological and geochemical characteristics of lake sediments (two records).

(1) Pollen data were presented for the Lakes Sayram²⁴, Aibi²⁵, Wulungu (Ulungur²⁶), Bosten²⁸, Balikun²⁷ and the Eastern Juyan Lake²⁹. The *Artemisia*/Chenopodiaceae ratio was used as climate proxy for the records from Sayram, Bosten and Balikun with higher ratios indicating steppe vegetation and wetter conditions in the arid to semi-arid region of Central Asia⁴⁷. Higher pollen concentrations in the lake sediments were used as a proxy of higher vegetation density and wetter conditions around Lake Aibi²⁵. Pollen and geochemical data of lake sediments were both assessed for Lake Wulungu and used to reconstruct climate conditions in the region and the lake level history²⁶ which was later on confirmed through salinity and lake level inferences based on the distribution of micro-crustaceans in the modern and Holocene sediments of Lake Wulungu⁴⁸. Holocene pollen data for the Eastern Juyan Lake were compared with modern pollen assemblages and used for a quantitative reconstruction

of past precipitation in the region²⁹. The lake desiccated 1700 years ago and aeolian sands were accumulated in the former lake basin²⁹.

(2) Stable isotope records were provided from Lakes Karakul³¹, Karakuli³², Issyk-Kul³³ and Manas³⁴. The Lakes Karakul, Issyk-Kul and Manas are closed basin lakes; and $\delta^{18}\text{O}$ values of lake carbonates represent the general hydrological budget of the lake with higher $\delta^{18}\text{O}$ values indicating more arid conditions and lower values representing more humid conditions⁴⁹. Lake Karakuli is an open-basin lake and lower δD values for plant leaf wax from the lake sediments were interpreted to indicate cooler and wetter climate conditions³².

(3) Ancient shorelines above current lake levels were dated at Lakes Hurleg³⁵, Qinghai³⁶ and Zhuyeze³⁸, providing direct evidence for the lake level changes and climate conditions in the past. Optically stimulated luminescence dating was applied at these three lakes due to the absence of organic matter commonly used for radiocarbon dating of Holocene lake deposits. The obtained age data of the shorelines are generally in good agreement with independent lake-level inferences based on analyses of sediments from these lakes^{50–52}.

(4) Sedimentological and geochemical proxies such as grain size, carbonate and total organic carbon content, and Sr/Ca ratios (a salinity and indirect lake-level proxy) were used for lake-level inferences from Lakes Bosten³⁰ and Hala³⁷. Pollen data are available for Lake Bosten and a comparison of the lake level inference based on the sedimentological and geochemical characteristics of the sediments and the A/C ratio shows a good agreement especially for the last 2500 years relevant for this study (Fig. 3).

Data availability. The presented data are accessible in the PANGAEA database, <https://doi.pangaea.de/10.1594/PANGAEA.871173>.

References

1. Qin, X. *et al.* New evidence of agricultural activity and environmental change associated with the ancient Loulan kingdom, China, around 1500 years ago. *Holocene* **22**, 53–61 (2012).
2. Zhang, H., Wu, J. W., Zheng, Q. H. & Yu, Y. J. A preliminary study of oasis evolution in the Tarim basin, Xinjiang, China. *J. Arid Environ.* **55**, 545–553 (2003).
3. Hao, X., Chen, Y., Xu, C. & Li, W. Impacts of climate change and human activities on the surface runoff in the Tarim River basin over the last fifty years. *Water Resour. Manage.* **22**, 1159–1171 (2008).
4. Hörner, N. G. Lop-nor. Topographical and Geological Summary. *Geografiska Annaler* **14**, 297–321 (1932).
5. Yang, X., Liu, Z., Zhang, F., White, P. D. & Wang, X. Hydrological changes and land degradation in the southern and eastern Tarim Basin, Xinjiang, China. *Land Degrad. Dev.* **17**, 381–392 (2006).
6. Lü, H. *et al.* A preliminary study of chronology for a newly discovered ancient city and five archaeological sites in Lop Nur, China. *Chin. Sci. Bull.* **55**, 63–71 (2009).
7. Zhang, F. *et al.* Hydrological changes and settlement migrations in the Keriya River delta in central Tarim Basin ca. 2.7–1.6 ka BP: Inferred from ^{14}C and OSL chronology. *Sci. China Earth Sci.* **54**, 1971–1980 (2011).
8. Ban, G. Han Dynasty (originally published in 92, reprinted by Zhonghua Book Company, Beijing, 1962).
9. Dong, Z. *et al.* Research progress in China's Lop Nur. *Earth-Sci. Rev.* **111**, 142–153 (2012).
10. Mallory, J. P. & Mair, V. H. *The Tarim Mummies: Ancient China and the Mystery of the Earliest Peoples from the West* (Thames & Hudson, London, 2000).
11. Xie, Y., Chen, F. & Qi, J. Past desertification processes of Minqin Oasis in arid China. *Int. J. Sustainable Dev. World Ecol.* **16**, 417–426 (2009).
12. Zhang, Y. Basic characteristics of the military reclamation in the western regions during the Han Dynasty. *Western Reg. Stud.* **3**, 6–12 (2007).
13. Gao, X. The development of irrigation farming of the Hexi Corridor in the West Han Dynasty and its impact on ecological environment. *J. Shihezi Univ.* **24**, 90–92 (2010).
14. Wang, J. & Duan, J. Qin-Han's developing west policies: Gains and losses. *J. Nanjing Agric. Univ.* **5**, 71–76 (2005).
15. Dong, J. Probe into the source of ancient China's troop stationing and land reclamation. *J. Shihezi Univ.* **21**, 30–32 (2007).
16. Wang, W. On the policy of stationing garrison troops to open up wasteland in border regions in the western Han Dynasty. *J. Urumqi Vocational Univ.* **4**, 22–24 (2009).
17. Zhou, X., Zhu, F. & Li, S. The formation and evolution of oasis in the Keriya River valley. *Quat. Sci.* **3**, 249–255 (1994).
18. Xuanzang. *Datangxiyuji* (Tang Dynasty, originally published in 645, reprinted by Zhonghua Book Company, Beijing, 1985).
19. Zhang, J. F., Liu, C. L., Wu, X. H., Liu, K. X. & Zhou, L. P. Optically stimulated luminescence and radiocarbon dating of sediments from Lop Nur (Lop Nur), China. *Quat. Geochron.* **10**, 150–155 (2012).
20. Liu, C., Zhang, J.-F., Jiao, P. & Mischke, S. The Holocene history of Lop Nur and its palaeoclimate implications. *Quat. Sci. Rev.* **148**, 163–175 (2016).
21. Xia, X. & Zhao, Y. Some latest achievements in research on environment and its evolution in Lop Nur region. *Bull. Nat. Sci. Found. China* **19**, 196–200 (2005).
22. Zhang, D. Station troops to open up wasteland of western region in the Han Dynasty based on the Han slips from Xuanquan and its significance. *Dunhuang Res.* **3**, 113–121, (in Chin) (2001).
23. Zhu, H. The role of stationed troops opening up wasteland of western regions during Han Dynasty in agriculture, technology and cultural exchange between China and the West. *Chin. Hist.* **22**, 21–27, (in Chin) (2003).
24. Jiang, Q. F. *et al.* Holocene vegetational and climatic variation in westerly dominated areas of Central Asia inferred from the Sayram Lake in northern Xinjiang, China. *Sci. China Earth Sci.* **56**, 339–353 (2013).
25. Wang, W., Feng, Z., Ran, M. & Zhang, C. Holocene climate and vegetation changes inferred from pollen records of Lake Aibi, northern Xinjiang, China: A potential contribution to understanding of Holocene climate pattern in East-central Asia. *Quat. Int.* **311**, 54–62 (2013).
26. Liu, X., Herzschuh, U., Shen, J., Jiang, Q. & Xiao, X. Holocene environmental and climatic changes inferred from Wulungu Lake in northern Xinjiang, China. *Quat. Res.* **70**, 412–425 (2008).
27. Tao, S. C. *et al.* Pollen-inferred vegetation and environmental changes since 16.7 ka BP at Balikun Lake, Xinjiang. *Chin. Sci. Bull.* **55**, 2449–2457 (2010).
28. Huang, X. Z., Chen, F. H., Fan, Y. X. & Yang, M. L. Dry late-glacial and early Holocene climate in arid central Asia indicated by lithological and palynological evidence from Bosten Lake, China. *Quat. Int.* **194**, 19–27 (2009).
29. Herzschuh, U., Tarasov, P., Wünnemann, B. & Hartmann, K. Holocene vegetation and climate of the Alashan Plateau, NW China, reconstructed from pollen data. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **211**, 1–17 (2004).
30. Wünnemann, B., Mischke, S. & Chen, F. H. Holocene sedimentary record from Bosten Lake, China. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **234**, 223–238 (2006).

31. Mischke, S. *et al.* Modern hydrology and late Holocene history of Lake Karakul, eastern Pamirs (Tajikistan): A reconnaissance study. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **289**, 10–24 (2010).
32. Aichner, B., Feakins, S. J., Lee, J. E., Herzschuh, U. & Liu, X. High-resolution leaf wax carbon and hydrogen isotopic record of the late Holocene paleoclimate in arid Central Asia. *Clim. Past* **11**, 619–633 (2015).
33. Ricketts, R. D., Johnson, T. C., Brown, E. T., Rasmussen, K. A. & Romanovsky, V. V. The Holocene paleolimnology of Lake Issyk-Kul, Kyrgyzstan: Trace element and stable isotope composition of ostracodes. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **176**, 207–227 (2001).
34. Rhodes, T. E. *et al.* A Late Pleistocene–Holocene lacustrine record from Lake Manas, Zunggar (northern Xinjiang, western China). *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **120**, 105–121 (1996).
35. Fan, Q. S., Ma, H. Z., Wei, H. C. & An, F. Y. Holocene lake-level changes of Hurlig Lake on northeastern Qinghai–Tibetan Plateau and possible forcing mechanism. *Holocene* **24**, 274–283 (2014).
36. Liu, X. J., Lai, Z., Madsen, D. & Zeng, F. Last deglacial and Holocene lake level variations of Qinghai Lake, north-eastern Qinghai–Tibetan Plateau. *J. Quat. Sci.* **30**, 245–257 (2015).
37. Wünnemann, B. *et al.* Implications of diverse sedimentation patterns in Hala Lake, Qinghai Province, China for reconstructing Late Quaternary climate. *J. Paleolimnol.* **48**, 725–749 (2012).
38. Long, H., Lai, Z. P., Fuchs, M., Zhang, J. R. & Li, Y. Timing of Late Quaternary palaeolake evolution in Tengger Desert of northern China and its possible forcing mechanisms. *Global Planet. Change* **92–93**, 119–129 (2012).
39. Yang, B., Braeuning, A., Shi, Y. & Chen, F. Evidence for a late Holocene warm and humid climate period and environmental characteristics in the arid zones of northwest China during 2.2–1.8 kyr B.P. *J. Geophys. Res.* **109**, D02105, doi: 10.1029/2003JD003787 (2004).
40. Lauterbach, S. *et al.* Climatic imprint of the mid-latitude Westerlies in the Central Tian Shan of Kyrgyzstan and teleconnections to North Atlantic climate variability during the last 6000 years. *Holocene* **24**, 970–984 (2014).
41. Yonenobu, H. & Itoh, T. ¹⁴C dating of the Niya site in the Tarim Basin. *Nucl. Instrum. Methods Phys. Res. B* **172**, 741–744 (2000).
42. Tang, Z., Chen, D., Wu, X. & Mu, G. Redistribution of prehistoric Tarim people in response to climate change. *Quat. Int.* **308–309**, 36–41 (2013).
43. Flad, R., Li, S., Wu, X. & Zhao, Z. Early wheat in China: Results from new studies at Donghuishan in the Hexi Corridor. *Holocene* **20**, 955–965 (2010).
44. Blott, S. J. & Pye, K. GRADISTAT: A grain size distribution and statistics package for the analysis of unconsolidated sediments. *Earth Surf. Process. Landf.* **26**, 1237–1248 (2001).
45. Turpen, J. B. & Angell, R. W. Aspects of moulting and calcification in the ostracod *Heterocypris*. *Biol. Bull.* **140**, 331–338 (1971).
46. von Grafenstein, U., Erlenkeuser, H. & Trumborn, P. Oxygen and carbon isotopes in modern fresh-water ostracod valves: Assessing vital offsets and autoecological effects of interest for palaeoclimate studies. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **148**, 133–152 (1999).
47. El-Moslimany, A. P. Ecological significance of common nonaraboreal pollen: Examples from drylands of the Middle East. *Rev. Palaeobot. Palynol.* **64**, 343–350 (1990).
48. Mischke, S. & Zhang, C. Ostracod distribution in Ulungur Lake (Xinjiang, China) and a reassessed Holocene record. *Ecol. Res.* **26**, 133–145 (2011).
49. Talbot, M. R. A review of the palaeohydrological interpretation of carbon and oxygen isotopic ratios in primary lacustrine carbonates. *Chem. Geol.* **80**, 261–279 (1990).
50. Zhao, C. *et al.* Holocene millennial-scale climate variations documented by multiple lake-level proxies in sediment cores from Hurlig Lake, Northwest China. *J. Paleolimnol.* **44**, 995–1008 (2010).
51. Shen, J., Liu, X., Wang, S. & Matsumoto, R. Palaeoclimatic changes in the Qinghai Lake area during the last 18,000 years. *Quat. Int.* **136**, 131–140 (2005).
52. Mischke, S., Lai, Z., Long, H. & Tian, F. Holocene climate and landscape change in the northeastern Tibetan Plateau foreland inferred from the Zhuyeze Lake record. *Holocene* **26**, 643–654 (2016).

Acknowledgements

Funding was provided by China's NSF projects (40830420, 41471003), the State key project (2003BA612A-06–15) of the Ministry of Science and Technology of China and the German Research Foundation (DFG grant Mi 730/16-1). We thank two anonymous reviewers who provided very constructive comments on an earlier version of this paper.

Author Contributions

S.M., C.L. and J.Z. designed research; S.M., C.L., J.Z., C.Z., P.J., B.P. performed research; and S.M., C.L., J.Z., H.Z. and B.P. analyzed data and wrote the paper.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Mischke, S. *et al.* The world's earliest Aral-Sea type disaster: the decline of the Loulan Kingdom in the Tarim Basin. *Sci. Rep.* **7**, 43102; doi: 10.1038/srep43102 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017