

MS# ECY17-0787

The China Plant Trait Database: towards a comprehensive regional compilation of functional traits for land plants

HAN WANG,¹ SANDY P. HARRISON,^{1,2} I. COLIN PRENTICE,^{1,3} YANZHENG YANG,^{4,1} FAN BAI,⁵ HENRIQUE FURSTENAU TOGASHI,^{6,7} MENG WANG,¹ SHUANGXI ZHOU,⁶ JIAN NI^{8,9}

¹ *State Key Laboratory of Soil Erosion and Dryland Farming on the Loess Plateau, College of Forestry, Northwest A&F University, Yangling 712100, China*

² *School of Archaeology, Geography and Environmental Sciences (SAGES), Reading University, Reading, RG6 6AB, UK*

³ *AXA Chair of Biosphere and Climate Impacts, Imperial College London, Silwood Park Campus, Buckhurst Road, Ascot SL5 7PY, UK*

⁴ *Ministry of Education Key Laboratory for Earth System Modelling, Department of Earth System Science, Tsinghua University, Beijing 100084, China*

⁵ *State Key Laboratory of Vegetation and Environmental Change, Institute of Botany, Chinese Academy of Science, Beijing 100093, China*

⁶ *Department of Biological Sciences, Macquarie University, North Ryde, NSW 2109, Australia*

⁷ *The Ecosystem Modelling and Scaling Infrastructure Facility, Macquarie University, North Ryde, NSW 2109, Australia*

⁸ *College of Chemistry and Life Sciences, Zhejiang Normal University, Yingbin Avenue 688, 321004 Jinhua, China*

⁹ *State Key Laboratory of Environmental Geochemistry, Institute of Geochemistry, Chinese Academy of Sciences, Lincheng West Road 99, 550081 Guiyang, China*

Abstract.

Plant functional traits provide information about adaptations to climate and environmental conditions, and can be used to explore the existence of alternative plant strategies within ecosystems. Trait data are also increasingly being used to provide parameter estimates for vegetation models. Here we present a new database of plant functional traits from China. Most global climate and vegetation types can be found in China, and thus the database is relevant for global modelling. The China Plant Trait Database contains information on morphometric, physical, chemical and photosynthetic traits from 122 sites spanning the range from boreal to tropical, and from deserts and steppes through woodlands and forests, including montane vegetation. Data collection at each site was based either on sampling the dominant species or on a stratified sampling of each ecosystem layer. The database contains information on 1215 unique species, though many species have been sampled at multiple sites. The original field identifications have been taxonomically standardized to the Flora of China. Similarly, derived photosynthetic traits, such as electron-transport and carboxylation capacities, were calculated using a standardized method. To facilitate trait-environment analyses, the database also contains detailed climate and vegetation information for each site.

Key words: plant traits, leaf morphometry, leaf economics, leaf chemistry, photosynthetic properties, J_{max} , V_{cmax}

The complete data set is available online at: [*to be completed at proof stage*].

Corresponding Editor: W. K. Michener.

¹E-mail: wanghan_sci@yahoo.com