



THE AGA KHAN UNIVERSITY

eCommons@AKU

Department of Medicine

Department of Medicine

January 1995

Severe thrombocytopenia in a man with prostatic cancer

Saeed Hamid

Aga Khan University, saeed.hamid@aku.edu

Ata Khan

Aga Khan University

Haleem Khan

Aga Khan University

Roger Sutton

Aga Khan University

Ikram A. Burney

Aga Khan University

See next page for additional authors

Follow this and additional works at: http://ecommons.aku.edu/pakistan_fhs_mc_med_med

Recommended Citation

Hamid, S., Khan, A., Khan, H., Sutton, R., Burney, I., Khurshid, M. (1995). Severe thrombocytopenia in a man with prostatic cancer. *Journal of Pakistan Medical Association*, 252-254.

Available at: http://ecommons.aku.edu/pakistan_fhs_mc_med_med/547

Authors

Saeed Hamid, Ata Khan, Haleem Khan, Roger Sutton, Ikram A. Burney, and Mohammad Khurshid

Severe Thrombocytopenia in a Man with Prostatic Cancer

Pages with reference to book, From 252 To 254

Saeed Hamid, Ata Khan, Haleem Khan, Roger Sutton, Ikram A. Burney, Mohammed Khurshid (Department of Medicine and Pathology, Aga Khan University Hospital, Karachi.)

A 66 year old gentleman was diagnosed to have metastatic adenocarcinoma of the prostate in 1990, following a CT scan of the abdomen and trans-urethral resection of prostate (TLJRP). He was started on estrogen therapy (Fosfesterol sodium). Two years later he presented to the emergency room with breathlessness and edema of the left arm of a week's duration. He admitted to persistence of symptoms of pmstatism and was found to have bilateral axillary lymphadenopathy and pitting edema of the left arm. A doppler scan revealed left subclavian vein thrombosis. The patient was started on i.v. heparin and was maintained on it until the resolution of arm edema and was subsequently switched to oral anticoagulation. The serum Prostate Specific Antigen (PSA) level was found to be 1460 ng/ml. Fosfesterol was discontinued and the patient was started on antiandrogen therapy (Cyproterone acetate). The patient was sent home on anticoagulation, but had to be readmitted shortly thereafter because of the sudden onset of shortness of breath. This time he was found to have bilateral pitting pedal edema and :tenderness in the right calf. Intravenous heparin was reinstated. An inferior venacavagram revealed extensive thrombosis of the right popliteal vein and the left common iliac vein. A clinical diagnosis of pulmonary embolism was made. Despite continuous i.v. heparin in adequate doses, the patient experienced recurrent episodes of shortness of breath. A greenfield filter was placed in the inferior vena cava to prevent further episodes of pulmonary embolism. The patient remained heparinized for more than two weeks, until complete resolution of symptoms had occurred. Subsequently oral anticoagulation was started and the dose of coumarin was adjusted to maintain an international normalized ratio (INR) between 1.8 and 2.0.

Over the next several months, the patient was followed up in the oncology clinic with serial measurements of PSA. Despite being on maximum doses of cyproterone acetate, the PSA level. did not normalize. Cyproterone was discontinued and the patient was started on the non-progestational anti-androgen flutamide. However, the PSA values continued to fluctuate and subsequently started to increase. The serum testosterone level was found to be high (235 ng/dl). The corresponding PSA level was 936 ng/ml. The patient was advised to have a subcapsular orchidectomy. He was admitted to the hospital, where the following laboratory values were obtained: Hb 10.1 gm/dl; TLC $5.7 \times 10^9/l$; platelets $147 \times 10^9/L$; PT 19/12 seconds (INR 1.9); and APTF 34/3 2 seconds. Coumann was stopped and heparin was started. On the 4th day, the platelet count was found to be $16 \times 10^9/L$. This was rechecked and was found to be the same. The PT, APT!' and FDP were all found to be within normal !imits. At this time, them was no purpura. The next day, ecchymoses had appeared and the platelet count had dropped to $7 \times 10^9/L$ and subsequently to $146 \times 10^9/L$ after another two days. The plan for subcapsular orchidectomy was cancelled and the LHRH analogue (Buserelin) was injected subcutaneously. Because of the relationship of the onset of thrombocytopenia and its resolution, to the start and discontinuation of heparin, a diagnosis of heparin-induced thrombocytopenia was entertained. Two weeks later, the patient was readmitted with calf tenderness and left leg edema. He was again found to have deep venous thrombosis, extending from the popliteal to the external iliac vein. Since the patient was already taking coumarin, which proved ineffective in preventing thrombus formation, the patient was started on low molecular weight heparm (LMWH) and aspirin. Presently, the patient is doing well, the symptoms have largely resolved and the platelet counts, repeated on several occasions, have remained within normal limits.

Discussion

Thrombocytopenia is one of the most common and important immunological complications of heparin therapy'. The attendant morbidity and mortality is significant^{2,3}. The incidence of the complication is 1-5%; higher with bovine than with porcine heparin⁴. Two clinical types have been described; a mild, early onset type and a severe, delayed onset type⁵. In the former type, the platelet count does not fall below $100 \times 10^9/L$ and the patient is usually asymptomatic. The situation is easily reversible without any specific therapy; the mechanism of thrombocytopenia is unknown. The second type is characterized by a reduction 'in the platelet count to usually less than $50 \times 10^9/L$, and it occurs 4-9 days after the onset of heparin therapy. This type is associated with an increased risk of thrombo-embolic complications and an immune mechanism is implicated in the pathogenesis^{6,7,8}. Of the various drug-induced thrombocytopenias, that caused by heparin is different as it is associated with relatively little risk of bleeding, but a high risk of paradoxical thrombosis involving major vessels. The thrombo-embolic complications may manifest as cerebral thrombosis, acute myocardial infarction, recurrent pulmonary emboli, or arterial thrombosis resulting in limb gangrene with a 20% risk of amputation⁹. The proposed mechanism of heparin induced thrombocytopenia is as follows: IgG antibody forms after several days of exposure to heparin. The antibodies are not specific for heparin and interact with heparin only in the presence of platelet factor 4 (PF4)¹⁰. The IgG/heparin PF4 immune complexes are formed on the platelet surface and activate platelets by binding to Fc receptors^{1,11,12,13}. Thrombosis is secondary to platelet activation and endothelial cell injury^{4,13}. Although the diagnosis of heparin induced thrombocytopenia is usually clinical, different laboratory tests can be employed, not only for the confirmation of the diagnosis, but also for screening agents which could be used as alternatives to heparin. In-vitro platelet aggregation is a rapid and simple test^{9,14,15}. ¹⁴C- serotonin release test is more sensitive and specific^{4,16}. Other assays have been developed but have little practical value for routine diagnosis⁴. Recognition of the condition requires immediate discontinuation of heparin. Platelet transfusion should be avoided, as these may precipitate arterial thrombosis. As alternatives to i.v. heparin for anticoagulation, different agents have been tried. Low molecular weight heparin (LMWH) has been used frequently, generally with a favourable clinical course. Although there may be 25-60% cross reactivity between unfractionated heparin and LMWH¹⁷, the incidence of heparin induced thrombocytopenia is significantly low^{18,19,20}. An in-vitro platelet aggregation test may help in selecting the type of LMWH. Besides LMWH, other treatment options include the use of Heparinoid²¹, Heparan sulphate and dermatan sulphate²², Ancrod, and the placement of inferior vena caval filters²³. For established thrombosis, aspirin, dipyridamole, prostacyclin analogues, fibrinolytic therapy and plasmapheresis all have been tried^{4,24} with variable success. Our patient developed severe thrombocytopenia on the fourth day of heparin therapy and this reversed four days after cessation of the drug without any specific intervention. The diagnosis was established clinically. The patient had been heparinized on two previous occasions, and retrospective analysis of records revealed that on one occasion the platelet count had dropped to $84 \times 10^9/L$, but reverted spontaneously. With as high an incidence of HIT as that reported in the literature (1-5%), together with the frequency of heparin use in clinical practice, physicians need to be aware of this important and potentially life-threatening complication of heparin therapy. LMWH could be usefully employed instead of unfractionated heparin and in such cases the probability of cross-reactivity could be decreased by performing a simple in-vitro platelet aggregation test.

References

1. Kelton, J. O., Smith, W. W., Warkentin, T. E. et al. Immunoglobulin U from patients with heparin-induced thrombocytopenia binds to a complex of heparin and platelet factor 4. *Blood* 1994;83:3232-3239.
2. Demasi, R., Bode, A. P., Krupp, C., et al. Heparin-induced thrombocytopenia. *Am. J. Surg.* 1994;60:26-29.
3. King, D. I., Kelton, S. G. Heparin-associated, thrombocytopenia. *Ann. Intern. Med.* 1994;100:536-541.
4. Warkentin, T. E., Kelton, J. G. Heparin and platelets. *Hematol. Oncol. Clin. North Am.* 1990;4:243-264.
5. Chong, B. H. Heparin-induced thrombocytopenia. *Aust. N. J. Med.* 1992;22:145-152.
6. Rhodes, G. R., Dixon, R. H., Silver, D. Heparin-induced thrombocytopenia with thrombotic and haemorrhagic manifestations. *Surg. Gynaecol. Obstet.* 1973;136:409-414.
7. Kelton, S. G., Sheridan, D., Santos, A. et al. Heparin-induced thrombocytopenia: Laboratory studies. *Blood* 1988;72:925-930.
8. Chong, B. H., Fawaz, I., Chesterman, C. N. Heparin-induced thrombocytopenia: mechanism of action of the heparin-dependent antibody with platelets. *Br. J. Hematol.* 1989;73:235-241.
9. Chong, B. H., Pilgrim, R. L., Cooley, M. A. et al. increased expression of platelet IgG Fc receptor in immune heparin-induced thrombocytopenia. *Blood* 1993;81:988-993.
10. Amiral, J., Bridey, F., Dreyfus, M. et al. Platelet factor 4 complexes to heparin is the target for antibodies generated in heparin-induced thrombocytopenia. *Thromb. hemost* 1992;68:95-96.
11. Adelman, B., Sobel, M., Fijimura, T. et al. Heparin associated thrombocytopenia: Observations on the mechanism of platelet aggregation. *3. Lab. Clin. Med.* 1989;113:204-210.
12. Anderson, G. P., Insights into heparin-induced thrombocytopenia. *Br. J. Hematol.* 1992;80:504-508.
13. Visentin, G. P., Ford, S. E., Scott, J. P. et al. Antibodies from patients with heparin-induced thrombocytopenia/thrombosis are specific for platelet factor 4 complexed with heparin or bound to endothelial cells. *J. Clin. Invest.* 1994;93:81-88.
14. Isenhardt, C. E., Brandt, J. T. Platelet aggregation studies for the diagnosis of heparin-induced thrombocytopenia. *Am. J. Clin Pathol.* 1993;99:324-330.
15. Keeling, D. M., Richards, E. M., Baglin, T. P. Platelet aggregation in response to four low molecular weight heparins and the heparinoid ORG 10172 in patients with heparin-induced thrombocytopenia. *Br. J. Hematol.* 1994;86:425-426.
16. Favalaro, E. J., Bernal-Hoyos, E., Exner, T. et al. Heparin-induced thrombocytopenia: laboratory investigation and confirmation of diagnosis. *Pathology* 1992;24: 177-183.
17. Kikta, M. J., Keller, M., P., Humphery, P. W. et al. Can low molecular weight heparins and heparinoids be safely given to patients with heparin-induced thrombocytopenia syndrome? *Surgery* 1993;114:705-710.
18. Mammen, E. F. Why low molecular weight heparin? *Semin Thromb Hemost* 1990;165: 1-4.
19. Green, D., Hirsh, J., Heit, J. et al. Low molecular weight heparin: A critical analysis of clinical trials. *Pharmacol Rev* 1994;46:89-109.
20. Warkentin, T. E., Hayward, C. P., Smith, C. A. et al. Determinants of donor platelet variability when testing for heparin-induced thrombocytopenia. *3. Lab. Clin Med.* 1992;120:371-379.
21. Magnani, H. N. Heparin-induced thrombocytopenia (HIT): an overview of 230 patients treated with organan (ORG 10172). *Thromb Haemost* 1993;70:554-561.
22. Hoppensteadt, D. A., Walenga, S. M., Fareed, J. Effect of dermatan sulfate and heparan sulfate on platelet activity compared to heparin. *Semin Thromb Hemost.* 1991 ;175:60-64.
23. Calligaro, K. D., Bergen, W. S., Haut, M. S., et al. Thromboembolic complications in patients with advanced cancer: anticoagulation versus green-field filter placement. *Ann Vase Surg* 1991 ;5: 186-189.
24. Brady, J., Riccio, J. A., Yumen, O. H. et al. Plasmapheresis: A therapeutic option in the management of heparin-induced thrombocytopenia with thrombosis. *Am. J. Clin. Pathol.* 1991;96:394-397.