

Fluorometabolite biosynthesis: Isotopically labelled glycerol incorporations into the antibiotic nucleocidin in *Streptomyces calvus*.


Axel Bartholomé,^a Jeffrey E. Janso,^b Usa Reilly^b and David O'Hagan.*^a

5 **Receipt/Acceptance Data [DO NOT ALTER/DELETE THIS TEXT]**

Publication data [DO NOT ALTER/DELETE THIS TEXT]

DOI: 10.1039/b000000x [DO NOT ALTER/DELETE THIS TEXT]

Deuterium and carbon-13 labelled glycerols have been fed to *Streptomyces calvus* fermentations and isotope incorporation into the fluorine containing antibiotic nucleocidin have been evaluated by ¹⁹F-NMR. A single deuterium atom was incorporated from [²H₅]- and (R)-[²H₂]- glycerol into C-5' of the antibiotic, suggesting that an oxidation occurs at this carbon after ribose ring assembly from glycerol (pentose phosphate 15 pathway), during nucleocidin biosynthesis.



Scheme 1. Fluorinase reaction of AdoMet **3** to give FDA **4** and then the relationship between FDA and FAc **1** and 4-FT **2**. Incorporation of (R)-[²H₂]- glycerol determined that both fluoromethyl hydrogens at C5' of FDA **4** are retained in the fluorometabolites.⁹ Nucleocidin **7** is a metabolite of *S. calvus*.

Fluorine containing natural products are exceedingly rare in nature.¹ The toxin fluoroacetate (FAc) **1** is the most widely distributed natural

product of this class where it has been identified in many plants² and a few bacteria.^{3,4} The mode of fluoroacetate biosynthesis in plants has not been established, however its biosynthesis in bacteria is known.⁵ *Streptomyces cattleya* is a bacterium that co-produces FAc **1** and 4-fluorothreonine (4-FT) **2**.³ Studies carried out with *S. cattleya* determined that the fluorination enzyme (fluorinase) catalyses a nucleophilic substitution reaction between fluoride ion and *S*-adenosyl-L-methionine (Ado-Met) **3** to generate 5'-fluorodeoxyadenosine (FDA) **4**.⁶ This is the only biosynthetic fluorination enzyme known. It has subsequently been identified by gene mining in several other bacteria.⁷ FDA **4** is then processed to FAc **1** and 4-FT **2**. A combination of isotopic labelling studies, including isotopically labelled glycerol **5**, and an understanding of the individual steps in the biosynthesis of **1** and **2** have determined that the C5' fluoromethyl and C4' carbon of FDA **4** are converted to FAc **1** and C3 and C4 of 4-FT **2**, the circled regions in Scheme 1.⁸ It has also been established that both hydrogens of the fluoromethyl group of FDA **4** are retained in the conversion to the fluoromethyl groups of **1** and **2**.⁹

The antibiotic nucleocidin **7** produced by the actinomycete soil bacterium, *Streptomyces calvus* is another example of the rare fluorometabolites.¹⁰ Its biosynthesis is unknown; however, the presence of the fluorine atom at the C4' position of the ribose ring¹¹ in **7** is inconsistent with the involvement of the fluorinase found in *S. cattleya*. Full genome sequencing of the nucleocidin producer *S. calvus* ATCC 13382 has not identified a fluorinase gene,¹² thus *S. calvus* may employ a very different process for enzymatic C-F bond formation relative to bacterial FAc **1** production.

Ever since its isolation, from *S. calvus* in 1956,¹⁰ nucleocidin **7** production has proved problematic, with consistently low titres and unpredictable production. Publically deposited strains appear to have lost the ability to produce nucleocidin.¹²⁻¹⁴ Some progress has been made to understand the inconsistent production, and it has been observed that a mutation of the *bldA* gene, encoding a Leu-tRNA, is apparent in the publically available strains. Correction of this mutation was recently shown¹² to re-establish production in *S. calvus*. In another study¹³ mutation of the *rpoB* gene (rifamycin resistance) led to an apparent increase in nucleocidin production. In this study we use an in-house strain of *S. calvus* T-3018 held by Pfizer. This strain does not have a *bldA* mutation, although production is still fickle with very low titres. Nucleocidin production can however be monitored, even at low production levels, directly from extracts by ¹⁹F{¹H}-NMR. As a first step towards elucidating the biosynthetic pathway to fluorometabolite **7** we now report the incorporation of various isotopically labelled glycerols **5 a-d**.

For this study we have supplemented fermentation cultures of *S. calvus* with glycerols [²-¹³C]- **5a**, [²H₅]- **5b**, (R)-[²H₁]- **5c** and (S)-[²H₁]- **5d**. Despite the low titres, we envisaged that the incorporation of deuterium or carbon-13 isotopes close to the fluorine

^a University of St Andrews, School of Chemistry and Centre for Biomolecular Sciences, North Haugh, St Andrews, Fife, KY16 9ST, UK Fax: 01334 463800; Tel: 01334 467171; E-mail: do1@st-andrews.ac.uk

^b Pfizer, 220-3529, 445 Eastern Point Rd., Groton, CT 06340, 860-715-6375, USA.

† Electronic Supplementary Information (ESI) available: See <http://dx.doi.org/10.1039/b000000x/>

atom of the antibiotic could be reported visually by $^{19}\text{F}\{^1\text{H}\}$ -NMR due to heavy atom induced chemical shifts of the fluorine resonance,¹¹⁰ and/or ^{13}C - ^{19}F -spin-spin coupling in the case of carbon-13 incorporation.

A first experiment explored pulse feeding of glycerol [^{13}C]- **5a** to a final concentration of 8mM, in a shake flask fermentation of *S. calvus*. Cultures were worked up after 20 days as previously¹¹⁵ described.¹³ Nucleocidin is excreted from the bacterial cells and thus the cells were spun down in a centrifuge and the supernatant was extracted into butanol. The extract was concentrated and was then analysed by $^{19}\text{F}\{^1\text{H}\}$ -NMR. The resultant spectrum is shown in Figure 1a. It is clear that the fluorine signal for nucleocidin has¹²⁰ accompanying satellites due to $^1J_{\text{CF}}$ (232Hz) coupling, consistent with a population (~25%) of nucleocidin **7** molecules enriched with carbon-13 at C4'. The satellite signals are off-centre relative to the natural abundance ^{19}F -NMR signal due to a heavy atom α -shift.⁸ This incorporation is entirely consistent with pentose¹²⁵ phosphate pathway involvement in ribose biosynthesis where carbons C3–C5 of D-ribose derive from the exogenously added glycerol, feeding into the C₃ metabolite pool *via* *sn*-glycerol-3-phosphate **6**.¹⁵





Figure 1. $^{19}\text{F}\{^1\text{H}\}$ NMR of nucleocidin **7** (-118.8ppm) after glycerol **5** feeding experiments to *S. calvus* fermentations. An unknown metabolite (-118.1ppm) is also present in some spectra. (a) [^{13}C]-Glycerol **5a**; (b) [$^2\text{H}_3$]-glycerol **5b**; (c) (*R*)-[$^2\text{H}_2$]-glycerol **5c**; (d) (*S*)-[$^2\text{H}_2$]-glycerol **5d**. A pictorial summary of the isotopic labelling outcome is given in each case.

Addition of perdeuterated glycerol **5b** (final conc 8 mM) to a *S. calvus* fermentation led to the incorporation of deuterium (~5%)¹⁵⁰ into nucleocidin **7** as determined by an upfield shift in the resultant $\{^1\text{H}\}^{19}\text{F}$ -NMR spectrum as shown in Figure 1b. The magnitude of this upfield shift at 0.22ppm is consistent with a single vicinal deuterium incorporated β - to the fluorine atom.¹⁵⁵ There is a F-C-C-D angular dependence on the magnitude of the shift ranging from 0.15ppm (0°) to 0.35ppm (180°) as determined

in classic studies by Lambert and Greifenstein.¹⁶ The observed value is intermediate between these extremes and consistent with a single vicinal C-D (γ -shift) bond approximately at 90° to the C-F bond. This experiment does not however distinguish the location of the incorporation between C3' or C5' of the ribose ring of **7**. To achieve such a distinction, comparative incorporation experiments were explored with the enantiomeric glycerols (*R*)-[$^2\text{H}_2$]- **5c** and (*S*)-[$^2\text{H}_2$]- **5d**. It is already established that the *pro-R* arm of glycerol delivers C5' of ribose through the pentose phosphate pathway, and that the *pro-S* arm delivers C3' of ribose. Glycerols **5c** and **5d** were prepared synthetically with enantiomeric purities > 95% *e.e.*, as previously described.⁹ The resultant $^{19}\text{F}\{^1\text{H}\}$ -NMR spectra from each of these glycerol feeding experiments are shown in Figure 1c and Figure 1d. It is clear that only glycerol (*R*)-[$^2\text{H}_2$]- **5c** results in the incorporation (~8%) of deuterium into nucleocidin **7**, and again only a single deuterium atom survives from the double labelled precursor. It follows from the known stereochemistry of glycerol processing through the pentose phosphate pathway, that this deuterium is located at C5'.^{15,17}

In this experiment a second fluorometabolite was observed in the $^{19}\text{F}\{^1\text{H}\}$ -NMR spectrum with a signal at -118.1ppm. This metabolite is detected periodically in nucleocidin extracts although its occurrence and level is unpredictable in our experience. Such a fluoro metabolite was also observed in the recent study where nucleocidin production was elicited by reversing the *bldA* mutation.¹² Its structure is unknown although given the similar level of deuterium incorporation from glycerol **5c**, it appears to be metabolically related to nucleocidin either as a biosynthetic precursor, or as a metabolite. For isomer (*S*)-[$^2\text{H}_2$]- **5d** the absence of any deuterium at C3' in **7** is consistent with *sn*-glycerol-3-phosphate **6** being processed to ribulose-5-phosphate *via* xylulose-5-phosphate, and the action of D-ribulose-5-phosphate 3-epimerase, with exchange of the surviving deuterium with bulk solvent.¹⁸ The predicted labelling pattern of ribose-5-phosphate from glycerol is illustrated in Scheme 2.



Scheme 2. The pentose phosphate pathway showing the fate of glycerol through to ribose-5-phosphate. Glycerol **5e** has a virtual labelling pattern which is a composite of the experimentally administered glycerols **5a-d**, and also highlights the stereochemical fate of the (*pro-R*) hydroxymethyl carbon (■). This pathway predicts incorporation of a C₃ glycerol unit (bold C-C bonds) into C-3-C-5 of ribose-5-phosphate, with the retention of two deuteriums at C-5 and the loss of deuteriums at C-3 and C-4.

The outcome where both **5b** and **5c** glycerols contribute only one deuterium to C5' of nucleocidin can be contrasted with the biosynthesis of FAc, **1** and 4-FT, **2** in *S. cattleya* discussed above²²⁵ and shown in Scheme 1. In those cases two deuterium atoms were incorporated into the fluorometabolites, and by implication into the C5' ribose carbon of FDA **4**.^{9a} For nucleocidin only one deuterium is incorporated.²³⁰

In summary, this study provides the first biosynthetic data on nucleocidin **7** assembly from isotope labelling studies. It is shown that glycerol is incorporated into the ribose ring moiety of nucleocidin, consistent with expectation *via* the pentose phosphate pathway. However, the presence of a single deuterium only at the C5' position of nucleocidin **7** after feeding experiments with glycerols **5b** and **5c** suggests that a hydrogen is lost from this carbon after ribose ring assembly but prior to, or concomitant with, fluorine introduction. This observation places a constraint on working hypotheses addressing nucleocidin biosynthesis.

Acknowledgement: We are grateful to Dr Alessandra Eustáquio for helping to establish this research project.

References

1. D. O'Hagan, H. Deng, *Chem. Rev.*, 2015, **115**, 634-649.
2. (a) T. McEwan, T. *Nature* 1964, **202**, 827; (b) P. B. Oelrichs, T. McEwan, *Nature*, 1961, **190**, 586-587.
3. M. Sanada, T. Miyano, S. Iwadare, J. M. Williamson, B. H. Arison, J. L. Smith, A. W. Douglas, J. M. Liesch, E. Inamine, E., *J. Antibiotic*, 1986, **39**, 259-265.
4. (a) S. Huang, L. Ma, M. H Tong, Y Yu, D. O'Hagan, H. Deng, *Org. Biol. Chem.*, 2014, **12**, 4828-4831; (b) L. Ma, A. Bartholomé, M. H. Tong, Z. Qin, Y. Yu, T. Shepherd, K. Kyeremeh, H. Deng, D. O'Hagan, *Chem. Sci.*, 2015, **6**, 1414 - 1419.
5. H. Deng, S. M. Cross, R. P McGlinchey, J. T. G Hamilton, D. O'Hagan, *Chem & Biol.*, 2008, **15**, 1268 -1276.
6. X. M. Zhi, S. Hackl, M. N. Thaker, L. Kalan, C. Weber, D. S. Urgast, E. M. Krupp, A. Brewer, S. Vanner, A. Szawiola, G. Yim, J. Feldmann, A. Bechtold, S. Vanner, D. L. Zechel, *ChemBioChem*, 2015, **16**, 2498 - 2506.
7. (a) D. O Hagan, C. Schaffrath, S. L. Cobb, J. T. G. Hamilton, C. D .Murphy, *Nature*, 2002, **416**, 279; (b) C. Dong, F. L. Huang, H. Deng, C. Schaffrath, J. B. Spencer, D. O'Hagan, J. H. Naismith, *Nature*, 2004, **427**, 561 - 565.
8. H. Deng, L. Ma, N. Bandaranayaka, Z. Qin, G. Mann, K. Kyeremeh, Y. Yu, T. Shepherd, J. H. Naismith, D. O'Hagan, *ChemBioChem*, 2014, **15**, 364-368.
9. J. T. G. Hamilton, C. D. Murphy, M. R Amin, D. O'Hagan, D. B. Harper, *J. Chem. Soc. Perkin Trans. I.*, 1998, 759-767.
10. (a) J. Nieschalk, J. T. G. Hamilton, C. D. Murphy, D. B. Harper, D. O'Hagan, *Chem. Comm.*, 1997, 799-800; (b) R. E. Hill, A. Iwanow, B. G. Sayer, W. Wysocka and I. D. Spencer, *J. Biol. Chem.*, 1987, **262**, 7463-7471.
11. S. O Thomas, V. L. Singleton, J. A. Lowry, R. W. Sharpe, L. M. Pruess, J. N. Porter, J. H. Mowat, N. Bohonos, *Antibiotics Annu.*, 1956, **1956-1957**, 716-721.
12. G. O Morton, J. E. Lancaster, G. E. Van Lear, W. Fulmor, W. E. Meyer, *J. Am. Chem. Soc.*, 1969, **91**, 1535-1537.
13. X. M. Zhu, S. Hackl, M. N. Thaker, L. Kalan, C. Weber, D. S. Urgast, E. M. Krupp, A. Brewer, S. Vanner, A. Szawiola, G. Yim, J. Feldmann, A. Bechtold, G. D. Wright, D. L. Zechel, *ChemBioChem*, 2015, **16**, 2498 - 2506.
14. K. Fukuda, T. Tamura, Y. Segawa, Y. Mutaguchi, K. Inagaki, *Actinomycetologica*, 2009, **23**, 51 - 55.
15. A. R. Maguire, W. D., Meng, S. M. Roberts, A. J. Willets, *J. Chem. Soc. Perkin Trans. I*, 1993, 1795-1808.
16. D. C. Crans, G. M. Whitesides, *J. Am. Chem.Soc.*, 1985, **107**, 7008 -7018.

16. (a) J. B. Lambert, L. G. Greifenstein, *J. Am. Chem. Soc.*, 1974, **96**, 5120 - 5124; (b) J. B. Lambert, L. G. Greifenstein, *J. Am. Chem. Soc.*, 1973, **95**, 6150 - 6152.
17. W. Van Winden, P. Verheijen, S. Heijnen, *Metab. Eng.*, 2001, **3**, 151 - 162.
18. (a) Y-R. Chen, F. W. Larimer, E. H. Serpersu, F. C. Hartman, *J. Biol. Chem.*, 1999, **22**, 2132 - 2136; (b) M. W. McDonough, W. A. Wood, *J. Biol. Chem.*, 1961, **236**, 1220 - 1224.