



Matsubara, V. H., Wang, Y., Bandara, H. M. H. N., Mayer, M. P. A., & Samaranayake, L. P. (2016). Probiotic lactobacilli inhibit early stages of *Candida albicans* biofilm development by reducing their growth, cell adhesion, and filamentation. *Applied Microbiology and Biotechnology*, *100*(14), 6415-6426. <https://doi.org/10.1007/s00253-016-7527-3>

Peer reviewed version

Link to published version (if available):
[10.1007/s00253-016-7527-3](https://doi.org/10.1007/s00253-016-7527-3)

[Link to publication record in Explore Bristol Research](#)
PDF-document

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/pure/about/ebr-terms>

Probiotic strain	Biofilm viable cells reduction (%)							
	<i>C. albicans</i> ATCC SC5314				<i>C. albicans</i> 75			
	Initial Colonization		Maturation		Initial Colonization		Maturation	
	Mean	SD	Mean	SD	Mean	SD	Mean	SD
<i>L. rhamnosus</i> LR32	32.0	6.4	61.8*	8.3	31.8	12.2	39.8	6.2
<i>L. casei</i> L324m	59.0	6.9	54.7	4.6	27.7	5.0	36.6	3.3
<i>L. acidophilus</i> NCFM	56.3	4.2	34.2*	7.0	27.4	5.5	25.5	9.1

* Significant difference between initial colonization and maturation (p<0.05).

Table 1. Percentage of reduction of viable *C. albicans* cells on biofilms after treatment with probiotic bacteria suspension (1×10^7 probiotic cells/ml) during initial colonization (24 h) and maturation (48 h) phases.