



**McMillan, Kathryn Anne and Kirk, Alison and Hewitt, Allan and MacRury, Sandra (2017) Objective, continuous measurement of sedentary behaviour and glucose in people with type 2 diabetes. In: Diabetes UK Professional Conference 2017, 2017-03-08 - 2017-03-10, Manchester Central. ,**

This version is available at <https://strathprints.strath.ac.uk/62241/>

**Strathprints** is designed to allow users to access the research output of the University of Strathclyde. Unless otherwise explicitly stated on the manuscript, Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Please check the manuscript for details of any other licences that may have been applied. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<https://strathprints.strath.ac.uk/>) and the content of this paper for research or private study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to the Strathprints administrator: [strathprints@strath.ac.uk](mailto:strathprints@strath.ac.uk)

# Objective, continuous measurement of sedentary behaviour and glucose in people with Type 2 Diabetes

K.A. McMillan<sup>1</sup>, A. Kirk<sup>1</sup>, A. Hewitt<sup>1</sup> & S. MacRury<sup>2</sup>

<sup>1</sup>Physical Activity for Health Research Group, School of Psychological Sciences and Health, University of Strathclyde, Glasgow

<sup>2</sup>Highland Diabetes Institute, University of Highlands and Islands, Inverness

## Introduction

- Substantial benefits of leading an active lifestyle following diagnosis of Type 2 diabetes have been identified<sup>1</sup>
- Reducing sedentary time has been shown to be effective in the management of blood glucose levels irrespective of physical activity levels<sup>2</sup>
- Regular breaks in sedentary behaviour with light intensity walking are associated with reduced variability in glucose<sup>3</sup>
- Little research has examined daily levels of glucose and sedentary behaviour in those with Type 2 diabetes using objective and continuous measurements in a free living context

## Aims

To explore levels of sedentary behaviour in those with Type 2 diabetes, and investigate the relationship between daily levels of sedentary behaviour and daily mean glucose in people with Type 2 diabetes using objective and continuous measurements

## Methods

- Ten participants with Type 2 diabetes managed with diet, Metformin or DPP4 inhibitors were recruited
- Participants completed a demographic questionnaire and wore an activPAL accelerometer and a FreeStyle Libre continuous glucose monitor for 3-14 days
- Participants were also required to document their sleep, food and medication in diaries
- Average proportion of time spent sedentary and daily mean glucose during the waking day were calculated.
- A multiple linear regression was calculated to explore the relationship between sedentary time, age, BMI, and daily mean glucose

## Example of activPAL and FreeStyle Libre Devices



## References

<sup>1</sup>Umpierre et al. (2011). Physical activity advice only or structured exercise training and association with HbA1c levels in type 2 diabetes: a systematic review and meta-analysis. *Jama*. 305.17,1790-1799.


<sup>2</sup>Dunstan et al. (2012). Breaking up prolonged sitting reduces postprandial glucose and insulin responses. *Diabetes care*. 35.5, 976-983.

<sup>3</sup>Dempsey et al. (2016). Benefits for type 2 diabetes of interrupting prolonged sitting with brief bouts of light walking or simple resistance activities. *Diabetes Care*. 39(6), 964-972.

## Results

- 10 adults with a mean age of 63.5±9.4 years participated
- Mean BMI was calculated as 30.8±6.9 kg/cm<sup>2</sup>
- Participants spent 67.5% of their waking day sitting/ lying (Figure 1)
- Daily mean glucose was calculated as 7.7±1.8mmol/l
- Regression analysis suggested a significant effect (F(3,105) = 16.52, *p* <0.01) for sitting/lying time, age and BMI on daily mean glucose (R<sup>2</sup> = 0.30)
- Daily mean glucose increased significantly (*p* <0.01) with increased sedentary time, BMI and age. With daily mean glucose increasing by 5.14mmol/l for each percent of daily sitting/lying, 0.16mmol/l for each increasing year of age and 0.21mmol/l for each increasing BMI unit

## Proportion of time spent sitting/ lying, standing and stepping



## Discussion and Conclusions

- Average sitting/lying time was higher than has previously been reported
- Percentage sitting/lying time, age and BMI were identified as significant predictors of higher daily mean glucose in people with Type 2 diabetes
- Results suggest that increased sedentariness is associated with increased daily mean glucose in those with Type 2 diabetes

## Recommendations

- Future analysis should focus on examining the relationship between daily sedentary behaviour and glucose in a larger sample size
- Patterns in sedentary behaviour and resultant patterns in glucose should also be examined

## Author's contact details:

[kathryn.mcmillan@strath.ac.uk](mailto:kathryn.mcmillan@strath.ac.uk)

Twitter:

KathrynMc91