



de Souza, W. M. et al. (2018) Discovery of novel anelloviruses in small mammals expands the host range and diversity of the Anelloviridae. *Virology*, 514, pp. 9-17. (doi:[10.1016/j.virol.2017.11.001](https://doi.org/10.1016/j.virol.2017.11.001))

This is the author's final accepted version.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/151586/>

Deposited on: 09 February 2018

Enlighten – Research publications by members of the University of Glasgow  
<http://eprints.gla.ac.uk>

**Running Title:** Novel anelloviruses in small mammals

**Title:** Discovery of novel anelloviruses in small mammals expands the host range and diversity of the *Anelloviridae*

**Authors:** William Marciel de Souza<sup>1,2†</sup>, Marcílio Jorge Fumagalli<sup>1†</sup>, Jansen de Araujo<sup>3</sup>, Gilberto Sabino-Santos Jr<sup>1</sup>, Felipe Gonçalves Motta Maia<sup>1,4</sup>, Marilia Farignoli Romeiro<sup>1</sup>, Sejal Modha<sup>2</sup>, Marcello Schiavo Nardi<sup>5</sup>, Luzia Helena Queiroz<sup>6</sup>, Edison Luiz Durigon<sup>3</sup>, Márcio Roberto Teixeira Nunes<sup>7,8</sup>, Pablo Ramiro Murcia<sup>2</sup> and Luiz Tadeu Moraes Figueiredo<sup>1</sup>.

†These first authors contributed equally to this article.

**Affiliations:**

<sup>1</sup> Virology Research Center, Ribeirão Preto Medical School, University of São Paulo, Ribeirão Preto, Brazil

<sup>2</sup> MRC-University of Glasgow Centre for Virus Research, Glasgow, United Kingdom.

<sup>3</sup> Laboratory Institute of Biomedical Sciences, University of São Paulo, São Paulo, Brazil.

<sup>4</sup> Department of Microbiology, Institute of Biomedical Sciences, University of São Paulo, São Paulo, Brazil.

<sup>5</sup> Divisão Técnica de Medicina Veterinária e Manejo da Fauna Silvestre, Prefeitura de São Paulo, Brazil.

<sup>6</sup> Faculty of Veterinary Medicine, São Paulo State University, Araçatuba, Brazil.

<sup>7</sup> Center for Technological Innovations, Evandro Chagas Institute, Ministry of Health, Ananindeua, Pará, Brazil.

<sup>8</sup> Department of Pathology, University of Texas Medical Branch, Galveston, TX, USA.

**# Corresponding author:** William Marciel de Souza, Virology Research Center, School of Medicine of Ribeirão Preto of University of São Paulo. Av. Bandeirantes, 3900, Monte Alegre, 14049-900, Ribeirão Preto, SP Brazil. E-mail: wmarciel@usp.br or wmarciel@hotmail.com Phone: +55 16 3315-4580.

## **ABSTRACT**

The *Anelloviridae* comprises single-stranded DNA viruses currently grouped in **sixty-eight** species classified in twelve genera. They have been found in **many** vertebrate hosts including primates. In this study, we describe the application of the high-throughput sequencing to examine the frequency and diversity of anelloviruses in rodents, bats and opossums captured in São Paulo State, Brazil. We report a total of twenty-six anelloviruses with sixteen nearly complete genomes and ten partial genomes, which include eleven **potential** novel species identified in rodents (Cricetidae), bats (Molossidae and Phyllostomidae), and opossums (Didelphidae). We also propose the inclusion of two **potential** new genera within the *Anelloviridae* family, **provisionally** named *Omegatorquevirus* and *Sigmatorquevirus*, including six and three novel species of anelloviruses, respectively. In summary, this study expands the diversity and the host range of the known anelloviruses.

**Keywords:** Anellovirus; rodent-borne virus; bat-borne virus; *Anelloviridae*; virome, and ssDNA viruses.

## Introduction

*Torque teno virus* (TTV) was discovered as a possible cause of post-transfusion hepatitis in humans in 1997 in Japan (Nishizawa et al., 1997). Currently, the *Anelloviridae* family comprises about 68 species, which are non-enveloped, circular, single-stranded DNA (ssDNA) viruses with a genome of 2.1 to 3.9 kb in length classified into 12 genera (Biagini, 2009; Biagini, 2017; Biagini et al., 2012).

Anelloviruses are associated with chronic infection without causing overt diseases and their prevalence in humans ranges from 5% to 90% (Spandole et al., 2015). Many studies have associated anelloviruses with a broad spectrum of diseases including hepatitis, multiple sclerosis, hepatocellular carcinomas, respiratory infections, blood disorders and autoimmune diseases, but these studies have not been confirmed that anelloviruses are the etiological agent of these diseases (Garcia-Alvarez et al., 2013; Mancuso et al., 2013; Masouridi-Levrat et al., 2016; Spandole et al., 2015; Tyschik et al., 2017). On the other hand, anelloviruses have been showed to be present on healthy individuals, and are commonly detected with high prevalence in several types of samples, such as blood, respiratory tract, gut, skin and human genitourinary tract (Afkari et al., 2012; Bernardin et al., 2010; Biagini et al., 1998; Blazsek et al., 2008; Breitbart and Rohwer, 2005; Burian et al., 2011; Chan et al., 2001; Lin et al., 2000; Okamura et al., 1999; Popgeorgiev et al., 2013; Pride et al., 2012). In addition, the viral load of anelloviruses in transplant recipients has been showed as one marker of the overall state of immunosuppression (De Vlaminck et al., 2013).

Anelloviruses are spread primarily through fecal-oral transmission and saliva (Ali et al., 2002; Maggi et al., 2003; Nishiyama et al., 2014). The lack of suitable cell culture systems and animal models impairs the study of anelloviruses in laboratory settings and thus we know very little about the specific function of these viral genes and proteins in pathogenesis (Hrazdilova et al., 2016).

Currently, anelloviruses have been identified in several mammals, such as primates, dogs, cats, pigs, rodents, bats and seals (Cibulski et al., 2014; Hrazdilova et al., 2016; Ng et al., 2011a; Nishiyama et al., 2015; Nishiyama et al., 2014; Okamoto, 2009; Shi et al., 2015). They have also been detected in invertebrates (i.e. mosquitoes), but probably due to blood feeding of mosquitoes obtained from vertebrate hosts (Ng et al., 2011b). Also, the Chicken anemia virus is a pathogen of chickens and has a worldwide distribution, which was included in *Anelloviridae* family (Schat, 2009).

In rodents, studies are limited to members of the subfamily Arvicolinae family Cricetidae and Muridae family (Nishiyama et al., 2015; Nishiyama et al., 2014). A comprehensive characterization of the natural hosts of anelloviruses is central to understanding their host range and transmission dynamics in nature. For this reason it is necessary to perform surveillance studies in other members of the Cricetidae rodents as well as in other mammalian species. In this study, we have applied the high-throughput sequencing (HTS) approach to determine the presence of the anelloviruses in small mammals, including Sigmodontinae rodents (Cricetidae family), bats of Molossidae and Phyllostomidae families, and opossums of Didelphidae family from various geographical locations of Southeastern Brazil.

## Materials and Methods

## Samples

A total of 919 specimens obtained from 13 different animal species were collected from 2008 to 2013 from rural areas of São Paulo State, South East Brazil. Individual specimens were pooled (n=40) based on the species, sample type (i.e. tissue, blood or sera), date and place of collection (**Supplementary Table 1**). To visualize the geographical distribution of the pools, we used the Tableau software v.10.1.1 (Tableau, USA) with referential geographical position for map generation, and the total of host pools were represented in clusters (**Figure 1**). The species of small mammals were identified using morphological characteristics keys as previously described (Bonvicino et al., 2008; Goes et al., 2016; Sikes and Gannon, 2011).

## Preparation of pools

Pool samples were processed using TissueLyser II (Qiagen, USA). Each sample tube was added a 5-mm stainless steel bead (Qiagen, USA) and the samples were processed for tissue disruption during 10 min at 30 Hz using TissueLyser II (Qiagen, USA). Then, the samples were centrifuge during 5 min at 10,000g. Subsequently, the supernatant was collected and filtered using 0.22 µm-pore-size polyvinylidene difluoride filters (Millipore, USA) to minimize the presence of eukaryotic cells and bacteria. Then, the pools were ultracentrifuged at 180,000xg for three hours at 4°C in 30% sucrose solution (Sorvall WX90 Ultracentrifuge, Thermo, USA), and pellets resuspended in 500µl of Hanks' balanced salt solution. To remove the naked DNA and RNA, 200µl of the resuspended pellet from each pooled sample were digested in a cocktail with 20U of Turbo DNase (Life Technologies, USA), 25U of benzonase (Sigma-Aldrich, USA), and 0.1 mg/ml of RNase A (Life Technologies,

USA) at 37°C for 2 hours in 20 µl of 10X DNase buffer (Life Technologies, USA). The viral genomes were extracted with a QIAamp viral RNA mini kit (Qiagen, USA).

### **Genome sequencing and assembly**

The DNAs were prepared for high-throughput sequencing using RAPID module with the TruSeq Universal adapter (Illumina, USA) protocols and standard multiplex adaptors. A paired-end, 150-base-read protocol in RAPID module was used for sequencing on an Illumina HiSeq 2500 instrument as recommended by the manufacturer. Sequencing was performed in Life Sciences Core Facility (LaCTAD) from State University of Campinas (UNICAMP), Brazil. A total of 7,059,398 to 94,508,748 paired end reads per pool were generated with 64.85% to 91.13% of bases  $\geq$  Q30 with a base call accuracy of 99.9% (**Supplementary Table 1**). The resulting sequencing reads were cleaned and *de novo* assembled using the MetaViC pipeline (Available on <https://github.com/sejmodha/MetaViC>).

### **Genome characterization**

Viral genomes were assessed for the genome size and ORF prediction with Geneious 9.1.2 (Biomatters, New Zealand) and CLC sequence viewer 7.7.1 (Qiagen, Germany). The ORFs were predicted using BLASTX database and based on position and nucleotide sequence similarities on NCBI database. GPMiner and Poly(A) signal miner webservers were used to predict the presence of TATA box regions and polyadenylation signals, respectively (Lee et al., 2012; Liu et al., 2003). GC-Rich regions and the molecular weight of putative proteins were predicted using Snapgene software (GSL biotech, USA) and ProtParam database, respectively (Artimo et al.,

2012). The nucleotide sequences determined in this study were deposited in GenBank under the accession numbers as described in **Table 1**.

### **Phylogenetic analysis**

Maximum likelihood (ML) phylogenetic trees were inferred using alignments on nucleotides level of predicted ORF1 identified in the present study with ORF1 of representative members of *Anelloviridae* family. Multiple sequence alignments (MSA) were generated using MAFFT v7.309 in RevTrans 2.0 (Wernersson and Pedersen, 2003). ML trees were inferred using IQ-TREE version 1.4.3 software using GTR+I+G4 nucleotides substitution model with 1,000 ultrafast bootstraps, as determined by the best-fit model based on Bayesian Information Criterion considered 88 reversible nucleotide substitution models (Kalyaanamoorthy et al., 2017; Nguyen et al., 2015). Statistical support for individual nodes was estimated using the bootstrap value. The phylogenetic trees were visualized using the FigTree software v.1.4.2. In addition, nucleotide and amino acids identifies were performed based on same alignments described above using the UGENE version 1.24.1 (Okonechnikov et al., 2012).

### **Frequency and diversity**

We calculated the frequency and diversity of the anelloviruses based on the host species and the sample type. The frequency was calculated based on the number of positive pooled samples containing anelloviruses divided by total number of pools for their respective host, and clustered by sample type. The diversity of anelloviruses species for each host was calculated based on the amount of different anelloviruses species identified by samples type and divided by the total anelloviruses species



identified reported in this study. The data were calculated using Tableau software v.10.1.1 (Tableau, USA) and the relative proportion was normalized to 100% for visualization purposes.

## Results

### Genome characterization of novel anelloviruses

The anelloviruses assemblies represent 1 to 33 % of viral assemblies, and 0.01 to 2% of total of contigs assemblies. From this result, we identified 26 anellovirus contigs that were classified into eleven novel species. The viruses were identified in samples derived from rodents, bats and opossums from São Paulo State (**Figure 1**), which included sixteen nearly complete genome and ten partial genomes (**Table 1**). The viruses identified in our study possess a ssDNA genome of 2.3 to 2.7 kb in length. Most anellovirus sequences had an untranslated region (UTR) with a predicted GC-rich region (>60% GC), a putative TATA box region and a polyadenylation tail signal (**Figure 2**).

All anellovirus sequences included *ORF1*, which encodes the putative ORF1 protein of 541 to 580 amino acids (aa) in length. The only exception is *Torque teno calomys tener virus* (TTCtV), which displayed a putative ORF1 of 337 aa, which is homologous to strains of *Rodent Torque teno virus 1* classified into same genus, the *Deltatorquevirus* genus. We also identified the *ORF2* that encodes a putative protein of 103 to 168 aa in length in *Opossum torque teno virus* (OTTV) and *Torque Teno didelphis albiventris virus* (TTDaV) (**Figure 2**). *ORF3* encodes a putative protein of 127 to 154 aa in length and it was identified in TTCtV, TTDaV, *Torque teno desmodus rotundus* (TTDrV) and *Torque teno carollia perspicillata* (TTCpV). In addition, we identified previously uncharacterized putative ORFs called “ORF?”,

potentially encoding proteins of 109 to 152 aa in length in *Rodent Torque teno virus* 3, 4, 6 and 7 (RoTTV), TTD<sub>r</sub>V and TTC<sub>t</sub>V (**Figure 2**). “ORF?” did not display homology with other anellovirus proteins, or with proteins of other viral families.

### **Phylogenetic analysis reveals two new genera into *Anelloviridae* family**

To determine the phylogenetic relationships amongst the novel anelloviruses sequences and those previously described, we inferred a phylogenetic tree that included 123 complete ORF1 nucleotide sequences (**Figure 3**). All known anelloviruses were split in two major groups, the primates and non-primates anelloviruses, which could be placed into 14 monophyletic clades (bootstrap values >70%). Twelve clades corresponded to established genera within the *Anelloviridae*, which include 68 species of anelloviruses recognized by ICTV into *Alphatorquevirus*, *Betatorquevirus*, *Deltatorquevirus*, *Epsilontorquevirus*, *Etatorquevirus*, *Gammatorquevirus*, *Gyrovirus*, *Iotatorquevirus*, *Kappatorquevirus*, *Lambdatorquevirus*, *Thetatorquevirus* and *Zetatorquevirus* genus (Biagini, 2017). Furthermore, we identified two clades that were distantly related to the aforementioned genera (**Figure 3**). To determine if the sequences identified in host two clades belong to previously uncharacterized genera, we calculated the pairwise genetic identity based on nucleotide level of ORF1 of all anelloviruses sequenced in our study combined with anelloviruses available in Genbank (**Supplementary Figure 1**). Using the taxonomic classification criteria of ICTV (i.e cut-off values for nucleotide sequence distance: isolates 20%, species 20 to 55%) (Biagini et al., 2012), we confirmed that eleven sequences constituted potential novel species of anelloviruses in four genera, and that two divergent clades did indeed belong to potential two novel genera within the *Anelloviridae*, which we proposed as

provisional named of *Omegatorquevirus* and *Sigmatorquevirus* (**Supplementary Figure 1**).

The genus *Omegatorquevirus* is constituted by the **RoTTV 4 to 8** identified in our study, which is composed of four anelloviruses that were previously reported, the *Rodent Torque teno virus* strain RN and AS 1014, as well as the *Mosquito VEM Anellovirus SDRB A and B* (Nishiyama et al., 2014; Shi et al., 2015) (**Figure 3**). On the other hand, the genus *Sigmatorquevirus* that we propose here is formed by four viruses that represent three novel species, the new species TTDaV and TTCpV, as well as TTDrV that is related with *Torque teno tadarida brasiliensis virus* (TTTbV) (Cibulski et al., 2014). Interestingly, TTDrV and TTTbV were found in different hosts despite exhibiting 91% of nucleotide identity, therefore they are considered the same viral species. In addition, we have identified two novel anelloviruses species, the TTCtV and **OTTV** classified into the *Deltatorquevirus* and *Alphatorquevirus* genera, respectively (**Figure 3**).

### **Frequency and diversity of anelloviruses**

Eighteen of forty-five pools were positive to anelloviruses, which includes nine of the thirteen species of small mammals species tested. We observed a high frequency of anelloviruses, including more than seven different species of small mammals with frequencies higher than 40% (**Figure 4a**). The most prevalent host species positive for anelloviruses were *Necromys lasiurus* and *Mus musculus* with 100% (**5/5 and 1/1**) of detection in the pools, followed by *Calomys tener* with 70% (**2/3**), all of which were detected in blood samples (**Figure 4a**). On the other hand, the greatest diversity of anelloviruses species were observed in *Oligoryzomys nigripes* and *Akodon montesis* with 36% (**4/11**) and 27% (**3/11**) of the total novel anelloviruses

species, respectively. Interestingly, different anelloviruses species were found in blood, liver and in both samples of *Oligoryzomys nigripes*, showing the greatest diversity when compared to the other species (Figure 4b).

## Discussion

The *Anelloviridae* have been detected in blood or serum of a broad spectrum of mammals, including primates, pigs, horses, cats, dogs, rodents and bats (Cibulski et al., 2014; Hrazdilova et al., 2016; Nishiyama et al., 2014; Okamoto, 2009; Okamoto et al., 2002). Our finding, which include the discovery of eleven novel species in previously unrecognized host species and the identification of two probably new anellovirus genera highlights the diversity of this viral family.

The genome of anelloviruses can vary in length depending on the host species (Manzin et al., 2015). The size of the genomes identified in this study were consistent with previous findings, being the only exception OTTV identified in *Didelphis albiventris*. This genome was larger than expected, and it was similar in size to those of *Alphatorqueviruses* described in primates. The UTRs of the viruses were conserved similar to those of previously reported anelloviruses. This suggests that UTRs must play important regulatory roles during viral replication (Spandole et al., 2015). Features such as the TATA box and polyadenylation signal are important during transcription of viral genes (Paran et al., 2000). We were able to map the TATA box and polyadenylation signals based on ORFs position for most of the viruses identified. However, experimental work is necessary to elucidate the function of these regulators.

By combining our phylogenetic tree with pairwise distance analysis and using the classification criteria of ICTV, we assigned eleven sequences to novel anelloviruses species (Biagini, 2009; Biagini, 2017; Biagini et al., 2012). Also, we

identified two [potential](#) new genera within the *Anelloviridae*, which we proposed to [provisional names of](#) *Omegatorquevirus* and *Sigmatorquevirus*. These two [potential](#) genera form divergent the clades with a nucleotide distance greater than 50% to the other twelve genera. The identification of two new genera into *Anelloviridae* family increases the diversity of the described anelloviruses. Also, while the inferred phylogenetic tree indicates an origin of the anelloviruses from non-primates, there is no clear evidence to state whether anelloviruses originated as a result of co-evolution with their host during millions of years or whether anelloviruses have originated and diversified over a much shorter time scale than their hosts. Such interesting question should be answered in future studies (Okamoto et al., 2001).

Sequences belonging to the *Omegatorquevirus* genus were the most frequently identified in our study. The most common species were [RoTTV-4](#) (eleven strains) and [RoTTV-5](#) (six strains) detected in *Akodon montesis*, *Akodon cursor*, *Calomys tener*, *Necromys lasiurus*, *Mus musculus* and *Oligoryzomys nigripes*. Almost all anelloviruses were identified in the rodent subfamily *Sigmodontinae* (*Cricetidae* family), which is predominantly South American (Parada et al., 2015). Interestingly, the *Omegatorquevirus* genus form a monophyletic clade with viruses most commonly found in rodents of the *Cricetidae* family (Nishiyama et al., 2014). The exceptions are Mos VEM SDRB A and B viruses [detected in mosquitoes from USA, which was suspected to be due the feed from rodents blood that potentially may be the original viral source](#) (Ng et al., 2011b) and the [RoTTV-4](#) strain 117 that was identified in *Mus musculus* (*Muridae* family), which have a wide distribution worldwide and anthropophilic behavior (Pocock et al., 2004). [Contrasting with Rodent TTV screening study in rodents captured in United Kingdom, which showed that \*Mus musculus\* mouse collected were absent to the presence of anelloviruses in samples of](#)

spleen and liver (Nishiyama et al., 2014)

Bats (order *Chiroptera*) have been recognized as important sources of viruses that can potentially cause human and veterinary diseases (Brook and Dobson, 2015). However, the bat anellovirus is restricted to a single species reported in the insectivorous phyllostomide *Tadarida brasiliensis* in Southern Brazil (Cibulski et al., 2014). We have also identified two novel anelloviruses species in *Carollia perspicillata* and *Desmodus rotundus*. These viruses form a unique monophyletic clade that we propose as *Sigmatorquevirus* genus. Another sequence belonging to the *Sigmatorquevirus* genus is TTDaV, which was derived from an opossum. Interestingly, OTTV, classified into *Alphatorquevirus* genus, was also obtained from an opossum. This is surprising because all members of *Alphatorquevirus* were identified only in primates (Spandole et al., 2015). We speculate that OTTV could be the result of anthroozoonosis, as recently proposed to Torque Teno viruses of human and swine (Ssemadaali et al., 2016). Finally, we identified in *Calomys tener* new specie within the *Deltatorquervirus* genus. This sequence clustered with other anelloviruses previously reported in rodents, shrew and mosquitos (Ng et al., 2011a; Nishiyama et al., 2014; Okamoto et al., 2001).

In sum, this is the first epidemiological study of anelloviruses in wild rodents and marsupials in the Americas, and it provides new information on the diversity and host range of anelloviruses in nature.

### **Acknowledgements**

We thank Luiz Gustavo Betim Góes, Cristiano de Carvalho and Wagner André Pedro for the help with the capture and identification of bats in Araçatuba region, São Paulo. We thank the field team of University of São Paulo: Fábio Maués,

Tatiana Ometto, Renata Hurtado and Felipe Alves Morais. We also appreciate the help in field campaigns of Márcio Schaefer, Mario Figueiredo, Felipe Morais, Jaqueline R. Silva, Paulo Paulosso and Dercílio Pavanelli. We are grateful to Edison Montilha, the Jataí Ecological Station manager for all the help, and Armando F. A. Nascimento who provide us a site for research and help on his farm Santa Gabriela in Batatais county, as well as José Teotônio (Zezinho) and the Secretary of Health from Cajuru county.

**Funding:** This work was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo, Brazil (Grant number. 13/14929-1, and Scholarships N°. 12/24150-9; 15/05778-5; 16/01414-1; 14/20851-8; 06/00572-0; 08/06411-4; 11/06810-9; 11/22663-6; 16/02568-2, 06/00572-0). PRM was supported by the Medical Research Council of the United Kingdom (Grant MC\_UU\_120/14/9).

## References

- Afkari, R., Pirouzi, A., Mohsenzadeh, M., Azadi, M., Jafari, M., 2012. Molecular detection of TT virus and SEN virus infections in hemodialysed patients and blood donors in south of Iran. *Indian journal of pathology & microbiology* 55, 478-480.
- Ali, S., Fevery, J., Peerlinck, K., Verslype, C., Schelstraete, R., Gyselinck, F., Emonds, M.P., Vermylen, J., Yap, S.H., 2002. TTV infection and its relation to serum transaminases in apparently healthy blood donors and in patients with clotting disorders who have been investigated previously for hepatitis C virus and GBV-C/HGV infection in Belgium. *J Med Virol* 66, 561-566.
- Artimo, P., Jonnalagedda, M., Arnold, K., Baratin, D., Csardi, G., de Castro, E., Duvaud, S., Flegel, V., Fortier, A., Gasteiger, E., Grosdidier, A., Hernandez, C.,

Ioannidis, V., Kuznetsov, D., Liechti, R., Moretti, S., Mostaguir, K., Redaschi, N., Rossier, G., Xenarios, I., Stockinger, H., 2012. ExPASy: SIB bioinformatics resource portal. *Nucleic acids research* 40, W597-603.

Bernardin, F., Operskalski, E., Busch, M., Delwart, E., 2010. Transfusion transmission of highly prevalent commensal human viruses. *Transfusion* 50, 2474-2483.

Biagini, P., 2009. Classification of TTV and related viruses (anelloviruses). *Current topics in microbiology and immunology* 331, 21-33.

Biagini, P., 2017. Create the family Anelloviridae containing 9 new genera and 47 new species. *International Committee on Taxonomy of Viruses*, <https://talk.ictvonline.org/taxonomy/>, pp. Code assigned: 2007.2075a-xxV.

Biagini, P., Bendinelli, M., Hino, S., Kakkola, L., Mankertz, A., Niel, C., Okamoto, H., Raidal, S., Teo, C.G., Todd, D., 2012. Family Anelloviridae in: King, A.M.Q., Adams, M.J., Carstens, E.B., Lefkowitz, E.J. (Eds.), *Virus Taxonomy - Ninth Report of the International Committee on Taxonomy of Viruses*. Elsevier pp. 331-341.

Biagini, P., Gallian, P., Cantaloube, J.F., De Micco, P., de Lamballerie, X., 1998. Presence of TT virus in French blood donors and intravenous drug users. *Journal of hepatology* 29, 684-685.

Blazsek, A., Sillo, P., Ishii, N., Gergely, P., Jr., Poor, G., Preisz, K., Hashimoto, T., Medvecz, M., Karpati, S., 2008. Searching for foreign antigens as possible triggering factors of autoimmunity: Torque Teno virus DNA prevalence is elevated in sera of patients with bullous pemphigoid. *Experimental dermatology* 17, 446-454.

Bonvicino, C.R., Oliveira, J.A., D'Andrea, P.S., 2008. *Guia dos roedores do Brasil, com chaves para gêneros baseadas em caracteres externos*, 1st ed. Centro Pan-



Americano de Febre Aftosa-Pan-American Health Organization//World Health Organization, Rio de Janeiro, Brazil.

Breitbart, M., Rohwer, F., 2005. Method for discovering novel DNA viruses in blood using viral particle selection and shotgun sequencing. *BioTechniques* 39, 729-736.

Brook, C.E., Dobson, A.P., 2015. Bats as 'special' reservoirs for emerging zoonotic pathogens. *Trends in microbiology* 23, 172-180.

Burian, Z., Szabo, H., Szekely, G., Gyurkovits, K., Pankovics, P., Farkas, T., Reuter, G., 2011. Detection and follow-up of torque teno midi virus ("small anelloviruses") in nasopharyngeal aspirates and three other human body fluids in children. *Archives of virology* 156, 1537-1541.

Chan, P.K., Tam, W.H., Yeo, W., Cheung, J.L., Zhong, S., Cheng, A.F., 2001. High carriage rate of TT virus in the cervixes of pregnant women. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* 32, 1376-1377.

Cibulski, S.P., Teixeira, T.F., de Sales Lima, F.E., do Santos, H.F., Franco, A.C., Roehle, P.M., 2014. A Novel Anelloviridae Species Detected in *Tadarida brasiliensis* Bats: First Sequence of a Chiropteran Anellovirus. *Genome announcements* 2.

De Vlamincq, I., Khush, K.K., Strehl, C., Kohli, B., Luikart, H., Neff, N.F., Okamoto, J., Snyder, T.M., Cornfield, D.N., Nicolls, M.R., Weill, D., Bernstein, D., Valantine, H.A., Quake, S.R., 2013. Temporal response of the human virome to immunosuppression and antiviral therapy. *Cell* 155, 1178-1187.

Garcia-Alvarez, M., Berenguer, J., Alvarez, E., Guzman-Fulgencio, M., Cosin, J., Miralles, P., Catalan, P., Lopez, J.C., Rodriguez, J.M., Micheloud, D., Munoz-Fernandez, M.A., Resino, S., 2013. Association of torque teno virus (TTV) and torque teno mini virus (TTMV) with liver disease among patients coinfecting with human

immunodeficiency virus and hepatitis C virus. *European journal of clinical microbiology & infectious diseases* : official publication of the European Society of Clinical Microbiology 32, 289-297.

Goes, L.G., Campos, A.C., Carvalho, C., Ambar, G., Queiroz, L.H., Cruz-Neto, A.P., Munir, M., Durigon, E.L., 2016. Genetic diversity of bats coronaviruses in the Atlantic Forest hotspot biome, Brazil. *Infection, genetics and evolution : journal of molecular epidemiology and evolutionary genetics in infectious diseases* 44, 510-513.

Hrazdilova, K., Slaninkova, E., Brozova, K., Modry, D., Vodicka, R., Celer, V., 2016. New species of Torque Teno miniviruses infecting gorillas and chimpanzees. *Virology* 487, 207-214.

[Kalyaanamoorthy, S., Minh, B.Q., Wong, T.K.F., von Haeseler, A., Jermini, L.S., 2017. ModelFinder: fast model selection for accurate phylogenetic estimates. \*Nature methods\* 14, 587-589.](#)

Lee, T.Y., Chang, W.C., Hsu, J.B., Chang, T.H., Shien, D.M., 2012. GPMIner: an integrated system for mining combinatorial cis-regulatory elements in mammalian gene group. *BMC genomics* 13 Suppl 1, S3.

[Lin, C.L., Kyono, W., Tongson, J., Chua, P.K., Easa, D., Yanagihara, R., Nerurkar, V.R., 2000. Fecal excretion of a novel human circovirus, TT virus, in healthy children. \*Clinical and diagnostic laboratory immunology\* 7, 960-963.](#)

Liu, H., Han, H., Li, J., Wong, L., 2003. An in-silico method for prediction of polyadenylation signals in human sequences. *Genome informatics. International Conference on Genome Informatics* 14, 84-93.

Maggi, F., Pifferi, M., Fornai, C., Andreoli, E., Tempestini, E., Vatteroni, M., Presciuttini, S., Marchi, S., Pietrobelli, A., Boner, A., Pistello, M., Bendinelli, M.,

2003. TT virus in the nasal secretions of children with acute respiratory diseases: relations to viremia and disease severity. *J Virol* 77, 2418-2425.

Mancuso, R., Saresella, M., Hernis, A., Agostini, S., Piancone, F., Caputo, D., Maggi, F., Clerici, M., 2013. Torque teno virus (TTV) in multiple sclerosis patients with different patterns of disease. *Journal of medical virology* 85, 2176-2183.

Manzin, A., Mallus, F., Macera, L., Maggi, F., Blois, S., 2015. Global impact of Torque teno virus infection in wild and domesticated animals. *Journal of infection in developing countries* 9, 562-570.

Masouridi-Levrat, S., Pradier, A., Simonetta, F., Kaiser, L., Chalandon, Y., Roosnek, E., 2016. Torque teno virus in patients undergoing allogeneic hematopoietic stem cell transplantation for hematological malignancies. *Bone marrow transplantation* 51, 440-442.

Ng, T.F., Wheeler, E., Greig, D., Waltzek, T.B., Gulland, F., Breitbart, M., 2011a. Metagenomic identification of a novel anellovirus in Pacific harbor seal (*Phoca vitulina richardsii*) lung samples and its detection in samples from multiple years. *The Journal of general virology* 92, 1318-1323.

Ng, T.F., Willner, D.L., Lim, Y.W., Schmieder, R., Chau, B., Nilsson, C., Anthony, S., Ruan, Y., Rohwer, F., Breitbart, M., 2011b. Broad surveys of DNA viral diversity obtained through viral metagenomics of mosquitoes. *PloS one* 6, e20579.

Nguyen, L.T., Schmidt, H.A., von Haeseler, A., Minh, B.Q., 2015. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular biology and evolution* 32, 268-274.

Nishiyama, S., Dutia, B.M., Sharp, C.P., 2015. Complete genome sequences of novel anelloviruses from laboratory rats. *Genome announcements* 3.

Nishiyama, S., Dutia, B.M., Stewart, J.P., Meredith, A.L., Shaw, D.J., Simmonds, P., Sharp, C.P., 2014. Identification of novel anelloviruses with broad diversity in UK rodents. *The Journal of general virology* 95, 1544-1553.

Nishizawa, T., Okamoto, H., Konishi, K., Yoshizawa, H., Miyakawa, Y., Mayumi, M., 1997. A novel DNA virus (TTV) associated with elevated transaminase levels in posttransfusion hepatitis of unknown etiology. *Biochemical and biophysical research communications* 241, 92-97.

Okamoto, H., 2009. TT viruses in animals. *Current topics in microbiology and immunology* 331, 35-52.

Okamoto, H., Nishizawa, T., Takahashi, M., Tawara, A., Peng, Y., Kishimoto, J., Wang, Y., 2001. Genomic and evolutionary characterization of TT virus (TTV) in tupaias and comparison with species-specific TTVs in humans and non-human primates. *The Journal of general virology* 82, 2041-2050.

Okamoto, H., Takahashi, M., Nishizawa, T., Tawara, A., Fukai, K., Muramatsu, U., Naito, Y., Yoshikawa, A., 2002. Genomic characterization of TT viruses (TTVs) in pigs, cats and dogs and their relatedness with species-specific TTVs in primates and tupaias. *The Journal of general virology* 83, 1291-1297.

Okamura, A., Yoshioka, M., Kubota, M., Kikuta, H., Ishiko, H., Kobayashi, K., 1999. Detection of a novel DNA virus (TTV) sequence in peripheral blood mononuclear cells. *Journal of medical virology* 58, 174-177.

[Okonechnikov, K., Golosova, O., Fursov, M., team, U., 2012. Unipro UGENE: a unified bioinformatics toolkit. \*Bioinformatics\* 28, 1166-1167.](#)

Parada, A., D'Elia, G., Palma, R.E., 2015. The influence of ecological and geographical context in the radiation of Neotropical sigmodontine rodents. *BMC evolutionary biology* 15, 172.

- Paran, N., Ori, A., Haviv, I., Shaul, Y., 2000. A composite polyadenylation signal with TATA box function. *Molecular and cellular biology* 20, 834-841.
- Pocock, M.J.O., Searle, J.B., White, P.C.L., 2004. Adaptations of animals to commensal habitats: population dynamics of house mice *Mus musculus domesticus* on farms. *Journal of Animal Ecology* 73, 878-888.
- Popgeorgiev, N., Boyer, M., Fancello, L., Monteil, S., Robert, C., Rivet, R., Nappez, C., Azza, S., Chiaroni, J., Raoult, D., Desnues, C., 2013. Marseillevirus-like virus recovered from blood donated by asymptomatic humans. *The Journal of infectious diseases* 208, 1042-1050.
- Pride, D.T., Salzman, J., Haynes, M., Rohwer, F., Davis-Long, C., White, R.A., 3rd, Loomer, P., Armitage, G.C., Relman, D.A., 2012. Evidence of a robust resident bacteriophage population revealed through analysis of the human salivary virome. *The ISME journal* 6, 915-926.
- Schat, K.A., 2009. Chicken anemia virus. *Current topics in microbiology and immunology* 331, 151-183.
- Shi, C., Liu, Y., Hu, X., Xiong, J., Zhang, B., Yuan, Z., 2015. A metagenomic survey of viral abundance and diversity in mosquitoes from Hubei province. *PloS one* 10, e0129845.
- Sikes, R.S., Gannon, W.L., 2011. Guidelines of the American Society of Mammalogists for the use of wild mammals in research. *Journal of Mammalogy* 92.
- Spandole, S., Cimponeriu, D., Berca, L.M., Mihaescu, G., 2015. Human anelloviruses: an update of molecular, epidemiological and clinical aspects. *Archives of virology* 160, 893-908.

Ssemadaali, M.A., Effertz, K., Singh, P., Kolyvushko, O., Ramamoorthy, S., 2016. Identification of heterologous Torque Teno Viruses in humans and swine. *Scientific reports* 6, 26655.

Tyschik, E.A., Shcherbakova, S.M., Ibragimov, R.R., Rebrikov, D.V., 2017. Transplacental transmission of torque teno virus. *Virology journal* 14, 92.

Wernersson, R., Pedersen, A.G., 2003. RevTrans: Multiple alignment of coding DNA from aligned amino acid sequences. *Nucleic acids research* 31, 3537-3539.

### Figure Legends

**Figure 1.** Map of São Paulo State with representative graphs of the numbers of pools per host in the geographic regions of capture.

**Figure 2.** Predicted genetic organization of newly discovered anelloviruses. The name and length of the determined nucleotide sequences of the viral species are shown in the center of the genomic scheme. The putative open reading frames 1 to 3 (ORFs) are showed in yellow, green and gray color, respectively, and the putative undetermined ORFs (ORF?) are indicated in red color. Putative TATA box region and poly A tail signal are presented with box in orange and green color. Dashed-lined indicate incomplete sequence.

**Figure 3.** Maximum likelihood phylogenies based on nucleotide level of ORF 1 showing the relationships of novel viruses identified into *Anelloviridae* family. Phylogenies are midpoint rooted for clarity of presentation. The Chicken anemia virus, an representative genome of Gyrovirus genus was used as outgroup. The scale

bar indicates evolutionary distance in numbers of substitutions per nucleotide site. Branches are colored according to genus into *Anelloviridae* family. Black circle indicate the main nodes with maximum likelihood bootstrap support levels above 75% bootstrap replicates. The viruses described in this study are highlighted in green color and genus proposed highlighted in pink color.

**Figure 4.** Frequency and diversity of anelloviruses identified in this study. (a) Frequency of anelloviruses based on the number of positive sample pools containing anelloviruses divided by total number of pools for their respective host. (b) Diversity of anelloviruses based on the amount of different anelloviruses species identified by samples type and divided by the total anelloviruses species identified. The relative proportion was normalized to 100% for visualization purposes.


**Table 1.** Viruses reported in this study, including the strain name, size in nucleotides, host, ID, sample type, number of animal samples per pool (N), collection region and year. \*Bold denotes prototype genome sequences.












**Figure Supplementary 1.** Heatmap of pairwise amino acid and nucleotides identities of the ORF1 of anelloviruses identified in this study and representative members of *Anelloviridae* family based on ICTV criteria. The amino acids sequences and nucleotide sequences are shown in the upper-right and lower-left of the table, respectively. The viruses described in this study are highlighted with bold.

**Table Supplementary 1.** Samples used in this study. Pools description of host specie, sample type, number of animals per pool (N), sample collection region and

year, the number of generated reads and the sequence quality score  $\geq Q30$  for each pool. Green color denote presence of anelloviruses in the pool and red color denote no presence of anelloviruses.







TTRoV - Torque teno rodent virus  
TTDrV - Torque teno desmodus rotundus virus

TTDaV - Torque teno didelphis albiventris virus  
TTCpV - Torque teno carollia perspicillata virus

TTV - Torque teno virus  
TTCrV - Torque teno calomys tener




**Rodents**



**Bats**



**Opossum**



Alphatorquevirus

Gammatorquevirus

Betatorquevirus

Zetatorquevirus  
Epsilontorquevirus

Deltatorquevirus

Iotatorquevirus

Lambdatorquevirus

Etatorquevirus



Thetatorquevirus

"Sigmatorquevirus"

Kappatorquevirus



"Omegatorquevirus"

NC\_001427 Chicken anemia virus



Genus	Species/Abbreviation	Strain	ORF1	Size	Host	Host Species	ID	Sample	N	Location	Date	Genbank accession number
<i>Deltatorquevirus</i>	<b><i>Torque teno calomys tener virus</i> (TTcTv)</b>	138	Partial	1,865	Rodent	<i>Calomys tener</i>	5B	Blood	38	Ribeirao Preto, SP	2008	MF541391
<i>Sigmatorquevirus</i>	<b><i>Torque teno carollia perspicillata virus</i> (TTCpV)</b>	2	Partial	1,728	Bat	<i>Carollia perspicillata</i>	18	Liver	18	Araçatuba, SP	2010	MF541393
<i>Sigmatorquevirus</i>	<b><i>Torque teno desmodus rotundus virus</i> (TDrV)</b>	182	Complete	2,529	Bat	<i>Desmodus rotundus</i>	24	Liver	8	Araçatuba, SP	2010	MF541386
<i>Sigmatorquevirus</i>	<b><i>Torque teno didelphis albiventris virus</i> (TDDaV)</b>	3470	Complete	2,605	Opossum	<i>Didelphis albiventris</i>	16	Serum	14	Teodoro Sampaio, SP	2009	MF541378
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 3</i> (RoTTV-3)</b>	250	Complete	2,538	Rodent	<i>Necromys lasiurus</i>	10B	Blood	58	Ribeirao Preto, SP	2008	MF541384
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 4</i> (RoTTV-4)</b>	188	Complete	2,561	Rodent	<i>Calomys tener</i>	6B	Blood	37	Ribeirao Preto, SP	2008	MF541383
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 4</i> (RoTTV-4)</b>	117	Complete	2,577	Rodent	<i>Mus musculus</i>	16B	Blood	24	Ribeirao Preto, SP	2008 - 2009	MF541382
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 4</i> (RoTTV-4)</b>	332	Complete	2,684	Rodent	<i>Necromys lasiurus</i>	10B	Blood	58	Ribeirao Preto, SP	2008	MF541376
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 4</i> (RoTTV-4)</b>	4269	Complete	2,594	Rodent	<i>Necromys lasiurus</i>	11B	Blood	52	Ribeirao Preto, SP	2009	MF541379
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 4</i> (RoTTV-4)</b>	1259	Complete	2,588	Rodent	<i>Necromys lasiurus</i>	9B	Blood	59	Ribeirao Preto, SP	2008	MF541380
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 4</i> (RoTTV-4)</b>	108	Complete	2,577	Rodent	<i>Necromys lasiurus</i>	8B	Blood	59	Ribeirao Preto, SP	2008	MF541381
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 4</i> (RoTTV-4)</b>	319	Complete	2,536	Rodent	<i>Necromys lasiurus</i>	12B	Blood	24	Ribeirao Preto, SP	2012 - 2013	MF541385
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 4</i> (RoTTV-4)</b>	67	Partial	1,340	Rodent	<i>Akodon montensis</i>	2B	Blood	55	Ribeirao Preto, SP	2008	MF541395
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 4</i> (RoTTV-4)</b>	152	Partial	1,099	Rodent	<i>Calomys tener</i>	6B	Blood	37	Ribeirao Preto, SP	2008	MF541397
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 4</i> (RoTTV-4)</b>	2216	Partial	1,047	Rodent	<i>Oligoryzomys nigripes</i>	13B	Blood	43	Ribeirao Preto, SP	2008 - 2009	MF541399
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 4</i> (RoTTV-4)</b>	1458	Partial	1,087	Rodent	<i>Oligoryzomys nigripes</i>	35	Liver	10	Vale do Ribeira, SP	2007 - 2008	MF541398
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 5</i> (RoTTV-5)</b>	2	Complete	3,433	Rodent	<i>Akodon montensis</i>	4B	Blood	48	Ribeirao Preto, SP	2012 - 2013	MF541374
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 5</i> (RoTTV-5)</b>	22	Complete	2,617	Rodent	<i>Akodon montensis</i>	4B	Blood	48	Ribeirao Preto, SP	2012 - 2013	MF541377
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 5</i> (RoTTV-5)</b>	2192	Complete	2,689	Rodent	<i>Oligoryzomys nigripes</i>	35	Liver	10	Vale do Ribeira, SP	2007 - 2008	MF541375
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 5</i> (RoTTV-5)</b>	49	Partial	1,830	Rodent	<i>Akodon cursor</i>	33	Lung	4	Nazare Paulista, SP	2009	MF541392
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 5</i> (RoTTV-5)</b>	1012	Partial	1,111	Rodent	<i>Akodon cursor</i>	31	Serum	8	Nazaré Paulista, SP	2009	MF541396
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 5</i> (RoTTV-5)</b>	1594	Partial	1,406	Rodent	<i>Necromys lasiurus</i>	11B	Blood	52	Ribeirao Preto, SP	2009	MF541394
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 6</i> (RoTTV-6)</b>	2404	Complete	2,505	Rodent	<i>Oligoryzomys nigripes</i>	13B	Blood	43	Ribeirao Preto, SP	2008 - 2009	MF541387
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 7</i> (RoTTV-7)</b>	15	Complete	2,379	Rodent	<i>Akodon montensis</i>	4B	Blood	48	Ribeirao Preto, SP	2012 - 2013	MF541388
<i>Omegatorquevirus</i>	<b><i>Rodent Torque teno virus 8</i> (RoTTV-8)</b>	2252	Complete	2,325	Rodent	<i>Oligoryzomys nigripes</i>	35	Liver	10	Vale do Ribeira, SP	2007 - 2008	MF541389
<i>Alphatorquevirus</i>	<b><i>Opossum torque teno virus</i> (OTTV)</b>	13255	Partial	2,220	Opossum	<i>Didelphis albiventris</i>	29	Serum	17	Teodoro Sampaio, SP	2009	MF541390

A large list of taxonomic identifiers (Accession Numbers, GenBank IDs, and other identifiers) is provided on the left side of the image, corresponding to the rows in the heatmap.



**Amino acids**

