

Ecosystem Heterogeneity Determines the Ecological Resilience of the Amazon to Climate Change

Naomi M. Levine^{1,2}
Ke Zhang^{1,3}
Marcos Longo⁴
Alessandro Baccini⁵
Oliver L. Phillips⁶
Simon L. Lewis^{6,7}
Esteban Alvarez-Dávila⁸
Ana Cristina Segalin de Andrade⁹
Roel Brienen⁶
Terry Erwin¹⁰
Ted R. Feldpausch¹¹
Abel Lorenzo Monteagudo Mendoza¹²
Percy Nuñez Vargas¹³
Adriana Prieto¹⁴
Javier Eduardo Silva Espejo¹³
Yadvinder Malhi¹⁵
Paul R. Moorcroft¹

For submission to *Proceedings of the National Academy of Sciences*
Major category: *Biological Sciences*, Minor category: *Sustainability Science*

- ¹ Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA
² Department of Biological Sciences, University of Southern California, Los Angeles, CA
³ Cooperative Institute for Mesoscale Meteorological Studies, University of Oklahoma, Norman, OK
⁴ Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA
⁵ The Woods Hole Research Center, 149 Woods Hole Road, Falmouth, MA
⁶ School of Geography, University of Leeds, Leeds, UK
⁷ Department of Geography, University College London, London, UK
⁸ Jardín Botánico de Medellín, Grupo de Investigación en Servicios Ecosistémicos y Cambio Climático, Medellín, Colombia.
⁹ Instituto Nacional de Pesquisas da Amazônia, Manaus, Brazil
¹⁰ Department of Entomology, Smithsonian Institution, Washington DC
¹¹ College of Life and Environmental Sciences, University of Exeter, Exeter, UK
¹² Jardín Botánico de Missouri, Oxapampa, Perú
¹³ Universidad Nacional de San Antonio Abad del Cusco, Cusco, Perú
¹⁴ Instituto de Ciencias Naturales, Bogotá, Colombia
¹⁵ School of Geography and the Environment, University of Oxford, Oxford, UK

Corresponding author: *Paul R. Moorcroft*

26 Oxford St., Cambridge, MA 02138, USA
paul_moorcroft@harvard.edu, (617) 496-6744

1 **ABSTRACT**

2 Amazon Forests, which store ~50% of tropical forest carbon and play a vital role in global water,
3 energy and carbon cycling, are predicted to experience both longer and more intensive dry
4 seasons by the end of the 21st century. However, the climate sensitivity of this region remains
5 uncertain: several studies have predicted large-scale die-back of the Amazon, while several more
6 recent studies predict that the biome will remain largely intact. Combining remote-sensing and
7 ground-based observations with a size-and-age structured terrestrial ecosystem model, we
8 explore the sensitivity and ecological resilience of these forests to changes in climate. We
9 demonstrate that water stress operating at the scale of individual plants, combined with spatial
10 variation in soil texture, explains observed patterns of variation in ecosystem biomass,
11 composition, and dynamics across the region, and strongly influences the ecosystem's resilience
12 to changes in dry season length. Specifically, our analysis suggests that, in contrast to existing
13 predictions of either stability or catastrophic biomass loss, the Amazon forest's response to a
14 drying regional climate is likely to be an immediate, graded, heterogeneous transition from high
15 biomass moist forests to transitional dry forests and woody savannah-like states. Fire, logging
16 and other anthropogenic disturbances may, however, accelerate this transition.

17

18 **Significance Statement**

19 **Understanding how changes in climate will impact terrestrial ecosystems is particularly**
20 **important in tropical forest regions, which store large amounts of carbon and exert**
21 **important feedbacks onto regional and global climate. By combining multiple types of**
22 **observations with a state-of-the-art terrestrial ecosystem model, we demonstrate that the**
23 **sensitivity of tropical forests to changes in climate is dependent on the length of the dry**

24 **season, soil type, but also importantly, the dynamics of individual-level competition within**
25 **plant canopies. These interactions result in ecosystems that are more sensitive to changes**
26 **in climate than has been predicted by traditional models, but that transition from one**
27 **ecosystem type to another in a continuous, non-tipping point, manner.**

28 \body

29 Amazonia consists of 815 million hectares of rainforest, transitional forest and savannah,
30 stores approximately half of tropical forest carbon (1), and plays a vital role in global water,
31 energy, and carbon cycling (2). While uncertainties in climate predictions for the region remain
32 large (3), recent analyses imply that significant portions of the basin will experience both longer
33 and more intense dry seasons by the end of the 21st century (3-6). There is particular concern
34 about southern Amazonian forests that already experience longer dry seasons than forests in
35 central and western Amazonia (3), and where a trend of increasing dry season length and
36 intensity has already been observed (7). Despite the importance of this region for the global
37 climate, the climate sensitivity of the Amazon forests remains uncertain: model predictions range
38 from a large-scale die-back of the Amazon (8, 9) to predictions that the biome will remain
39 largely intact, and may even increase in biomass (10-12). While some of these differences can
40 be attributed to differences in the predicted future climate forcing of the region (13, 14), accurate
41 predictions of how changes in climate will impact Amazonian forests also rely on an accurate
42 characterization of how the ecosystem is affected by a given change in climate forcing. In this
43 study, we examine the climate sensitivity of the Amazon ecosystem focusing on the mechanisms
44 underpinning changes in forest dynamics and their implications for the timing and nature of
45 basin-wide shifts in biomass in response to a drying climate.

46 Variation in forest biomass across the Amazon basin (15-17) has been shown to correlate
47 with dry season length (16-18) (Fig. 1), soil texture (16), shifts in stem turnover rate (19) and
48 forest composition (20). In general, high-biomass moist tropical forests occur where dry season
49 length (DSL, defined here as the number of months where precipitation is < 100 mm (6, 9)), is
50 short, and low-biomass savannah-like ecosystems are primarily found when DSLs are long (Fig.
51 1a). In addition, a significant relationship is observed between regional-scale spatial
52 heterogeneity in above-ground biomass ($AGB > 2 \text{ kg C m}^{-2}$) and DSL, with drier places having
53 greater spatial heterogeneity: this pattern is seen at both the scale of 1° (Fig. 1c, $r^2=0.88$, $p<0.01$
54 for remote-sensing based AGB estimates), but also at smaller spatial scales (see *SI Appendix S1*).
55 In other words, in moist areas, where DSL is short, forests have relatively homogenous levels of
56 AGB while, in drier areas, forests are increasingly heterogeneous. As we show below, this
57 observed heterogeneity in response to increasing DSL has important implications for how the
58 structure, composition, and dynamics of Amazon forests will be impacted by changes in climate.

59 The Ecosystem Demography model (ED2), a process-based terrestrial biosphere model
60 that represents individual plant-level dynamics including competition for light and water (21, 22),
61 was used to investigate the impact of ecosystem heterogeneity on the Amazon forest's ecological
62 resilience to climate perturbations (*SI Appendix S3*). Here the term ecological resilience is used
63 to describe the ability of a forest to maintain fundamental characteristics, such as carbon pools,
64 composition, and structure, despite changes in climate (23). ED2 simulations for the Amazon
65 region, forced with a regional climate dataset derived from *in situ* measurements and remote-
66 sensing observations, correctly reproduce the observed pattern of AGB variability as a function
67 of DSL and soil texture (*Fig. 1, SI Appendix S4*). This is also true for ED2 model simulations for

68 sites where ground-based soil texture, and forest structure, turnover, and composition
69 measurements are available (*Fig. S1, SI Appendix S4*).

70 An ensemble of model simulations with varying soil texture was used to investigate the
71 mechanisms that underpin the observed variable response to increasing DSL (*SI Appendix S3*). In
72 the model, individual plant productivity is modified by a measure of plant water stress (γ_{WS}) that
73 integrates soil texture, precipitation, and plant transpiration demand such that, as γ_{WS} increases,
74 the plants close their stomata to reduce water loss. In the ED2 ensemble simulations, plot
75 biomass is highly correlated with the average γ_{WS} for the forested sites (defined here as $AGB > 3$
76 $kg\ C\ m^{-2}$) (*Fig. 2c, $r^2=0.96-0.99, p<0.01, SI Appendix S5$*). Associated with changes in AGB that
77 occur as water stress increases are correlated changes in the productivity and composition of the
78 plant canopy (*SI Appendix S6*).

79 The important role that water stress operating at the scale of individual plants plays in
80 generating these responses is illustrated by comparing the native ED2 model predictions to
81 output from a horizontally and vertically averaged version of the model (ED2-BL), analogous to
82 a conventional ‘big leaf’ terrestrial biosphere model that represents the canopy in an aggregated
83 manner (*SI Appendix S3*). In the ED2-BL simulations, there is no significant relationship
84 between the spatial heterogeneity of forested sites and DSL ($r^2=0.24\ p=0.32, Fig. 1a,c$). The
85 absence of individual-level plant dynamics in ED2-BL results in a markedly different response to
86 variations in soil texture and dry season length than the native model formulation: biomass
87 initially declines as a function of increasing water stress, but then a tipping point is reached,
88 beyond which the high biomass forest is no longer stable and is replaced by a low biomass
89 savannah (*Fig. 2*). The result is a bimodal distribution of AGB across the basin in the ED2-BL
90 simulations, in contrast to the continuous distribution seen in the native model formulation and

91 the observations (Fig. 1b). This response mirrors the response seen in other ‘big-leaf’ type
92 ecosystem models (9). In native ED2 simulations, when water stress is prevented from
93 influencing plant productivity, DSL and soil texture no longer impact AGB (*SI Appendix S5* and
94 *Fig. S5*). Taken together, these simulations indicate that the driving mechanism behind the
95 observed heterogeneous response to changes in DSL is the differential performance of
96 individuals within the canopy to declining water availability, and how this response is modulated
97 by soils with different hydrologic properties. Specifically, the size-and-age structure of the ED2
98 plant canopy results in individuals’ differential access to both light and soil water, influencing
99 the dynamics of individual plant growth and mortality (*SI Appendix S6*). Due to the non-linear
100 nature of functions governing plant growth, mortality, and recruitment, this heterogeneity results
101 in a more continuous, graded response to changes in water stress than the big leaf (ED2-BL)
102 formulation (Fig. 2). The consequence of this heterogeneity in plant-level responses to changes
103 in soil moisture is that soil texture is likely to become increasingly important for controlling
104 AGB as dry season length increases. Soil fertility gradients also influence Amazonian AGB (16-
105 18); however, as we show in *SI Appendix S2*, it does not account for the observed regional-scale
106 pattern of increasing biomass heterogeneity with increasing DSL.

107 The ED2 biosphere model was used to investigate the expected patterns and time-scales
108 of Amazonian ecosystem response to a one-to-four month change in DSL over the 21st century
109 (6). Earlier analyses have suggested that, by accurately representing the dynamics of individual
110 trees, models such as ED2 that incorporate plant-level dynamics are likely to provide more
111 realistic estimates of forest successional change (21). Forests with a 4-month dry season (24% of
112 the Amazon basin) are projected to lose ~20% of their biomass with a 2-month increase in DSL
113 (range of 11 - 58% loss of AGB dependent on clay content), while drier forests (6-month DSL)

114 respond more rapidly to changes in climate, losing ~29% (20 – 37% loss dependent on clay
115 content) of their biomass with a one month increase in DSL (Fig. 3a, *SI Appendix S7*). As the
116 forests adjust to the new climate regime, the spatial heterogeneity of the forest structure,
117 composition and biomass across the range of soil textures gradually increases. As seen in Fig. 3b,
118 the model predicts that forests in soils with low clay content will be relatively unaffected;
119 however, in high clay content soils, the increase in levels of water stress caused by the onset of a
120 longer dry season will result in marked changes in forest AGB and composition, beginning
121 approximately three years after the perturbation (Fig. 3c). The timescale of the predicted initial
122 ecosystem response is consistent with the results from two field-based through-fall exclusion
123 experiments, which showed declining biomass 3-4 years after a drought was introduced (24, 25).
124 Underlying these predicted changes in AGB and canopy composition are reductions in plant
125 growth and increases in mortality rates (Figs. S14 and S15). While the majority of the change in
126 AGB occurs in the first hundred years, the composition and structure of the forest continues to
127 reorganize for more than two-hundred years after the perturbation (Fig. 3c). Specifically, the
128 simulations predict a substantial decline in the abundance of late-successional trees in high clay
129 content soils. This prediction arises as a simple consequence of the slower rate of growth of late-
130 successional trees that makes them more vulnerable to water stress-induced increases in
131 mortality rates, and less competitive against mid-successional species that are favored by
132 drought-induced increases in understory light levels. This prediction of increased vulnerability of
133 late-successional trees to increases in water stress is, as-yet, untested; however, more generally,
134 our analysis highlights how shifts in climate forcing are likely to drive significant shifts in
135 tropical forest composition and structure over decadal and centennial timescales.

136 Recent work has hypothesized that two stable ecosystem states may exist along the
137 boundaries of tropical forests and that a tipping point may occur once a climatological moisture
138 threshold is passed (26, 27). Instead, by combining field observations, remote-sensing estimates,
139 and a terrestrial biosphere model, we find no evidence either that an irreversible rapid transition
140 or dieback of Amazon forests will occur in response to a drying climate (8, 9), or that forests will
141 be unresponsive (11, 12). Rather, our results suggest that, at least in the case of Amazonian
142 forests, the ecosystem will exhibit an immediate but heterogeneous response to changes in its
143 climate forcing and that a continuum of transitional forest ecosystem states exists. These
144 conclusions are consistent with experimental observations across Amazonia of short-term
145 drought impacts (28). Furthermore, we find that future climate-induced shifts between a moist
146 tropical forest and dry forest will be a more graded transition accompanied by increasing spatial
147 heterogeneity in forest above-ground biomass, composition, and dynamics across gradients in
148 soil texture. The ability of Amazonian forests to undergo reorganization of their structure and
149 composition in response to climate-induced changes in levels of plant stress acts as an important
150 buffer against more drastic threshold changes in vegetation state that would otherwise occur;
151 however, it also means that the forests are more sensitive to smaller magnitude changes in their
152 climate forcing than previous studies have suggested.

153 The analysis conducted here intentionally focused on the direct impacts of changes in
154 climate forcing on vegetation, and did not incorporate the effects of soil nutrients, climate driven
155 changes in fire frequency, the effects of increasing atmospheric carbon-dioxide concentrations,
156 the impacts of land transformation, and biosphere-atmosphere feedbacks. With regard to soil
157 nutrients, at the basin-scale, analyses indicate that forest composition, structure, biomass, and
158 dynamics also vary across a gradient in soil fertility (16, 17) with the younger, more fertile soils

159 of western Amazonian supporting forests with lower above-ground biomass and higher rates of
160 biomass productivity and stem turnover relative to the forests of the central Amazon and
161 Guianan Shield, which are located on older, more nutrient-poor soils. Meanwhile landscape-scale
162 studies in the central (29) and northwestern (30) Amazonia have found that more fertile clay
163 soils have higher above-ground biomass than nutrient-poor sandy soils. Further discussion of the
164 impact of soil nutrients can be found in *SI Appendix S2*).

165 Plant water availability is affected by both the hydraulic properties of soils and plant
166 hydraulic architecture. Our findings of the importance of individual plant water stress on forest
167 response to changes in climate highlight the need for additional studies into both of these
168 important, but relatively under-studied, properties of tropical forests. With regard to soil
169 hydraulic properties, recent studies suggest that the relationship between a soil's texture and its
170 hydraulic properties may differ significantly between tropical and temperate soils (31, 32).
171 However, the impact of these differences on plant water availability remains uncertain. With
172 regard to plant hydraulic architecture, while some measurements exist on rooting properties and
173 vascular architecture of tropical trees (33-36), the above and below-ground hydraulic attributes
174 of tropical trees remain poorly characterized, especially compared to that of temperate trees.

175 In some areas, particularly those with long dry seasons, increasing water stress is likely to
176 be accompanied by increases in fire frequency, which may act to generate more rapid transitions
177 from a higher biomass forested state to a more savannah-like biome (26, 27). Since these two
178 mechanisms have distinct impacts on forest composition, structure, and function, both must be
179 considered when predicting future responses to changes in climate. The potential impacts of fire
180 on patterns of ecosystem change are discussed in *SI Appendix S1*. Recent modeling studies
181 indicate that CO₂ fertilization may mitigate the impact of increasing water stress (37); however,

182 experimental studies are needed to better quantify the impact of elevated CO₂ concentrations on
183 the physiological functioning of Amazon trees.

184 While regional patterns of Amazonian above-ground biomass (AGB) are complex,
185 reflecting the impact of multiple factors, our results suggest that plant-level responses to soil
186 texture heterogeneity and changes in DSL are important in explaining the observed basin-wide
187 pattern of variation in Amazonian AGB, providing a mechanistic explanation for the observed
188 correlations between DSL, above-ground biomass, and changes in stand structure and
189 composition (16, 17) . These conclusions may also apply to African and Asian tropical forests;
190 however, important differences exist in the future climate predictions for these regions (38) and
191 their soil edaphic and nutrient characteristics, and historical fire regimes (39-41).

192 The response of forests to changes in their climate forcing is an emergent ecosystem-
193 level response that is ultimately driven by individual trees responding to changes in their local
194 environments. Non-linearities in the performance of individual plants, such as their rates of
195 photosynthetic assimilation and mortality, as environmental conditions change imply that
196 terrestrial biosphere models need to represent these differential responses of individuals in order
197 to accurately capture emergent ecosystem properties (42). This analysis demonstrates that the
198 conventional approach of modeling average plants in average environments within
199 climatological grid cells underestimates the direct, near-term response of tropical forests to
200 climatological change, but overestimates the direct impacts of larger-scale changes in forcing.
201 Consequently, accurate predictions for the timing and nature of forest responses to changes in
202 climate require consideration of how climate and soils affect the performance of individuals
203 within plant canopies. As we have shown here, models that incorporate plant-level dynamics are
204 able to more accurately characterize observed extant patterns of variation in the structure,

205 composition and dynamics of Amazonian ecosystems, and that accounting for these patterns has
206 important implications for the sensitivity and ecological resilience of Amazon forests to different
207 levels of climatological perturbation.

208

209 **METHODS**

210 The Ecosystem Demography Biosphere Model (ED2) is an integrated terrestrial
211 biosphere model that incorporates hydrology, land-surface biophysics, vegetation dynamics and
212 soil carbon biogeochemistry and uses a size- and age-structured system of partial differential
213 equations (PDEs) to approximate the individual level dynamics of plant canopies (21, 22, 43).
214 The horizontally and vertically averaged Ecosystem Demography Model (ED2-BL) represents
215 exactly the same biophysical and biogeochemical processes as ED2, but the size- and age-
216 structured canopy is replaced with a horizontally- and vertically- averaged canopy akin to those
217 used by conventional terrestrial biosphere models (*see SI Appendix S3 for further information on*
218 *the model formulation*).

219 ED2 and ED2-BL models were run for the entire Amazon basin forced with a rescaled
220 NCEP reanalysis product (44) and observation-based soil maps (45, 46) at 1° resolution, and
221 increasing atmospheric CO₂ (47) (*SI Appendix S3*). The model results were compared against
222 remote-sensing estimates (1, 48) aggregated to the same resolution as the model simulations.
223 Plot-based observations were made on the scale of 0.4-10 ha (20), and were compared against
224 model simulations forced with site-specific inputs (Table S4).

225 The water stress factor (γ_{WS}) was used in both the ED2 and ED2-BL models to scale
226 photosynthesis in response to water stress. γ_{WS} was calculated for each individual (i) as:

$$227 \quad \gamma_{WS}^i = 1 - \frac{K \int_0^{RD} (\theta(z) - \theta_{WP}) dz C_{root}^i}{K \int_0^{RD} (\theta(z) - \theta_{WP}) dz C_{root}^i + T_{max}^i} \quad (Eq. 1)$$

228 where γ_{WS} ranges from 0 (unstressed) to 1 (stressed). T_{\max}^i is the maximum transpiration (kg H₂O
229 yr⁻¹) for individual i , C_{root} is the root biomass (kgC) for individual i , $\theta(z)$ is the soil moisture (kg
230 H₂O m⁻³) at soil depth z , K is the root conductance (m² kgC⁻¹ yr⁻¹), and θ_{WP} is the soil wilting
231 point (kg H₂O m⁻³). The available soil water (kgH₂O m⁻²), $\theta(z) - \theta_{\text{WP}}$, is integrated over the rooting
232 depth (RD) of the individual.

233 Spatial heterogeneity ($\frac{\sigma}{\mu}$) was calculated over 1-month DSL intervals for model
234 simulations and remote-sensing based estimates. These calculations were done at 1-ha resolution
235 for plot-based observations and model simulations for these locations, 500 m and 1-km
236 resolution for remote-sensing based estimates, and 1° resolution for the regional model
237 simulations and remote-sensing based estimates. Due to the relatively low number of plots, the
238 spatial heterogeneity of the plot-based observations was calculated for three DSL categories: 0-2
239 months, 2-5 months, and 5-8 months.

240

241 **Acknowledgements:**

242 Funding was provided by the Gordon and Betty Moore Foundation and a NOAA Climate and
243 Global Change fellowship. We acknowledge RAINFOR partners for generously providing plot
244 data, in particular T. Baker, G. López-Gonzalez, and the late S. Almeida, A. Gentry, and S.
245 Patiño. RAINFOR inventories have been funded by NERC and the Gordon and Betty Moore
246 Foundation. OP is supported by an ERC Advanced Grant and is a Royal Society-Wolfson
247 Research Merit Award holder.

248

249

250 **Author Contributions:**

251 N.M.L. was involved in study design, model development, ran model simulations, performed
252 data analysis, and wrote the manuscript; K.Z. was involved with model development, ran model
253 simulations and commented on the manuscript; M.L. was involved with model development and
254 provided comments on the manuscript; A.B. provided remote-sensing data and comments on the
255 manuscript; O.P., S.L.L., T.E., T.R.F., and Y.M. provided unpublished field-based observations
256 and comments on the manuscript; E.A., A.A., R.B., A.L.M.M., P.N.V., A.P., and J.E.S.E.
257 provided unpublished field-based observations; P.R.M. was involved in study design, model
258 development, and data analysis, and provided comments on the manuscript.

REFERENCES:

1. Baccini A, *et al.* (2012) Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps. *Nature Climate Change* 2(3):182-185.
2. Werth D & Avissar R (2002) The local and global effects of Amazon deforestation. *Journal of Geophysical Research: Atmospheres* 107(D20):LBA 55-51-LBA 55-58.
3. Boisier JP, Ciais P, Ducharne A, & Guimberteau M (2015) Projected strengthening of Amazonian dry season by constrained climate model simulations. *Nature Clim. Change* 5(7):656-660.
4. Malhi Y, *et al.* (2008) Climate change, deforestation, and the fate of the Amazon. *Science* 319(5860):169-172.
5. Chou C & Neelin JD (2004) Mechanisms of global warming impacts on regional tropical precipitation. *Journal of Climate* 17(13):2688-2701.
6. Malhi Y, *et al.* (2009) Exploring the likelihood and mechanism of a climate-change-induced dieback of the Amazon rainforest. *Proceedings of the National Academy of Sciences* 106(49):20610-20615.
7. Fu R, *et al.* (2013) Increased dry-season length over southern Amazonia in recent decades and its implication for future climate projection. *Proceedings of the National Academy of Sciences*.
8. Cox PM, *et al.* (2004) Amazonian forest dieback under climate-carbon cycle projections for the 21st century. *Theoretical and Applied Climatology* 78(1-3):137-156.
9. Good P, *et al.* (2011) Quantifying Environmental Drivers of Future Tropical Forest Extent. *Journal of Climate* 24(5):1337-1349.
10. Thompson SL, *et al.* (2004) Quantifying the effects of CO₂-fertilized vegetation on future global climate and carbon dynamics. *Geophys. Res. Lett.* 31(23):L23211.
11. Cox PM, *et al.* (2013) Sensitivity of tropical carbon to climate change constrained by carbon dioxide variability. *Nature* 494(7437):341-344.
12. Huntingford C, *et al.* (2013) Simulated resilience of tropical rainforests to CO₂-induced climate change. *Nature Geosci* 6(4):268-273.
13. Zhang K, *et al.* (2015) The Fate of Amazonian Ecosystems over the Coming Century Arising from Changes in Climate, Land-use and CO₂. *Global Change Biology*.

14. Galbraith D, *et al.* (2010) Multiple mechanisms of Amazonian forest biomass losses in three dynamic global vegetation models under climate change. *New Phytologist* 187(3):647-665.
15. Malhi Y, *et al.* (2004) The above-ground coarse wood productivity of 104 Neotropical forest plots. *Global Change Biology* 10(5):563-591.
16. Quesada CA, *et al.* (2012) Basin-wide variations in Amazon forest structure and function are mediated by both soils and climate. *Biogeosciences* 9(6):2203-2246.
17. ter Steege H, *et al.* (2006) Continental-scale patterns of canopy tree composition and function across Amazonia. *Nature* 443(7110):444-447.
18. Baraloto C, *et al.* (2011) Disentangling stand and environmental correlates of aboveground biomass in Amazonian forests. *Global Change Biology* 17(8):2677-2688.
19. Phillips OL, *et al.* (2004) Pattern and process in Amazon tree turnover, 1976-2001. *Philosophical Transactions of the Royal Society of London Series B-Biological Sciences* 359(1443):381-407.
20. Baker TR, *et al.* (2004) Variation in wood density determines spatial patterns in Amazonian forest biomass. *Global Change Biology* 10(5):545-562.
21. Moorcroft PR, Hurtt GC, & Pacala SW (2001) A method for scaling vegetation dynamics: The ecosystem demography model (ED). *Ecological Monographs* 71(4):557-585.
22. Medvigy D, Wofsy SC, Munger JW, Hollinger DY, & Moorcroft PR (2009) Mechanistic scaling of ecosystem function and dynamics in space and time: Ecosystem Demography model version 2. *Journal of Geophysical Research-Biogeosciences* 114.
23. Walker B, Holling CS, Carpenter SR, & Kinzig A (2004) Resilience, adaptability and transformability in social--ecological systems. *Ecology and society* 9(2):5.
24. Nepstad DC, Tohver IM, Ray D, Moutinho P, & Cardinot G (2007) Mortality of large trees and lianas following experimental drought in an amazon forest. *Ecology* 88(9):2259-2269.
25. da Costa ACL, *et al.* (2010) Effect of 7 yr of experimental drought on vegetation dynamics and biomass storage of an eastern Amazonian rainforest. *New Phytologist* 187(3):579-591.

26. Staver AC, Archibald S, & Levin SA (2011) The Global Extent and Determinants of Savanna and Forest as Alternative Biome States. *Science* 334(6053):230-232.
27. Hirota M, Holmgren M, Van Nes EH, & Scheffer M (2011) Global Resilience of Tropical Forest and Savanna to Critical Transitions. *Science* 334(6053):232-235.
28. Phillips OL, *et al.* (2009) Drought Sensitivity of the Amazon Rainforest. *Science* 323(5919):1344-1347.
29. Laurance WF, *et al.* (1999) Relationship between soils and Amazon forest biomass: a landscape-scale study. *Forest Ecology and Management* 118(1–3):127-138.
30. Jimenez EM, *et al.* (2014) Edaphic controls on ecosystem-level carbon allocation in two contrasting Amazon forests. *Journal of Geophysical Research-Biogeosciences* 119(9):1820-1830.
31. Hodnett MG & Tomasella J (2002) Marked differences between van Genuchten soil water-retention parameters for temperate and tropical soils: a new water-retention pedo-transfer functions developed for tropical soils. *Geoderma* 108(3–4):155-180.
32. Marthews TR, *et al.* (2014) High-resolution hydraulic parameter maps for surface soils in tropical South America. *Geosci. Model Dev.* 7(3):711-723.
33. Nepstad DC, *et al.* (1994) The role of deep roots in the hydrological and carbon cycles of Amazonian forests and pastures.
34. Nepstad DC, *et al.* (2002) The effects of partial throughfall exclusion on canopy processes, aboveground production, and biogeochemistry of an Amazon forest. *Journal of Geophysical Research-Atmospheres* 107(D20).
35. Schuldt B, Leuschner C, Brock N, & Horna V (2013) Changes in wood density, wood anatomy and hydraulic properties of the xylem along the root-to-shoot flow path in tropical rainforest trees. *Tree physiology* 33(2):161-174.
36. Oliveira RS, Dawson TE, Burgess SS, & Nepstad DC (2005) Hydraulic redistribution in three Amazonian trees. *Oecologia* 145(3):354-363.
37. Rammig A, *et al.* (2010) Estimating the risk of Amazonian forest dieback. *New Phytologist* 187(3):694-706.
38. Chadwick R, Boutle I, & Martin G (2012) Spatial Patterns of Precipitation Change in CMIP5: Why the Rich Do Not Get Richer in the Tropics. *Journal of Climate* 26(11):3803-3822.

39. Dwyer E, Pereira JMC, Gregoire JM, & DaCamara CC (2000) Characterization of the spatio-temporal patterns of global fire activity using satellite imagery for the period April 1992 to March 1993. *Journal of Biogeography* 27(1):57-69.
40. Baillie IC (1996) Soils of the humid tropics. *The tropical rainforest (Second edition)* ed Richards PW (Cambridge University Press, Cambridge, UK), pp 256-286.
41. Vitousek PM & Sanford RL (1986) Nutrient cycling in moist tropical forest. *Annual Review of Ecology and Systematics* 17:137-167.
42. Levin SA, Grenfell B, Hastings A, & Perelson AS (1997) Mathematical and computational challenges in population biology and ecosystems science. *Science* 275(5298):334-343.
43. Hurtt GC, Moorcroft PR, Pacala SW, & Levin SA (1998) Terrestrial models and global change: challenges for the future. *Global Change Biology* 4(5):581-590.
44. Sheffield J, Goteti G, & Wood EF (2006) Development of a 50-year high-resolution global dataset of meteorological forcings for land surface modeling. *Journal of Climate* 19(13):3088-3111.
45. Quesada CA, *et al.* (2010) Variations in chemical and physical properties of Amazon forest soils in relation to their genesis. *Biogeosciences* 7(5):1515-1541.
46. Anonymous (Global Soil Task 2000, Global Soil Data Products CD_ROM. ed IGBP-DIS (<http://www.daac.ornl.gov/SOILS/igbp.html>).
47. Nakicenovic N, *et al.* (2000) IPCC Special report on emissions scenarios. (Cambridge, UK & New York, NY, USA), p 599.
48. Saatchi SS, *et al.* (2011) Benchmark map of forest carbon stocks in tropical regions across three continents. *Proceedings of the National Academy of Sciences of the United States of America* 108(24):9899-9904.

Figure captions:

Fig. 1: Change in above-ground biomass with dry season length for remote-sensing based estimates (black and gray circles), ground-based plot measurements (blue triangles), ED2 model output (green circles), and ED2-BL model output (purple circles). Panel b displays the distribution of AGB in the observations and the two models and panel c displays the change in percent biomass variability, coefficient of variation (CV) defined as $1 \sigma / \text{mean}$. Results are for undisturbed primary vegetation forests.

Fig. 2: Impact of changes in soil clay fraction (panels a and b) and plant water stress (panels c and d) on above-ground biomass (AGB) in the ED2 (panels a and c) and ED2-BL (panels b and d) model simulations. Four climatological conditions are shown, a 2-month dry season, a 4-month dry season, a 6-month dry season and an 8-month dry season.

Fig. 3: Predicted response of forest above-ground biomass and composition to an increase in dry season length (DSL). Panel a shows the change in above-ground biomass after 100 years as a result of increasing dry season length for forests with historic dry season lengths of 2, 4, and 6 months for the range of soil-textures simulated in the ensemble model simulations ($N=30$). The magnitude of the change in AGB is influenced by soil clay fraction: the mean (solid line), 1σ deviation (shaded region), and minimum and maximum values (dashed lines) are shown. Panels b and c: barplots illustrating the impact of a 2 month increase in dry season (from 4 to 6 months) on a forest situated on a low clay content soil and a forest situated on a high clay content soil. The color of the bars indicates the contribution of mid- and late-successional trees, illustrating the shift in composition caused by the increase in DSL.