

IMAGES IN NEONATAL MEDICINE

Aplasia cutis congenita of the scalp with sagittal venous sinus exposure

Aplasia cutis congenita is an uncommon condition (estimated global incidence 0.5–1 in 10 000 neonates)¹ characterised by a localised defect of all skin layers and subcutaneous tissue, sometimes extending to the underlying bone and meningeal layers.² Some of the possible complications include haemorrhage, trauma, venous sinus thrombosis and infection (including meningitis).³ The management is controversial, and includes surgical and conservative treatment options.⁴

We present the case of a term male newborn, weighing 2615 g with normal delivery by a 24-year-old primipara. Prenatal ultrasounds were unremarkable as was prenatal history regarding infections and drug or medication use. Family history



Figure 1 Aplasia cutis congenita of the scalp measuring 5×5 cm with sagittal venous sinus exposure. Photo taken in the first hours of life (Israel Macedo, Maternidade Alfredo da Costa).

was also unremarkable. At the time of the delivery it was noted a large lesion of the scalp measuring 5×5 cm, involving all the layers of the skin, underlying subcutaneous tissue, bone and dura mater through which the sagittal venous sinus was visible (figure 1). There was no visible leak of cerebrospinal fluid. No other associated malformations were noted. The lesion was dressed initially twice a day with sterile fatty gauze bandage and after 2 weeks with soft silicone dressings (Mepitel) to minimise trauma and drying and to promote a re-epithelialisation of the lesion. Prophylactic antibiotics (flucloxacillin and gentamicin) were initiated. Progressive closure of the lesion issued without complications over the course of the next weeks.

Rapid identification, clinical assessment and institution of appropriate protection of the lesion are necessary to achieve the best outcome. A multidisciplinary approach is recommended.

Jenny F Goncalves,¹ Tiago Milheiro Silva,¹ Israel Macedo²

¹Departamento de Pediatria Médica, Hospital Dona Estefania, Lisboa, Portugal

²Maternidade Dr Alfredo da Costa, Lisboa, Lisboa, Portugal

Correspondence to Dr Jenny F Goncalves, Hospital Dona Estefania, Rua Jacinta Marto, Lisboa 1169-045, Portugal, jennycfg@gmail.com

Competing interests None declared.

Provenance and peer review Not commissioned; internally peer reviewed.



CrossMark

To cite Goncalves JF, Silva TM, Macedo I. *Arch Dis Child Fetal Neonatal Ed* 2016;**101**:F283.

Received 17 January 2016

Accepted 10 February 2016

Published Online First 1 March 2016

Arch Dis Child Fetal Neonatal Ed 2016;**101**:F283.

doi:10.1136/archdischild-2016-310480

REFERENCES

- 1 Browning JA. Aplasia cutis congenital: approach to evaluation and management. *Dermatol Ther* 2013;**26**:439–44.
- 2 Puvabanditsin S, February M, Garrow E, *et al*. Our experience with a severe case of aplasia cutis congenita with a large skull defect. *Int J Dermatol* 2015.
- 3 Duan X, Yang GE, Yu D, *et al*. Aplasia cutis congenital: a case report and literature review. *Exp Ther Med* 2015;**10**:1893–5.
- 4 Brzezinski P, Pinteala T, Chiriac AE, *et al*. Aplasia cutis congenital of the scalp—what are the steps to be followed? Case report and review of the literature. *An Bras Dermatol* 2015;**90**:100–3.