



University of Pennsylvania  
**ScholarlyCommons**

---

Statistics Papers

Wharton Faculty Research

---

1977

# Families of Sample Means Converge Slowly

J Michael Steele  
*University of Pennsylvania*

Follow this and additional works at: [http://repository.upenn.edu/statistics\\_papers](http://repository.upenn.edu/statistics_papers)

 Part of the [Physical Sciences and Mathematics Commons](#)

---

## Recommended Citation

Steele, J. M. (1977). Families of Sample Means Converge Slowly. *The Annals of Probability*, 5 (6), 1036-1038. <http://dx.doi.org/10.1214/aop/1176995672>

At the time of publication, author J. Michael Steele was affiliated with University of British Columbia. Currently, he is a faculty member at the Statistics Department at the University of Pennsylvania.

This paper is posted at ScholarlyCommons. [http://repository.upenn.edu/statistics\\_papers/21](http://repository.upenn.edu/statistics_papers/21)  
For more information, please contact [repository@pobox.upenn.edu](mailto:repository@pobox.upenn.edu).

---

# Families of Sample Means Converge Slowly

**Abstract**

The uniform empirical integral differences of Sethuraman's large deviation theorem are proved to converge arbitrarily slowly.

**Keywords**

empirical measures, slow convergence, uniform convergence

**Disciplines**

Physical Sciences and Mathematics

**Comments**

At the time of publication, author J. Michael Steele was affiliated with University of British Columbia. Currently, he is a faculty member at the Statistics Department at the University of Pennsylvania.

## FAMILIES OF SAMPLE MEANS CONVERGE SLOWLY

BY J. MICHAEL STEELE

University of British Columbia

The uniform empirical integral differences of Sethuraman's large deviation theorem are proved to converge arbitrarily slowly.

**1. Introduction.** The purpose of this note is to answer a question of N. Glick [2, page 1380] concerning the rate of convergence to zero of a family of sample means as introduced by Sethuraman [1].

To introduce Glick's question we let  $(\Omega, X, P)$  be a probability space, and let  $X_i, i = 1, 2, \dots$  be a sequence of  $M$ -valued Borel measurable random variables which are independent and identically distributed. We will suppose also that  $M$  is a complete separable metric space. The distribution of the  $X_i(\omega)$  will be denoted by  $\mu(\cdot)$  and the sample probability measure of  $X_1(\omega), X_2(\omega), \dots, X_n(\omega)$  will be denoted by  $\mu(n, \omega, \cdot)$ . (That is,  $\mu(n, \omega, \cdot)$  is the measure which places mass  $1/n$  at each of the points  $X_1(\omega), X_2(\omega), \dots, X_n(\omega)$ .)

Numerous authors have considered properties of the random variable defined by

$$(1.1) \quad \sup_{f \in \mathcal{F}} |\int f(x) \mu(n, \omega, dx) - \int f(x) \mu(dx)|$$

where  $\mathcal{F}$  is a suitable class of functions on  $M$ . Noteworthy among such results both for its precision and generality is the large deviation theorem of Sethuraman [1] and [2]. This result says, under certain restrictions on  $\mathcal{F}$ , one has for  $\varepsilon > 0$  that

$$(1.2) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \log P(\sup_{f \in \mathcal{F}} |\int f(x) \mu(n, \omega, dx) - \int f(x) \mu(dx)| \geq \varepsilon) \\ = \log \rho(\mathcal{F}, \varepsilon)$$

where  $0 < \rho(\mathcal{F}, \varepsilon) < 1$ . The conditions which Sethuraman requires of  $\mathcal{F}$  in (1.2) are only two:

(S<sub>1</sub>)  $\mathcal{F}$  is a class of functions from  $M$  to the real line which is pre-compact in the topology of uniform convergence on compact sets;

(S<sub>2</sub>) There is a function  $g(x)$  such that  $|f(x)| \leq g(x)$  for all  $f \in \mathcal{F}$  and such that  $\int \exp(tg(x)) \mu(dx) < \infty$  for all  $t$ .

The question posed by N. Glick [2, page 1330] is to ascertain the truth or falsity of

$$(1.3) \quad n^{\frac{1}{2}} \sup_{f \in \mathcal{F}} |\int f(x) \mu(n, \omega, dx) - \int f(x) \mu(dx)| = O_p(1)$$

Received February 28, 1977; revised May 16, 1977.

AMS 1970 subject classifications. 60F15, 60F10, 62E20.

Key words and phrases. Empirical measures, slow convergence, uniform convergence.

where  $\mathcal{F}$  satisfies the Sethuraman conditions  $S_1$  and  $S_2$ . (Here one writes  $Z_n = O_p(1)$  to mean, as usual, that  $\max_n P(Z_n \geq \lambda) \rightarrow 0$  as  $\lambda \rightarrow \infty$ .)

The fact that (1.3) is false is one immediate consequence of the main theorem of this note. Much more pointedly, it is proved that the random variables (1.1) converge to zero arbitrarily slowly.

**2. Main result.**

**THEOREM 1.** *Let  $X_i, i = 1, 2, \dots$  be i.i.d. and uniformly distributed on  $[0, 1]$ . For any sequence  $c_n$  such that  $c_n \rightarrow \infty$  there is a class of functions  $\mathcal{F}$  which is (a) uniformly bounded by 1 and (b) compact in the topology of uniform convergence such that*

$$(1.4) \quad c_n \sup_{f \in \mathcal{F}} |\int f(x)\mu(dx) - \int f(x)\mu(n, \omega, dx)| \rightarrow \infty$$

with probability one.

**PROOF.** To construct the class  $\mathcal{F}$  suppose that  $0 < h_n \leq 1$  is a sequence of reals such that  $h_n \rightarrow 0$  and  $k_n$  is a sequence of integers such that  $k_n > n$ . For each  $n$  a finite set  $\mathcal{F}_n$  of functions is defined by the following procedure.

First  $[0, 1]$  is divided into  $k_n$  intervals of equal length. Next a function  $f$  on  $[0, 1]$  is defined by (1) choosing  $n$  of the  $k_n$  intervals and defining  $f$  to be zero on the chosen intervals; (2) defining  $f$  to be zero on the first and the last of the intervals; (3) defining  $f$  to be  $h_n$  on the intervals where  $f$  is not yet defined and which are not adjacent to intervals where  $f$  has been defined, and (4) defining  $f$  by linear interpolation on all remaining intervals.

Finally  $\mathcal{F}_n$  is taken to be the set of all  $f$  defined by the preceding process. We note the elements of  $\mathcal{F}_n$  are bounded by  $h_n$  and that  $\mathcal{F}_n$  contains at most  $\binom{k_n}{n}$  distinct elements.

The class  $\mathcal{F}$  is defined to be the union of all the  $\mathcal{F}_n$  and the function which is identically zero. Since  $h_n \rightarrow 0$  and since each  $\mathcal{F}_n$  is finite, any infinite sequence from  $\mathcal{F}$  is seen to contain a uniformly convergent subsequence. Since the uniform topology is metric, this shows  $\mathcal{F}$  is compact.

To provide estimates for (1.4) we suppose that  $k_n$  is chosen so large that for  $A_n = \{\omega : |X_i - X_j| \leq 1/k_n \text{ for some } i \neq j, i, j \leq n\}$  we have  $P(A_n) \leq 1/n^2$ . It will also be supposed that  $k_n$  is taken so large that  $2(n + 2)/k_n \leq h_n/2$  for our future convenience.

Now for each  $\omega \in A_n^c$  one can select an  $f^* \in \mathcal{F}_n$  such that  $f^*$  is zero at each of the points  $X_1(\omega), X_2(\omega), \dots, X_n(\omega)$ . Consequently one has  $\int f^*(x)\mu(n, \omega, dx) = 0$ . On the other hand, by elementary geometry one has

$$\int f\mu(dx) \geq h_n - 2(n + 2)/k_n$$

for any  $f \in \mathcal{F}_n$ . Hence by the choice which was made for  $k_n$ , one has for  $\omega \in A_n^c$ ,

$$(1.5) \quad \sup_{f \in \mathcal{F}} |\int f(x)\mu(dx) - \int f(x)\mu(n, \omega, dx)| \geq |\int f^*(x)\mu(dx) - \int f^*(x)\mu(n, \omega, dx)| \geq h_n/2.$$

Since  $P(A_n \text{ i.o.}) = 0$  by the Borel–Cantelli lemma we have for a.e.  $\omega$  an  $N(\omega)$  such that

$$(1.6) \quad c_n \sup_{f \in \mathcal{F}} |\int f(x)\mu(dx) - \int f(x)\mu(n, \omega, dx)| \geq c_n h_n/2$$

for all  $n \geq N(\omega)$ . Finally, since the only property of  $h_n$  required in the construction is that  $h_n \rightarrow 0$ , the  $h_n$  can be selected such that  $c_n h_n \rightarrow \infty$ . Such a choice completes the proof of the theorem.

**3. Applications and extentions.** One should note that the class constructed in Theorem 1 certainly satisfies the Sethuraman conditions  $S_1$  and  $S_2$ . Consequently, the question posed by N. Glick is completely answered. In the style of that question one now has that

$$\sup_{f \in \mathcal{F}} |\int f(x)\mu(dx) - \int f(x)\mu(n, \omega, dx)| = o_p(1)$$

is best possible under the conditions  $S_1$  and  $S_2$ .

The construction used in Theorem 1 obviously can be applied to yield more general results. In particular one can prove

**THEOREM 2.** *Let  $X_i, i = 1, 2, \dots$  be i.i.d. random variables which take values in  $R^d$  and which have distribution  $\mu$ . Suppose also that  $\mu$  is not concentrated on any countable set. Then for any  $c_n \rightarrow \infty$  there is a uniformly bounded class  $\mathcal{F}$  which is compact under uniform convergence such that (1.4) holds with probability one.*

The proof of Theorem 2 requires only mild modifications of the technique used previously so it will not be given.

**4. A conjecture.** The two previous results should not make one overly doubtful of relationships such as suggested by (1.3). In particular, if one considers a class  $G$  of functions on  $[0, 1]$  which are uniformly bounded and have uniformly bounded derivatives, the techniques of the present note do nothing to discredit the conjecture that

$$(4.1) \quad n^{\frac{1}{2}} \sup_{f \in G} |\int f(x)\mu(dx) - \int f(x)\mu(n, \omega, dx)| = O_p(1).$$

There are reasons to believe that (4.1) is in fact true. If (4.1) is true this fact would have several uses, and if (4.1) is false this too should be learned.

#### REFERENCES

- [1] SETHURAMAN, J. (1964). On the probability of large deviations of families of sample means. *Ann. Math. Statist.* **35** 1304–1316.
- [2] SETHURAMAN, J. (1970). Corrections to “On the probability of large deviations of families of sample means”. *Ann. Math. Statist.* **41** 1376–1380.

DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF BRITISH COLUMBIA  
2075 WEBBROOK PLACE  
VANCOUVER, B. C.  
CANADA V6T 1W5