

Tumor suppressive Ca²⁺ signaling is driven by IP₃ receptor fitness

Geert Bultynck^{1,*} and Michelangelo Campanella^{2,3,*}

¹ KU Leuven, Lab. Molecular and Cellular Signaling, Dep. Cellular and Molecular Medicine and Leuven Kanker Instituut (LKI), Campus Gasthuisberg O/N-I bus 802, Herestraat 49, BE-3000 Leuven, Belgium.

² Department of Comparative Biomedical Sciences, Royal Veterinary College, NW1 0TU, London, United Kingdom.

³ University College London Consortium for Mitochondrial Research, University College London, WC1 6BT, London, United Kingdom.

* Corresponding Authors:

Geert Bultynck, KU Leuven, Lab. Molecular and Cellular Signaling, Dep. Cellular and Molecular Medicine and Leuven Kanker Instituut (LKI), Campus Gasthuisberg O/N-I bus 802, Herestraat 49, BE-3000 Leuven, Belgium; E-mail: geert.bultynck@kuleuven.be;

Michelangelo Campanella, Hobday Building, RVC Camden Campus, Royal College Street, NW1 0TU, London, United Kingdom; E-mail: mcampanella@rvc.ac.uk

Intracellular Ca²⁺ signals critically control a plethora of cellular functions, of which many impact cellular death and/or survival, processes often dysregulated in cancer [1]. In healthy cells, Ca²⁺ signaling is employed for normal cell physiology and survival functions [2]. Yet, when a cell is exposed to toxic stimuli or suffering from enduring cell stress, like irreparable DNA damage, the Ca²⁺-signaling toolbox can be rapidly switched from a “pro-survival” modus into a “pro-death” modus, thereby initiating demise pathways [3]. The highly dynamic nature of Ca²⁺ signaling allows cells to swiftly respond to stress and damage, preventing the survival of damaged cells and malignant transformation that eventually results in tumor formation. Alterations in the expression, activity and regulation of Ca²⁺-transport systems both at the plasma membrane and at organelles like the endoplasmic reticulum (ER) and mitochondria have been implicated in oncogenesis and neoplasia [1, 4]. These changes result in aberrant Ca²⁺-signaling events that could favor resistance to cell death, migration or senescence escape [5].

Over the last decade, we learnt that tight contacts and functional connections involving Ca²⁺ exchanges between the ER, the main intracellular Ca²⁺-storage organelle, and

the mitochondria are pivotal for cell-fate decisions [3, 6-8]. These contact sites contain chaperone-coupled Ca²⁺-flux systems: the IP₃ receptors (IP₃Rs) at the ER side and the voltage-dependent anion channels (VDACs) at the mitochondrial outer membrane side [9]. These are controlled/exploited by several cellular factors and regulatory proteins, including oncogenes and tumor suppressors [10-12]. Basal Ca²⁺ fluxes between ER and mitochondria sustain anabolic pathways for mitochondrial metabolism, ensuring proper cell cycle progression [13]. Yet, continued elevated ER-mitochondrial Ca²⁺ transfers result in loss of mitochondrial membrane integrity and release of apoptogenic factors [14]. Tuned ER-mitochondrial Ca²⁺ transfer is therefore key to cells' response to pro-apoptotic stimuli: the failure of which results in cell death resistance, as often observed in cancer cells [15, 16]. In fact, the efficacy of chemotherapeutic agents and photodynamic therapy depends on the ability of these agents to elicit ER-mitochondrial Ca²⁺ exchanges [15]. In the context of apoptosis, previous work proposed unique roles for the type 3 IP₃R isoform (IP₃R3) [17] and type 1 VDAC isoform (VDAC1) [18] even though other IP₃R isoforms can contribute to the initiation of cell death programs [19, 20].

Comment on Kuchay S, Giorgi C, Simoneschi D, Pagan J, Missiroli S, Saraf A, Florens L, Washburn MP, Collazo-Lorduy A, Castillo-Martin M, Cordon-Cardo C, Sebti SM, Pinton P, Pagano M (2017). PTEN counteracts FBXL2 to promote IP3R3- and Ca2+-mediated apoptosis limiting tumour growth. *Nature* 546(7659): 554-558. doi: 10.1038/nature22965.

AND

Bononi A, Giorgi C, Patergnani S, Larson D, Verbruggen K, Tanji M, Pellegrini L, Signorato V, Olivetto F, Pastorino S, Nasu M, Napolitano A, Gaudino G, Morris P, Sakamoto G, Ferris LK, Danese A, Raimondi A, Tacchetti C, Kuchay S, Pass HI, Affar EB, Yang H, Pinton P, Carbone M (2017). BAP1 regulates IP3R3-mediated Ca2+ flux to mitochondria suppressing cell transformation. *Nature* 546(7659): 549-553. doi: 10.1038/nature22798

doi: 10.15698/cst2017.11.109

Received: 12.10.2017, Accepted 20.10.2017, Published 01.11.2017.

Keywords: cancer, oncogenesis, tumor suppression, IP₃R3, calcium signaling, PTEN, FBXL2, BAP1.

In some cells, the role played by the IP₃R3 in pro-apoptotic Ca²⁺ transfers from ER to mitochondria might relate to its ability to preferentially partner with the VDAC1 complex [18]. Notably, several tumor suppressors and oncogenes located at the ER membranes dodge ER Ca²⁺ homeostasis and dynamics [10, 21]. In general, tumor suppressors increase ER-mitochondrial Ca²⁺ fluxes, whereas oncogenes suppress ER-mitochondrial Ca²⁺ fluxes. The actions of both tumor suppressors and oncogenes at the ER can involve changes of the steady-state ER Ca²⁺-filling state through modulation of sarco/endoplasmic reticulum Ca²⁺ ATPases (SERCA) and/or ER Ca²⁺-leak channels. For instance, during stress, the tumor suppressor p53 accumulates at the ER and enhances the Ca²⁺-pump activity SERCA, causing ER Ca²⁺ overload and the likelihood for pro-apoptotic ER-mitochondrial Ca²⁺ fluxes [22-24]. The anti-apoptotic protein Bcl-2 increases IP₃R phosphorylation and its sensitivity to IP₃, enhancing the passive Ca²⁺ leak from the ER, lowering ER Ca²⁺ levels and so pro-apoptotic ER-mitochondrial Ca²⁺ fluxes [25]. Several tumor suppressors and oncogenes have been identified as direct regulators of the IP₃R, whereby tumor suppressors (like BRCA1, PTEN, PML) and oncogenes (like Bcl-2, PKB/Akt) that respectively promote and suppress the activity of IP₃R channels by impacting their gating and consequently their open probability [10, 21]. Besides this direct regulation of IP₃R gating, it is clear that total IP₃R-protein levels impact Ca²⁺ flux from the ER to the mitochondria and in turn cellular sensitivity to death as well. Insights are now available on IP₃R degradation by ER-assisted and 26S proteasomal turnover after cell stimulation and IP₃R activation [26]. In such conditions, IP₃Rs become ubiquitinated due to recruitment of the erlin1/2 complex and RNF170, an E3 ubiquitin ligase [27-29].

Hitherto, not much was known about the molecular mechanisms impacting basal IP₃R turn-over and controlling their steady-state in stressed cells; equally, whether dysregulation of these mechanisms was involved in oncogenesis and/or tumor progression. Nevertheless, it is clear that IP₃R levels do impact apoptotic sensitivity [20, 30-32], and hence cell death and survival proteins were found implicated in regulating IP₃R levels [33, 34].

Recent work from Kuchay *et al.* revealed an unexpected role for the tumor suppressor lipid/protein phosphatase PTEN, an allele frequently lost in cancer [35] and well-known negative regulator of PKB/Akt signaling, stabilizing IP₃R channels by protecting them from proteasomal degradation [36] (**Fig. 1**). This novel function adds to its recently discovered presence at the mitochondria-associated membranes (MAMs), where it contributes to cell death sensitivity by suppressing IP₃R3-mediated Ca²⁺ fluxes [37]. Independently of its catalytic activity, PTEN competes with the F-box protein FBXL2 (the receptor subunit of one of 69 human SCF (SKP1, CUL1, F-box protein) [38, 39] to bind to IP₃R3 channels, in particular to a region in the ligand-binding domain. Normal cells that express PTEN will have a low level of the IP₃R3/FBXL2-complex formation, preventing the ubiquitination of IP₃R3 channels and subsequent targeting to the proteasome. Consistently with previous

observations [27, 40], activation of cells with agonists increase FBXL2/IP₃R3-complex formation and subsequent IP₃R3 post-transcriptional regulation via ubiquitination similarly to RNF170-mediated ubiquitination of IP₃Rs [27]. Interestingly, FBXL2 activity itself is Ca²⁺ dependent, but antagonized by calmodulin [41]. Hence, activation of IP₃Rs may not only make these channels more susceptible for degradation by increased interaction with FBXL2 but release of Ca²⁺ itself through IP₃Rs may trigger local activation of FBXL2 associated with IP₃R3 leading to IP₃R3 degradation. Notably, FBXL2 binding to its substrates (like cyclin D3) was found to occur via canonical calmodulin-binding motif thereby preventing ubiquitination. It is nonetheless likely that FBXL2 binding to IP₃Rs does not occur by targeting its calmodulin-binding motif even though a role for calmodulin in regulating IP₃R3/FBXL2-complex formation may not be ruled out. In any case, access to the degron region in the ligand-binding core of IP₃R3 for FBXL2 is facilitated by deletion of the suppressor domain and loss of PTEN is associated with increased FBXL2 binding to IP₃R3 and degradation of IP₃R3, contributing to the apoptotic resistance of cells [36]. In cancer cells lacking PTEN, FBXL2 knockdown could therefore restore IP₃R3 levels and apoptotic sensitivity.

Notably, authors show that by mutating the degron region in the ligand-binding domain in IP₃R3 the FBXL2 binding and IP₃R3 degradation were both prevented. Using a knockin approach in which wild-type IP₃R3 was altered into a non-degradable IP₃R3 mutant version (IP₃R3^{Q550A/F553A/R554A}), the authors could restore the rise of Ca²⁺ induced by photodynamic therapy and apoptosis in PTEN-negative cancer cells. This correlation was eloquently observed in human tumor samples beyond the convincing *in vivo* xenograft models, in which tumor cells expressing a non-degradable IP₃R3 version or treated with geranylgeranyltransferase inhibitor (that prevents FBXL2 accumulation at ER membranes and activity, which depends on its geranylgeranylation [42]) were greatly sensitized towards photodynamic therapy compared to tumor cells expressing wild-type (degradable) IP₃R3 or untreated tumors.

It is important to note that FBXL2 has been previously implicated in cancer, but rather acting via the suppression of cell cycle progression and proliferation, as observed in lung tumors [43], leukemic cells [44], gastric cancer cells [45], and prevalently of tumor suppressive nature. Instead, in this work, the effects of FBXL2 on IP₃R3 are tumor promoting by increasing IP₃R3 degradation and making the cells more resistant to cell death [36]. These effects are neutralized by PTEN, which prevents FBXL2 binding to IP₃R3 channels. This interference by PTEN is likely selective for IP₃R3 and not for other FBXL2 targets regulating the cell cycle. Nevertheless, the anti-cancer properties of FBXL2 activators like the small molecule BC-1258 [46] will be adversely impacted by FBXL2-mediated IP₃R3 degradation, likely limiting their application to PTEN-positive cancers. Vice versa, the previously discovered tumor suppressive properties of FBXL2 in cancer might be further boosted if FBXL2 could be selectively/subcellularly activated to pro-




FIGURE 1: IP₃R3 channels are targets for ubiquitination and deubiquitylation enzymes, whose activities are deregulated in cancer cells, thereby impacting IP₃R3 stability and ultimately cell death sensitivity. (A) IP₃R3, located at the ER-mitochondrial interface, here particularly IP₃R3, promote ER-mitochondrial Ca²⁺ fluxes, thereby enabling adequate cell death sensitivity in healthy cells undergoing genotoxic stress and cell damage, an important feature that prevents oncogenesis. The levels of IP₃R3 channels are critical for this process and are dynamically regulated via ubiquitination. BAP1, a tumor suppressor with deubiquitylation enzyme activities and frequently mutated in cancers including mesothelioma, deubiquitylates IP₃R3, preventing its proteasomal degradation and resulting in IP₃R3 stabilization. In contrast, FBXL2, an F-Box protein with ubiquitin ligase activity, ubiquitinates IP₃R3, directing it for proteasomal degradation and resulting in a decline in IP₃R3 levels. Importantly, PTEN, another tumor suppressor that antagonizes Akt signaling and frequently mutated in a variety of cancers including prostate cancer, competes with FBXL2 for IP₃R3 binding and thus antagonizes FBXL2-mediated ubiquitination of IP₃R3, stabilizing IP₃R3. In healthy cells, it is anticipated that the balance between deubiquitylation and ubiquitination of IP₃R3 channels is favored towards deubiquitylated IP₃R3, enabling sufficient IP₃R3 channels to sustain adequate ER-mitochondrial Ca²⁺ fluxes in cells undergoing genotoxic stress and cell damage to engage cell death and eradicate pre-malignant cells with DNA damage. **(B)** Several cancer types display chromosomal aberrations with defects in the expression of several tumor suppressors, including a reduction in BAP1 levels and a loss in PTEN expression. As a consequence, both alterations will favor IP₃R3 ubiquitination and degradation, since reduced BAP1 levels will lead to reduced deubiquitylation of IP₃R3 and loss of PTEN will enable FBXL2 to mediate IP₃R3 ubiquitination. As a consequence, the balance between ubiquitination and deubiquitylation of IP₃R3s will shift towards ubiquitinated IP₃R3, impairing ER-mitochondrial Ca²⁺ fluxes that are needed to engage cell death programs in response to genotoxic stress and cell damage. As such, cells will be able to withstand this cellular stress, resulting in their survival and proliferation despite accumulated DNA damage. This phenomenon is an early event in oncogenesis, enabling malignant cell formation and tumoral behavior.

mote its cell-cycle targets while shielding IP₃R3 for degradation.

In a separate study, Bononi *et al.* revealed a novel deubiquitylating enzyme that actually counteracts the level of IP₃R3 ubiquitination, namely BRCA-associated protein 1 (BAP1) [47] (**Fig. 1**). BAP1 is a potent tumor suppressor, which protects against environmental stress and damage [48]. Loss of one BAP1 allele either inherited or acquired during life has been associated with environmental stress-induced carcinogenesis, like UV light for uveal melanoma and asbestos for mesothelioma. Germline mutations in BAP1 resulting in aberrant/loss of BAP1 expression were associated with a high incidence of familial mesothelioma and uveal melanoma, while somatic mutations in BAP1 were found in sporadic mesotheliomas [49]. Germline mutations in BAP1 greatly enhanced the sensitivity of mice to develop mesothelioma when exposed to asbestos [50]. Recently, it has been shown in patient fibroblasts that loss of BAP1 displayed a metabolic rewiring towards aerobic glycolysis and reduced mitochondrial respiration associated with malignancy and carcinogenesis [51]. BAP1 was therefore identified as a novel IP₃R3-interacting protein that impacts its post-translational modification. BAP1 causes IP₃R3 stabilization and prevents the channel to be degraded by the proteasome. Loss of only 1 allele in BAP1 is sufficient to protect cells from undergoing apoptotic cell death via suppressed Ca²⁺ release triggered by apoptotic stimuli like H₂O₂. An effect operated via the IP₃R3 degradation due to a decreased BAP1-mediated deubiquitylation of IP₃R3. As a consequence, exposure of cells being BAP1^{+/-} to DNA-damaging conditions will result in a higher percentage of cells surviving despite having damaged DNA. Such cells bearing genomic aberrations are at high risk for neoplastic behavior and oncogenesis, resulting in malignant cell growth and tumor formation. However, restoring IP₃R3 in these cells could be an attractive strategy to reinstate cell death sensitivity of BAP1^{+/-} cells.

Based on these recent developments, it will be therefore important to examine the interplay between BAP1 and FBXL2. It is indeed not clear whether the increased IP₃R3 ubiquitination observed when BAP1 levels are declined is mediated through FBXL2 and whether lack of BAP1 results in increased FBXL2 association with IP₃R3 channels. In any case, by inhibiting FBXL2's action on IP₃R3 without inhibiting its action on cell cycle targets could be an attractive avenue to restore IP₃R3 levels in these cancers. Another strategy can be boosting ER Ca²⁺-store loading in tumor cells by activating the SERCA pumps, thereby increasing the likelihood for ER-mitochondrial Ca²⁺ fluxes and restoring cellular sensitivity to apoptosis [22, 52]. Tumor-selective SERCA modulation is challenging but not impossible though, as evidenced by the ability to locally release thapsigargin in

the vicinity of pancreatic tumor cells using peptide-coupled prodrugs that are enzymatically cleaved by prostate-specific factors [53]. SERCA-activating approaches may be based on p53, recently proposed to activate SERCA activity in response to chemotherapy [22-24], although p53 is very frequently mutated in cancer. In wild-type p53 tumors, direct p53 activators might be of use. Other solace may come from SERCA-activating small molecules like CDN1163 [54], provided these agents can be delivered to tumor cells while sparing healthy cells [52].

These recent papers highlight that deregulation of IP₃R3 ubiquitination homeostasis not only impacts death and survival of cells but also contributes to the oncogenic behavior of cells with dysfunctional tumor suppressors by either (i) lacking PTEN [36] or by (ii) displaying deficiencies in BAP1 [47]. They underpin the emerging role of altered Ca²⁺ signaling at the MAM level as a key event in apoptosis resistance that contributes early events associated with oncogenesis and tumor formation. Challenging will be the translation of these insights into anti-cancer therapies for which not only tumor-selective applications will be required but also further understanding in the selective targeting at the level of the IP₃R3.

ACKNOWLEDGEMENTS

The research activities led by M.C. are supported by the following funders, who are gratefully acknowledged: The Biotechnology and Biological Sciences Research Council; the London Interdisciplinary Doctoral Training Programme (LiDO); The Petplan Charitable Trust and the LAM-Bighi Grant Initiative. The research activities led by G.B. are supported by the following funders, who are gratefully acknowledged: Scientific Research – Flanders (FWO), Research Council – KU Leuven and Stichting Tegen Kanker.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

COPYRIGHT

© 2017 Bultynck and Campanella. This is an open-access article released under the terms of the Creative Commons Attribution (CC BY) license, which allows the unrestricted use, distribution, and reproduction in any medium, provided the original author and source are acknowledged.

Please cite this article as: Geert Bultynck and Michelangelo Campanella (2017). Tumor suppressive Ca²⁺ signaling is driven by IP₃ receptor fitness. *Cell Stress* 1(2): 73-78. doi: 10.15698/cst2017.11.109

REFERENCES

1. Monteith GR, Prevarskaya N, Roberts-Thomson SJ (2017). The calcium-cancer signalling nexus. *Nat Rev Cancer* 17(6): 367-380. doi: 10.1038/nrc.2017.18
2. Berridge MJ, Bootman MD, Roderick HL (2003). Calcium signalling: dynamics, homeostasis and remodelling. *Nat Rev Mol Cell Biol* 4(7): 517-529. doi: 10.1038/nrm1155

3. Marchi S, Patergnani S, Missiroli S, Morciano G, Rimessi A, Wieckowski MR, Giorgi C, Pinton P (2017). Mitochondrial and endoplasmic reticulum calcium homeostasis and cell death. **Cell Calcium**. doi: 10.1016/j.ceca.2017.05.003
4. Bultynck G, Parys JB (2017). Ca²⁺ signaling and cell death: Focus on Ca²⁺-transport systems and their implication in cell death and survival. **Cell Calcium**. doi: 10.1016/j.ceca.2017.09.001
5. Ivanova H, Kerkhofs M, La Rovere RM, Bultynck G (2017). Endoplasmic Reticulum-Mitochondrial Ca²⁺ Fluxes Underlying Cancer Cell Survival. **Front Oncol** 7(70). doi: 10.3389/fonc.2017.00070
6. Marchi S, Patergnani S, Pinton P (2014). The endoplasmic reticulum-mitochondria connection: one touch, multiple functions. **Biochim Biophys Acta** 1837(4): 461-469. doi: 10.1016/j.bbabi.2013.10.015
7. Marchi S, Bittremieux M, Missiroli S, Morganti C, Patergnani S, Sbrano L, Rimessi A, Kerkhofs M, Parys JB, Bultynck G, Giorgi C, Pinton P (2017). Endoplasmic Reticulum-Mitochondria Communication Through Ca²⁺ Signaling: The Importance of Mitochondria-Associated Membranes (MAMs). **Adv Exp Med Biol** 997: 49-67. doi: 10.1007/978-981-10-4567-7_4
8. La Rovere RM, Roest G, Bultynck G, Parys JB (2016). Intracellular Ca²⁺ signaling and Ca²⁺ microdomains in the control of cell survival, apoptosis and autophagy. **Cell Calcium** 60(2): 74-87. doi: 10.1016/j.ceca.2016.04.005
9. Giorgi C, Missiroli S, Patergnani S, Duszynski J, Wieckowski MR, Pinton P (2015). Mitochondria-associated membranes: composition, molecular mechanisms, and physiopathological implications. **Antioxid Redox Signal** 22(12): 995-1019. doi: 10.1089/ars.2014.6223
10. Bittremieux M, Parys JB, Pinton P, Bultynck G (2016). ER functions of oncogenes and tumor suppressors: Modulators of intracellular Ca signaling. **Biochim Biophys Acta** 1863(6 Pt B): 1364-78. doi: 10.1016/j.bbamcr.2016.01.002
11. Vervliet T, Clerix E, Seitaj B, Ivanova H, Monaco G, Bultynck G (2017). Modulation of Ca²⁺ signaling by anti-apoptotic Bcl-2 proteins at the ER-mitochondrial interface. **Frontiers in Oncology** 7: 75. doi: 10.3389/fonc.2017.00075
12. Marchi S, Giorgi C, Oparka M, Duszynski J, Wieckowski MR, Pinton P (2014). Oncogenic and oncosuppressive signal transduction at mitochondria-associated endoplasmic reticulum membranes. **Mol Cell Oncol** 1(2): e956469. doi: 10.4161/23723548.2014.956469
13. Cardenas C, Foskett JK (2012). Mitochondrial Ca²⁺ signals in autophagy. **Cell Calcium** 52(1): 44-51. doi: 10.1016/j.ceca.2012.03.001
14. Giorgi C, Baldassari F, Bononi A, Bonora M, De Marchi E, Marchi S, Missiroli S, Patergnani S, Rimessi A, Suski JM, Wieckowski MR, Pinton P (2012). Mitochondrial Ca²⁺ and apoptosis. **Cell Calcium** 52(1): 36-43. doi: 10.1016/j.ceca.2012.02.008
15. Pedriali G, Rimessi A, Sbrano L, Giorgi C, Wieckowski MR, Previati M, Pinton P (2017). Regulation of Endoplasmic Reticulum-Mitochondria Ca²⁺ Transfer and Its Importance for Anti-Cancer Therapies. **Front Oncol** 7: 180. doi: 10.3389/fonc.2017.00180
16. Kerkhofs M, Giorgi C, Marchi S, Seitaj B, Parys JB, Pinton P, Bultynck G, Bittremieux M (2017). Alterations in Ca²⁺ Signalling via ER-Mitochondria Contact Site Remodelling in Cancer. **Adv Exp Med Biol** 997(225-254). doi: 10.1007/978-981-10-4567-7_17
17. Marchi S, Marinello M, Bononi A, Bonora M, Giorgi C, Rimessi A, Pinton P (2012). Selective modulation of subtype III IP₃R by Akt regulates ER Ca²⁺ release and apoptosis. **Cell Death Dis** 3: e304. doi: 10.1038/cddis.2012.45
18. De Stefani D, Bononi A, Romagnoli A, Messina A, De Pinto V, Pinton P, Rizzuto R (2012). VDAC1 selectively transfers apoptotic Ca²⁺ signals to mitochondria. **Cell Death Differ** 19(2): 267-273. doi: 10.1038/cdd.2011.92
19. Ivanova H, Vervliet T, Missiaen L, Parys JB, De Smedt H, Bultynck G (2014). Inositol 1,4,5-trisphosphate receptor-isoform diversity in cell death and survival. **Biochim Biophys Acta** 1843(10): 2164-2183. doi: 10.1016/j.bbamcr.2014.03.007
20. Jayaraman T, Marks AR (1997). T cells deficient in inositol 1,4,5-trisphosphate receptor are resistant to apoptosis. **Mol Cell Biol** 17(6): 3005-3012. PMID: 9154798
21. Akl H, Bultynck G (2013). Altered Ca²⁺ signaling in cancer cells: proto-oncogenes and tumor suppressors targeting IP₃ receptors. **Biochim Biophys Acta** 1835(2): 180-193. doi: 10.1016/j.bbcan.2012.12.001
22. Bittremieux M, Bultynck G (2015). p53 and Ca²⁺ signaling from the endoplasmic reticulum: partners in anti-cancer therapies. **Oncoscience** 2(3): 233-238. doi: 10.18632/oncoscience.139
23. Giorgi C, Bonora M, Missiroli S, Poletti F, Ramirez FG, Morciano G, Morganti C, Pandolfi PP, Mammano F, Pinton P (2015). Intravital imaging reveals p53-dependent cancer cell death induced by phototherapy via calcium signaling. **Oncotarget** 6(3): 1435-1445. doi: 10.18632/oncotarget.2935
24. Giorgi C, Bonora M, Sorrentino G, Missiroli S, Poletti F, Suski JM, Galindo Ramirez F, Rizzuto R, Di Virgilio F, Zito E, Pandolfi PP, Wieckowski MR, Mammano F, Del Sal G, Pinton P (2015). p53 at the endoplasmic reticulum regulates apoptosis in a Ca²⁺-dependent manner. **Proc Natl Acad Sci U S A** 112(6): 1779-1784. doi: 10.1073/pnas.1410723112
25. Oakes SA, Scorrano L, Opferman JT, Bassik MC, Nishino M, Pozzan T, Korsmeyer SJ (2005). Proapoptotic BAX and BAK regulate the type 1 inositol trisphosphate receptor and calcium leak from the endoplasmic reticulum. **Proc Natl Acad Sci USA** 102(1): 105-110. doi: 10.1073/pnas.0408352102
26. Wright FA, Wojcikiewicz RJ (2016). Chapter 4 - Inositol 1,4,5-Trisphosphate Receptor Ubiquitination. **Prog Mol Biol Transl Sci** 141(141-159). doi: 10.1016/bs.pmbts.2016.02.004
27. Lu JP, Wang Y, Sliter DA, Pearce MM, Wojcikiewicz RJ (2011). RNF170 protein, an endoplasmic reticulum membrane ubiquitin ligase, mediates inositol 1,4,5-trisphosphate receptor ubiquitination and degradation. **J Biol Chem** 286(27): 24426-24433. doi: 10.1074/jbc.M111.251983
28. Pearce MM, Wormer DB, Wilkens S, Wojcikiewicz RJ (2009). An endoplasmic reticulum (ER) membrane complex composed of SPFH1 and SPFH2 mediates the ER-associated degradation of inositol 1,4,5-trisphosphate receptors. **J Biol Chem** 284(16): 10433-10445. doi: 10.1074/jbc.M809801200
29. Wright FA, Lu JP, Sliter DA, Dupre N, Rouleau GA, Wojcikiewicz RJ (2015). A Point Mutation in the Ubiquitin Ligase RNF170 That Causes Autosomal Dominant Sensory Ataxia Destabilizes the Protein and Impairs Inositol 1,4,5-Trisphosphate Receptor-mediated Ca²⁺ Signaling. **J Biol Chem** 290(22): 13948-13957. doi: 10.1074/jbc.M115.655043
30. Madesh M, Hawkins BJ, Milovanova T, Bhanumathy CD, Joseph SK, Ramachandrarao SP, Sharma K, Kurosaki T, Fisher AB (2005). Selective role for superoxide in InsP₃ receptor-mediated mitochondrial dysfunction and endothelial apoptosis. **J Cell Biol** 170(7): 1079-1090. doi: 10.1083/jcb.200505022
31. Khan MT, Bhanumathy CD, Schug ZT, Joseph SK (2007). Role of inositol 1,4,5-trisphosphate receptors in apoptosis in DT40 lymphocytes. **J Biol Chem** 282(45): 32983-32990. doi: 10.1074/jbc.M705183200

32. Assefa Z, Bultynck G, Szlufcik K, Nadif Kasri N, Vermassen E, Goris J, Missiaen L, Callewaert G, Parys JB, De Smedt H (2004). Caspase-3-induced truncation of type 1 inositol trisphosphate receptor accelerates apoptotic cell death and induces inositol trisphosphate-independent calcium release during apoptosis. *J Biol Chem* 279(41): 43227-43236. doi: 10.1074/jbc.M403872200
33. Schulman JJ, Wright FA, Kaufmann T, Wojcikiewicz RJ (2013). The Bcl-2 protein family member Bok binds to the coupling domain of inositol 1,4,5-trisphosphate receptors and protects them from proteolytic cleavage. *J Biol Chem* 288(35): 25340-25349. doi: 10.1074/jbc.M113.496570
34. Pierro C, Cook SJ, Foets TC, Bootman MD, Roderick HL (2014). Oncogenic K-Ras suppresses IP₃-dependent Ca²⁺ release through remodelling of the isoform composition of IP₃Rs and ER luminal Ca²⁺ levels in colorectal cancer cell lines. *J Cell Sci* 127(Pt 7): 1607-1619. doi: 10.1242/jcs.141408
35. Beroukhi R, Mermel CH, Porter D, Wei G, Raychaudhuri S, Donovan J, Barretina J, Boehm JS, Dobson J, Urashima M, Mc Henry KT, Pinchback RM, Ligon AH, Cho YJ, Haery L, Greulich H, Reich M, Winkler W, Lawrence MS, Weir BA, Tanaka KE, Chiang DY, Bass AJ, Loo A, Hoffman C, Prensner J, Liefeld T, Gao Q, Yecies D, Signoretti S, et al. (2010). The landscape of somatic copy-number alteration across human cancers. *Nature* 463(7283): 899-905. doi: 10.1038/nature08822
36. Kuchay S, Giorgi C, Simoneschi D, Pagan J, Missiroli S, Saraf A, Florens L, Washburn MP, Collazo-Lorduy A, Castillo-Martin M, Cordon-Cardo C, Sebt SM, Pinton P, Pagano M (2017). PTEN counteracts FBXL2 to promote IP₃R3- and Ca²⁺-mediated apoptosis limiting tumour growth. *Nature* 546(7659): 554-558. doi: 10.1038/nature22965
37. Bononi A, Bonora M, Marchi S, Missiroli S, Poletti F, Giorgi C, Pandolfi PP, Pinton P (2013). Identification of PTEN at the ER and MAMs and its regulation of Ca²⁺ signaling and apoptosis in a protein phosphatase-dependent manner. *Cell Death Differ* 20(12): 1631-1643. doi: 10.1038/cdd.2013.77
38. Skaar JR, Pagan JK, Pagano M (2009). SnapShot: F box proteins I. *Cell* 137(6): 1160-1160 e1161. doi: 10.1016/j.cell.2009.05.039
39. Ilyin GP, Riialand M, Glaise D, Guguen-Guillouzo C (1999). Identification of a novel Skp2-like mammalian protein containing F-box and leucine-rich repeats. *FEBS Lett* 459(1): 75-79. PMID: 10508920
40. Zhu CC, Wojcikiewicz RJ (2000). Ligand binding directly stimulates ubiquitination of the inositol 1, 4,5-trisphosphate receptor. *Biochem J* 348 (Pt 3): 551-556. PMID: 10839985
41. Chen BB, Coon TA, Glasser JR, Mallampalli RK (2011). Calmodulin antagonizes a calcium-activated SCF ubiquitin E3 ligase subunit, FBXL2, to regulate surfactant homeostasis. *Mol Cell Biol* 31(9): 1905-1920. doi: 10.1128/MCB.00723-10
42. Wang C, Gale M, Jr., Keller BC, Huang H, Brown MS, Goldstein JL, Ye J (2005). Identification of FBL2 as a geranylgeranylated cellular protein required for hepatitis C virus RNA replication. *Mol Cell* 18(4): 425-434. doi: 10.1016/j.molcel.2005.04.004
43. Chen BB, Glasser JR, Coon TA, Mallampalli RK (2012). F-box protein FBXL2 exerts human lung tumor suppressor-like activity by ubiquitin-mediated degradation of cyclin D3 resulting in cell cycle arrest. *Oncogene* 31(20): 2566-2579. doi: 10.1038/onc.2011.432
44. Chen BB, Glasser JR, Coon TA, Zou C, Miller HL, Fenton M, McDyer JF, Boyiadzis M, Mallampalli RK (2012). F-box protein FBXL2 targets cyclin D2 for ubiquitination and degradation to inhibit leukemic cell proliferation. *Blood* 119(13): 3132-3141. doi: 10.1182/blood-2011-06-358911
45. Li LQ, Pan D, Chen H, Zhang L, Xie WJ (2016). F-box protein FBXL2 inhibits gastric cancer proliferation by ubiquitin-mediated degradation of forkhead box M1. *FEBS Lett* 590(4): 445-452. doi: 10.1002/1873-3468.12071
46. Chen BB, Glasser JR, Coon TA, Mallampalli RK (2013). Skp-cullin-F box E3 ligase component FBXL2 ubiquitinates Aurora B to inhibit tumorigenesis. *Cell Death Dis* 4: e759. doi: 10.1038/cddis.2013.271
47. Bononi A, Giorgi C, Patergnani S, Larson D, Verbruggen K, Tanji M, Pellegrini L, Signorato V, Olivetto F, Pastorino S, Nasu M, Napolitano A, Gaudino G, Morris P, Sakamoto G, Ferris LK, Danese A, Raimondi A, Tacchetti C, Kuchay S, Pass HI, Affar EB, Yang H, Pinton P, Carbone M (2017). BAP1 regulates IP₃R3-mediated Ca²⁺ flux to mitochondria suppressing cell transformation. *Nature* 546(7659): 549-553. doi: 10.1038/nature22798
48. Jensen DE, Proctor M, Marquis ST, Gardner HP, Ha SI, Chodosh LA, Ishov AM, Tommerup N, Vissing H, Sekido Y, Minna J, Borodovsky A, Schultz DC, Wilkinson KD, Maul GG, Barlev N, Berger SL, Prendergast GC, Rauscher FJ, 3rd (1998). BAP1: a novel ubiquitin hydrolase which binds to the BRCA1 RING finger and enhances BRCA1-mediated cell growth suppression. *Oncogene* 16(9): 1097-1112. PMID: 9528852
49. Testa JR, Cheung M, Pei J, Below JE, Tan Y, Sementino E, Cox NJ, Dogan AU, Pass HI, Trusa S, Hesdorffer M, Nasu M, Powers A, Rivera Z, Comertpay S, Tanji M, Gaudino G, Yang H, Carbone M (2011). Germline BAP1 mutations predispose to malignant mesothelioma. *Nat Genet* 43(10): 1022-1025. doi: 10.1038/ng.912
50. Xu J, Kadariya Y, Cheung M, Pei J, Talarchek J, Sementino E, Tan Y, Menges CW, Cai KQ, Litwin S, Peng H, Karar J, Rauscher FJ, Testa JR (2014). Germline mutation of Bap1 accelerates development of asbestos-induced malignant mesothelioma. *Cancer Res* 74(16): 4388-4397. doi: 10.1158/0008-5472.CAN-14-1328
51. Bononi A, Yang H, Giorgi C, Patergnani S, Pellegrini L, Su M, Xie G, Signorato V, Pastorino S, Morris P, Sakamoto G, Kuchay S, Gaudino G, Pass HI, Napolitano A, Pinton P, Jia W, Carbone M (2017). Germline BAP1 mutations induce a Warburg effect. *Cell Death Differ* 24(10): 1694-1704. doi: 10.1038/cdd.2017.95
52. Chemaly ER, Troncone L, Lebeche D (2017). SERCA control of cell death and survival. *Cell Calcium* doi: 10.1016/j.ceca.2017.07.001
53. Doan NT, Paulsen ES, Sehgal P, Moller JV, Nissen P, Denmeade SR, Isaacs JT, Dionne CA, Christensen SB (2015). Targeting thapsigargin towards tumors. *Steroids* 97:2-7. doi: 10.1016/j.steroids.2014.07.009
54. Kang S, Dahl R, Hsieh W, Shin A, Zsebo KM, Buettner C, Hajjar RJ, Lebeche D (2016). Small Molecular Allosteric Activator of the Sarco/Endoplasmic Reticulum Ca²⁺-ATPase (SERCA) Attenuates Diabetes and Metabolic Disorders. *J Biol Chem* 291(10): 5185-5198. doi: 10.1074/jbc.M115.705012