

Classes of Ordinary Differential Equations Obtained for the Probability Functions of Exponentiated Pareto Distribution

Hilary I. Okagbue, Olusunmbo O. Agboola, *Member, IAENG*, Paulinus O. Ugwoke and Abiodun A. Opanuga

Abstract— In this paper, the differential calculus was used to obtain some classes of ordinary differential equations (ODEs) for the probability density function, quantile function, survival function, inverse survival function, hazard function and reversed hazard function of the exponentiated Pareto distribution. The stated necessary conditions required for the existence of the ODEs are consistent with the various parameters that defined the distribution. Solutions of these ODEs by using numerous available methods are new ways of understanding the nature of the probability functions that characterize the distribution. The method can be extended to other probability distributions and can serve as an alternative to approximation.

Index Terms— Exponentiated Pareto distribution, reversed hazard function, calculus, differentiation, inverse survival function

I. INTRODUCTION

THE distribution was introduced by Gupta et al. [1] for modeling failure time data but was proposed formally as a probability model by [2]. Afify [3] obtained the Bayes estimates for the distribution. Ali and Woo [4] obtained the maximum likelihood estimates of the tail probability of the distribution. Nooghabi and Nooghabi [5] derived the improved moments for the distribution. The distribution is a submodel of the extended Pareto distribution proposed by Mead [6] and the new Weibull –Pareto distribution by Tahir et al. [7].

Other examples of exponentiated distributions obtained by different researchers include: exponentiated Weibull distribution [8-10], exponentiated generalized inverted exponential distribution [11], exponentiated generalized inverse Gaussian distribution [12], exponentiated generalized inverse Weibull distribution [13-14], gamma-exponentiated exponential distribution [15], exponentiated Gompertz distribution [16-17], beta exponentiated

Mukherjee-Islam Distribution [18], transmuted exponentiated Pareto-i distribution [19], gamma exponentiated exponential–Weibull distribution [20], exponentiated gamma distribution [21], exponentiated Gumbel distribution [22], exponentiated uniform distribution [23] and beta exponentiated Weibull distribution [24-25]. Others are: exponentiated log-logistic distribution [26], McDonald exponentiated gamma distribution [27], exponentiated Generalized Weibull Distribution [28], beta exponentiated gamma distribution [29], exponentiated gamma distribution [30], exponentiated Pareto distribution [31], exponentiated Kumaraswamy distribution [32], exponentiated modified Weibull extension distribution [33], exponentiated Weibull-Pareto distribution [34], exponentiated lognormal distribution [35], exponentiated Perks distribution [36], Kumaraswamy-transmuted exponentiated modified Weibull distribution [37], exponentiated power Lindley–Poisson distribution [38] and exponentiated Chen distribution [39]. Also available in scientific are: exponentiated reduced Kies distribution [40], exponentiated inverse Weibull geometric distribution [41], exponentiated geometric distribution [42-43], exponentiated Weibull geometric distribution [44], exponentiated transmuted Weibull geometric distribution [45], exponentiated half logistic distribution [46], transmuted exponentiated Gumbel distribution [47], exponentiated Kumaraswamy-power function distribution [48], exponentiated-log-logistic geometric distribution [49] and bivariate exponentiated generalized Weibull-Gompertz distribution [50].

The aim of this research is to develop ordinary differential equations (ODE) for the probability density function (PDF), Quantile function (QF), survival function (SF), inverse survival function (ISF), hazard function (HF) and reversed hazard function (RHF) of exponentiated Pareto distribution by the use of differential calculus. Calculus is a very key tool in the determination of mode of a given probability distribution and in estimation of parameters of probability distributions, amongst other uses. The research is an extension of the ODE to other probability functions other than the PDF. Similar works done where the PDF of probability distributions was expressed as ODE whose solution is the PDF are available. They include: Laplace distribution [51], beta distribution [52], raised cosine distribution [53], Lomax distribution [54], beta prime distribution or inverted beta distribution [55].

Manuscript received June 30, 2017; revised July 25, 2017. This work was sponsored by Covenant University, Ota, Nigeria.

H. I. Okagbue, O. O. Agboola and A. A. Opanuga are with the Department of Mathematics, Covenant University, Ota, Nigeria (hilary.okagbue@covenantuniversity.edu.ng; ola.agboola@covenantuniversity.edu.ng; abiodun.opanuga@covenantuniversity.edu.ng)

P. O. Ugwoke is with the Department of Computer, University of Nigeria, Nsukka, Nigeria and Digital Bridge Institute, International Centre for Information & Communications Technology Studies, Abuja, Nigeria.

II. PROBABILITY DENSITY FUNCTION

The probability density function of the exponentiated Pareto distribution is given as;

$$f(x) = \alpha\theta(1+x)^{-(\alpha+1)}[1-(1+x)^{-\alpha}]^{\theta-1} \quad (1)$$

To obtain the first order ordinary differential equation for the probability density function of the exponentiated Pareto distribution, differentiate equation (1), to obtain;

$$f'(x) = \left\{ \begin{array}{l} -\frac{(\alpha+1)(1+x)^{-(\alpha+2)}}{(1+x)^{-(\alpha+1)}} \\ +\frac{\alpha(\theta-1)(1+x)^{-(\alpha+1)}[1-(1+x)^{-\alpha}]^{\theta-2}}{[1-(1+x)^{-\alpha}]^{\theta-1}} \end{array} \right\} f(x) \quad (2)$$

$$f'(x) = \left\{ -\frac{(\alpha+1)}{(1+x)} + \frac{\alpha(\theta-1)(1+x)^{-(\alpha+1)}}{1-(1+x)^{-\alpha}} \right\} f(x) \quad (3)$$

The condition necessary to the existence of equation is $\alpha, \theta, x > 0$.

The ordinary differential equations can be obtained for given values of the parameters. Some cases are considered and can be seen in Table I.

TABLE I
CLASSES OF DIFFERENTIAL EQUATIONS OBTAINED FOR THE PROBABILITY DENSITY FUNCTION OF EXPONENTIATED PARETO DISTRIBUTION FOR DIFFERENT PARAMETERS

α	θ	Ordinary differential equations
1	1	$(1+x)f'(x) + 2f(x) = 0$
1	2	$x^3(1+x)f'(x) + (2x^3 - (1+x)^2)f(x) = 0$
1	3	$x^3(1+x)f'(x) + 2(x^3 - (1+x)^2)f(x) = 0$

To obtain an ordinary differential equation that is independent of the powers of the parameters, differentiate equation (3);

$$f''(x) = \left\{ \begin{array}{l} \frac{\alpha+1}{(1+x)^2} - \frac{(\theta-1)\alpha^2((1+x)^{-(\alpha+1)})^2}{(1-(1+x)^{-\alpha})^2} \\ -\frac{\alpha(\theta-1)(\alpha+1)(1+x)^{-(\alpha+2)}}{1-(1+x)^{-\alpha}} \end{array} \right\} f(x) + \left\{ -\frac{(\alpha+1)}{(1+x)} + \frac{\alpha(\theta-1)(1+x)^{-(\alpha+1)}}{1-(1+x)^{-\alpha}} \right\} f'(x) \quad (4)$$

The following equations obtained from equation (3) are required in the simplification of equation (4). From equation (3);

$$-\frac{(\alpha+1)}{(1+x)} + \frac{\alpha(\theta-1)(1+x)^{-(\alpha+1)}}{1-(1+x)^{-\alpha}} = \frac{f'(x)}{f(x)} \quad (5)$$

$$\frac{\alpha(\theta-1)(1+x)^{-(\alpha+1)}}{1-(1+x)^{-\alpha}} = \frac{f'(x)}{f(x)} + \frac{\alpha+1}{1+x} \quad (6)$$

$$\left(\frac{\alpha(\theta-1)(1+x)^{-(\alpha+1)}}{1-(1+x)^{-\alpha}} \right)^2 = \left(\frac{f'(x)}{f(x)} + \frac{\alpha+1}{1+x} \right)^2 \quad (7)$$

$$\frac{(\theta-1)\alpha^2((1+x)^{-(\alpha+1)})^2}{(1-(1+x)^{-\alpha})^2} = \frac{1}{\theta-1} \left(\frac{f'(x)}{f(x)} + \frac{\alpha+1}{1+x} \right)^2 \quad (8)$$

$$\frac{\alpha(\theta-1)(\alpha+1)(1+x)^{-(\alpha+1)}}{1-(1+x)^{-\alpha}} = \alpha+1 \left(\frac{f'(x)}{f(x)} + \frac{\alpha+1}{1+x} \right) \quad (9)$$

$$\frac{\alpha(\theta-1)(\alpha+1)(1+x)^{-(\alpha+2)}}{1-(1+x)^{-\alpha}} = \frac{\alpha+1}{x+1} \left(\frac{f'(x)}{f(x)} + \frac{\alpha+1}{1+x} \right) \quad (10)$$

Substituting equations (5), (8) and (10) into equation (4) gives

$$f''(x) = \frac{f'^2(x)}{f(x)} + \left\{ \begin{array}{l} \frac{\alpha+1}{(1+x)^2} - \frac{1}{\theta-1} \left(\frac{f'(x)}{f(x)} + \frac{\alpha+1}{1+x} \right) \\ 2 - \frac{\alpha+1}{x+1} \left(\frac{f'(x)}{f(x)} + \frac{\alpha+1}{1+x} \right) \end{array} \right\} f(x) \quad (11)$$

The condition necessary to the existence of equation is $\alpha, x > 0, \theta > 1$.

The ordinary differential equations can be obtained for the particular values of the parameters.

III. QUANTILE FUNCTION

The Quantile function of the exponentiated Pareto distribution is given as;

$$Q(p) = \frac{1}{(1-p^{\frac{1}{\theta}})^{\alpha}} - 1 \quad (12)$$

To obtain the first order ordinary differential equation for the quantile function of the exponentiated Pareto distribution, differentiate equation (12), to obtain;

$$Q'(p) = \frac{1}{\alpha\theta} p^{\frac{1}{\theta}-1} (1-p^{\frac{1}{\theta}})^{-\frac{1}{\alpha}(\frac{1}{\theta}+1)} \quad (13)$$

The condition necessary to the existence of equation is $\alpha, \theta > 0, 0 < p < 1$.

Equation (12) can be written as;

$$\frac{1}{(1-p^{\frac{1}{\theta}})^{\alpha}} = Q(p) + 1 \quad (14)$$

Substituting equation (14) into equation (13) yields

$$Q'(p) = \frac{p^{\frac{1}{\theta}}(Q(p)+1)}{\alpha\theta p(1-p^{\frac{1}{\theta}})} \quad (15)$$

$$\alpha\theta p(1-p^{\frac{1}{\theta}})Q'(p) - p^{\frac{1}{\theta}}(Q(p)+1) = 0 \quad (16)$$

The ordinary differential equations can be obtained for given values of the parameters. Some cases are considered and can be seen in Table II.

TABLE II
CLASSES OF DIFFERENTIAL EQUATIONS OBTAINED FOR THE
QUANTILE FUNCTION OF EXPONENTIATED PARETO
DISTRIBUTION FOR DIFFERENT PARAMETERS

α	θ	ordinary differential equations
1	1	$(1-p)Q'(p) - Q(p) - 1 = 0$
1	2	$2(1-p)Q'(p) - Q(p) - 1 = 0$
1	3	$3(1-p)Q'(p) - Q(p) - 1 = 0$
2	1	$2p(1-\sqrt{p})Q'(p) - \sqrt{p}Q(p) - \sqrt{p} = 0$
2	2	$4p(1-\sqrt{p})Q'(p) - \sqrt{p}Q(p) - \sqrt{p} = 0$

IV. SURVIVAL FUNCTION

The survival function of the exponentiated Pareto distribution is given as;

$$S(t) = 1 - [1 - (1+t)^{-\alpha}]^\theta \quad (17)$$

To obtain the first order ordinary differential equation for the survival function of the exponentiated Pareto distribution, differentiate equation (17), to obtain;

$$S'(t) = -\alpha\theta(1+t)^{-(\alpha+1)}[1 - (1+t)^{-\alpha}]^{\theta-1} \quad (18)$$

$$S'(t) = -\frac{\alpha\theta(1+t)^{-\alpha}[1 - (1+t)^{-\alpha}]^\theta}{(1+t)[1 - (1+t)^{-\alpha}]} \quad (19)$$

The condition necessary to the existence of equation is $\alpha, \theta, t > 0$.

Equation (17) can be written as;

$$[1 - (1+t)^{-\alpha}]^\theta = 1 - S(t) \quad (20)$$

Substituting equation (20) into equation (19) one has

$$S'(t) = -\frac{\alpha\theta(1+t)^{-\alpha}(1 - S(t))}{(1+t)[1 - (1+t)^{-\alpha}]} \quad (21)$$

$$(1+t)(1 - (1+t)^{-\alpha})S'(t) + \alpha\theta(1+t)^{-\alpha}(1 - S(t)) = 0 \quad (22)$$

The ordinary differential equations can be obtained for given values of the parameters. Some cases are considered and are shown in Table III.

TABLE III
CLASSES OF DIFFERENTIAL EQUATIONS OBTAINED FOR THE
SURVIVAL FUNCTION OF EXPONENTIATED PARETO
DISTRIBUTION FOR DIFFERENT PARAMETERS

α	θ	ordinary differential equations
1	1	$t(1+t)S'(t) - S(t) + 1 = 0$
1	2	$t(1+t)S'(t) - 2S(t) + 2 = 0$
1	3	$t(1+t)S'(t) - 3S(t) + 3 = 0$
2	1	$t(1+t)(2+t)S'(t) - 2S(t) + 2 = 0$
2	2	$t(1+t)(2+t)S'(t) - 4S(t) + 4 = 0$
2	3	$t(1+t)(2+t)S'(t) - 6S(t) + 6 = 0$

Using equation (3) and (18), it can be seen that;

$$S'(t) = -f(t) \Rightarrow S''(t) = -f'(t) \quad (23)$$

$$S''(t) = \left\{ -\frac{(\alpha+1)}{(1+t)} + \frac{\alpha(\theta-1)(1+t)^{-(\alpha+1)}}{1 - (1+t)^{-\alpha}} \right\} S'(t) \quad (24)$$

The condition necessary to the existence of equation is $\alpha, \theta, t > 0$.

The ordinary differential equations can be obtained for given values of the parameters. Some cases are considered and shown in Table IV.

TABLE IV:
CLASSES OF SECOND ORDER DIFFERENTIAL EQUATIONS
OBTAINED FOR THE PROBABILITY SURVIVAL FUNCTION OF
EXPONENTIATED PARETO DISTRIBUTION FOR DIFFERENT
PARAMETERS

α	θ	ordinary differential equations
1	1	$(1+t)S''(t) + 2S'(t) = 0$
1	2	$t^3(1+t)S''(t) + (2t^3 - (1+t)^2)S'(t) = 0$
1	3	$t^3(1+t)S''(t) + 2(t^3 - (1+t)^2)S'(t) = 0$

V. INVERSE SURVIVAL FUNCTION

The inverse survival function of the exponentiated Pareto distribution is given as;

$$Q(p) = \frac{1}{(1 - (1-p)^\theta)^{\frac{1}{\alpha}}} - 1 \quad (25)$$

To obtain the first order ordinary differential equation for the inverse survival function of the exponentiated Pareto distribution, differentiate equation (25), to obtain;

$$Q'(p) = -\frac{(1-p)^{\frac{1}{\theta}-1} (1 - (1-p)^\theta)^{-\frac{1}{\alpha}}}{\alpha\theta} \quad (26)$$

The condition necessary to the existence of equation is $\alpha, \theta > 0, 0 < p < 1$.

$$Q'(p) = -\frac{(1-p)^{\frac{1}{\theta}} (1 - (1-p)^\theta)^{\frac{1}{\alpha}}}{\alpha\theta(1-p)(1 - (1-p)^\theta)} \quad (27)$$

Equation (25) can also be written as;

$$\frac{1}{(1 - (1-p)^\theta)^{\frac{1}{\alpha}}} = Q(p) + 1 \quad (28)$$

Substituting equation (28) into equation (27), one obtains

$$Q'(p) = -\frac{(1-p)^{\frac{1}{\theta}}(Q(p)+1)}{\alpha\theta(1-p)(1 - (1-p)^\theta)} \quad (29)$$

$$\alpha\theta(1-p)(1 - (1-p)^\theta)Q'(p) + (1-p)^{\frac{1}{\theta}}(Q(p)+1) = 0 \quad (30)$$

The ordinary differential equations can be obtained for given values of the parameters. Some cases are considered and shown in Table V.

TABLE V
CLASSES OF DIFFERENTIAL EQUATIONS OBTAINED FOR THE
INVERSE SURVIVAL FUNCTION OF EXPONENTIATED PARETO
DISTRIBUTION FOR DIFFERENT PARAMETERS

θ	α	Ordinary differential equations
1	1	$pQ'(p) + Q(p) + 1 = 0$
1	2	$2pQ'(p) + Q(p) + 1 = 0$
1	3	$3pQ'(p) + Q(p) + 1 = 0$
2	1	$2(1-p)(1-\sqrt{1-p})Q'(p) + \sqrt{1-p}(Q'(p)+1) = 0$
2	2	$4(1-p)(1-\sqrt{1-p})Q'(p) + \sqrt{1-p}(Q'(p)+1) = 0$
2	3	$6(1-p)(1-\sqrt{1-p})Q'(p) + \sqrt{1-p}(Q'(p)+1) = 0$

VI. HAZARD FUNCTION

The hazard function of the exponentiated Pareto distribution is:

$$h(t) = \frac{\alpha\theta(1+t)^{-(\alpha+1)}[1-(1+t)^{-\alpha}]^{\theta-1}}{1-[1-(1+t)^{-\alpha}]^{\theta}} \quad (31)$$

To obtain the first order ordinary differential equation for the hazard function of the exponentiated Pareto distribution, differentiate equation (31), to obtain;

$$h'(t) = -\frac{(\alpha+1)(1+t)^{-(\alpha+2)}}{(1+t)^{-(\alpha+1)}}h(t) + \frac{\alpha(\theta-1)(1+t)^{-(\alpha+1)}[1-(1+t)^{-\alpha}]^{\theta-2}}{[1-(1+t)^{-\alpha}]^{\theta-1}}h(t) + \frac{\alpha\theta(1+t)^{-(\alpha+1)}[1-(1+t)^{-\alpha}]^{\theta-1}(1-[1-(1+t)^{-\alpha}]^{\theta})^{-2}h(t)}{(1-[1-(1+t)^{-\alpha}]^{\theta})^{-1}} \quad (32)$$

The condition necessary to the existence of equation is $\alpha, \theta, t > 0$.

$$h'(t) = \left\{ \begin{aligned} &-\frac{(\alpha+1)}{1+t} + \frac{\alpha(\theta-1)(1+t)^{-(\alpha+1)}}{1-(1+t)^{-\alpha}} \\ &+ \frac{\alpha\theta(1+t)^{-(\alpha+1)}[1-(1+t)^{-\alpha}]^{\theta-1}}{(1-[1-(1+t)^{-\alpha}]^{\theta})} \end{aligned} \right\} h(t) \quad (33)$$

$$h'(t) = \left\{ -\frac{(\alpha+1)}{1+t} + \frac{\alpha(\theta-1)(1+t)^{-(\alpha+1)}}{1-(1+t)^{-\alpha}} + h(t) \right\} h(t) \quad (34)$$

The ordinary differential equations can be obtained for given values of the parameters. To obtain an ordinary differential equation that is independent of the powers of the parameters,

differentiate equation (34);

$$h''(t) = \left\{ \begin{aligned} &\frac{\alpha+1}{(1+t)^2} - \frac{(\theta-1)\alpha^2((1+t)^{-(\alpha+1)})^2}{(1-(1+t)^{-\alpha})^2} \\ &-\frac{\alpha(\theta-1)(\alpha+1)(1+t)^{-(\alpha+2)}}{1-(1+t)^{-\alpha}} + h'(t) \end{aligned} \right\} h(t) \quad (35)$$

$$+ \left\{ -\frac{(\alpha+1)}{(1+t)} + \frac{\alpha(\theta-1)(1+t)^{-(\alpha+1)}}{1-(1+t)^{-\alpha}} + h(t) \right\} h'(t)$$

The condition necessary to the existence of equation is $\alpha, \theta, t > 0$.

The following equations obtained from equation (34) are required in the simplification of equation (35). From equation (34), we have

$$-\frac{(\alpha+1)}{(1+t)} + \frac{\alpha(\theta-1)(1+t)^{-(\alpha+1)}}{1-(1+t)^{-\alpha}} + h(t) = \frac{h'(t)}{h(t)} \quad (36)$$

$$\frac{\alpha(\theta-1)(1+t)^{-(\alpha+1)}}{1-(1+t)^{-\alpha}} = \frac{h'(t)}{h(t)} + \frac{\alpha+1}{1+t} - h(t) \quad (37)$$

$$\left(\frac{\alpha(\theta-1)(1+t)^{-(\alpha+1)}}{1-(1+t)^{-\alpha}} \right)^2 = \left(\frac{h'(t)}{h(t)} + \frac{\alpha+1}{1+t} - h(t) \right)^2 \quad (38)$$

$$\frac{(\theta-1)\alpha^2((1+t)^{-(\alpha+1)})^2}{(1-(1+t)^{-\alpha})^2} = \frac{1}{\theta-1} \left(\frac{h'(t)}{h(t)} + \frac{\alpha+1}{1+t} - h(t) \right)^2 \quad (39)$$

$$\frac{\alpha(\theta-1)(\alpha+1)(1+t)^{-(\alpha+1)}}{1-(1+t)^{-\alpha}} = \alpha+1 \left(\frac{h'(t)}{h(t)} + \frac{\alpha+1}{1+t} - h(t) \right) \quad (40)$$

$$\frac{\alpha(\theta-1)(\alpha+1)(1+t)^{-(\alpha+2)}}{1-(1+t)^{-\alpha}} = \frac{\alpha+1}{t+1} \left(\frac{h'(t)}{h(t)} + \frac{\alpha+1}{1+t} - h(t) \right) \quad (41)$$

Substitute equations (36), (39) and (41) into equation (35) to get

$$h''(t) = \frac{h'^2(t)}{h(t)} + \left\{ \frac{\alpha+1}{(1+t)^2} - \frac{1}{\theta-1} \left(\frac{h'(t)}{h(t)} + \frac{\alpha+1}{1+t} - h(t) \right)^2 \right\} h(t) + \left\{ -\frac{\alpha+1}{t+1} \left(\frac{h'(t)}{h(t)} + \frac{\alpha+1}{1+t} - h(t) \right) + h'(t) \right\} h(t) \quad (42)$$

The condition necessary to the existence of equation (42) is $\alpha, x > 0, \theta > 1$.

The ordinary differential equations can be obtained for the particular values of the parameters.

VII. REVERSED HAZARD FUNCTION

The reversed hazard function of the exponentiated Pareto distribution is given as;

$$j(t) = \frac{\alpha\theta(1+t)^{-(\alpha+1)}}{1-(1+t)^{-\alpha}} \quad (43)$$

To obtain the first order ordinary differential equation for the reversed hazard function of the exponentiated Pareto

distribution, differentiate equation (43), to obtain;

$$j'(t) = \left\{ \begin{array}{l} \frac{(\alpha+1)(1+t)^{-(\alpha+2)}}{(1+t)^{-(\alpha+1)}} \\ \frac{\alpha(1+t)^{-(\alpha+1)}(1-(1+t)^{-\alpha})^{-2}}{(1-(1+t)^{-\alpha})^{-1}} \end{array} \right\} j(t) \quad (44)$$

The condition necessary to the existence of equation is $\alpha, \theta, t > 0$.

$$j'(t) = - \left\{ \frac{\alpha+1}{1+t} + \frac{\alpha(1+t)^{-(\alpha+1)}}{1-(1+t)^{-\alpha}} \right\} j(t) \quad (45)$$

$$j'(t) = - \left\{ \frac{\alpha+1}{1+t} + \frac{j(t)}{\theta} \right\} j(t) \quad (46)$$

The first order ordinary differential equations for the reversed hazard function of the exponentiated Pareto distribution is given by;

$$\theta(t+1)j'(t) + \theta(\alpha+1)j(t) + (t+1)j^2(t) = 0 \quad (47)$$

$$j(1) = \frac{\alpha\theta 2^{-(\alpha+1)}}{1-2^{-\alpha}} = \frac{\alpha\theta 2^{-\alpha}}{2(1-2^{-\alpha})} = \frac{\alpha\theta}{2(2^\alpha - 1)} \quad (48)$$

The ODEs of all the probability functions considered can be obtained for the particular values of the distribution. Several analytic, semi-analytic and numerical methods can be applied to obtain the solutions of the respective differential equations [56-69]. Also, comparison with two or more solution methods is useful in understanding the link between ODEs and the probability distributions.

VIII. CONCLUDING REMARKS

In this work, differentiation was used to obtain some classes of ordinary differential equations for the probability density function (PDF), quantile function (QF), survival function (SF), inverse survival function (ISF), hazard function (HF) and reversed hazard function (RHF) of the exponentiated Pareto distribution. In all, the parameters that define the distribution determine the nature of the respective ODEs and the range determines the existence of the ODEs. Furthermore, the complexity of the ODEs depends largely on the values of the parameters.

ACKNOWLEDGMENT

The authors are unanimous in appreciation of financial sponsorship from Covenant University. The constructive suggestions of the reviewers are greatly appreciated.

REFERENCES

[1] R.C. Gupta, P.L. Gupta and R.D. Gupta, "Modeling failure time data by Lehman alternatives", *Comm. Stat. Theo. Meth.*, vol. 27, no. 4, pp. 887-904, 1998.
[2] S. Nadarajah, "Exponentiated Pareto distributions", *Statistics*, vol. 39, no. 3, pp. 255-260, 2005.
[3] W.M. Afify, "On estimation of the exponentiated Pareto distribution under different sample schemes", *Stat. Methodol.*, vol. 7, no. 2, pp. 77-83, 2010.
[4] M.M. Ali and J. Woo, "Estimation of tail-probability and reliability in exponentiated Pareto case", *Pak. J. Stat.*, vol. 26, no. 1, pp. 39-47, 2010.

[5] M.J. Nooghabi and H.J. Nooghabi, "Efficient estimation of pdf, CDF and rth moment for the exponentiated Pareto distribution in the presence of outliers", *Statistics*, vol. 45, no. 2, pp. 121-122, 2011.
[6] M.E. Mead, "An extended Pareto distribution", *Pak. J. Stat. Oper. Res.*, vol. 10, no. 3, pp. 313-329, 2014.
[7] M.H. Tahir, G.M. Cordeiro, A. Alzaatreh, M. Mansoor and M. Zubair, "A new Weibull-Pareto distribution: properties and applications", *Comm. Stat. Simul. Comput.*, vol. 45, no. 10, pp. 3548-3567, 2016.
[8] M. Pal, M.M. Ali and J. Woo, "Exponentiated Weibull distribution", *Statistica*, vol. 66, no. 2, pp. 139-147, 2006.
[9] G.S. Mudholkar and D.K. Srivastava, "Exponentiated Weibull family for analyzing bathtub failure-rate data", *IEEE Trans. Relia.*, vol. 42, no. 2, pp. 299-302, 1993.
[10] M.N. Nassar and F.H. Eissa, "On the exponentiated Weibull distribution", *Comm. Stat. Theo. Meth.*, vol. 32, no. 7, pp. 1317-1336, 2003.
[11] P.E. Oguntunde, A.O. Adejumo and O.S. Balogun, "Statistical properties of the exponentiated generalized inverted exponential distribution", *Appl. Math.*, vol. 4, no. 2, pp. 47-55, 2014.
[12] A.J. Lemonte and G.M. Cordeiro, "The exponentiated generalized inverse Gaussian distribution", *Stat. Prob. Lett.*, vol. 81, no. 4, pp. 506-517, 2011.
[13] A. Flaih, H. Elsalloukh, E. Mendi and M. Milanova, "The exponentiated inverted Weibull distribution", *Appl. Math. Inf. Sci.*, vol. 6, no. 2, pp. 167-171, 2012.
[14] I. Elbatal and H.Z. Muhammed, "Exponentiated generalized inverse Weibull distribution", *Appl. Math. Sci.*, vol. 8, no. 81, pp. 3997-4012, 2014.
[15] M.M. Ristić and N. Balakrishnan, "The gamma-exponentiated exponential distribution", *J. Stat. Comput. Simul.*, vol. 82, no. 8, pp. 1191-1206, 2012.
[16] H.H. Abu-Zinadah and A.S. Aloufi, "Some characterizations of the exponentiated Gompertz distribution", *Int. Math. Forum*, vol. 9, no. 30, pp. 1427-1439, 2014.
[17] H.H. Abu-Zinadah, "Six method of estimations for the shape parameter of exponentiated Gompertz distribution", *Appl. Math. Sci.*, vol. 8, no. 85-88, pp. 4349-4359, 2014.
[18] S.A. Siddiqui, S. Dwivedi, P. Dwivedi and M. Alam, "Beta exponentiated Mukherjee-Islam distribution: Mathematical study of different properties", *Global J. Pure Appl. Math.*, vol. 12, no. 1, pp. 951-964, 2016.
[19] A. Fatima and A. Roohi, "Transmuted exponentiated Pareto-I distribution", *Pak. J. Statist.*, vol. 32, no. 1, pp. 63-80, 2015.
[20] T.K. Pogány and A. Saboor, "The Gamma exponentiated exponential-Weibull distribution", *Filomat*, vol. 30, no. 12, pp. 3159-3170, 2016.
[21] S. Nadarajah and A.K. Gupta, "The exponentiated gamma distribution with application to drought data", *Calcutta Stat. Assoc. Bul.*, vol. 59, no. 1-2, pp. 29-54, 2007.
[22] S. Nadarajah, "The exponentiated Gumbel distribution with climate application", *Environmetrics*, vol. 17, no. 1, pp. 13-23, 2006.
[23] C.S. Lee and H.Y. Won, "Inference on reliability in an exponentiated uniform distribution", *J. Korean Data Info. Sci. Soc.*, vol. 17, no. 2, pp. 507-513, 2006.
[24] G.M. Cordeiro, A.E. Gomes, C.Q. da-Silva and E.M. Ortega, "The beta exponentiated Weibull distribution", *J. Stat. Comput. Simul.*, vol. 83, no. 1, pp. 114-138, 2013.
[25] S. Hashmi and A.Z. Memon, "Beta exponentiated Weibull distribution (its shape and other salient characteristics)", *Pak. J. Stat.*, vol. 32, no. 4, pp. 301-327, 2016.
[26] K. Rosaiah, R.R.L. Kantam and S. Kumar, "Reliability test plans for exponentiated log-logistic distribution", *Econ. Qual. Control*, vol. 21, no. 2, pp. 279-289, 2006.
[27] A.A. Al-Babtain, F. Merovci and I. Elbatal, "The McDonald exponentiated gamma distribution and its statistical properties", *SpringerPlus*, vol. 4, no. 1, art. 2, 2015.
[28] P.E. Oguntunde, O.A. Odetunmbi and A.O. Adejumo, "On the Exponentiated Generalized Weibull Distribution: A Generalization of the Weibull Distribution", *Indian J. Sci. Tech.*, vol. 8, no. 35, 2015.
[29] N. Feroze and I. Elbatal, "Beta exponentiated gamma distribution: some properties and estimation", *Pak. J. Stat. Oper. Res.*, vol. 12, no. 1, pp. 141-154, 2016.
[30] A.I. Shawky and R.A. Bakoban, "Exponentiated gamma distribution: Different methods of estimations", *J. Appl. Math.*, vol. 2012, art. no. 284296, 2012.

- [31] A.I. Shawky and H.H. Abu-Zinadah, Exponentiated Pareto distribution: Different method of estimations, *Int. J. Contem. Math. Sci.*, vol. 4, no. 14, pp. 677- 693, 2009.
- [32] A.J. Lemonte, W. Barreto-Souza and G.M. Cordeiro, The exponentiated Kumaraswamy distribution and its log-transform, *Braz. J. Prob. Stat.*, vol. 27, no. 1, pp. 31-53, 2013.
- [33] A.M. Sarhan and J. Apaloo, "Exponentiated modified Weibull extension distribution", *Relia. Engine. Syst. Safety*, vol. 112, pp. 137-144, 2013.
- [34] A.Z. Afify, H.M. Yousof, G.G. Hamedani and G. Aryal, "The exponentiated Weibull-Pareto distribution with application", *J. Stat. Theory Appl.*, vol. 15, pp. 328-346, 2016.
- [35] C.S. Kakde and D.T. Shirke, "On exponentiated lognormal distribution", *Int. J. Agric. Stat. Sci.*, vol. 2, pp. 319-326, 2006.
- [36] B. Singh and N. Choudhary, "The exponentiated Perks distribution", *Int. J. Syst. Assur. Engine. Magt.*, vol. 8, no. 2, pp. 468-478, 2017.
- [37] A. Al-Babtain, A.A. Fattah, A.H.N. Ahmed and F. Merovci, "The Kumaraswamy-transmuted exponentiated modified Weibull distribution", *Comm. Stat. Simul. Comput.*, vol. 46, no. 5, pp. 3812-3832, 2017.
- [38] M. Pararai, G. Warahena-Liyanaage and B.O. Oluyede, "Exponentiated power Lindley-Poisson distribution: Properties and applications", *Comm. Stat. Theo. Meth.*, vol. 46, no. 10, pp. 4726-4755, 2017.
- [39] S. Dey, D. Kumar, P.L. Ramos and F. Louzada, "Exponentiated Chen distribution: Properties and estimation", *Comm. Stat. Simul. Comput.*, To appear, 2017.
- [40] C.S. Kumar and S.H.S. Dharmaja, "The exponentiated reduced Kies distribution: Properties and applications", *Comm. Stat. Theo. Meth.*, To appear, 2017.
- [41] Y. Chung, D.K. Dey and M. Jung, "The exponentiated inverse Weibull geometric distribution", *Pak. J. Stat.*, vol. 33, no. 3, pp. 161-178.
- [42] S. Nadarajah and S.A.A. Bakar, "An exponentiated geometric distribution", *Appl. Math. Model.*, vol. 40, no. 13, pp. 6775-6784, 2016.
- [43] V. Nekoukhou, M.H. Alamatsaz and H. Bidram, "A note on exponentiated geometric distribution: Another generalization of geometric distribution", *Comm. Stat. Theo. Meth.*, vol. 45, no. 5, pp. 1575-1575, 2016.
- [44] Y. Chung and Y. Kang, "The exponentiated Weibull geometric distribution: Properties and Estimations", *Comm. Stat. Appl. Meth.*, vol. 21, no. 2, pp. 147-160, 2014.
- [45] A.A. Fattah, S. Nadarajah and A.H.N. Ahmed, "The exponentiated transmuted Weibull geometric distribution with application in survival analysis", *Comm. Stat. Simul. Comput.*, To appear, 2017.
- [46] W. Gui, "Exponentiated half logistic distribution: different estimation methods and joint confidence regions", *Comm. Stat. Simul. Comput.*, To appear, 2017.
- [47] D. Deka, B. Das and B.K. Baruah, B. "Transmuted exponentiated Gumbel distribution (TEGD) and its application to water quality data", *Pak. J. Stat. Oper. Res.*, vol. 13, no. 1, pp. 115-126, 2017.
- [48] N. Bursa and G. Ozel, "The exponentiated Kumaraswamy-power function distribution", *Hacettepe J. Math. Stat.*, vol. 46, no. 2, pp. 277-292, 2017.
- [49] N.V. Mendoza, E.M. Ortega and G.M. Cordeiro, "The exponentiated-log-logistic geometric distribution: Dual activation", *Comm. Stat. Theo. Meth.*, vol. 45, no. 13, pp. 3838-3859, 2016.
- [50] A.H. El-Bassiouny, M.A., EL-Damcese, A. Mustafa and M.S. Eliwa, "Bivariate exponentiated generalized Weibull-Gompertz distribution", *J. Appl. Prob.*, vol. 11, no. 1, pp. 25-46, 2016.
- [51] N.L. Johnson, S. Kotz and N. Balakrishnan, Continuous univariate distributions, Wiley New York. ISBN: 0-471-58495-9, 1994.
- [52] W.P. Elderton, Frequency curves and correlation, Charles and Edwin Layton. London, 1906.
- [53] H. Rinne, Location scale distributions, linear estimation and probability plotting using MATLAB, 2010.
- [54] N. Balakrishnan and C.D. Lai, Continuous bivariate distributions, 2nd edition, Springer New York, London, 2009.
- [55] N.L. Johnson, S. Kotz and N. Balakrishnan, Continuous Univariate Distributions, Volume 2. 2nd edition, Wiley, 1995.
- [56] S.O. Edeki, H.I. Okagbue, A.A. Opanuga and S.A. Adeosun, "A semi - analytical method for solutions of a certain class of second order ordinary differential equations", *Applied Mathematics*, vol. 5, no. 13, pp. 2034 - 2041, 2014.
- [57] S.O. Edeki, A.A. Opanuga and H.I. Okagbue, "On iterative techniques for numerical solutions of linear and nonlinear differential equations", *J. Math. Computational Sci.*, vol. 4, no. 4, pp. 716-727, 2014.
- [58] A.A. Opanuga, S.O. Edeki, H.I. Okagbue, G.O. Akinlabi, A.S. Osheku and B. Ajayi, "On numerical solutions of systems of ordinary differential equations by numerical-analytical method", *Appl. Math. Sciences*, vol. 8, no. 164, pp. 8199 - 8207, 2014.
- [59] S.O. Edeki, A.A. Opanuga, H.I. Okagbue, G.O. Akinlabi, S.A. Adeosun and A.S. Osheku, "A Numerical-computational technique for solving transformed Cauchy-Euler equidimensional equations of homogenous type. *Adv. Studies Theo. Physics*, vol. 9, no. 2, pp. 85 - 92, 2015.
- [60] S.O. Edeki, E.A. Owoloko, A.S. Osheku, A.A. Opanuga, H.I. Okagbue and G.O. Akinlabi, "Numerical solutions of nonlinear biochemical model using a hybrid numerical-analytical technique", *Int. J. Math. Analysis*, vol. 9, no. 8, pp. 403-416, 2015.
- [61] A.A. Opanuga, S.O. Edeki, H.I. Okagbue and G.O. Akinlabi, "Numerical solution of two-point boundary value problems via differential transform method", *Global J. Pure Appl. Math.*, vol. 11, no. 2, pp. 801-806, 2015.
- [62] A.A. Opanuga, S.O. Edeki, H.I. Okagbue and G. O. Akinlabi, "A novel approach for solving quadratic Riccati differential equations", *Int. J. Appl. Engine. Res.*, vol. 10, no. 11, pp. 29121-29126, 2015.
- [63] A.A. Opanuga, O.O. Agboola and H.I. Okagbue, "Approximate solution of multipoint boundary value problems", *J. Engine. Appl. Sci.*, vol. 10, no. 4, pp. 85-89, 2015.
- [64] A.A. Opanuga, O.O. Agboola, H.I. Okagbue and J.G. Oghonyon, "Solution of differential equations by three semi-analytical techniques", *Int. J. Appl. Engine. Res.*, vol. 10, no. 18, pp. 39168-39174, 2015.
- [65] A.A. Opanuga, H.I. Okagbue, S.O. Edeki and O.O. Agboola, "Differential transform technique for higher order boundary value problems", *Modern Appl. Sci.*, vol. 9, no. 13, pp. 224-230, 2015.
- [66] A.A. Opanuga, S.O. Edeki, H.I. Okagbue, S.A. Adeosun and M.E. Adeosun, "Some Methods of Numerical Solutions of Singular System of Transistor Circuits", *J. Comp. Theo. Nanosci.*, vol. 12, no. 10, pp. 3285-3289, 2015.
- [67] A.A. Opanuga, E.A. Owoloko, H.I. Okagbue and O.O. Agboola, "Finite difference method and Laplace transform for boundary value problems", Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering 2017, 5-7 July, 2017, London, U.K., pp. 65-69.
- [68] A.A. Opanuga, E.A. Owoloko, O.O. Agboola and H.I. Okagbue, "Application of homotopy perturbation and modified Adomian decomposition methods for higher order boundary value problems", Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering 2017, 5-7 July, 2017, London, U.K., pp. 130-134.
- [69] O.O. Agboola, A.A. Opanuga and J.A. Gbadeyan, "Solution of third order ordinary differential equations using differential transform method", *Global J. Pure Appl. Math.*, vol. 11, no. 4, pp. 2511-2516, 2015.