

Classes of Ordinary Differential Equations Obtained for the Probability Functions of Harris Extended Exponential Distribution

Hilary I. Okagbue, *Member, IAENG*, Muminu O. Adamu, Enahoro A. Owoloko,
Esivue A. Suleiman

Abstract— In this paper, the differential calculus was used to obtain some classes of ordinary differential equations (ODE) for the probability density function, quantile function, survival function, inverse survival function and hazard function of the Harris extended exponential distribution. The case of reversed hazard function was excluded because of its complexity. The stated necessary conditions required for the existence of the ODEs are consistent with the various parameters that defined the distribution. Solutions of these ODEs by using numerous available methods are new ways of understanding the nature of the probability functions that characterize the distribution. The method can be extended to other probability distributions, functions and can serve an alternative to estimation and approximation.

Index Terms— Harris, exponential, differentiation, quantile function, survival function.

I. INTRODUCTION

THIS is a lifetime probability model proposed by [1]. The distribution is as a result of mixture of Harris [2] and exponential distributions and applications of the distribution as a flexible model was shown by [3]. The distribution is a lifetime with decreasing failure rate [4-5]. Alternatively the distribution can be regarded as an extension or modification of the exponential distribution.

The aim of this paper is to develop ordinary differential equations (ODE) for the probability density function (PDF), Quantile function (QF), survival function (SF), inverse survival function (ISF) and hazard function (HF) of Harris extended exponential distribution by the use of differential calculus. Calculus is a very key tool in the determination of mode of a given probability distribution and in estimation of parameters of probability distributions, amongst other uses. The research is an extension of the ODE to other probability functions other than the PDF. Similar works done where the PDF of probability distributions was expressed as ODE whose solution is the PDF are available. They include: Laplace distribution [6], beta distribution [7], raised cosine

Manuscript received July 30, 2017. This work was sponsored by Covenant University, Ota, Nigeria.

H. I. Okagbue, E. A. Owoloko and E. A. Suleiman are with the Department of Mathematics, Covenant University, Ota, Nigeria.

hilary.okagbue@covenantuniversity.edu.ng

Alfred.owoloko@covenantuniversity.edu.ng

M. O. Adamu is with the Department of Mathematics, University of Lagos, Akoka, Nigeria.

distribution [8], Lomax distribution [9], beta prime distribution or inverted beta distribution [10].

II. PROBABILITY DENSITY FUNCTION

The probability density function of the Harris extended exponential distribution (HEE) is given as;

$$f(x) = \frac{\lambda \theta^{\frac{1}{k}} e^{-\lambda x}}{(1 - \bar{\theta} e^{-\lambda k x})^{1 + \frac{1}{k}}} \quad (1)$$

Where $\bar{\theta} = 1 - \theta$, $x, \theta, \bar{\theta}, \lambda, k > 0$

To obtain the first order ordinary differential equation for the probability density function of the HEE distribution, differentiate equation (1), to obtain;

$$f'(x) = \left\{ \frac{\lambda e^{-\lambda x}}{e^{-\lambda x}} - \frac{\bar{\theta} \lambda (k+1) e^{-\lambda k x} (1 - \bar{\theta} e^{-\lambda k x})^{-(2 + \frac{1}{k})}}{(1 - \bar{\theta} e^{-\lambda k x})^{-(1 + \frac{1}{k})}} \right\} f(x) \quad (2)$$

$$f'(x) = \left\{ -\lambda - \frac{\bar{\theta} \lambda (k+1) e^{-\lambda k x}}{(1 - \bar{\theta} e^{-\lambda k x})} \right\} f(x) \quad (3)$$

The condition necessary for the existence of equation is $x, \theta, \bar{\theta}, \lambda, k > 0$.

Differentiate equation (3);

$$f''(x) = \left\{ -\lambda - \frac{\bar{\theta} \lambda (k+1) e^{-\lambda k x}}{(1 - \bar{\theta} e^{-\lambda k x})} \right\} f'(x) + \left\{ \frac{(k+1)k \bar{\theta}^2 \lambda^2 (e^{-\lambda k x})^2}{(1 - \bar{\theta} e^{-\lambda k x})^2} + \frac{(k+1)k \bar{\theta} \lambda^2 e^{-\lambda k x}}{(1 - \bar{\theta} e^{-\lambda k x})} \right\} f(x) \quad (4)$$

The condition necessary for the existence of equation is $x, \theta, \bar{\theta}, \lambda, k > 0$.

The following equations obtained from (3) are required in the simplification of equation (4);

$$\frac{f'(x)}{f(x)} = -\lambda - \frac{\bar{\theta} \lambda (k+1) e^{-\lambda k x}}{(1 - \bar{\theta} e^{-\lambda k x})} \quad (5)$$

$$\frac{\bar{\theta}\lambda(k+1)e^{-\lambda kx}}{(1-\bar{\theta}e^{-\lambda kx})} = -\left(\frac{f'(x)}{f(x)} + \lambda\right) \quad (6)$$

$$\frac{\bar{\theta}^2\lambda^2(k+1)^2(e^{-\lambda kx})^2}{(1-\bar{\theta}e^{-\lambda kx})^2} = \left(\frac{f'(x)}{f(x)} + \lambda\right)^2 \quad (7)$$

$$\frac{(k+1)\bar{\theta}^2\lambda^2(e^{-\lambda kx})^2}{(1-\bar{\theta}e^{-\lambda kx})^2} = \frac{1}{k+1}\left(\frac{f'(x)}{f(x)} + \lambda\right)^2 \quad (8)$$

$$\frac{k(k+1)(\bar{\theta}\lambda(e^{-\lambda kx})^2)}{(1-\bar{\theta}e^{-\lambda kx})^2} = \frac{k}{k+1}\left(\frac{f'(x)}{f(x)} + \lambda\right)^2 \quad (9)$$

$$\frac{\bar{\theta}\lambda^2(k+1)e^{-\lambda kx}}{(1-\bar{\theta}e^{-\lambda kx})} = -\lambda\left(\frac{f'(x)}{f(x)} + \lambda\right) \quad (10)$$

$$\frac{\bar{\theta}\lambda^2k(k+1)e^{-\lambda kx}}{(1-\bar{\theta}e^{-\lambda kx})} = -k\lambda\left(\frac{f'(x)}{f(x)} + \lambda\right) \quad (11)$$

Substitute equations (5), (9) and (11) into equation (4);

$$f''(x) = \frac{f'^2(x)}{f(x)} + \left\{ \begin{array}{l} \frac{k}{k+1}\left(\frac{f'(x)}{f(x)} + \lambda\right)^2 \\ -k\lambda\left(\frac{f'(x)}{f(x)} + \lambda\right) \end{array} \right\} f(x) \quad (12)$$

$$f(1) = \frac{\lambda\theta^{\frac{1}{k}}e^{-\lambda}}{(1-\bar{\theta}e^{-\lambda k})^{1+\frac{1}{k}}} \quad (13)$$

$$f'(1) = -\lambda\theta^{\frac{1}{k}}e^{-\lambda} \left\{ \frac{\lambda(1-\bar{\theta}e^{-\lambda k}) + \bar{\theta}\lambda(k+1)e^{-\lambda k}}{(1-\bar{\theta}e^{-\lambda k})^{2+\frac{1}{k}}} \right\} \quad (14)$$

The second order ordinary differential equation for the probability density function of the HEE distribution can be obtained and evaluated for particular values of k and λ. When k = 1, equation (12), (13) and (14) become;

$$f''(x) = \frac{f'^2(x)}{f(x)} + \left\{ \begin{array}{l} \frac{1}{2}\left(\frac{f'(x)}{f(x)} + \lambda\right)^2 \\ -\lambda\left(\frac{f'(x)}{f(x)} + \lambda\right) \end{array} \right\} f(x) \quad (15)$$

$$f''(x) = \frac{3f'^2(x)}{2f(x)} - \frac{\lambda^2 f(x)}{2} \quad (16)$$

$$2f(x)f''(x) - 3f'^2(x) + \lambda^2 f^2(x) = 0 \quad (17)$$

$$f(1) = \frac{\lambda\theta e^{-\lambda}}{(1-\bar{\theta}e^{-\lambda})^2} \quad (18)$$

$$f'(1) = -\lambda\theta e^{-\lambda} \left\{ \frac{\lambda(1+\bar{\theta}e^{-\lambda})}{(1-\bar{\theta}e^{-\lambda})^3} \right\} \quad (19)$$

III. QUANTILE FUNCTION

The Quantile function of the Harris extended exponential distribution (HEE) is given as;

$$Q(p) = -\frac{1}{k\lambda} \ln\left(\frac{(1-p)^k}{\theta + \bar{\theta}(1-p)^k}\right) \quad (20)$$

Differentiate equation (20);

$$Q'(p) = -\frac{1}{k\lambda} \left[\frac{\theta + \bar{\theta}(1-p)^k}{(1-p)^k} \left(\frac{k\bar{\theta}(1-p)^k(1-p)^{k-1}}{(\theta + \bar{\theta}(1-p)^k)^2} - \frac{k(1-p)^{k-1}}{\theta + \bar{\theta}(1-p)^k} \right) \right] \quad (21)$$

$$Q'(p) = \frac{1}{\lambda(1-p)} - \frac{\bar{\theta}(1-p)^{k-1}}{\lambda(\theta + \bar{\theta}(1-p)^k)} \quad (22)$$

The condition necessary for the existence of equation is $\theta, \bar{\theta}, \lambda, k > 0, 0 \leq p < 1$.

The first order ordinary differential equation for the quantile function of the HEE distribution can be obtained and evaluated for particular values of all the parameters especially k, p and λ. This is summarized in the **Table 1**.

Table 1: Classes of differential equations obtained for the quantile function of the HEE distribution for different parameters.

k	λ	θ	θ̄	Ordinary differential equation
1	1	1	1	(1-p)(2-p)Q'(p)-1=0
1	1	1	2	(1-p)(3-2p)Q'(p)-1=0
1	1	2	1	(1-p)(3-p)Q'(p)-2=0
1	2	1	1	2(1-p)(2-p)Q'(p)-1=0
1	1	2	2	(1-p)(2-p)Q'(p)-1=0
1	2	2	2	2(1-p)(2-p)Q'(p)-1=0
1	2	2	1	(1-p)(3-p)Q'(p)-1=0
1	2	1	2	2(1-p)(3-2p)Q'(p)-1=0
2	1	1	1	(1-p)(1+(1-p)^2)Q'(p)-1=0
2	2	1	1	2(1-p)(1+(1-p)^2)Q'(p)-1=0
2	2	2	1	(1-p)(2+(1-p)^2)Q'(p)-1=0
2	2	2	2	(1-p)(2+2(1-p)^2)Q'(p)-1=0
2	1	2	1	(1-p)(2+(1-p)^2)Q'(p)-2=0
2	1	1	2	(1-p)(1+2(1-p)^2)Q'(p)-1=0
2	1	2	2	(1-p)(2+2(1-p)^2)Q'(p)-2=0
2	2	1	2	2(1-p)(1+2(1-p)^2)Q'(p)-1=0

In order to obtain a more simplified differential equation that will be the function of the quantile function, differentiate equation (22);

$$Q''(p) = -\frac{1}{\lambda(1-p)^2} - \frac{\bar{\theta}^2 k((1-p)^{k-1})^2}{\lambda(\theta + \bar{\theta}(1-p)^k)^2} + \frac{\bar{\theta}(k-1)(1-p)^{k-2}}{\lambda(\theta + \bar{\theta}(1-p)^k)} \quad (23)$$

The condition necessary for the existence of equation is that $\theta, \bar{\theta}, \lambda, k > 0, 0 \leq p < 1$.

The following equations obtained from (22) are needed in the simplification of (23).

$$\frac{\bar{\theta}(1-p)^{k-1}}{\lambda(\theta + \bar{\theta}(1-p)^k)} = \frac{1}{\lambda(1-p)} - Q'(p) \quad (24)$$

$$\frac{\bar{\theta}^2((1-p)^{k-1})^2}{\lambda^2(\theta + \bar{\theta}(1-p)^k)^2} = \left(\frac{1}{\lambda(1-p)} - Q'(p) \right)^2 \quad (25)$$

$$\frac{\bar{\theta}^2 k((1-p)^{k-1})^2}{\lambda(\theta + \bar{\theta}(1-p)^k)^2} = \lambda k \left(\frac{1}{\lambda(1-p)} - Q'(p) \right)^2 \quad (26)$$

$$\frac{\bar{\theta}(k-1)(1-p)^{k-1}}{\lambda(\theta + \bar{\theta}(1-p)^k)} = k-1 \left(\frac{1}{\lambda(1-p)} - Q'(p) \right) \quad (27)$$

$$\frac{\bar{\theta}(k-1)(1-p)^{k-2}}{\lambda(\theta + \bar{\theta}(1-p)^k)} = \frac{k-1}{1-p} \left(\frac{1}{\lambda(1-p)} - Q'(p) \right) \quad (28)$$

Substitute equations (26) and (28) into equation (23);

$$Q''(p) = -\frac{1}{\lambda(1-p)^2} - \lambda k \left(\frac{1}{\lambda(1-p)} - Q'(p) \right)^2 + \frac{k-1}{1-p} \left(\frac{1}{\lambda(1-p)} - Q'(p) \right) \quad (29)$$

$$Q(0) = -\frac{1}{k\lambda} \ln \left(\frac{1}{\theta + \bar{\theta}} \right) \quad (30)$$

$$Q'(0) = \frac{1}{\lambda} - \frac{\bar{\theta}}{\lambda(\theta + \bar{\theta})} = \frac{\theta}{\lambda(\theta + \bar{\theta})} \quad (31)$$

The ordinary differential equations can be obtained for the particular values of k and λ .

When k = 1, equations (29)-(31) become;

$$Q''(p) = -\frac{1}{\lambda(1-p)^2} - \lambda \left(\frac{1}{\lambda(1-p)} - Q'(p) \right)^2 \quad (32)$$

$$Q''(p) = -\frac{2}{\lambda(1-p)^2} + \frac{2Q'(p)}{(1-p)} - \lambda Q'^2(p) \quad (33)$$

$$\lambda(1-p)^2 Q''(p) + \lambda^2(1-p)^2 Q'^2(p) - 2\lambda(1-p)Q'(p) + 2 = 0 \quad (34)$$

$$Q(0) = -\frac{1}{\lambda} \ln \left(\frac{1}{\theta + \bar{\theta}} \right) \quad (35)$$

$$Q'(0) = \frac{\theta}{\lambda(\theta + \bar{\theta})} \quad (36)$$

IV. SURVIVAL FUNCTION

The survival function of the Harris extended exponential distribution (HEE) is given as;

$$S(t) = \left[\frac{\theta e^{-\lambda kt}}{1 - \bar{\theta} e^{-\lambda kt}} \right]^{\frac{1}{k}} \quad (37)$$

Differentiate equation (37);

$$S'(t) = \left\{ \begin{array}{l} \frac{\theta \lambda e^{-\lambda kt} (\theta e^{-\lambda kt})^{\frac{1}{k}-1}}{(\theta e^{-\lambda kt})^{\frac{1}{k}}} \\ - \frac{\bar{\theta} \lambda e^{-\lambda kt} (1 - \bar{\theta} e^{-\lambda kt})^{-\left(\frac{1}{k}+1\right)}}{(1 - \bar{\theta} e^{-\lambda kt})^{\frac{1}{k}}} \end{array} \right\} S(t) \quad (38)$$

$$S'(t) = - \left\{ \frac{\theta \lambda e^{-\lambda kt}}{\theta e^{-\lambda kt}} + \frac{\bar{\theta} \lambda e^{-\lambda kt}}{(1 - \bar{\theta} e^{-\lambda kt})} \right\} S(t) \quad (39)$$

$$S'(t) = - \left\{ \lambda + \frac{\bar{\theta} \lambda e^{-\lambda kt}}{(1 - \bar{\theta} e^{-\lambda kt})} \right\} S(t) \quad (40)$$

The condition necessary for the existence of equation is that $t, \theta, \bar{\theta}, \lambda, k > 0$.

Equation (37) can also be written as;

$$S^k(t) = \frac{\theta e^{-\lambda kt}}{1 - \bar{\theta} e^{-\lambda kt}} \quad (41)$$

$$\frac{S^k(t)}{\theta} = \frac{e^{-\lambda kt}}{1 - \bar{\theta} e^{-\lambda kt}} \quad (42)$$

$$\frac{\bar{\theta} S^k(t)}{\theta} = \frac{\bar{\theta} e^{-\lambda kt}}{1 - \bar{\theta} e^{-\lambda kt}} \quad (43)$$

Substitute equation (43) into equation (40);

$$S'(t) = - \left\{ \lambda + \frac{\bar{\theta} S^k(t)}{\theta} \right\} S(t) \quad (44)$$

The first order ordinary differential equation for the survival function of the HEE distribution is given by;

$$\theta S'(t) + \theta \lambda S(t) + \bar{\theta} S^{k+1}(t) = 0 \quad (45)$$

$$S(1) = \left[\frac{\theta e^{-\lambda k}}{1 - \bar{\theta} e^{-\lambda k}} \right]^{\frac{1}{k}} \quad (46)$$

The ordinary differential equations can be obtained for the particular values of k and λ .

When k = 1, equations (45) and (46) become;

$$\theta S'(t) + \theta \lambda S(t) + \bar{\theta} S^2(t) = 0 \quad (47)$$

$$S(1) = \frac{\theta e^{-\lambda}}{1 - \bar{\theta} e^{-\lambda}} = \frac{\theta}{e^{\lambda} - \bar{\theta}} \quad (48)$$

In order to obtain the second order differential equation, differentiate equation (40);

$$S''(t) = \left\{ -\lambda - \frac{\bar{\theta}\lambda e^{-\lambda kt}}{(1-\bar{\theta}e^{-\lambda kt})} \right\} S'(t) + \left\{ \frac{k(\bar{\theta}\lambda e^{-\lambda kt})^2}{(1-\bar{\theta}e^{-\lambda kt})^2} + \frac{\bar{\theta}k\lambda^2 e^{-\lambda kt}}{(1-\bar{\theta}e^{-\lambda kt})} \right\} S(t) \quad (49)$$

The condition necessary for the existence of equation is that $t, \theta, \bar{\theta}, \lambda, k > 0$.

The following equations obtained from equation (40) are needed in the simplification of equation (49);

$$\frac{S'(t)}{S(t)} = -\lambda - \frac{\bar{\theta}\lambda e^{-\lambda kt}}{(1-\bar{\theta}e^{-\lambda kt})} \quad (50)$$

$$\frac{\bar{\theta}\lambda e^{-\lambda kt}}{(1-\bar{\theta}e^{-\lambda kt})} = -\left(\frac{S'(t)}{S(t)} + \lambda \right)$$

$$\left(\frac{\bar{\theta}\lambda e^{-\lambda kt}}{(1-\bar{\theta}e^{-\lambda kt})} \right)^2 = \left(\frac{S'(t)}{S(t)} + \lambda \right)^2 \quad (51)$$

$$\frac{k(\bar{\theta}\lambda e^{-\lambda kt})^2}{(1-\bar{\theta}e^{-\lambda kt})^2} = k \left(\frac{S'(t)}{S(t)} + \lambda \right)^2 \quad (52)$$

$$\frac{k\bar{\theta}\lambda e^{-\lambda kt}}{(1-\bar{\theta}e^{-\lambda kt})} = -k \left(\frac{S'(t)}{S(t)} + \lambda \right) \quad (53)$$

$$\frac{k\bar{\theta}\lambda^2 e^{-\lambda kt}}{(1-\bar{\theta}e^{-\lambda kt})} = -k\lambda \left(\frac{S'(t)}{S(t)} + \lambda \right) \quad (54)$$

(55) Substitute equations (50), (53) and (55) into equation (49);

$$S''(t) = \frac{S'^2(t)}{S(t)} + \left\{ \begin{array}{l} k \left(\frac{S'(t)}{S(t)} + \lambda \right)^2 \\ -k\lambda \left(\frac{S'(t)}{S(t)} + \lambda \right) \end{array} \right\} S(t) \quad (56)$$

Simplify equation (56);

$$S''(t) = \frac{(k+1)S'^2(t)}{S(t)} + k\lambda S'(t) \quad (57)$$

The second order ordinary differential equation for the survival function of the HEE distribution is given by;

$$S(t)S''(t) - (k+1)S'^2(t) - k\lambda S(t)S'(t) = 0 \quad (58)$$

$$S'(1) = -\left\{ \lambda + \frac{\bar{\theta}\lambda e^{-\lambda k}}{(1-\bar{\theta}e^{-\lambda k})} \right\} S(1) = -\left(\lambda + \frac{\bar{\theta}\lambda e^{-\lambda k}}{(1-\bar{\theta}e^{-\lambda k})} \right) \left(\frac{\theta e^{-\lambda k}}{1-\bar{\theta}e^{-\lambda k}} \right)^{\frac{1}{k}} \quad (59)$$

One case of equation (58) is considered, that is when $k = 1$, equations (58), (46) and (59) become;

$$S(t)S''(t) - 2S'^2(t) - \lambda S(t)S'(t) = 0 \quad (60)$$

$$S(1) = \left(\frac{\theta e^{-\lambda}}{1-\bar{\theta}e^{-\lambda}} \right) \quad (61)$$

$$S'(1) = -\left(\lambda + \frac{\bar{\theta}\lambda e^{-\lambda}}{(1-\bar{\theta}e^{-\lambda})} \right) \left(\frac{\theta e^{-\lambda}}{1-\bar{\theta}e^{-\lambda}} \right) = -\frac{(\lambda - \bar{\theta}e^{-\lambda}(\lambda-1))\bar{\theta}e^{-\lambda}}{(1-\bar{\theta}e^{-\lambda})^2} \quad (62)$$

V. INVERSE SURVIVAL FUNCTION

The inverse survival function of the Harris extended exponential distribution (HEE) can be obtained from the survival function;

$$\left[\frac{\theta e^{-\lambda k Q(p)}}{1-\bar{\theta}e^{-\lambda k Q(p)}} \right]^{\frac{1}{k}} = p \quad (63)$$

$$\frac{\theta}{e^{\lambda k Q(p)} - \bar{\theta}} = p^k \quad (64)$$

$$e^{\lambda k Q(p)} = \frac{\theta + \bar{\theta}p^k}{p^k} \quad (65)$$

Taking logarithm;

$$\lambda k Q(p) = \ln(\theta + \bar{\theta}p^k) - \ln p^k \quad (66)$$

Differentiating;

$$\lambda k Q'(p) = \frac{k\bar{\theta}p^{k-1}}{\theta + \bar{\theta}p^k} - \frac{kp^{k-1}}{p^k} \quad (67)$$

$$\lambda Q'(p) = \frac{\bar{\theta}p^{k-1}}{\theta + \bar{\theta}p^k} - \frac{1}{p} \quad (68)$$

The condition necessary for the existence of equation (68) is that $\theta, \bar{\theta}, \lambda, k > 0, 0 \leq p < 1$.

The first order ordinary differential equation for the inverse survival function of the HEE distribution can be obtained and evaluated for particular values of all the parameters especially k, p and λ . This is summarized in the **Table 2**.

Table 2: Classes of differential equations obtained for the inverse survival function of the HEE distribution for different parameters.

k	λ	θ	$\bar{\theta}$	ordinary differential equation
1	1	1	1	$p(p+1)Q'(p) + 1 = 0$
1	1	1	2	$p(2p+1)Q'(p) + 1 = 0$
1	1	2	1	$p(p+2)Q'(p) + 2 = 0$
1	1	2	2	$p(p+1)Q'(p) + 1 = 0$
1	2	1	1	$2p(p+1)Q'(p) + 1 = 0$
1	2	1	2	$2p(2p+1)Q'(p) + 1 = 0$
1	2	2	1	$p(p+2)Q'(p) + 1 = 0$
1	2	2	2	$p(p+1)Q'(p) + 1 = 0$
2	1	1	1	$p(p^2+1)Q'(p) + 1 = 0$
2	1	1	2	$p(2p^2+1)Q'(p) + 1 = 0$

Table 2 continued

k	λ	θ	$\bar{\theta}$	ordinary differential equation
2	1	2	1	$p(p^2 + 2)Q'(p) + 2 = 0$
2	1	2	2	$p(p^2 + 1)Q'(p) + 1 = 0$
2	2	1	1	$2p(p^2 + 1)Q'(p) + 1 = 0$
2	2	1	2	$2p(2p^2 + 1)Q'(p) + 1 = 0$
2	2	2	1	$p(p^2 + 2)Q'(p) + 2 = 0$
2	2	2	2	$p(p^2 + 1)Q'(p) + 1 = 0$

VI. HAZARD FUNCTION

The hazard function of the Harris extended exponential distribution (HEE) is given as;

$$h(t) = \frac{\lambda}{1 - \bar{\theta}e^{-\lambda kt}} \tag{69}$$

Differentiate equation (69);

$$h'(t) = -\frac{\bar{\theta}\lambda^2 k e^{-\lambda kt}}{(1 - \bar{\theta}e^{-\lambda kt})^2} \tag{70}$$

The condition necessary for the existence of equation (70) is $t, \theta, \bar{\theta}, \lambda, k > 0$.

Use equation (69) in (70);

$$h'(t) = -\bar{\theta}k e^{-\lambda kt} h^2(t) \tag{71}$$

Equation (69) can also be written as;

$$(1 - \bar{\theta}e^{-\lambda kt})h(t) = \lambda \tag{72}$$

$$\bar{\theta}e^{-\lambda kt} = \frac{h(t) - \lambda}{h(t)} \tag{73}$$

Substitute equation (73) into equation (71);

$$h'(t) = -k(h(t) - \lambda)h(t) \tag{74}$$

The first order ordinary differential equation for the hazard function of the HEE distribution is given by;

$$h'(t) + kh^2(t) - k\lambda h(t) = 0 \tag{75}$$

$$h(1) = \frac{\lambda}{1 - \bar{\theta}e^{-\lambda k}} \tag{76}$$

One case of equation (75) is considered, that is when $k = 1$, equations (75) and (76) become;

$$h'(t) + h^2(t) - \lambda h(t) = 0 \tag{77}$$

$$h(1) = \frac{\lambda}{1 - \bar{\theta}e^{-\lambda}} \tag{78}$$

The ODEs of all the probability functions considered can be obtained for the particular values of the distribution. Several analytic, semi-analytic and numerical methods can be applied to obtain the solutions of the respective differential equations. Also comparison with two or more solution methods is useful in understanding the link between ODEs and the probability distributions. Several semi-analytical methods can be applied to solve the ODEs for a given parameter of chosen probability function. The ODEs can be

solved using any of the semi-analytical methods as applied in [11-26].

VII. CONCLUDING REMARKS

In this work, differentiation was used to obtain some classes of ordinary differential equations for the probability density function (PDF), quantile function (QF), survival function (SF), inverse survival function (ISF) and hazard function (HF) of the Harris extended exponential distribution. The result of the reversed hazard function (RHF) was not obtained because of its complexity. The work was simplified by the application of simple algebraic procedures. The inverse survival function and quantile function generated several ODEs based on their different parameters. In all, the parameters that define the distribution determine the nature of the respective ODEs and the range determines the existence of the ODEs.

ACKNOWLEDGMENT

The authors are unanimous in appreciation of financial sponsorship from Covenant University. The constructive suggestions of the reviewers are greatly appreciated.

REFERENCES

- [1] L.G.B. Pinho, G.M. Cordeiro and J.S. Nohre, "The Harris extended exponential distribution", *Comm. Stat. Theo. Meth.*, vol. 44, no. 16, pp. 3486-3502, 2015.
- [2] T.E. Harris, "Branching processes", *Ann. Math. Stat.*, vol. 19, pp. 474-494, 1948.
- [3] L.G.B. Pinho, G.M. Cordeiro and J.S. Nohre, "On the Harris-G class of distributions: General results and application", *Braz. J. Prob. Stat.*, vol. 29, no. 4, pp. 813-832, 2015.
- [4] W. Barreto-Souza and H.S. Bakouch, "A new lifetime model with decreasing failure rate", *Statistics*, vol. 47, no. 2, pp. 465-476, 2013.
- [5] M. Chahkandi and M.Ganjali, "On some lifetime distributions with decreasing failure rate", *Comput. Stat. Data Analy.*, vol. 53, no. 12, pp. 4433-4440, 2009.
- [6] N.L. Johnson, S. Kotz and N. Balakrishnan, *Continuous univariate distributions*, Wiley New York. ISBN: 0-471-58495-9, 1994.
- [7] W.P. Elderton, *Frequency curves and correlation*, Charles and Edwin Layton. London, 1906.
- [8] H. Rinne, *Location scale distributions, linear estimation and probability plotting using MATLAB*, 2010.
- [9] N. Balakrishnan and C.D. Lai, *Continuous bivariate distributions*, 2nd edition, Springer New York, London, 2009.
- [10] N.L. Johnson, S. Kotz and N. Balakrishnan, *Continuous Univariate Distributions*, Volume 2. 2nd edition, Wiley, 1995.
- [11] S.O. Edeki, H.I. Okagbue, A.A. Opanuga and S.A. Adeosun, "A semi - analytical method for solutions of a certain class of second order ordinary differential equations", *Applied Mathematics*, vol. 5, no. 13, pp. 2034 - 2041, 2014.
- [12] S.O. Edeki, A.A Opanuga and H.I Okagbue, "On iterative techniques for numerical solutions of linear and nonlinear differential equations", *J. Math. Computational Sci.*, vol. 4, no. 4, pp. 716-727, 2014.
- [13] A.A. Opanuga, S.O. Edeki, H.I. Okagbue, G.O. Akinlabi, A.S. Osheku and B. Ajayi, "On numerical solutions of systems of ordinary differential equations by numerical-analytical method", *Appl. Math. Sciences*, vol. 8, no. 164, pp. 8199 - 8207, 2014.
- [14] S.O. Edeki, A.A. Opanuga, H.I. Okagbue, G.O. Akinlabi, S.A. Adeosun and A.S. Osheku, "A Numerical-computational technique for solving transformed Cauchy-Euler equidimensional equations of homogenous type. *Adv. Studies Theo. Physics*, vol. 9, no. 2, pp. 85 - 92, 2015.
- [15] S.O. Edeki, E.A. Owoloko, A.S. Osheku, A.A. Opanuga, H.I. Okagbue and G.O. Akinlabi, "Numerical solutions of nonlinear biochemical model using a hybrid numerical-analytical technique", *Int. J. Math. Analysis*, vol. 9, no. 8, pp. 403-416, 2015.
- [16] A.A. Opanuga, S.O. Edeki, H.I. Okagbue and G.O. Akinlabi, "Numerical solution of two-point boundary value problems via

- differential transform method”, *Global J. Pure Appl. Math.*, vol. 11, no. 2, pp. 801-806, 2015.
- [17] A.A. Opanuga, S.O. Edeki, H.I. Okagbue and G. O. Akinlabi, “A novel approach for solving quadratic Riccati differential equations”, *Int. J. Appl. Engine. Res.*, vol. 10, no. 11, pp. 29121-29126, 2015.
- [18] A.A Opanuga, O.O. Agboola and H.I. Okagbue, “Approximate solution of multipoint boundary value problems”, *J. Engine. Appl. Sci.*, vol. 10, no. 4, pp. 85-89, 2015.
- [19] A.A. Opanuga, O.O. Agboola, H.I. Okagbue and J.G. Oghonyon, “Solution of differential equations by three semi-analytical techniques”, *Int. J. Appl. Engine. Res.*, vol. 10, no. 18, pp. 39168-39174, 2015.
- [20] A.A. Opanuga, H.I. Okagbue, S.O. Edeki and O.O. Agboola, “Differential transform technique for higher order boundary value problems”, *Modern Appl. Sci.*, vol. 9, no. 13, pp. 224-230, 2015.
- [21] A.A. Opanuga, S.O. Edeki, H.I. Okagbue, S.A. Adeosun and M.E. Adeosun, “Some Methods of Numerical Solutions of Singular System of Transistor Circuits”, *J. Comp. Theo. Nanosci.*, vol. 12, no. 10, pp. 3285-3289, 2015.
- [22] O.O. Agboola, A.A. Opanuga and J.A. Gbadeyan, “Solution of third order ordinary differential equations using differential transform method”, *Global J. Pure Appl. Math.*, vol. 11, no. 4, pp. 2511-2516, 2015.
- [23] A.A. Opanuga, H.I. Okagbue, O.O. Agboola, "Irreversibility Analysis of a Radiative MHD Poiseuille Flow through Porous Medium with Slip Condition," *Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering 2017, 5-7 July, 2017, London, U.K.*, pp. 167-171.
- [24] A.A. Opanuga, E.A. Owoloko, O.O. Agboola, H.I. Okagbue, "Application of Homotopy Perturbation and Modified Adomian Decomposition Methods for Higher Order Boundary Value Problems," *Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering 2017, 5-7 July, 2017, London, U.K.*, pp. 130-134.
- [25] A.A. Opanuga, E.A. Owoloko, H. I. Okagbue, O.O. Agboola, "Finite Difference Method and Laplace Transform for Boundary Value Problems," *Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering 2017, 5-7 July, 2017, London, U.K.*, pp. 65-69.
- [26] A.A. Opanuga, E.A. Owoloko, H.I. Okagbue, “Comparison Homotopy Perturbation and Adomian Decomposition Techniques for Parabolic Equations,” *Lecture Notes in Engineering and Computer Science: Proceedings of The World Congress on Engineering 2017, 5-7 July, 2017, London, U.K.*, pp. 24-27.