

# Draft Genome Sequence of the Commercial Biocontrol Strain *Pantoea agglomerans* P10c

Theo H. M. Smits,<sup>a</sup> Fabio Rezzonico,<sup>a</sup> Jochen Blom,<sup>b</sup> Alexander Goesmann,<sup>b</sup> Azzurra Abelli,<sup>c</sup> Roberto Kron Morelli,<sup>c</sup> Joël L. Vanneste,<sup>d</sup> Brion Duffy<sup>a</sup>

Environmental Genomics and Systems Biology Research Group, Institute for Natural Resource Sciences, Zürich University of Applied Sciences ZHAW, Wädenswil, Switzerland<sup>a</sup>; Bioinformatics and Systems Biology, Justus-Liebig-Universität, Giessen, Germany<sup>b</sup>; Agrifutur, Alfianello (Brescia), Italy<sup>c</sup>; The New Zealand Institute for Plant & Food Research Limited, Ruakura, Hamilton, New Zealand<sup>d</sup>

**We report here the draft genome sequence of the biocontrol strain *Pantoea agglomerans* P10c, composed of a draft chromosome and two plasmids: the 559-kb large *Pantoea* plasmid 1 (pPag3) and a 182-kb plasmid (pPag1). A genomic island containing pantocin A biosynthesis genes was identified.**

Received 19 October 2015 Accepted 23 October 2015 Published 10 December 2015

**Citation** Smits THM, Rezzonico F, Blom J, Goesmann A, Abelli A, Kron Morelli R, Vanneste JL, Duffy B. 2015. Draft genome sequence of the commercial biocontrol strain *Pantoea agglomerans* P10c. *Genome Announc* 3(6):e01448-15. doi:10.1128/genomeA.01448-15.

**Copyright** © 2015 Smits et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Theo H. M. Smits, [theo.smits@zhaw.ch](mailto:theo.smits@zhaw.ch).

*Pantoea agglomerans* is a ubiquitous bacterial species often found in association with plants. Several strains of *P. agglomerans* have been found to control fungal or bacterial postharvest diseases (1, 2) and to reduce disease incidence in the orchards and fields, for example, basal kernel blight of barley caused by *Pseudomonas syringae* pv. *syringae* (3) or fire blight caused by *Erwinia amylovora* (2, 4, 5). *P. agglomerans* is one of the most common bacteria isolated from fire blight-susceptible plant species (4). The mechanisms by which *P. agglomerans* suppresses plant pathogens include nutritional competition and preemptive exclusion in combination with a variety of antibacterial organic acids and peptide antibiotics (5–9).

*P. agglomerans* P10c was isolated from pear blossoms in New Zealand. Selected for its exceptional ability to rapidly colonize apple and pear flowers and to suppress fire blight disease (10), it has been developed as a commercial biocontrol agent (Blossom Bless; Agrinova NZ Limited). Whole-genome sequencing (Illumina MiSeq 2 × 300-bp shotgun sequencing) yielded 2,693,476 reads representing ~160× genome coverage. The genomes were assembled using a combination of *de novo* assembly with NGen version 4 (DNASTar, Madison, WI) and comparisons with available *P. agglomerans* genomes (11–13) using Mauve version 2.3.1 (14). The final assembly consists of a draft chromosome (16 contigs for a total 4,034,977 bp) and two closed plasmids, pPag1 (182,134 bp) and pPag3 (558,805 bp). All sequences were annotated automatically using GenDB (15), with manual optimization (16).

The plasmids pPag1 and pPag3 are relatives of the respective *Pantoea vagans* C9-1 plasmids (16). Plasmid pPag3 is a group I member of the large *Pantoea* plasmid 1 family (LPP-1) of plasmids, most closely related to the plasmids of other sequenced *P. agglomerans* strains (11). Plasmid pPag1 is present in all *P. agglomerans* genomes studied to date but also shows some variation between strains.

Comparative analysis using EDGAR (17) confirmed that the genomes of *P. agglomerans* strains are highly collinear. A distinct

genomic island integrated in the *mutS* N-terminus was identified containing the pantocin A biosynthetic genes (18), one potential mechanism of action in biocontrol. Overall, this genomic island is related to the pantocin A genomic island of *P. vagans* C9-1, with slight variations (16).

The genome sequence of *P. agglomerans* P10c will give us the opportunity to improve strain-specific fingerprinting important for registration and intellectual property protection of active biological agents in commercial biocontrol products. Moreover, this provides a foundation to elucidate mechanisms of action and the regulation of antimicrobial compound biosynthesis, which in turn can be exploited to improve practical and commercial value through the optimization of formulation technology, performance reliability, and efficacy of this biocontrol strain.

**Nucleotide sequence accession numbers.** The whole-genome shotgun project of *P. agglomerans* P10c has been deposited at DDBJ/EMBL/GenBank under the accession no. LIME00000000. The version described in this paper is version LIME01000000. The finished plasmid sequences received accession numbers LIME01000017 (P10c pPag1) and LIME01000018 (P10c pPag3).

## ACKNOWLEDGMENTS

This genome project was supported by the European Union's Seventh Framework Programme for Research, Technological Development and Demonstration under grant agreement no. 613678 (DROPSA), the EUPHRESO Era-Net pilot project "PhytFire," and the Department of Life Sciences and Facility Management of ZHAW.

## REFERENCES

- Bonaterra A, Mari M, Casalini L, Montesinos E. 2003. Biological control of *Monilinia laxa* and *Rhizopus stolonifer* in postharvest of stone fruit by *Pantoea agglomerans* EPS125 and putative mechanisms of antagonism. *Int J Food Microbiol* 84:93–104. [http://dx.doi.org/10.1016/S0168-1605\(02\)00403-8](http://dx.doi.org/10.1016/S0168-1605(02)00403-8).
- Vanneste JL, Cornish DA, Yu J, Voyle MD. 1998. A microcin produced by a strain of *Erwinia herbicola* is involved in biological control of fire

- blight and soft rot caused by *Erwinia* sp. *Acta Hort* 513:39–46. <http://dx.doi.org/10.17660/ActaHortic.1998.513.3>.
3. Braun-Kiewnick A, Jacobsen BJ, Sands DC. 2000. Biological control of *Pseudomonas syringae* pv. *syringae*, the causal agent of basal kernel blight of barley, by antagonistic *Pantoea agglomerans*. *Phytopathology* 90: 368–375. <http://dx.doi.org/10.1094/PHYTO.2000.90.4.368>.
  4. Johnson KB, Stockwell VO. 1998. Management of fire blight: a case study in microbial ecology. *Annu Rev Phytopathol* 36:227–248. <http://dx.doi.org/10.1146/annurev.phyto.36.1.227>.
  5. Vanneste JL. 1996. Honey bees and epiphytic bacteria to control fire blight, a bacterial disease of apple and pear. *Biocontrol News Inform* 17: 67N–78N.
  6. Vanneste JL, Yu J, Beer SV. 1992. Role of antibiotic production by *Erwinia herbicola* Eh252 in biological control of *Erwinia amylovora*. *J Bacteriol* 174:2785–2796.
  7. Stockwell VO, Johnson KB, Sugar D, Loper JE. 2010. Control of fire blight by *Pseudomonas fluorescens* A506 and *Pantoea vagans* C9-1 applied as single strains and mixed inocula. *Phytopathology* 100:1330–1339. <http://dx.doi.org/10.1094/PHYTO-03-10-0097>.
  8. Wilson M, Lindow SE. 1994. Coexistence among epiphytic bacterial populations mediated through nutritional resource partitioning. *Appl Environ Microbiol* 60:4468–4477.
  9. Stockwell VO, Johnson KB, Sugar D, Loper JE. 2002. Antibiosis contributes to biological control of fire blight by *Pantoea agglomerans* strain Eh252 in orchards. *Phytopathology* 92:1202–1209. <http://dx.doi.org/10.1094/PHYTO.2002.92.11.1202>.
  10. Vanneste JL, Cornish DA, Yu J, Voyle MD. 2002. P10c: a new biological control agent for control of fire blight which can be sprayed or distributed using honey bees. *Acta Hort* 590:231–236. <http://dx.doi.org/10.17660/ActaHortic.2002.590.33>.
  11. De Maayer P, Chan W, Blom J, Venter SN, Duffy B, Smits THM, Coutinho TA. 2012. The large universal *Pantoea* plasmid LPP-1 plays a major role in biological and ecological diversification. *BMC Genomics* 13:625. <http://dx.doi.org/10.1186/1471-2164-13-625>.
  12. Matsuzawa T, Mori K, Kadowaki T, Shimada M, Tashiro K, Kuhara S, Inagawa H, Soma G-I, Takegawa K. 2012. Genome sequence of *Pantoea agglomerans* IG1. *J Bacteriol* 194:1258–1259. <http://dx.doi.org/10.1128/JB.06674-11>.
  13. Smith DDN, Kirzinger MWB, Stavriniades J. 2013. Draft genome sequence of the antibiotic-producing cystic fibrosis isolate *Pantoea agglomerans* Tx10. *Genome Announc* 1(5):e00904-13. <http://dx.doi.org/10.1128/genomeA.00904-13>.
  14. Darling ACE, Mau B, Blattner FR, Perna NT. 2004. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res* 14:1394–1403. <http://dx.doi.org/10.1101/gr.2289704>.
  15. Meyer F, Goesmann A, McHardy AC, Bartels D, Bekel T, Clausen J, Kalinowski J, Linke B, Rupp O, Giegerich R, Pühler A. 2003. GenDB—an open source genome annotation system for prokaryote genomes. *Nucleic Acids Res* 31:2187–2195. <http://dx.doi.org/10.1093/nar/kg312>.
  16. Smits THM, Rezzonico F, Kamber T, Goesmann A, Ishimaru CA, Frey JE, Stockwell VO, Duffy B. 2011. Metabolic versatility and antibacterial metabolite biosynthesis are distinguishing genomic features of the fire blight antagonist *Pantoea vagans* C9-1. *PLoS One* 6:e22247. <http://dx.doi.org/10.1371/journal.pone.0022247>.
  17. Blom J, Albaum SP, Doppmeier D, Pühler A, Vorhölter F, Zakrzewski M, Goesmann A. 2009. Edgar: a software framework for the comparative analysis of prokaryotic genomes. *BMC Bioinformatics* 10:154. <http://dx.doi.org/10.1186/1471-2105-10-154>.
  18. Jin M, Liu L, Wright SAI, Beer SV, Clardy J. 2003. Structural and functional analysis of pantocin A: an antibiotic from *Pantoea agglomerans* discovered by heterologous expression of cloned genes. *Angew Chem Int Ed* 42:2898–2901. <http://dx.doi.org/10.1002/anie.200351053>.