

5-10-2014

A Thermal Window for Yawning in Humans: Yawning as a Brain Cooling Mechanism

Jorg J. M. Massen
University of Vienna

Kim Dusch
University of Vienna

Omar T. Eldakar
Nova Southeastern University, oeldakar@nova.edu

Andrew C. Gallup
State University of New York

Follow this and additional works at: https://nsuworks.nova.edu/cnso_bio_facarticles

 Part of the [Biology Commons](#)

NSUWorks Citation

Massen, Jorg J. M.; Kim Dusch; Omar T. Eldakar; and Andrew C. Gallup. 2014. "A Thermal Window for Yawning in Humans: Yawning as a Brain Cooling Mechanism." *Physiology & Behavior* 130, (): 145-148. doi:10.1016/j.physbeh.2014.03.032.

This Article is brought to you for free and open access by the Department of Biological Sciences at NSUWorks. It has been accepted for inclusion in Biology Faculty Articles by an authorized administrator of NSUWorks. For more information, please contact nsuworks@nova.edu.



A thermal window for yawning in humans: Yawning as a brain cooling mechanism



Jorg J.M. Massen^{a,*}, Kim Dusch^b, Omar Tonsi Eldakar^c, Andrew C. Gallup^{d,*}

^a Department of Cognitive Biology, University of Vienna, Vienna, Austria

^b Department of Education, University of Vienna, Vienna, Austria

^c Farquhar College of Arts and Sciences, Nova Southeastern University, Ft. Lauderdale, USA

^d Psychology Department, SUNY College at Oneonta, USA

HIGHLIGHTS

- The thermoregulatory theory of yawning posits that yawns function in brain cooling.
- Yawning is constrained to an optimal thermal zone of ambient temperature.
- This theory explains basic features of both spontaneous and contagious yawning.
- Applications include improved treatment of patients with thermoregulatory problems.

ARTICLE INFO

Article history:

Received 15 November 2013

Received in revised form 24 February 2014

Accepted 31 March 2014

Available online 12 April 2014

Keywords:

Yawning

Contagious yawning

Thermoregulation

Ambient temperature

ABSTRACT

The thermoregulatory theory of yawning posits that yawns function to cool the brain in part due to counter-current heat exchange with the deep inhalation of ambient air. Consequently, yawning should be constrained to an optimal thermal zone or range of temperature, i.e., a thermal window, in which we should expect a lower frequency at extreme temperatures. Previous research shows that yawn frequency diminishes as ambient temperatures rise and approach body temperature, but a lower bound to the thermal window has not been demonstrated. To test this, a total of 120 pedestrians were sampled for susceptibility to self-reported yawn contagion during distinct temperature ranges and seasons (winter: 1.4 °C, n = 60; summer: 19.4 °C, n = 60). As predicted, the proportion of pedestrians reporting yawning was significantly lower during winter than in summer (18.3% vs. 41.7%), with temperature being the only significant predictor of these differences across seasons. The underlying mechanism for yawning in humans, both spontaneous and contagious, appears to be involved in brain thermoregulation.

© 2014 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/3.0/>).

1. Introduction

Yawning occurs with an average duration of 4 to 7 s, and consists of three distinct phases: an active gaping of the jaw with inspiration, a brief period of acme corresponding with apnea and peak muscle contraction, and a passive closure of the jaw with shorter expiration [1]. In humans [2], as well as a handful of other social vertebrates [3–7], yawning can be categorized into two basic forms: spontaneous and contagious. Both forms include similar motor action patterns, but spontaneous yawns seem to be triggered by physiological mechanisms of homeostasis and arousal since they reliably occur during distinct behavioral contexts [8,9] and follow a consistent circadian pattern [10]. In contrast, contagious yawns are elicited simply by sensing or even

thinking about the action in others [11]. Unlike its spontaneous form, which appears evolutionarily older by its observed presence in all classes of vertebrates [12] and early onset in uterine development [13], contagious yawning appears to be a more recently derived behavior as evidenced by its presence in relatively few highly social species [2–7] and delayed ontogeny [14–18]. Research investigating contagious yawning has emphasized the influence of interpersonal and emotional-cognitive variables on its expression [4,5,19–28], but there have been few attempts to combine theoretical frameworks when explaining both contagious and spontaneous effects. Due to the potential multifunctionality of yawning across species [12,29], however, recent reports on social primates have highlighted potentially important differences in yawn morphology or intensity [5,30,31].

Although it is commonly believed that yawns serve a respiratory function, experimental procedures have shown yawn frequency is independent of brain/blood levels of O₂ and CO₂ [32]. A more recent theory, which posits that the motor action of yawning functions as a brain

* Corresponding authors.

E-mail addresses: jorgmassen@gmail.com (J.J.M. Massen), a.c.gallup@gmail.com (A.C. Gallup).




Fig. 1. (a) The proportion of participants reporting yawning, and (b) the mean \pm s.e.m. frequency of reported yawns in the two seasonal conditions in Vienna, Austria (light gray bars), as well as the conditions of an earlier study in Tucson, Arizona USA (dark gray bars). Average temperatures and sample sizes for each are in bold. The best-fit lines demonstrate a non-linear relationship, with (a) probability of yawning and (b) yawn frequency dropping at extreme ambient temperatures.

cooling mechanism [33,34], has received growing empirical support [reviewed by 35]. For example, research on both rats and humans shows that yawning is preceded by intermittent rises in brain temperature and localized mild hyperthermia and then followed by equivalent decreases in temperature immediately thereafter [36,37]. While various critiques have been proposed regarding the thermoregulatory theory [38–42], no study has found evidence contrary to its main predictions and all current arguments remain untenable [35,43].

According to the thermoregulatory theory, the cooling effects of yawns occur through thermoregulatory mechanisms of counter-current heat exchange, evaporative cooling and enhanced cerebral blood flow [44]. Consequently, the effectiveness of yawning is dependent on the ambient air temperature, and the expression of this behavior should be constrained to an optimal thermal zone or range of temperature, i.e., a thermal window. In particular, this theory posits that yawns should (1) increase in frequency with initial rises in ambient temperature, as this stimulates thermoregulatory mechanisms to control temperatures within a normal range, (2) decrease as ambient temperatures draw near or exceed body temperature, since taking a deep inhalation of air above one's body temperature would be counter-productive, and likewise (3) diminish when temperatures fall below a certain point, because thermoregulatory cooling responses are no longer necessary and countercurrent heat exchange could result in deviations below optimal thermal homeostasis. Since both spontaneous and contagious yawns are indistinguishable, aside from different triggers, the predictions of the thermal window hypothesis should apply to both forms.

Experimental and observational research reports of spontaneous yawning in non-human primates [9,45], birds [46,47], and rats [48] have confirmed the first two predictions of this model. Additionally, it

Table 1

Descriptive statistics: mean and s.d. for each variable between and across conditions. Independent t-test and χ^2 comparisons are between winter and summer conditions.

	Total	Winter	Summer	Test statistic	<i>p</i>
Sex (m:f)	48:72	25:35	23:37	$\chi^2 = 0.04$	0.852
Age (year)	28.6 ± 7.3	28.2 ± 7.3	29.0 ± 7.3	$t = -0.57$	0.567
Temp. (°C)	10.4 ± 10.1	1.4 ± 2.7	19.4 ± 5.9	$t = -21.42$	<0.001
Humidity (%)	54.8 ± 15.4	62.5 ± 16.0	47.0 ± 10.0	$t = 6.35$	<0.001
Time (min) ^a	63.8 ± 83.0	69.0 ± 74.6	59.8 ± 89.5	$t = 0.55$	0.586
Sleep (h)	7.1 ± 1.5	7.2 ± 1.5	7.1 ± 1.4	$t = 0.44$	0.661

^a Time represents the time spent outside prior to participating.

was recently discovered that self-reported contagious yawning frequency in humans varies with seasonal climate variation [49]. In particular, two independent groups of pedestrians were sampled in an arid desert climate (Tucson, AZ, USA): the first in summer (37 °C) and the other during 'winter' (22 °C). Contagious yawning frequency was significantly lower during the hot summer climate (24% vs. 45%), with temperature being the only significant factor contributing to this response after controlling for other variables, such as humidity, sleep and time spent outside.

Here we tested the lower bound of the thermal window hypothesis by investigating the frequency of self-reported contagious yawning in a climate with a colder winter season (Vienna, Austria). In this case the summer condition provided temperatures equivalent to those in winter months of Tucson, while the winter condition included temperatures at and slightly below freezing.

2. Methods

2.1. Participants

Participants were 120 random pedestrians recruited in and around the city of Vienna, Austria (Lat.: 48.21; Lon.: 16.37). The experiments were conducted during two distinct time frames: December 2012–March 2013 (winter; average temperature: 1.4 °C) and June 2013–October 2013 (summer; average temperature: 19.4 °C). In total, per season we recruited 60 participants (winter: 25 males, 35 females; summer: 23 males, 37 females). Participants were all over 18 years of

Table 2

Best-fitting models (GLMMs) showing the factors influencing a) whether an individual reported yawning (binomial distribution, logit link function) ($n = 120$) and b) how often they yawned ($n = 120$). Original models included sex, season (winter or summer), age, temperature, humidity, time spent outside and hours of sleep and all 2-way interactions between these variables.

Variable	denom. df	Beta \pm s.e.m.	<i>F</i>	<i>p</i>
<i>a) Dependent variable is yawn (y/n)</i>				
Temperature	61	0.078 ± 0.02	13.09	0.001
<i>b) Dependent variable is number of yawns</i>				
Temperature	116	0.113 ± 0.04	7.25	0.008
Age	116	-0.053 ± 0.03	4.09	0.045
Winter or summer	116	1.186 ± 0.84	1.98	0.162

age and gave verbal and written consent to participate in this study (see ESM). The Ethics Committee of the University of Vienna approved this research.

2.2. Procedure

The procedure was similar to [49]: an experimenter approached pedestrians in public places and asked them to participate in a survey about contagious yawning. The participants then were instructed to carefully look through a series of 18 images of people yawning, after which they took a small survey self-reporting on whether and how often they had yawned during the experiment [for validity in this approach, see 50] and if not whether they had the urge to do so, how long they had been outside before participating in the study, how long they slept the night before, and how old they were. The last item was included to replicate a recent effect of age-related declines in contagious yawning [51]. To further trigger contagious yawns, additional questions were included for participants to report on their yawning behavior outside of the study (see ESM).

While the participants were taking the survey, an experimenter recorded the relative humidity (%) and ambient air temperature (°C) using a digital thermometer/hygrometer (TFA®) that was placed in the shade. Experimenters avoided directing their attention towards the participants during the experiment, since research suggests that people are less likely to yawn when they are being observed [52]. To avoid potential effects of circadian rhythms, all surveys were conducted between 13:00 h and 15:00 h [49].

2.3. Analysis

We compared variables across season using independent t-tests, Mann–Whitney U tests (non-normally distributed scale data), or Chi-square tests with Yates correction (binomial data). To assess the influence of several variables on reported yawning in parallel, we ran General Linear Mixed Models (GLMMs). If our response variable was binomial (yawn: yes/no) we ran GLMMs with a binomial distribution and logit link function. The sex of the participant was entered as a fixed factor and age, temperature, humidity, time spent outside and hours of sleep as fixed covariates. In addition, we included all 2-way interactions of these variables. To achieve the best models we used a backward stepwise approach, and our model choices were based on comparisons of the corrected Akaike Information Criteria (cAIC). For reasons of clarity, here we present only the best fitting models. All analyses were performed in IBM SPSS Statistics v.20 for Mac OS, with α set to 0.05. All reported *p*-values are 2-tailed.

3. Results

Of the 120 participants, 36 (30.0%) reported yawning during the experiment. As predicted, there was a significant difference in self-reported yawn contagion between the two seasons, with only 11 out of 60 (18.3%) participants in winter reporting yawning, and 25 out of 60 (41.7%) participants in summer reporting yawning ($\text{Chi}^2 = 6.71$, $p = 0.010$) (Fig. 1a). Further analyses showed that participants also reported more total yawns during the summer (Mann–Whitney U = 2203, $n = 120$, $p = 0.009$) (Fig. 1b).

As several variables differed between the two seasons (Table 1), we ran GLMMs to assess which variables best predicted self-reported yawning. The best fitting model revealed that temperature was the only significant predictor, with an increased likelihood to report a yawn at higher temperatures (Table 2a); i.e., none of the other variables (sex, season, age, humidity, time spent outside and hours of sleep the night before) had a significant effect on the likelihood of reporting a yawn, and were therefore excluded from the best fitting model (Table 2a).

Similarly, we found that temperature was a significant predictor of reported yawn frequency (Table 2b), with a greater number of yawns

being reported at higher temperatures. Consistent with previous research [51], age also had a significantly negative effect on reported yawn frequency. Season was also included in the best fitting model, albeit as a non-significant effect. None of the other variables had a significant effect on the reported yawn frequency and were therefore excluded from the best fitting model (Table 2b).

4. Discussion

Overall, these results show that significantly fewer pedestrians reported contagious yawning during the cold winter (−4 to 7 °C), and that, similar to effects observed in an arid desert climate [49], temperature was the only significant predictor of this response when controlling for other variables. As predicted by the thermal window hypothesis, reports of yawning were constrained to an optimal thermal zone or range of ambient temperature (Fig. 1). Importantly, changes in daylight across the seasons cannot account for these results. First, a particular time frame was chosen for both studies (between 1 and 3 pm) whereby contagious yawning frequencies remain unchanged [53]. Second, the proportion of people that reported yawning in the summer in Vienna, Austria (current study) was comparable to that of the winter in Tucson, Arizona, USA [49], whereas there is a large difference in daylight hours between these samples (summer in Vienna: ± 16 h vs. winter in Tucson: ± 10 h). Lastly, an inverse seasonal pattern emerged between the two study locations; i.e., whereas in Tucson there was a high frequency of reported yawning in winter, and a low frequency of reported yawning in summer, in Vienna there was a high frequency of reported yawning in summer, and a low frequency of reported yawning in winter. Thus, it cannot be generalized that people yawn more or less in winter vs. summer, nor with greater or fewer hours of daylight. Instead, the ambient air temperature accompanying the season appears to determine reported yawn frequency.

This report adds to accumulating research suggesting that the underlying mechanism for yawning, both spontaneous and contagious, is involved in brain thermoregulation. The thermoregulatory theory provides clear predictions for both the primitive and derived features of this behavior. That is, the thermoregulatory benefits resulting from yawning provide the mechanism by which arousal or state change can be achieved [8,10], while the spreading of this behavior, i.e., yawn contagion, would therefore coordinate arousal in a group and enhance overall group vigilance [33]. In addition to enhancing the basic understanding of why we yawn, applications from this research include improved treatment and diagnosis of patients with thermoregulatory problems [34,37,54,55].

Acknowledgments

We would like to thank Bettina Janker for her help with approaching people on the street, and two anonymous reviewers for helpful comments to an earlier draft of this paper. This research was funded by a Lise Meitner grant of the Austrian Science Fund (FWF, grant nr.: M 1351-B17) to JJMM.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.physbeh.2014.03.032>.

References

- [1] Barbizet J. Yawning. *J Neurol Neurosurg Psychiatry* 1958;21:203–9.
- [2] Provine RR. Yawning as a stereotyped action pattern and releasing stimulus. *Ethology* 1986;72:109–22.
- [3] Anderson JR, Myowa-Yamakoshi M, Matsuzawa T. Contagious yawning in chimpanzees. *Proc R Soc B* 2004;271:S468–70.
- [4] Demuru E, Palagi E. In *Nobobos yawn contagion is higher among kin and friends*. *PLoS ONE* 2012;7:e49613.

- [5] Palagi E, Leone A, Mancini G, Ferrari PF. Contagious yawning in gelada baboons as a possible expression of empathy. *Proc Natl Acad Sci U S A* 2009;106:19262–7.
- [6] Miller ML, Gallup AC, Vogel AR, Vicario SM, Clark AB. Evidence for contagious behaviors in budgerigars (*Melopsittacus undulatus*): an observational study of yawning and stretching. *Behav Process* 2012;89:264–70.
- [7] Joly-Mascheroni RM, Senju A, Shepherd AJ. Dogs catch human yawns. *Biol Lett* 2008;4:446–8.
- [8] Baenninger R. On yawning and its functions. *Psychon Bull Rev* 1997;4:198–207.
- [9] Deputte BL. Ethological study of yawning in primates. I. Quantitative analysis and study of causation in two species of old world monkeys (*Cercocebus albigena* and *Macaca fascicularis*). *Ethology* 1994;98:221–45.
- [10] Provine RR, Hamernik HB, Curchack BC. Yawning: relation to sleeping and stretching in humans. *Ethology* 1987;76:152–60.
- [11] Provine RR. Yawning: the yawn is primal, unstoppable and contagious, revealing the evolutionary and neural basis of empathy and unconscious behavior. *Am Sci* 2005;93:532–9.
- [12] Baenninger R. Some comparative aspects of yawning in *Betta splendens*, *Homo sapiens*, *Panthera leo*, and *Papio sphinx*. *J Comp Psychol* 1987;101:349–54.
- [13] de Vries JJP, Visser GHA, Prechtl HF. The emergence of fetal behaviour. I. Qualitative aspects. *Early Hum Dev* 1982;7:301–22.
- [14] Anderson JR, Meno P. Psychological influences on yawning in children. *Curr Psychol Lett Behav Brain Cog* 2003;11:2–7.
- [15] Millen A, Anderson JR. Neither infants nor toddlers catch yawns from their mothers. *Biol Lett* 2010;7:440–2.
- [16] Hoogenhout M, van der Straaten K, Pileggi L-A, Malcolm-Smith S. Young children display contagious yawning when looking at the eyes. *J Child Adolesc Behav* 2013;1:1.
- [17] Madsen EA, Persson T. Contagious yawning in domestic dog puppies (*Canis lupus familiaris*): the effect of ontogeny and emotional closeness on low-level imitation in dogs. *Anim Cogn* 2012;16:233–40.
- [18] Madsen EA, Persson T, Sayehli S, Lenninger S, Sonesson G. Chimpanzees show a developmental increase in susceptibility to contagious yawning: a test of the effect of ontogeny and emotional closeness on yawn contagion. *PLoS ONE* 2013;8:e76266.
- [19] Campbell MW, de Waal FBM. Ingroup–outgroup bias in contagious yawning by chimpanzees supports link to empathy. *PLoS ONE* 2011;6:e18283.
- [20] Norscia I, Palagi E. Yawn contagion and empathy in *Homo sapiens*. *PLoS ONE* 2011;6:e28472.
- [21] de Waal FBM. Empathy in primates and other mammals. In: Decety J, editor. *Empathy from bench to bedside*. Cambridge: MIT Press; 2012. p. 87–106.
- [22] Platek SM, Critton SR, Myers TE, Gallup GG. Contagious yawning: the role of self-awareness and mental state attribution. *Cogn Brain Res* 2003;17:223–7.
- [23] Platek SM, Mohamed FB, Gallup Jr GG. Contagious yawning and the brain. *Cogn Brain Res* 2005;23:448–52.
- [24] Arnott SR, Singhal A, Goodale MA. An investigation of auditory contagious yawning. *Cogn Affect Behav Neurosci* 2009;9:335–42.
- [25] Nahab FB, Hattori N, Saad ZS, Hallett M. Contagious yawning and the frontal lobe: an fMRI study. *Hum Brain Mapp* 2009;30:1744–51.
- [26] Haker H, Kawohl W, Herwig U, Rössler W. Mirror neuron activity during contagious yawning — an fMRI study. *Brain Imaging Behav* 2013;7:28–34.
- [27] Cooper NR, Puzzo I, Pawley AD, Bowes-Mulligan RA, Kirkpatrick EV, Antoniou PA, et al. Bridging a yawning chasm: EEG investigations into the debate concerning the role of the human mirror neuron system in contagious yawning. *Cogn Affect Behav Neurosci* 2012;12:393–405.
- [28] Massen JJM, Vermunt DA, Sterck EHM. Male yawning is more contagious than female yawning among chimpanzees (*Pan troglodytes*). *PLoS ONE* 2012;7:e40697.
- [29] Gallup AC. Why do we yawn? Primitive versus derived features. *Neurosci Biobehav Rev* 2011;35:765–9.
- [30] Leone A, Ferrari PF, Palagi E. Different yawns, different functions? Testing social hypotheses on spontaneous yawning in *Theropithecus gelada*. *Sci Rep* 2014;4.
- [31] Vick SJ, Paukner A. Variation and context of yawns in captive chimpanzees (*Pan troglodytes*). *Am J Primatol* 2010;72:262–9.
- [32] Provine RR, Tate BC, Geldmacher LL. Yawning: no effect of 3–5% CO₂, 100% O₂, and exercise. *Behav Neural Biol* 1987;48:382–93.
- [33] Gallup AC, Gallup Jr GG. Yawning as a brain cooling mechanism: nasal breathing and forehead cooling diminish the incidence of contagious yawning. *Evol Psychol* 2007;5:92–101.
- [34] Gallup AC, Gallup Jr GG. Yawning and thermoregulation. *Physiol Behav* 2008;95:10–6.
- [35] Gallup AC, Eldakar OT. The thermoregulatory theory of yawning: what we know from five years of research. *Front Neurosci* 2013;6:1–13.
- [36] Shoup-Knox ML, Gallup AC, Gallup Jr GG, McNay EC. Yawning and stretching predict brain temperature changes in rats: support for the thermoregulatory hypothesis. *Front Evol Neurosci* 2010;2:1–5.
- [37] Gallup Jr GG, Gallup AC. Excessive yawning and thermoregulation: two case studies of chronic, debilitating bouts of yawning. *Sleep Breath* 2010;14:157–9.
- [38] Walusinski O. Why do we yawn? Past and current hypotheses. In: Shoja MM, Agutter PS, Tubbs RS, Ghanei M, Ghabili K, Harris A, Loukas M, editors. *Hypotheses in clinical medicine*. Hauppauge, NY: Nova Science Publishers; 2013. p. 245–56.
- [39] Elo H. Yawning and thermoregulation. *Sleep Breath* 2010;14:391–2.
- [40] Elo H. Yawning cannot cause significant temperature decreases in humans. *Sleep Med* 2011;12:102.
- [41] Guggisberg AG, Mathis J, Schnider A, Hess CW. Why do we yawn? *Neurosci Biobehav Rev* 2010;34:1267–76.
- [42] Guggisberg AG, Mathis J, Schnider A, Hess CW. Why do we yawn? The importance of evidence for specific yawn-induced effects. *Neurosci Biobehav Rev* 2011;35:1302–4.
- [43] Gallup AC. The thermoregulatory hypothesis of yawning: time to reconsider terms such as “impossible” and “cannot” and evaluate theories based on evidence. *Sleep Med* 2011;12:102–3.
- [44] Gallup AC, Hack GD. Human paranasal sinuses and selective brain cooling: a ventilation system activated by yawning? *Med Hypotheses* 2011;77:970–3.
- [45] Campos FA, Fedigan LM. Behavioural adaptations to heat stress and water scarcity in white-faced capuchins (*Cebus capucinus*) in Santa Rosa National Park, Costa Rica. *Am J Phys Anthropol* 2009;138:101–11.
- [46] Gallup AC, Miller ML, Clark AB. Yawning and thermoregulation in budgerigars (*Melopsittacus undulatus*). *Anim Behav* 2009;77:109–13.
- [47] Gallup AC, Miller ML, Clark AB. The direction and range of ambient temperature influences yawning in budgerigars (*Melopsittacus undulatus*). *J Comp Psychol* 2010;124:133–8.
- [48] Gallup AC, Miller RR, Clark AB. Changes in ambient temperature trigger yawning but not stretching in rats. *Ethology* 2011;117:145–53.
- [49] Gallup AC, Eldakar OT. Contagious yawning and seasonal climate variation. *Front Evol Neurosci* 2011;3(3):1–5.
- [50] Greco M, Baenninger R. Self-report as a valid measure of yawning in the laboratory. *Bull Psychonomic Soc* 1989;27:75–6.
- [51] Giganti F, Toselli M, Ramat S. Developmental trends in a social behaviour: contagious yawning in the elderly. *Eta Evolutiva* 2012;34:111–7.
- [52] Baenninger R, Greco M. Some antecedents and consequences of yawning. *Psychol Rec* 1991;41:453–60.
- [53] Giganti F, Zilli I. The daily time course of contagious and spontaneous yawning among humans. *J Ethol* 2011;29:215–9.
- [54] Gallup AC, Gallup Jr GG, Feo C. Yawning, sleep, and symptom relief in patients with multiple sclerosis. *Sleep Med* 2010;11:329–30.
- [55] Gallup AC, Gallup JA. Frequent yawning as an initial signal of fever relief. *Med Hypotheses* 2013;81:1034–5.