

GENERALIZED HÖRMANDER'S CONDITIONS AND WEIGHTED ENDPOINT ESTIMATES

MARÍA LORENTE, JOSÉ MARÍA MARTELL, CARLOS PÉREZ,
AND MARÍA SILVINA RIVEROS

ABSTRACT. We consider two-weight estimates for singular integral operators and their commutators with bounded mean oscillation functions. Hörmander type conditions in the scale of Orlicz spaces are assumed on the kernels. We prove weighted weak-type estimates for pairs of weights (u, Su) where u is an arbitrary nonnegative function and S is a maximal operator depending on the smoothness of the kernel. We also obtain sufficient conditions on a pair of weights (u, v) for the operators to be bounded from $L^p(v)$ to $L^{p,\infty}(u)$. One-sided singular integrals, as the differential transform operator, are under study. We also provide applications to Fourier multipliers and homogeneous singular integrals.

CONTENTS

1. Introduction	2
2. Preliminaries	5
2.1. Young functions and Orlicz spaces	5
2.2. Muckenhoupt weights	6
2.3. Singular Integral operators and Hörmander's type conditions	7
3. Statements of the main results	8
3.1. Singular integral operators	8
3.2. Commutators with BMO functions	10
3.3. One-sided operators	12
4. Applications	13
4.1. The differential transform operator	14
4.2. An example of a one-sided operator with $K \in H_\infty \cap H_{e^{t^{1/k}}, k}$	15
4.3. Multipliers	16
4.4. Kernels related to H_r and M_{L^r}	17
4.5. Homogeneous Singular Integrals	17
5. Proofs of the main results	18
6. Proofs in the one-sided case	25
References	29

2000 *Mathematics Subject Classification.* 42B20, 42B25.

Key words and phrases. Calderón-Zygmund operators, homogeneous singular integrals, multipliers, one-sided operators, commutators, BMO, Hörmander's condition of Young type, Muckenhoupt weights, two-weight estimates.

The first and last authors are partially supported by MCYT Grant MTM2005-08350-C03-02. The first author is also supported by Junta de Andalucía Grant FQM354. The second author is partially supported by MEC "Programa Ramón y Cajal, 2005", by MEC Grant MTM2007-60952, and by UAM-CM Grant CCG07-UAM/ESP-1664. The third author is partially supported by MTM2006-05622. The last author is partially supported by CONICET, Agencia Córdoba Ciencias and SECYT-UNC.

1. INTRODUCTION

The Calderón-Zygmund decomposition is a very powerful tool in Harmonic Analysis. Since its discovery in [6], many results have used it to derive boundedness properties of singular integral operators. For instance, using that the Hilbert or the Riesz transforms are bounded on L^2 , and by means of the Calderón-Zygmund decomposition, one proves that these classical operators are of weak-type $(1, 1)$. From this starting point, in the literature one can find many boundedness results for the Hilbert and the Riesz transforms: estimates on L^p , one-weight and two-weight norm inequalities,

The Calderón-Zygmund theory generalizes these ideas to provide a general framework allowing us to deal with singular integral operators. A typical Calderón-Zygmund convolution operator T is bounded on $L^2(\mathbb{R}^n)$ and has a kernel K on which different conditions are assumed. In the easiest case, K behaves as the kernel of the Hilbert or Riesz transforms. That is, K decays as $|x|^{-n}$ and its gradient as $|x|^{-n-1}$. It was already proved in [19] that these assumptions can be relaxed in order to show that T is of weak-type $(1, 1)$: it suffices to impose that K satisfies the so-called Hörmander condition (we write $K \in H_1$),

$$\int_{|x|>c|y|} |K(x-y) - K(x)| dx \leq C, \quad y \in \mathbb{R}^n, \quad c > 1.$$

From here, and by interpolation, T is bounded on $L^p(\mathbb{R}^n)$ with $1 < p < \infty$.

The underlying measure dx can be replaced by $w(x) dx$ where w is a Muckenhoupt A_p weight: The Hilbert and the Riesz transforms are bounded on $L^p(w) = L^p(w(x) dx)$ if and only if $w \in A_p$ for $1 < p < \infty$. For $p = 1$, the weak-type $(1, 1)$ with respect to w holds if and only if $w \in A_1$. The decay assumed before on the kernel and its gradient guarantees the same weighted estimates for the operator T . However, the Hörmander condition does not suffice to derive such estimates as it is proved in [18] (see also [26]). One can relax the decay conditions assumed on the kernel and still prove the previous weighted norm inequalities. Namely, it is enough to impose that K satisfies the following Lipschitz condition (we write $K \in H_\infty^*$):

$$|K(x-y) - K(x)| \leq C \frac{|y|^\alpha}{|x|^{\alpha+n}}, \quad |x| > c|y|.$$

With this condition in hand, one can show Coifman's estimate (see [7]): for any $0 < p < \infty$ and any $w \in A_\infty$

$$\int_{\mathbb{R}^n} |Tf(x)|^p w(x) dx \leq C \int_{\mathbb{R}^n} Mf(x)^p w(x) dx. \quad (1.1)$$

These estimates can be seen as a control of the operator T by the Hardy-Littlewood maximal function M and this allows one to show that T satisfies the most of the weighted estimates that M does (see [14] for more details).

When relaxing the H_∞^* condition, the operators become more singular and less smoother. Thus, the Coifman estimates to be expected will have a worse maximal operator on the righthand side. For instance, one has a scale of Hörmander's conditions based on the Lebesgue spaces L^r for $1 \leq r \leq \infty$ (see [22], [39] and [43]). A singular integral operator, with kernel satisfying the L^r -Hörmander condition, $1 < r \leq \infty$, satisfies a Coifman estimate with the maximal operator $M_{r'}$ in the righthand side (here $M_{r'}f(x) = M(|f|^{r'})(x)^{1/r'}$). These estimates are shown to be sharp in [26].

Let us notice that as r goes to 1 then r' goes to ∞ and the corresponding Coifman estimates get worse. In particular, when $K \in H_1$ one lacks of Coifman estimates (see [26]).

Sometimes, this scale of Hörmander conditions based on the Lebesgue spaces is too coarse and gives estimates that are not accurate enough. For instance, let us consider the differential transform operator studied in [20] and [4]:

$$T^+ f(x) = \sum_{j \in \mathbb{Z}} \nu_j (D_j f(x) - D_{j-1} f(x)), \quad (1.2)$$

where $\|\{\nu_j\}_j\|_\infty < \infty$ and

$$D_j f(x) = \frac{1}{2^j} \int_x^{x+2^j} f(t) dt.$$

We have that T^+ is a singular integral operator with kernel K supported in $(-\infty, 0)$, and therefore T^+ is a one-sided singular integral operator (that is the reason why we write T^+). In [4] it was shown that $K \in \cap_{r \geq 1} H_r$ (here H_r is the Hörmander condition associated with L^r , see the precise definition below). Thus one can show that T^+ satisfies a Coifman estimate with M_q in the righthand side for any $1 < q < \infty$. Indeed, exploiting the fact that T^+ is a one-sided operator one can do better: $M_q f$ can be replaced by the pointwise smaller operator $M_q^+ f$ (the corresponding one-sided maximal function) and A_∞ by the bigger class A_∞^+ (see the precise definitions and more details below). Notice that one can take any $1 < q < \infty$, with the case $q = 1$ remaining open (in general, $K \notin H_\infty$). Nevertheless, there are other maximal operators that one can write between M (or M^+) and M_q (or M_q^+): any iteration of the Hardy-Littlewood maximal function, or maximal operators associated with Orlicz spaces lying between L^1 and L^q as $L(\log L)^\alpha$, $\alpha > 0$.

These ideas motivated [24] on which new classes of Hörmander conditions based on Orlicz spaces were introduced. Roughly, given a Young function \mathcal{A} , associated with the Orlicz space $L^\mathcal{A}$ one can define a Hörmander class $H_\mathcal{A}$ (see Definition 2.3). Thus, a singular integral operator with kernel in $H_\mathcal{A}$, is controlled in the sense of Coifman by the maximal operator $M_{\overline{\mathcal{A}}}$ (which is the maximal function associated with the space $L^{\overline{\mathcal{A}}}$) where $\overline{\mathcal{A}}$ is the conjugate function of \mathcal{A} . This was obtained in [24] as well as the one-sided case (see Theorems 2.4 and 3.11 below).

For the differential transform introduced above one can show that $K \in H_{e^{t^{1/(1+\varepsilon)}}}$ for any $\varepsilon > 0$. Thus T^+ satisfies a Coifman type estimate with $M_{L(\log L)^{1+\varepsilon}}^+$ on the right hand side—in terms of iterations one can write $(M^+)^3$ —and this maximal operator is pointwise smaller than M_q^+ for any $1 < q < \infty$.

Coifman's estimates are important from the point of view of weighted norm inequalities since they encode a lot of information about the singularity of the operator T (see [10] and [14]). In some sense, T behaves as the maximal operator that controls it. For instance, one shows that T is bounded on $L^p(w)$ for $1 < p < \infty$, $w \in A_p$. Also one can see that T is of weak-type $(1, 1)$ for weights in A_1 , T is bounded on weighted rearrangement invariant function spaces, T satisfies weighted modular inequalities (see [14]), etc. All these one-weight estimates are based on the fact that (1.1) is valid for any weight in A_∞ and the weight always move within this class.

The situation changes when one works with two-weight inequalities. Let us focus on the end-point estimates for $p = 1$. In the one weight case, M is bounded from $L^1(w)$ to $L^{1,\infty}(w)$ for every $w \in A_1$. Also, there is a version of (1.1) in the sense of $L^{1,\infty}(w)$, that is, $\|Tf\|_{L^{1,\infty}(w)} \lesssim \|Mf\|_{L^{1,\infty}(w)}$ for every $w \in A_\infty$ (see [10]). These two facts imply at once that T is of weak-type $(1, 1)$ for weights in A_1 . In the two-weight case, Vitali's covering lemma easily gives that for every weight u (a weight is a non-negative locally integrable function)

$$u\{x \in \mathbb{R}^n : Mf(x) > \lambda\} \lesssim \frac{1}{\lambda} \int_{\mathbb{R}^n} |f(x)| Mu(x) dx.$$

However, this estimate is not known for the singular integral operators with smooth kernel. Even for the Hilbert or the Riesz transforms the validity of this estimate is an open question. Reasoning as above, one seeks for pairs of weight (u, Su) for which these operators are of weak-type $(1, 1)$, where S will be a maximal operator worse, in principle, than M . For instance, one can put $S = M_q$ for every $1 < q < \infty$: using that $M_q u \in A_1$ and Coifman's estimate (in $L^{1,\infty}$) one easily obtains the estimate proved in [9]

$$\|Tf\|_{L^{1,\infty}(u)} \leq \|Tf\|_{L^{1,\infty}(M_q u)} \lesssim \|Mf\|_{L^{1,\infty}(M_q u)} \lesssim \|f\|_{L^1(M_q u)}.$$

As observed before, there are some other maximal operators that lie between M and M_q as the iterations of M or $M_{L(\log L)^\alpha}$, $\alpha > 0$. In [32], by means of the Calderón-Zygmund decomposition, it was proved that if T is a singular integral operator with smooth kernel (say $K \in H_\infty^*$), as the Hilbert or Riesz transform, then for any $\varepsilon > 0$ and any weight u

$$u\{x \in \mathbb{R}^n : |Tf(x)| > \lambda\} \lesssim \frac{1}{\lambda} \int_{\mathbb{R}^n} |f(x)| M_{L(\log L)^\varepsilon} u(x) dx. \quad (1.3)$$

Note that in terms of iterations one can write M^2 .

The goal of this paper is to study estimates like (1.3) when the operator T has a less smoother kernel. That is, if we impose that the kernel of T satisfies a Hörmander condition in the scale of Orlicz spaces, we seek for a maximal operator S so that T is of weak-type $(1, 1)$ with respect to the pair of weights (u, Su) . The main technique to be used is the Calderón-Zygmund decomposition. The bad part is handled by using the smoothness of the kernel. For the good part, one needs a strong two-weight estimate that usually follows from a Coifman estimate (see Theorem 2.6). We also obtain weighted end-point estimates for the commutators of such operators with BMO functions. The corresponding Coifman estimates have been studied in [23]. As one of our main examples is the differential transform operator presented above, we also pay attention to the one-sided operators in which case one can obtain better estimates by replacing a maximal operator by its corresponding one-sided analog.

The paper is organized as follows. The following section contains some of the preliminaries and definitions that are needed to state our results. In Section 3 we state our main results on singular integral operators, their commutators with BMO functions and also we consider the one-sided case. Some applications, including the differential transform operator and multipliers, are given in Section 4. Finally Sections 5 and 6 contain the proof of our main results.

2. PRELIMINARIES

2.1. Young functions and Orlicz spaces. We recall some of the needed background for Orlicz spaces we refer the reader to [37] and [3] for a complete account of this topic. A function $\mathcal{A} : [0, \infty) \rightarrow [0, \infty)$ is a Young function if it is continuous, convex, increasing and satisfies $\mathcal{A}(0) = 0$, $\mathcal{A}(\infty) = \infty$. We will assume the Young functions are normalized so that $\mathcal{A}(1) = 1$. We introduce the following localized and averaged Luxemburg norm associated with the Orlicz space $L^{\mathcal{A}}$: given a cube Q

$$\|f\|_{\mathcal{A},Q} = \inf \left\{ \lambda > 0 : \frac{1}{|Q|} \int_Q \mathcal{A} \left(\frac{|f(x)|}{\lambda} \right) dx \leq 1 \right\}.$$

For instance, when $\mathcal{A}(t) = t^r$ with $r \geq 1$ then we have

$$\|f\|_{L^r,Q} = \left(\frac{1}{|Q|} \int_Q |f(x)|^r dx \right)^{\frac{1}{r}}.$$

It is well known that if $\mathcal{A}(t) \leq C\mathcal{B}(t)$ for $t \geq t_0$ then $\|f\|_{\mathcal{A},Q} \leq C\|f\|_{\mathcal{B},Q}$. Thus the behavior of $\mathcal{A}(t)$ for $t \leq t_0$ does not matter: if $\mathcal{A}(t) \approx \mathcal{B}(t)$ for $t \geq t_0$ the latter estimate implies that $\|f\|_{\mathcal{A},Q} \approx \|f\|_{\mathcal{B},Q}$. This means that in most of the cases we will not be concerned about the value of the Young functions for t small.

We can now define the Hardy-Littlewood maximal function associated with \mathcal{A} as

$$M_{\mathcal{A}}f(x) = \sup_{Q \ni x} \|f\|_{\mathcal{A},Q}.$$

When $\mathcal{A}(t) = t$ then $M_{\mathcal{A}} = M$ is the Hardy-Littlewood maximal function. For $\mathcal{A}(t) = t^r$ with $r > 1$ we have $M_{\mathcal{A}}f(x) = M(|f|^r)(x)^{1/r}$.

Given a Young function \mathcal{A} , we say that \mathcal{A} is doubling, we write $\mathcal{A} \in \Delta_2$, if $\mathcal{A}(2t) \leq C\mathcal{A}(t)$ for every $t \geq t_0 > 0$. For $1 < p < \infty$, \mathcal{A} belongs to B_p if there exists $c > 0$ such that

$$\int_c^\infty \frac{\mathcal{A}(t)}{t^p} \frac{dt}{t} < \infty.$$

This condition appears first in [34] and it was shown that $\mathcal{A} \in B_p$ if and only if $M_{\mathcal{A}}$ is bounded on $L^p(\mathbb{R}^n)$.

Abusing on the notation if $\mathcal{A}(t) = t^r$, $\mathcal{A}(t) = e^{t^\alpha} - 1$ or $\mathcal{A}(t) = t^r(1 + \log^+ t)^\alpha$, the Orlicz norms are respectively written as $\|\cdot\|_r = \|\cdot\|_{L^r}$, $\|\cdot\|_{\exp L^\alpha}$, $\|\cdot\|_{L^r(\log L)^\alpha}$ and the corresponding maximal operators as $M_r = M_{L^r}$, $M_{\exp L^\alpha}$ and $M_{L^r(\log L)^\alpha}$. For $k \geq 0$, it is known that $M_{L(\log L)^k} f(x) \approx M^{k+1} f(x)$ where M^k is the k -times iterated of M (see [33], [38] and [14]).

In \mathbb{R} , we can also define the one-sided maximal functions associated with a given Young function \mathcal{A} :

$$M_{\mathcal{A}}^+ f(x) = \sup_{b > x} \|f\|_{\mathcal{A},(x,b)} \quad \text{and} \quad M_{\mathcal{A}}^- f(x) = \sup_{a < x} \|f\|_{\mathcal{A},(a,x)}.$$

The one-sided Hardy-Littlewood maximal functions M^+ , M^- correspond to the case $\mathcal{A}(t) = t$.

Given a Young function \mathcal{A} , let $\overline{\mathcal{A}}$ denote its associate function: the Young function with the property that $t \leq \mathcal{A}^{-1}(t) \overline{\mathcal{A}}^{-1}(t) \leq 2t$, $t > 0$. If $\mathcal{A}(t) = t^r$ with $1 < r < \infty$, then $\overline{\mathcal{A}}(t) = t^{r'}$; if $\mathcal{A}(t) = t^r \log(e+t)^\alpha$, then $\overline{\mathcal{A}}(t) \approx t^{r'} \log(e+t)^{-\alpha(r'-1)}$.

One has the generalized Hölder's inequality

$$\frac{1}{|Q|} \int_Q |f g| \leq 2 \|f\|_{\mathcal{A},Q} \|g\|_{\overline{\mathcal{A}},Q}.$$

There is a further generalization that turns out to be useful for our purposes, see [31]: If $\mathcal{A}, \mathcal{B}, \mathcal{C}$ are Young functions such that $\mathcal{A}^{-1}(t) \mathcal{B}^{-1}(t) \mathcal{C}^{-1}(t) \leq t$, for all $t \geq t_0 > 0$ (in what follows we assume that $t_0 = 1$ for simplicity and clearness in the computations), —sometimes, we will equivalently write $\mathcal{A}^{-1}(t) \mathcal{B}^{-1}(t) \leq \overline{\mathcal{C}}^{-1}(t)$ — then

$$\|f g h\|_{L^1, Q} \leq C \|f\|_{\mathcal{A}, B} \|g\|_{\mathcal{B}, Q} \|h\|_{\mathcal{C}, Q}, \quad \|f g\|_{\overline{\mathcal{C}}, Q} \leq C \|f\|_{\mathcal{A}, Q} \|g\|_{\mathcal{B}, Q}. \quad (2.1)$$

Remark 2.1. Let us observe that when $\mathcal{D}(t) = t$, which gives L^1 , then $\overline{\mathcal{D}}(t) = 0$ if $s \leq 1$ and $\overline{\mathcal{D}}(t) = \infty$ otherwise. Although $\overline{\mathcal{D}}$ is not a Young function one can see that the space $L^{\overline{\mathcal{D}}}$ coincides with L^∞ . On the other hand, as the (generalized) inverse is $\overline{\mathcal{D}}^{-1}(t) \equiv 1$, the previous Hölder's inequalities make sense with the appropriate changes if one of the three functions is \mathcal{D} or $\overline{\mathcal{D}}$. We will use this throughout the paper.

Remark 2.2. The convexity of \mathcal{A} implies that $\mathcal{A}(t)/t$ is increasing and so $t \leq C \mathcal{A}(t)$ for all $t \geq 1$. This yields that $\|f\|_{L^1, B} \leq C \|f\|_{\mathcal{A}, B}$ for all Young functions \mathcal{A} .

2.2. Muckenhoupt weights. We recall the definition of the Muckenhoupt classes A_p , $1 \leq p \leq \infty$. Let w be a non-negative locally integrable function and $1 \leq p < \infty$. We say that $w \in A_p$ if there exists $C_p < \infty$ such that for every ball $B \subset \mathbb{R}^n$

$$\left(\frac{1}{|B|} \int_B w(x) dx \right) \left(\frac{1}{|B|} \int_B w(x)^{1-p'} dx \right)^{p-1} \leq C_p,$$

when $1 < p < \infty$, and for $p = 1$,

$$\frac{1}{|B|} \int_B w(y) dy \leq C_1 w(x), \quad \text{for a.e. } x \in B,$$

which can be equivalently written as $Mw(x) \leq C_1 w(x)$ for a.e. $x \in \mathbb{R}^n$. Finally we set $A_\infty = \cup_{p \geq 1} A_p$. It is well known that the Muckenhoupt classes characterize the boundedness of the Hardy-Littlewood maximal function on weighted Lebesgue spaces. Namely, $w \in A_p$, $1 < p < \infty$, if and only if M is bounded on $L^p(w)$; and $w \in A_1$ if and only if M maps $L^1(w)$ into $L^{1,\infty}(w)$.

In \mathbb{R} , the weighted estimates for the one-sided Hardy-Littlewood maximal function M^+ (and analogously for M^-) are modeled for the classes A_p^+ which are defined as follows. Given $1 < p < \infty$, $w \in A_p^+$, if there exists a constant $C_p < \infty$ such that for all $a < b < c$

$$\frac{1}{(c-a)^p} \left(\int_a^b w(x) dx \right) \left(\int_b^c w(x)^{1-p'} dx \right)^{p-1} \leq C_p.$$

We say that $w \in A_1^+$ if $M^-w(x) \leq C_1 w(x)$ for a.e. $x \in \mathbb{R}$. The class A_∞^+ is defined as the union of all the A_p^+ classes, $A_\infty^+ = \cup_{p \geq 1} A_p^+$. The classes A_p^- are defined in a similar way. It is interesting to note that $A_p = A_p^+ \cap A_p^-$, $A_p \subsetneq A_p^+$ and $A_p \subsetneq A_p^-$. See [41], [27], [28], [29] for more definitions and results.

2.3. Singular Integral operators and Hörmander's type conditions. Let T be a singular integral operator of convolution type, that is, T is bounded on $L^2(\mathbb{R}^n)$ and

$$Tf(x) = \text{p.v.} \int_{\mathbb{R}^n} K(x-y) f(y) dy$$

where K is a measurable function defined away from 0. Convolution operators are considered for simplicity, but the results presented here can be stated for variable kernels with the appropriate changes. The precise statements and the details are left to the reader.

When $n = 1$ and we further assumed that the kernel K is supported on $(-\infty, 0)$ we say that T is a one-sided singular integral and we write T^+ to emphasize it. The results that we present below for (regular) singular integrals apply to T^+ . However, taking advantage of the extra assumption on the kernel, one can be more precise and get better estimates (see Section 3.3).

We introduce the different Hörmander type conditions assumed on the kernel K . The weakest one is the so-called Hörmander condition H_1 (we simply say $K \in H_1$ or K satisfies the L^1 -Hörmander condition): there are constants $c > 1$ and $C > 0$ such that

$$\int_{|x|>c|y|} |K(x-y) - K(x)| dx \leq C, \quad y \in \mathbb{R}^n.$$

The strongest one is the classical Lipschitz condition called H_∞^* (this notation is not standard but we keep H_∞ for a weaker L^∞ -condition, see the definition below). We say that $K \in H_\infty^*$ if there are $\alpha, C > 0$ and $c > 1$ such that

$$|K(x-y) - K(x)| \leq C \frac{|y|^\alpha}{|x|^{\alpha+n}}, \quad |x| > c|y|.$$

Between H_1 and H_∞^* one finds the L^r -Hörmander conditions (which are called $H_{L^r} = H_r$ in the definition below). These classes appeared implicitly in the work [22] where it is shown that classical L^r -Dini condition for K implies $K \in H_r$ (see also [39] and [43]). However, there are examples of singular integrals like the differential transform operator from Ergodic Theory defined in (1.2), whose kernel $K \in H_r$ for all $1 \leq r < \infty$ but $K \notin H_\infty$. As it was obtained in [23], K satisfies a Hörmander condition in the scale of the Orlicz spaces that lies between the intersection of the classes H_r for $1 \leq r < \infty$ and H_∞ . The same happens with the one-sided discrete square function considered in [42] and [24]. All this has motivated the definition of the L^A -Hörmander conditions in [24]:

Definition 2.3. *The kernel K is said to satisfy the L^A -Hörmander condition, we write $K \in H_A$, if there exist $c \geq 1$, $C > 0$ such that for any $y \in \mathbb{R}^n$ and $R > c|y|$,*

$$\sum_{m=1}^{\infty} (2^m R)^n \|K(\cdot - y) - K(\cdot)\|_{\mathcal{A}, |x| \sim 2^m R} \leq C.$$

We say that $K \in H_\infty$ if K satisfies the previous condition with $\|\cdot\|_{L^\infty, |x| \sim 2^m R}$ in place of $\|\cdot\|_{\mathcal{A}, |x| \sim 2^m R}$.

We have used the notation: $|x| \sim s$ for $s < |x| \leq 2s$ and

$$\|f\|_{\mathcal{A}, |x| \sim s} = \|f \chi_{\{|x| \sim s\}}\|_{\mathcal{A}, B(0, 2s)}.$$

Note that if $\mathcal{A}(t) = t$ then $H_{\mathcal{A}} = H_1$. On the other hand, since $t \leq C \mathcal{A}(t)$ for $t \geq 1$ we have that $H_{\mathcal{A}} \subset H_1$ which implies that the classical unweighted Calderón-Zygmund theory can be applied to T . Also, it is easy to see that $H_{\infty}^* \subset H_{\infty} \subset H_{\mathcal{A}}$. For convenience thorough this paper we write $|\cdot| = |\cdot|_{\infty}$ so that everything is adapted to cubes in place of balls (with the appropriate changes everything can be written in terms of balls). For simplicity we also assume that $c = 1$.

Coifman type estimates were proved for kernels in these classes in [24]:

Theorem 2.4 ([24]). *Let \mathcal{A} be a Young function and let T be a singular integral operator with kernel $K \in H_{\mathcal{A}}$. Then for any $0 < p < \infty$ and $w \in A_{\infty}$,*

$$\int_{\mathbb{R}^n} |Tf(x)|^p w(x) dx \leq C \int_{\mathbb{R}^n} M_{\overline{\mathcal{A}}} f(x)^p w(x) dx, \quad f \in L_c^{\infty}, \quad (2.2)$$

whenever the left-hand side is finite.

Note that this improves the previous results in [22], [39] and [43] (for sharpness issues see also [26]). Similar results are also proved for vector-valued and one-sided operators (see [24]).

Remark 2.5. Abusing on the notation, as in Remark 2.1, if $K \in H_{\infty}$, then (2.2) holds with $M_{\overline{\mathcal{A}}} = M$, where $\overline{\mathcal{A}}(t) = t$. This was obtained in [26] improving the corresponding result for the smaller class H_{∞}^* .

The previous estimates are useful in applications as one has that T and $M_{\overline{\mathcal{A}}}$ have a similar behavior (see [14]). For instance, two-weight estimates can be proved in the following way:

Theorem 2.6 ([23]). *Let \mathcal{A} be a Young function and $1 < p < \infty$. Suppose that there exist Young functions \mathcal{D} , \mathcal{E} such that $\mathcal{E} \in B_{p'}$ and $\mathcal{D}^{-1}(t) \mathcal{E}^{-1}(t) \leq \overline{\mathcal{A}}^{-1}(t)$ for $t \geq t_0 > 0$. Set $\mathcal{D}_p(t) = \mathcal{D}(t^{1/p})$. Let T be a linear operator such that its adjoint T^* satisfies (2.2). Then for any weight u ,*

$$\int_{\mathbb{R}^n} |Tf(x)|^p u(x) dx \leq C \int_{\mathbb{R}^n} |f(x)|^p M_{\mathcal{D}_p} u(x) dx. \quad (2.3)$$

Remark 2.7. Abusing in the notation, the previous result contains the case $\overline{\mathcal{A}}(t) = t$ on which in (2.2) one has $M_{\overline{\mathcal{A}}} = M$. Then, \mathcal{D} and \mathcal{E} are conjugate functions and so (2.3) holds for any \mathcal{D}_p such that $\overline{\mathcal{D}} \in B_{p'}$. In particular, in (2.3) we can take the pair of weights $(u, M_{L(\log L)^{p-1+\delta}} u)$ for any $\delta > 0$: pick $\mathcal{D}(t) = t^p (1 + \log^+ t)^{p-1+\delta}$ whose conjugate function is $\overline{\mathcal{D}}(t) \approx t^{p'}/(1 + \log^+ t)^{1+\delta(p'-1)} \in B_{p'}$.

3. STATEMENTS OF THE MAIN RESULTS

3.1. Singular integral operators. We are going to obtain endpoint two-weight norm inequalities for singular integral operators where different Hörmander's conditions are assumed on the kernel. Namely, we seek for the following weak-type $(1, 1)$ estimates with pairs of weights (u, Su) where S will be a certain maximal function depending on the smoothness of the kernel:

$$u\{x \in \mathbb{R}^n : |Tf(x)| > \lambda\} \leq \frac{C}{\lambda} \int_{\mathbb{R}^n} |f(x)| Su(x) dx. \quad (3.1)$$

Theorem 3.1. *Let T be a singular integral operator with kernel K .*

- (a) *Let \mathcal{A} be a Young function such that its complementary function $\bar{\mathcal{A}} \in \Delta_2$ and assume that there exists $r > 1$ so that $\liminf_{t \rightarrow \infty} \bar{\mathcal{A}}(t)/t^r > 0$. If $K \in H_{\mathcal{A}}$ then (3.1) holds for the pairs of weights $(u, M_{\bar{\mathcal{A}}}u)$.*
- (b) *Let \mathcal{A} be a Young function and assume that there exist $1 < p < \infty$, and Young functions \mathcal{D} and \mathcal{E} such that $\mathcal{D}^{-1}(t)\mathcal{E}^{-1}(t) \leq \bar{\mathcal{A}}^{-1}(t)$ for $t \geq t_0 > 0$ with $\mathcal{E} \in B_{p'}$. If $K \in H_{\mathcal{A}}$ then, (3.1) holds for the pairs of weights $(u, M_{\mathcal{D}_p}u)$ with $\mathcal{D}_p(t) = \mathcal{D}(t^{1/p})$.*
- (c) *If $K \in H_{\infty}$, then (3.1) holds for the pairs of weights $(u, M_{L(\log L)^{\varepsilon}}u)$ for any $\varepsilon > 0$.*

Remark 3.2. In part (c) we improve [32], as we consider a wider class of kernels $H_{\infty}^* \subsetneq H_{\infty}$.

Remark 3.3. Let us notice that when $\liminf_{t \rightarrow \infty} \bar{\mathcal{A}}(t)/t^r > 0$, the pair of weights in (a) is better than the one in (b): one can see that $\bar{\mathcal{A}}(t) \lesssim \mathcal{D}_p(t)$ for $t \geq 1$. Take an arbitrary $t \geq 1$. The fact that $\mathcal{E} \in B_{p'}$ implies $\mathcal{E}(t) \lesssim t^{p'}$. Also, $\bar{\mathcal{A}}(t) \geq t$ as $\bar{\mathcal{A}}$ is a Young function. Then, from the condition assumed on $\bar{\mathcal{A}}$, \mathcal{D} and \mathcal{E} it follows that $\mathcal{D}^{-1}(t) \lesssim t^{1/p}$ and therefore

$$\bar{\mathcal{A}}^{-1}(t) \geq \mathcal{D}^{-1}(t)\mathcal{E}^{-1}(t) \gtrsim \mathcal{D}^{-1}(t)^p t^{1/p'} / \mathcal{D}^{-1}(t)^{p-1} \gtrsim \mathcal{D}^{-1}(t)^p = \mathcal{D}_p^{-1}(t).$$

Remark 3.4. We would like to emphasize that in part (a) the associated Coifman estimate in Theorem 2.4 tells us that T is controlled by $M_{\bar{\mathcal{A}}}$. Here we show that the pair of weights of the form $(u, M_{\bar{\mathcal{A}}}u)$ is valid. In the previous remark, we have observed that in (b) one gets a bigger maximal operator $M_{\mathcal{D}_p}$. In many applications, although we take p very close to 1, we always obtain a maximal operator pointwise greater than $M_{\bar{\mathcal{A}}}$. This is the case in (c) which covers the classical Hilbert and Riesz transforms. Here as these operators are controlled by M (in the sense of Coifman) one would wish to show that the pair of weights (u, Mu) is valid. However this remains as an open question and the best known result is $(u, M_{L(\log L)^{\varepsilon}}u)$.

There is a general extrapolation principle that allows one to pass from pairs of weights (u, Su) , with S being a maximal operator, to general pairs of weights (u, v) . The main ideas are implicit in [12], [13] and are further exploited in [11]. Below we present a proof in the one-sided case (see Theorem 3.14), that can be easily adapted to the present situation.

Theorem 3.5. *Let \mathcal{F} be a Young function and assume that a given operator T satisfies*

$$u\{x \in \mathbb{R}^n : |Tf(x)| > \lambda\} \leq \frac{C}{\lambda} \int_{\mathbb{R}^n} |f(x)| M_{\mathcal{F}}u(x) dx \quad (3.2)$$

for every weight u and $\lambda > 0$. Given $1 < p < \infty$, let \mathcal{G}, \mathcal{H} be Young functions such that $\mathcal{G}^{-1}(t)\mathcal{H}^{-1}(t) \leq \mathcal{F}^{-1}(t)$ for all $t \geq t_0 > 0$ and $\mathcal{H} \in B_{p'}$. Then for any pair of weights (u, v) satisfying

$$\|u^{1/p}\|_{\mathcal{G}, Q} \|v^{-1/p}\|_{L^{p'}, Q} \leq C \quad (3.3)$$

and for any and $\lambda > 0$ we have

$$u\{x \in \mathbb{R}^n : |Tf(x)| > \lambda\} \leq \frac{C}{\lambda^p} \int_{\mathbb{R}^n} |f(x)|^p v(x) dx. \quad (3.4)$$

3.2. Commutators with BMO functions. We are going to consider commutators of singular integral operators with BMO functions. Let us remind that a locally integrable functions b is in BMO if

$$\|b\|_{\text{BMO}} = \sup_Q \frac{1}{|Q|} \int_Q |b(x) - b_Q| dx < \infty,$$

where the sup runs over all cubes $Q \subset \mathbb{R}^n$ with the sides parallel to the coordinate axes and where b_Q stands for the average of b over Q .

We define the (first-order) commutator by

$$T_b^1 f(x) = [b, T]f(x) = b(x)Tf(x) - T(bf)(x).$$

The higher order commutators T_b^k are defined by induction as $T_b^k = [b, T_b^{k-1}]$ for $k \geq 2$. Note that for every $k \geq 1$

$$T_b^k f(x) = \text{p.v.} \int_{\mathbb{R}^n} (b(x) - b(y))^k K(x - y) f(y) dy.$$

For $k = 0$ we understand that $T_b^0 = T$.

In [23], Coifman type estimates were proved for commutators of singular integral operators with kernels in the following Hörmander classes that depend on the order of the commutator.

Definition 3.6. Let \mathcal{A} be a Young function and $k \in \mathbb{N}$. We say that the kernel K satisfies the $L^{\mathcal{A},k}$ -Hörmander condition, we write $K \in H_{\mathcal{A},k}$, if there exist $c \geq 1$ and $C > 0$ (depending on \mathcal{A} and k) such that for all $y \in \mathbb{R}^n$ and $R > c|y|$

$$\sum_{m=1}^{\infty} (2^m R)^n m^k \|K(\cdot - y) - K(\cdot)\|_{\mathcal{A},|x| \sim 2^m R} \leq C.$$

We say that $K \in H_{\infty,k}$ if K satisfies the previous condition with $\|\cdot\|_{L^\infty,|x| \sim 2^m R}$ in place of $\|\cdot\|_{\mathcal{A},|x| \sim 2^m R}$.

As before, for simplicity we will assume that $c = 1$. For these classes the following Coifman estimates are obtained:

Theorem 3.7 ([23]). Let $b \in \text{BMO}$ and $k \geq 0$.

(a) Let \mathcal{A}, \mathcal{B} be Young functions, such that $\overline{\mathcal{A}}^{-1}(t) \mathcal{B}^{-1}(t) \overline{\mathcal{C}}_k^{-1}(t) \leq t$ for $t \geq t_0 > 0$ with $\overline{\mathcal{C}}_k(t) = e^{t^{1/k}} - 1$. If T is a singular integral operator with kernel $K \in H_{\mathcal{B}} \cap H_{\mathcal{A},k}$ (or, in particular, $K \in H_{\mathcal{B},k}$), then for any $0 < p < \infty$, $w \in A_\infty$,

$$\int_{\mathbb{R}^n} |T_b^k f(x)|^p w(x) dx \leq C \|b\|_{\text{BMO}}^{pk} \int_{\mathbb{R}^n} M_{\overline{\mathcal{A}}} f(x)^p w(x) dx, \quad f \in L_c^\infty, \quad (3.5)$$

whenever the left-hand side is finite.

(b) If $K \in H_\infty \cap H_{e^{t^{1/k}},k}$ (or, in particular, $K \in H_{\infty,k}$) then (3.5) holds with M^{k+1} —the $k+1$ -iteration of M — in place of $M_{\overline{\mathcal{A}}}$.

This result and Theorem 2.6 can be used to derived endpoint estimates of the form

$$u\{x \in \mathbb{R}^n : |T_b^k f(x)| > \lambda\} \leq C \int_{\mathbb{R}^n} \mathcal{C}_k \left(\frac{\|b\|_{\text{BMO}}^k |f(x)|}{\lambda} \right) Su(x) dx, \quad (3.6)$$

where $\mathcal{C}_k(t) = t(1 + \log^+ t)^k$.

Theorem 3.8. *Let T be a singular integral operator with kernel K , $k \in \mathbb{N}$, $b \in \text{BMO}$ and T_b^k the k -th order commutator of T .*

(a) *Let \mathcal{A} , \mathcal{B} be Young functions such that $\overline{\mathcal{A}}^{-1}(t) \mathcal{B}^{-1}(t) \overline{\mathcal{C}}_k^{-1}(t) \leq t$ for $t \geq t_0 > 0$ with $\overline{\mathcal{C}}_k(t) = e^{t^{1/k}} - 1$. Let $K \in H_{\mathcal{B}} \cap H_{\mathcal{A},k}$ (or, in particular, $K \in H_{\mathcal{B},k}$).*

(a.1) *If $\overline{\mathcal{A}} \in \Delta_2$ and there exists $r > 1$ so that $\liminf_{t \rightarrow \infty} \overline{\mathcal{A}}(t)/t^r > 0$, then (3.6) holds for the pairs of weights $(u, M_{\overline{\mathcal{A}}}u)$.*

(a.2) *Assume that there exist $1 < p < \infty$, and Young functions \mathcal{D} and \mathcal{E} such that $\mathcal{D}^{-1}(t) \mathcal{E}^{-1}(t) \leq \overline{\mathcal{A}}^{-1}(t)$ for $t \geq t_0 > 0$ with $\mathcal{E} \in B_{p'}$. Then, (3.6) holds for the pairs of weights $(u, M_{\mathcal{D}_p}u)$ with $\mathcal{D}_p(t) = \mathcal{D}(t^{1/p})$.*

(b) *If $K \in H_{\infty} \cap H_{e^{t^{1/k}},k}$ (or, in particular, $K \in H_{\infty,k}$), then (3.6) holds for the pairs of weights $(u, M_{L(\log L)^{k+\varepsilon}}u)$ for any $\varepsilon > 0$.*

Remark 3.9. In part (b) we obtain the same result as in [35], but considering a weaker condition on the kernel K , since $H_{\infty}^* \subsetneq H_{\infty} \cap H_{e^{t^{1/k}},k}$.

Remark 3.10. Notice that we can understand (b) as an extension of (a) when \mathcal{B} corresponds to L^{∞} and so $\mathcal{A}(t) = \overline{\mathcal{C}}_k(t) = e^{t^{1/k}} - 1$. Observe that in that case we also have $\overline{\mathcal{A}}^{-1}(t) \mathcal{B}^{-1}(t) \overline{\mathcal{C}}_k^{-1}(t) \leq t$ (where $\mathcal{B}^{-1}(t) \equiv 1$).

This result can be extended to the multilinear commutators considered in [36]. Given $k \geq 1$, a singular integral operator T with kernel K and a vector $\vec{b} = (b_1, \dots, b_k)$ of locally integrable functions, the multilinear commutator is defined as

$$T_{\vec{b}}f(x) = \int_{\mathbb{R}^n} \left(\prod_{l=1}^k (b_l(x) - b_l(y)) \right) K(x, y) f(y) dy.$$

When $k = 0$ we understand that $T_{\vec{b}} = T$. Notice that if $k = 1$ and $\vec{b} = b$ then $T_{\vec{b}} = T_b^1$. For $k \geq 1$ if $b_1 = \dots = b_k = b$ then $T_{\vec{b}} = T_b^k$.

For standard commutators, one assumes that $b \in \text{BMO}$, and by John-Nirenberg's inequality we have that $\|b\|_{\text{BMO}} \approx \sup_Q \|b - b_Q\|_{\exp L}$. This can be seen as a supremum of the oscillations of b on the space $\exp L$.

As it was done in [36], when dealing with multilinear commutators, the symbols b_j are assumed to be in one of this oscillation spaces. Given $s \geq 1$ we set

$$\|f\|_{\text{Osc}(\exp L^s)} = \sup_Q \|f - f_Q\|_{\exp L^s}$$

and the space $\text{Osc}(\exp L^s)$ is the set of measurable functions $f \in L_{\text{loc}}^1(\mathbb{R}^n)$ such that $\|f\|_{\text{Osc}(\exp L^s)} < \infty$. Let us notice that $\text{Osc}(\exp L^s) \subset \text{Osc}(\exp L^1) = \text{BMO}$. We assume that for each $1 \leq l \leq k$, $b_l \in \text{Osc}(\exp L^{s_l})$ with $s_l \geq 1$. We set $\frac{1}{s} = \frac{1}{s_1} + \dots + \frac{1}{s_k}$,

For these commutators in [23, Theorem 7.1] it is shown that under the previous conditions if $K \in H_{\mathcal{B},k}$ and $\overline{\mathcal{A}}^{-1}(t)\mathcal{B}^{-1}(t)\overline{\mathcal{C}}_{1/s}^{-1}(t) \leq t$ with $\overline{\mathcal{C}}_{1/s}(t) = e^{t^s}$, then $T_{\overline{\mathcal{B}}}$ satisfies a Coifman estimate with $M_{\overline{\mathcal{A}}}$ on the righthand side. In the case $K \in H_{\infty,k}$, the maximal operator is $M_{L(\log L)^{1/s}}$. In this way, we can extend Theorem 3.8 to the multilinear commutators: in (a) we assume $K \in H_{\mathcal{B},k}$ and replace k by $1/s$, and in (b) we assume $K \in H_{\infty,k}$ and replace k by $1/s$. The precise formulation is left to the interest reader. The proof of this result follows the same scheme, see Remark 5.2 below.

3.3. One-sided operators. In \mathbb{R} we can consider a smaller class of operators and obtain estimates for the so-called one-sided operators. These are singular integral operators with kernels supported on $(-\infty, 0)$ and we write T^+ to emphasize it. One can also consider operators T^- with kernels supported on $(0, \infty)$, for simplicity we restrict ourselves to the first type.

Let us highlight that one-sided operators are singular integral operators, therefore the previous results can be applied to them. However, exploiting the fact that they are supported on $(-\infty, 0)$ one can obtain better estimates. For instance, if $K \in H_{\infty}^*$ (indeed $K \in H_{\infty}$ suffices) then T^+ can be controlled by M on $L^p(w)$ for every $0 < p < \infty$ and $w \in A_{\infty}$, consequently T^+ is bounded on $L^p(w)$ for every $w \in A_p$, $1 < p < \infty$. These follow from the classical theory for Calderón-Zygmund operators. Moreover, exploiting the fact that the kernel of T^+ is supported on $(-\infty, 0)$ one can do better: in the Coifman estimate we can write the pointwise smaller one-sided maximal operator M^+ and consider a bigger class of weights $w \in A_{\infty}^+$; thus T^+ is bounded on $L^p(w)$ for every $w \in A_p^+$, $1 < p < \infty$ (note that we have that $A_p \subsetneq A_p^+$).

The same happens with Theorems 2.4, 3.7 and 2.6:

Theorem 3.11 ([24], [23]). *Let T^+ be a one-sided singular integral operator with kernel K supported in $(-\infty, 0)$.*

- (i) *Under the assumptions of Theorem 2.4 or 3.7, one can improve (2.2) and (3.5): A_{∞} is replaced by the bigger class of weights A_{∞}^+ , and $M_{\overline{\mathcal{A}}}$ is replaced by the pointwise smaller operator $M_{\overline{\mathcal{A}}}^+$ —in (b) of Theorem 3.7 M^{k+1} is replaced by $(M^+)^{k+1}$ —.*
- (ii) *Under the assumptions of Theorem 2.6, if the adjoint of T^+ —which is a one-sided operator with kernel supported on $(0, \infty)$ — satisfies (2.2) for all $0 < p < \infty$, $w \in A_{\infty}^-$ and with $M_{\overline{\mathcal{A}}}^- f$ on the righthand side, then, for any weight u , it follows that T^+ verifies (2.3) with $M_{\overline{\mathcal{D}}_p}^- u$ in place of $M_{\overline{\mathcal{D}}_p} u$.*

Here, we can obtain one-sided versions of Theorems 3.1 and 3.8:

Theorem 3.12. *Let T^+ be a singular integral operator with kernel K supported in $(-\infty, 0)$.*

- (i) *Under the assumptions of Theorem 3.1, then T^+ satisfies (3.1) for the pairs of weights $(u, M_{\overline{\mathcal{A}}}^- u)$ in (a), $(u, M_{\overline{\mathcal{D}}_p}^- u)$ in (b), and $(u, M_{L(\log L)^{\varepsilon}}^- u)$ in (c).*

(ii) Under the assumptions of Theorem 3.8, then $T_b^{+,k}$ —which is the k -th order commutator of T^+ — satisfies (3.6) for the pairs of weights $(u, M_{\mathcal{A}}^- u)$ in (a.1), $(u, M_{\mathcal{D}_p}^- u)$ in (a.2), and $(u, M_{L(\log L)^{k+\varepsilon}}^- u)$ in (b).

Remark 3.13. In (i) when $K \in H_\infty$ we improve the results in [1] where the stronger condition $K \in H_\infty^*$ was assumed. For example of these kernels see that reference.

We can also get an improvement of the estimates in Theorem 3.5 when we start with pairs based on one-sided maximal functions:

Theorem 3.14. Let \mathcal{F} be a Young function and assume that a given operator T satisfies (3.2) with $M_{\mathcal{F}}^-$ in place of $M_{\mathcal{F}}$. Let $1 < p < \infty$, and \mathcal{G}, \mathcal{H} as in Theorem 3.5. If (u, v) is a pair of weights such that, for all $a < b < c$ with $b - a < c - b$,

$$\|u^{1/p}\|_{\mathcal{G},(a,b)} \|v^{-1/p}\|_{L^{p'},(b,c)} \leq C,$$

we have for all $\lambda > 0$

$$u\{x \in \mathbb{R} : |Tf(x)| > \lambda\} \leq \frac{C}{\lambda^p} \int_{\mathbb{R}} |f(x)|^p v(x) dx.$$

Let us notice that here one does not need to work with one-sided operators as this abstract result does not use any property of T but the initial two-weight estimate which involves the one-sided maximal function $M_{\mathcal{F}}^-$. Notice that when applying this result T will be T^+ or $T_b^{+,k}$ from Theorem 3.12.

To prove this Theorem 3.14 we need to find sufficient conditions on (u, v) that guarantee the boundedness of $M_{\mathcal{F}}^-$ from $L^p(v)$ to $L^p(u)$. This result with $M_{\mathcal{F}}$ appears in [12] and here we extend it to the one-sided case. For convenience we state it in terms of $M_{\mathcal{F}}^+$ and to pass to $M_{\mathcal{F}}^-$ one just switches the intervals of integration in the corresponding Muckenhoupt type condition.

Theorem 3.15. Let $1 < p < \infty$ and let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be Young functions such that $\mathcal{B}^{-1}(t)\mathcal{C}^{-1}(t) \leq \mathcal{A}^{-1}(t)$, for all $t \geq t_0 > 0$, with $\mathcal{C} \in B_p$. If (u, v) is a pair of weights such that, for all $a < b < c$ with $b - a < c - b$,

$$\|u^{1/p}\|_{L^p,(a,b)} \|v^{-1/p}\|_{\mathcal{B},(b,c)} \leq C, \quad (3.7)$$

then

$$\int_{\mathbb{R}} M_{\mathcal{A}}^+ f(x)^p u(x) dx \leq C \int_{\mathbb{R}} |f(x)|^p v(x) dx.$$

4. APPLICATIONS

In this section we present some applications. As we have already observed, our results include those in [32] and [35] for Calderón-Zygmund singular integrals operators with kernels in H_∞^* . We observed before that weaker conditions on the kernels, say H_∞ for T and $K \in H_\infty \cap H_{e^{t/k}, k}$ (or, in particular, $K \in H_{\infty, k}$) for T_b^k , lead us to the same conclusions.

4.1. The differential transform operator. Consider the differential transform operator studied in [20] and [4]

$$T^+ f(x) = \sum_{j \in \mathbb{Z}} \nu_j (D_j f(x) - D_{j-1} f(x)), \quad (4.1)$$

where $\|\{\nu_j\}_j\|_\infty < \infty$ and

$$D_j f(x) = \frac{1}{2^j} \int_x^{x+2^j} f(t) dt.$$

This operator appears when studying the rate of convergence of the averages $D_j f$. Let us observe that $D_j f \rightarrow f$ a.e. when $j \rightarrow -\infty$ and $D_j f \rightarrow 0$ when $j \rightarrow \infty$. Notice that $T^+ f(x) = K * f(x)$, where

$$K(x) = \sum_{j \in \mathbb{Z}} \nu_j \left(\frac{1}{2^j} \chi_{(-2^j, 0)}(x) - \frac{1}{2^{j-1}} \chi_{(-2^{j-1}, 0)}(x) \right).$$

Observe that K is supported in $(-\infty, 0)$, and therefore T^+ is a one-sided singular integral operator (so we write T^+). In [4] it is proved that, for appropriate f ,

$$T^+ f(x) = \lim_{(N_1, N_2) \rightarrow (-\infty, \infty)} \sum_{N_1}^{N_2} \nu_j (D_j f(x) - D_{j-1} f(x)) \quad \text{for a.e. } x \in \mathbb{R}.$$

It was shown that $K \in \cap_{r \geq 1} H_r$ and so T^+ is bounded on $L^p(w)$ for all $w \in A_p^+$, $1 < p < \infty$, and maps $L^1(w)$ into $L^{1, \infty}(w)$ for all $w \in A_1^+$.

When trying to prove Coifman type estimates for T^+ , one obtains that T^+ is controlled by M_s^+ for every $1 < s < \infty$. In general $K \notin H_\infty$ (see [23] for the case $\{\nu_j\} = \{(-1)^j\}$), thus it is not clear whether one can take $s = 1$, that is, whether T^+ behaves as a one-sided singular integral operator with smooth kernel. This motivates the new Hörmander's type conditions in [24], [23]: If one shows that its kernel belongs to some class near L^∞ then one would obtain a maximal operator near M^+ . In [23] it was shown that $K \in H_{e^{t^{1/(1+\varepsilon)}}}$ for any $\varepsilon > 0$ and that $K \in H_{e^{t^{1/(1+k+\varepsilon)}}}, k$, for any $\varepsilon > 0$ and $k \geq 1$. Thus, by Theorems 2.4 and 3.7 for any $k \geq 0$, $\varepsilon > 0$, $0 < p < \infty$, and $w \in A_\infty^+$

$$\int_{\mathbb{R}} |T_b^{+,k} f(x)|^p w(x) dx \leq C \int_{\mathbb{R}} M_{L(\log L)^{k+1+\varepsilon}}^+ f(x)^p w(x) dx.$$

Applying Theorem 3.12 we obtain the following end-point estimates:

Theorem 4.1. *Let $b \in \text{BMO}$ and $k \geq 0$. Let T^+ be the differential transform operator defined above, and let $T_b^{+,k}$ be its k -th order commutator. Then, for any $\varepsilon > 0$,*

$$u\{x \in \mathbb{R} : |T_b^{+,k} f(x)| > \lambda\} \leq C \int_{\mathbb{R}} \mathcal{C}_k \left(\frac{|f(x)|}{\lambda} \right) M_{L(\log L)^{k+1+\varepsilon}}^- u(x) dx,$$

for all $\lambda > 0$.

Note that this result includes the case $k = 0$ on which $T_b^{+,0} = T^+$.

Remark 4.2. One can write the last estimate in terms of iterations of M^- since $M_{L(\log L)^{k+1+\varepsilon}}^- u(x) \leq C(M^-)^{k+3} u(x)$, for $\varepsilon > 0$ small enough. Thus, the previous estimate holds for the pair of weights $(u, (M^-)^{k+3} u)$.

Proof of Theorem 4.1. Given $k \geq 0$ and $\varepsilon > 0$ we fix $\mathcal{D}(t) = t^p (1 + \log^+ t)^{k+1+\varepsilon}$ so that $\mathcal{D}_p(t) \approx t(1 + \log^+ t)^{k+1+\varepsilon}$. We take $1 < p < 1 + \varepsilon/(2(k+2))$, $\mathcal{A}(t) \approx \exp(t^{1/(1+k+\varepsilon/(2p))}) - 1$ and $\mathcal{B}(t) \approx \exp(t^{1/(1+\varepsilon/(2p))}) - 1$. Then, as mentioned before $K \in H_B \cap H_{\mathcal{A},k}$ (note that for $k = 0$, we just have $K \in H_A$). We also notice that for $k \geq 1$ it follows that $\overline{\mathcal{A}}^{-1}(t) \mathcal{B}^{-1}(t) \overline{\mathcal{C}}_k^{-1}(t) \lesssim t$ for $t \geq 1$. Next, we pick $\mathcal{E}(t) \approx t^{p'}/(1 + \log^+ t)^{\varepsilon(p'-1)/2-(k+1)}$ and observe that our choice of p guarantees that $\varepsilon(p'-1)/2-(k+1) > 1$, therefore $\mathcal{E} \in B_{p'}$. Besides, we have $\mathcal{D}^{-1}(t) \mathcal{E}^{-1}(t) \lesssim \overline{\mathcal{A}}^{-1}(t)$ for $t \geq 1$. Then applying Theorem 3.12, that is, the one-sided version of Theorem 3.1 part (b) when $k = 0$, and the one-sided version of Theorem 3.8 part (a.2) when $k \geq 1$, we conclude the desired estimate. \square

As a corollary of Theorem 4.1, applying Theorem 3.14 we get the following weak-type estimates for general pairs of weights (u, v) :

Corollary 4.3. *Let $b \in \text{BMO}$ and $k \geq 0$. Let T^+ be the differential transform operator defined above, and let $T_b^{+,k}$ be its k -th order commutator. Then, for any $\varepsilon > 0$, if (u, v) is a pair of weights such that, for all $a < b < c$ with $b - a < c - b$,*

$$\|u^{1/p}\|_{L^p(\log L)^{(k+2)p-1+\varepsilon, (a,b)}} \|v^{-1/p}\|_{L^{p', (b,c)}} \leq C,$$

we have for all $\lambda > 0$

$$u\{x \in \mathbb{R} : |T_b^{+,k} f(x)| > \lambda\} \leq \frac{C}{\lambda^p} \int_{\mathbb{R}} |f(x)|^p v(x) dx.$$

The proof of this result follows at once from Theorem 3.14. The starting estimate is given by Theorem 4.1, so $\mathcal{F}(t) = t(1 + \log^+ t)^{k+1+\varepsilon}$ for every $\varepsilon > 0$, and we take $\mathcal{H}(t) = t^{p'}/(1 + \log^+ t)^{1+\delta} \in B_{p'}$ for any $\delta > 0$. This leads to the desired function \mathcal{G} . Details are left to the interested reader.

4.2. An example of a one-sided operator with $K \in H_\infty \cap H_{e^{t^{1/k}}, k}$. We consider the one-sided operator

$$T^+ f(x) = \sum_{j \in \mathbb{Z}} \nu_j (D_j f(x) - D_{j-1} f(x)),$$

where $\|\{\nu_j\}_j\|_\infty < \infty$ and

$$D_j f(x) = \frac{1}{2^j(1+j^2)} \int_x^{x+2^j} f(t) dt.$$

Observe that

$$K(x) = \sum_{j \in \mathbb{Z}} \nu_j \left(\frac{1}{2^j(1+j^2)} \chi_{(-2^j, 0)}(x) - \frac{1}{2^{j-1}(1+(j-1)^2)} \chi_{(-2^{j-1}, 0)}(x) \right).$$

This operator is similar to the previous one. In [23] was proved that $K \in H_\infty \cap H_{e^{t^{1/k}}, k}$. Thus, by Theorems 2.4 and 3.7 for each $k \geq 0$, $0 < p < \infty$, and $w \in A_\infty^+$

$$\int_{\mathbb{R}} |T_b^{+,k} f(x)|^p w(x) dx \leq C \int_{\mathbb{R}} M_{L(\log L)^k}^+ f(x)^p w(x) dx. \quad (4.2)$$

Note that in the righthand side one can alternatively write $(M^+)^{k+1} f$ as $(M^+)^{k+1} f \approx M_{L(\log L)^k}^+ f$ a.e.. We apply Theorem 3.12, that is, when $k = 0$ we use the one-sided

version of Theorem 3.1 part (c), and when $k \geq 1$ we employ the one-sided version of Theorem 3.8 part (b). Thus, we conclude the following end-point estimates: given $b \in \text{BMO}$, for every $k \geq 0$ and for any $\varepsilon > 0$

$$u\{x \in \mathbb{R} : |T_b^{+,k} f(x)| > \lambda\} \leq C \int_{\mathbb{R}} \mathcal{C}_k \left(\frac{|f(x)|}{\lambda} \right) M_{L(\log L)^{k+\varepsilon}}^- u(x) dx. \quad (4.3)$$

Note that taking $\varepsilon > 0$ small enough, $M_{L(\log L)^{k+\varepsilon}}^- u(x) \leq C(M^-)^{k+2} u(x)$.

Remark 4.4. In terms of iterations of the one-sided Hardy-Littlewood maximal function, notice that in (4.2) we have $k + 1$ iterations and in (4.3) we have $k + 2$, so we obtain an extra iteration. This is because in (c) of Theorem 3.1, in (b) of Theorem 3.8 and in their corresponding versions for one-sided operators we lose a small power of the logarithm. This happens also with Calderón-Zygmund operators with smooth kernel as the Hilbert and Riesz transforms: they are controlled, in the sense of Coifman, by M , but the end-point estimate holds for the pair of weights $(u, M^2 u)$ —indeed one can write $(u, M_{L(\log L)^\varepsilon} u)$ for any $\varepsilon > 0$ —. It is not known, even for the Hilbert and Riesz transforms, whether the pair of weights (u, Mu) is valid for the corresponding weak-type estimate.

Notice that in the case of the differential transform operator in both the Coifman inequality and the end-point estimate the number of iterations for the k -th order commutator is $k + 3$. This happens as we already have a small power of the logarithm floating around.

From Theorem 3.14 proceeding as in Corollary 4.3 we obtain the following two-weight weak-type estimates: given $b \in \text{BMO}$, for every $k \geq 0$ and for any $\varepsilon > 0$ if (u, v) is a pair of weights such that, for all $a < b < c$ with $b - a < c - b$,

$$\|u^{1/p}\|_{L^p(\log L)^{(k+1)p-1+\varepsilon},(a,b)} \|v^{-1/p}\|_{L^{p'},(b,c)} \leq C,$$

then $T_b^{+,k} f$ maps $L^p(v)$ into $L^{p,\infty}(u)$. This extends the sharp results obtained in [12] for Calderón-Zygmund operators with smooth kernels to the setting of one-sided operators.

4.3. Multipliers. Let $m \in L^\infty(\mathbb{R}^n)$ and consider the multiplier operator T defined *a priori* for f in the Schwartz class by $\widehat{Tf}(\xi) = m(\xi) \widehat{f}(\xi)$. Given $1 < s \leq 2$ and $0 \leq l \in \mathbb{N}$ we say that $m \in M(s, l)$ if

$$\sup_{R>0} R^{|\alpha|} \|D^\alpha m\|_{L^s, |\xi| \sim R} < +\infty, \quad \text{for all } |\alpha| \leq l.$$

In [23] it was proved the following: let $m \in M(s, l)$, with $1 < s \leq 2$, $0 \leq l \leq n$ and $l > n/s$. Then for all $k \geq 0$ and any $\varepsilon > 0$ we have that for all $0 < p < \infty$ and $w \in A_\infty$,

$$\int_{\mathbb{R}^n} |T_b^k f(x)|^p w(x) dx \leq C \int_{\mathbb{R}^n} M_{n/l+\varepsilon} f(x)^p w(x) dx. \quad (4.4)$$

The proof of such estimates consists in obtaining that a family of truncations of the kernel $\{K^N\}_N$ are uniformly in $H_{L^r(\log L)^{kr},k}$ with $r' = n/l + \varepsilon$. Thus, taking $\mathcal{A}(t) = t^r$, $\mathcal{B}(t) = t^r (1 + \log^+ t)^{kr}$ we have $K^N \in H_{\mathcal{B}} \cap H_{\mathcal{A},k}$ (this follows easily from $K^N \in H_{L^r(\log L)^{kr},k}$). Notice that $\overline{\mathcal{A}}^{-1}(t) \mathcal{B}^{-1}(t) \overline{\mathcal{C}}_k^{-1}(t) \lesssim t$ for $t \geq 1$ and therefore (4.4) follows from Theorem 3.7 for the k -th order commutators of T^N (which is the

operator whose kernel is K^N) with constants that are independent of N . A standard approximation argument leads to the desired estimate for T_b^k . We refer the reader to [23] for more details.

The same argument allows us to apply Theorem 3.1 part (a) and Theorem 3.8 part (a.1) to T^N . Observe that $\overline{\mathcal{A}}(t) = t^{r'}$, then choosing $1 < s < r'$ we obtain $\liminf_{t \rightarrow \infty} \overline{\mathcal{A}}(t)/t^s = +\infty$. Therefore, taking limits we have the following result:

Theorem 4.5. *Let $m \in M(s, l)$ with $1 < s \leq 2$, $0 \leq l \leq n$ and $l > n/s$. Then for all $k \geq 0$ and any $\varepsilon > 0$ we have*

$$u\{x \in \mathbb{R}^n : |T_b^k f(x)| > \lambda\} \leq C \int_{\mathbb{R}^n} \mathcal{C}_k \left(\frac{|f(x)|}{\lambda} \right) M_{n/l+\varepsilon} u(x) dx.$$

From this estimate one can obtain weak-type estimates for general pairs of weights by using Theorem 3.5. The precise statements are left to the reader.

4.4. Kernels related to H_r and M_{L^r} . Implicit in [39] (see also [22], [43]) and as it was observed in [26] when $K \in H_{L^r}$, that is, when the kernel satisfies the L^r -Hörmander condition, then one obtains that T is controlled by $M_{L^{r'}}$. In [23] different extensions of that inequality for the higher order commutators were considered. Following the notation of Theorem 3.7 these are the different conditions and maximal operators obtained: for every $1 < r < \infty$ and $k \geq 0$, we have

$H_{\mathcal{B},k}$	$H_{\mathcal{B}} \cap H_{\mathcal{A},k}$	$M_{\overline{\mathcal{A}}} f$
$H_{L^r,k}$	$H_{L^r} \cap H_{L^r(\log L)^{-kr},k}$	$M_{L^{r'}(\log L)^{kr'}} f$
$H_{L^r(\log L)^{kr},k}$	$H_{L^r(\log L)^{kr}} \cap H_{L^r,k}$	$M_{L^{r'}} f$
$H_{L^r(\log L)^k,k}$	$H_{L^r(\log L)^k} \cap H_{L^r(\log L)^{-k(r-1),k}$	$M_{L^{r'}(\log L)^k} f \approx (M_{L^{r'}})^{k+1}$

TABLE 1. Examples of different H_r -conditions

Thus, applying Theorem 3.1 part (a) and Theorem 3.8 part (a.1) we obtain that T_b^k satisfies (3.6) with the different pairs of weights $(u, M_{\overline{\mathcal{A}}} u)$ in the previous table.

4.5. Homogeneous Singular Integrals. Denote by $\Sigma = \Sigma_{n-1}$ the unit sphere on \mathbb{R}^n . For $x \neq 0$, we write $x' = x/|x|$. Let us consider $\Omega \in L^1(\Sigma)$. This function can be extended to $\mathbb{R}^n \setminus \{0\}$ as $\Omega(x) = \Omega(x')$ (abusing on the notation we call both functions Ω). Thus Ω is a function homogeneous of degree 0. We assume that $\int_{\Sigma} \Omega(x') d\sigma(x') = 0$. Set $K(x) = \Omega(x)/|x|^n$ and let T be the operator associated with the kernel K .

Given a Young function \mathcal{A} we define the $L^{\mathcal{A}}$ -modulus of continuity of Ω as

$$\varpi_{\mathcal{A}}(t) = \sup_{|y| \leq t} \|\Omega(\cdot + y) - \Omega(\cdot)\|_{\mathcal{A}, \Sigma}.$$

Given $\Omega \in L^{\mathcal{B}}(\Sigma)$ and T be as above. Let $k \geq 0$ and \mathcal{A}, \mathcal{B} be Young functions such that $\overline{\mathcal{A}}^{-1}(t) \mathcal{B}^{-1}(t) \overline{\mathcal{C}}_k^{-1}(t) \leq t$ for all $t \geq 1$. If

$$\int_0^1 \varpi_{\mathcal{B}}(t) \frac{dt}{t} + \int_0^1 \left(1 + \log \frac{1}{t}\right)^k \varpi_{\mathcal{A}}(t) \frac{dt}{t} < \infty,$$

then it was proved in [23] that $K \in H_{\mathcal{B}} \cap H_{\mathcal{A},k}$ and therefore

$$\int_{\mathbb{R}^n} |T_b^k f(x)|^p w(x) dx \leq C \int_{\mathbb{R}^n} M_{\overline{\mathcal{A}}} f(x)^p w(x) dx,$$

for every $0 < p < \infty$ and $w \in A_{\infty}$.

Once it is known that $K \in H_{\mathcal{B}} \cap H_{\mathcal{A},k}$ one can apply Theorems 3.1 and 3.8 to derive the corresponding two-weight end-point estimates. The precise statements and further details are left to the interested reader.

5. PROOFS OF THE MAIN RESULTS

Proof of Theorem 3.1. Without loss of generality we can assume that u is bounded and has compact support (otherwise we prove the corresponding estimate for $u_N = \min\{u, N\} \chi_{B(0,N)}$ with bounds independent of N and apply the monotone convergence theorem). We assume that $0 \leq f \in L_c^\infty(\mathbb{R}^n)$ and consider the standard Calderón-Zygmund decomposition of f at level λ : there exists a collection of maximal (and so disjoint) dyadic cubes $\{Q_j\}_j$ (with center x_j and sidelength $2r_j$) such that

$$\lambda < \frac{1}{|Q_j|} \int_{Q_j} f \leq 2^n \lambda. \quad (5.1)$$

We write $f = g + h$ where

$$g = f \chi_{\mathbb{R}^n \setminus \cup_j Q_j} + \sum_j f_{Q_j} \chi_{Q_j}, \quad h = \sum_j h_j = \sum_j (f - f_{Q_j}) \chi_{Q_j}$$

where f_{Q_j} denotes the average of f over Q_j . Let us recall that $0 \leq g(x) \leq 2^n \lambda$ a.e. and also that each h_j has vanishing integral. We set $\tilde{Q}_j = 2Q_j$, $\tilde{\Omega} = \cup_j \tilde{Q}_j$, and $\tilde{u} = u \chi_{\mathbb{R}^n \setminus \tilde{\Omega}}$. Then,

$$\begin{aligned} u\{x \in \mathbb{R}^n : |Tf(x)| > \lambda\} &\leq u(\tilde{\Omega}) + u\{x \in \mathbb{R}^n \setminus \tilde{\Omega} : |Th(x)| > \lambda/2\} \\ &\quad + u\{x \in \mathbb{R}^n \setminus \tilde{\Omega} : |Tg(x)| > \lambda/2\} \\ &= I + II + III. \end{aligned}$$

We estimate each term separately. The estimates for I and II are obtained in the same way in the three cases (a), (b) and (c). We show that

$$I \lesssim \frac{1}{\lambda} \int_{\mathbb{R}^n} f(x) M u(x) dx, \quad II \lesssim \frac{1}{\lambda} \int_{\mathbb{R}^n} f(x) M_{\overline{\mathcal{A}}} u(x) dx, \quad (5.2)$$

where, in case (c), as $K \in H_{\infty} = H_{L^{\infty}}$ it is understood that $\overline{\mathcal{A}}(t) = t$ so $M_{\overline{\mathcal{A}}} = M_{L^1} = M$. Let us observe that both estimates lead us to the desired conclusions in the three cases (a), (b) and (c). Regarding I , Mu is controlled by $M_{\overline{\mathcal{A}}}u$ in (a) —as $\overline{\mathcal{A}}$ is a Young function—, by $M_{\mathcal{D}_p}u$ in (b) —as we pointed out in Remark 3.3 that $\mathcal{D}^{-1}(t) \lesssim t^{1/p}$ for $t \geq 1$ which yields $\mathcal{D}_p(t) \geq t$ for $t \geq 1$ — and by $M_{L(\log L)^\varepsilon}u$ in (c). For II , $M_{\overline{\mathcal{A}}}u$ is the desired weight in (a); in (b) we observed in Remark 3.3 that $M_{\overline{\mathcal{A}}}u \lesssim M_{\mathcal{D}_p}u$; and in (c) we have $M_{\overline{\mathcal{A}}}u = Mu \leq M_{L(\log L)^\varepsilon}u$.

Let us show the first estimate in (5.2). By (5.1) we have

$$I = u(\cup_j \tilde{Q}_j) \leq \sum_j u(\tilde{Q}_j) = 2^n \sum_j \frac{u(\tilde{Q}_j)}{|\tilde{Q}_j|} |\tilde{Q}_j| \leq \frac{2^n}{\lambda} \sum_j \frac{u(\tilde{Q}_j)}{|\tilde{Q}_j|} \int_{\tilde{Q}_j} f(x) dx$$

$$\leq \frac{2^n}{\lambda} \sum_j \int_{Q_j} f(x) M u(x) dx = \frac{2^n}{\lambda} \int_{\mathbb{R}^n} f(x) M u(x) dx.$$

Next, we estimate II : as the functions h_j has vanishing integral

$$\begin{aligned} II &= u \left\{ x \in \mathbb{R}^n \setminus \tilde{\Omega} : \left| \sum_j T h_j(x) \right| > \lambda/2 \right\} \leq \frac{2}{\lambda} \sum_j \int_{\mathbb{R}^n \setminus \tilde{\Omega}} |T h_j(x)| u(x) dx \\ &\leq \frac{2}{\lambda} \sum_j \int_{\mathbb{R}^n \setminus \tilde{\Omega}} \left| \int_{Q_j} (K(x-y) - K(x-x_{Q_j})) h_j(y) dy \right| u(x) dx \\ &\leq \frac{2}{\lambda} \sum_j \int_{Q_j} |h_j(y)| \int_{\mathbb{R}^n \setminus \tilde{Q}_j} |K(x-y) - K(x-x_{Q_j})| u(x) dx dy. \end{aligned}$$

We claim that for every $y \in Q_j$ we have

$$\int_{\mathbb{R}^n \setminus \tilde{Q}_j} |K(x-y) - K(x-x_{Q_j})| u(x) dx \lesssim \operatorname{ess\,inf}_{Q_j} M_{\bar{\mathcal{A}}} u. \quad (5.3)$$

This estimate drives us to

$$\begin{aligned} II &\lesssim \frac{1}{\lambda} \sum_j \operatorname{ess\,inf}_{Q_j} M_{\bar{\mathcal{A}}} u \int_{Q_j} |h_j(y)| dy \lesssim \frac{1}{\lambda} \sum_j \operatorname{ess\,inf}_{Q_j} M_{\bar{\mathcal{A}}} u \int_{Q_j} f(y) dy \\ &\leq \frac{1}{\lambda} \sum_j \int_{Q_j} f(y) M_{\bar{\mathcal{A}}} u(y) dy \leq \frac{1}{\lambda} \int_{\mathbb{R}^n} f(y) M_{\bar{\mathcal{A}}} u(y) dy. \end{aligned}$$

We obtain (5.3): using the generalized Hölder's inequality for \mathcal{A} and $\bar{\mathcal{A}}$ (when $K \in H_\infty$ we understand that $\bar{\mathcal{A}}(t) = t$ and so we have the corresponding $L^1 - L^\infty$ Hölder's estimate)

$$\begin{aligned} &\int_{\mathbb{R}^n \setminus \tilde{Q}_j} |K(x-y) - K(x-x_{Q_j})| u(x) dx \\ &\leq \sum_{k=1}^{\infty} \int_{|x-x_{Q_j}| \sim 2^k r_j} |K(x-y) - K(x-x_{Q_j})| u(x) dx \\ &\lesssim \sum_{k=1}^{\infty} (2^k r_j)^n \|K(\cdot - y) - K(\cdot - x_{Q_j})\|_{\mathcal{A}, |x-x_{Q_j}| \sim 2^k r_j} \|u\|_{\bar{\mathcal{A}}, |x-x_{Q_j}| \leq 2^{k+1} r_j} \\ &\leq C \operatorname{ess\,inf}_{Q_j} M_{\bar{\mathcal{A}}} u, \end{aligned}$$

where in the last estimate we have used that $K \in H_{\mathcal{A}}$.

To complete the proof, it remains to estimate III . Here, the proof changes in each of the cases. We start with (a). As $\liminf_{t \rightarrow \infty} \bar{\mathcal{A}}(t)/t^r > 0$ then there exists $c = c_r$ such that $\bar{\mathcal{A}}(t) \geq c t^r$ for every $t \geq 1$. On the other hand, using that $\bar{\mathcal{A}} \in \Delta_2$ there exist $1 < s < \infty$ (indeed we can take $s > r$) such that $\bar{\mathcal{A}}(t) \leq C t^s$ for every $t \geq 1$ (this follows by iterating the Δ_2 -condition). Then, taking $p > s$ we have

$$\begin{aligned} III &= u \{ x \in \mathbb{R}^n \setminus \tilde{\Omega} : |Tg(x)| > \lambda/2 \} \leq \frac{2^p}{\lambda^p} \int_{\mathbb{R}^n} |Tg(x)|^p \tilde{u}(x) dx \\ &\leq \frac{2^p}{\lambda^p} \int_{\mathbb{R}^n} |Tg(x)|^p M_r \tilde{u}(x) dx \lesssim \frac{1}{\lambda^p} \int_{\mathbb{R}^n} M_{\bar{\mathcal{A}}} g(x)^p M_r \tilde{u}(x) dx, \quad (5.4) \end{aligned}$$

where in the last inequality we have used Theorem 2.4 and the fact that $M_r \tilde{u} \in A_1 \subset A_\infty$ as $r > 1$. Notice that one has to check that the left-hand side of (2.2) is finite. Indeed, as we have assumed that $u \in L^\infty$ we have

$$\begin{aligned} \int_{\mathbb{R}^n} |Tg(x)|^p M_r \tilde{u}(x) dx &\leq \|u\|_{L^\infty} \int_{\mathbb{R}^n} |Tg(x)|^p dx \lesssim \|u\|_{L^\infty} \int_{\mathbb{R}^n} g(x)^p dx \\ &\lesssim \|u\|_{L^\infty} \lambda^{p-1} \int_{\mathbb{R}^n} g(x) dx = \|u\|_{L^\infty} \lambda^{p-1} \int_{\mathbb{R}^n} f(x) dx < \infty, \end{aligned}$$

where we have used that T is bounded on $L^p(\mathbb{R}^n)$ as $K \in H_{\mathcal{A}} \subset H_1$; and also that f and u are bounded with compact support. We can continue with the estimate of *III*: as $\bar{\mathcal{A}}(t) \leq C t^s$ for every $t \geq 1$ it follows that

$$\begin{aligned} III &\lesssim \frac{1}{\lambda^p} \int_{\mathbb{R}^n} M_s g(x)^p M_r \tilde{u}(x) dx = \frac{1}{\lambda^p} \int_{\mathbb{R}^n} M(g^s)(x)^{p/s} M_r \tilde{u}(x) dx \\ &\lesssim \frac{1}{\lambda^p} \int_{\mathbb{R}^n} g(x)^p M_r \tilde{u}(x) dx \lesssim \frac{1}{\lambda^p} \int_{\mathbb{R}^n} g(x)^p M_{\bar{\mathcal{A}}} \tilde{u}(x) dx, \end{aligned} \quad (5.5)$$

where we have used that $M_r \tilde{u} \in A_1$, therefore M is bounded on $L^{p/s}(M_r \tilde{u})$ and also that $\bar{\mathcal{A}}(t) \geq c t^r$ for every $t \geq 1$. We claim that

$$\int_{\cup_j Q_j} g(x) M_{\bar{\mathcal{A}}} \tilde{u}(x) dx \lesssim \int_{\cup_j Q_j} f(x) M_{\bar{\mathcal{A}}} \tilde{u}(x) dx. \quad (5.6)$$

From (5.5), this estimate and the fact that $0 \leq g(x) \leq 2^n \lambda$ a.e. yield

$$\begin{aligned} III &\lesssim \frac{1}{\lambda} \int_{\mathbb{R}^n} g(x) M_{\bar{\mathcal{A}}} \tilde{u}(x) dx = \frac{1}{\lambda} \int_{\mathbb{R}^n \setminus \cup_j Q_j} f(x) M_{\bar{\mathcal{A}}} \tilde{u}(x) dx + \frac{1}{\lambda} \int_{\cup_j Q_j} g(x) M_{\bar{\mathcal{A}}} \tilde{u}(x) dx \\ &\lesssim \frac{1}{\lambda} \int_{\mathbb{R}^n} f(x) M_{\bar{\mathcal{A}}} \tilde{u}(x) dx \leq \frac{1}{\lambda} \int_{\mathbb{R}^n} f(x) M_{\bar{\mathcal{A}}} u(x) dx, \end{aligned}$$

which is the desired estimate for *III*.

To complete the proof of (a) we need to show (5.6). We first obtain that for any Young function \mathcal{C} , any weight v with $M_{\mathcal{C}} v < \infty$ a.e, and any cube Q we have

$$M_{\mathcal{C}}(v \chi_{\mathbb{R}^n \setminus 2Q})(y) \approx \operatorname{ess\,inf}_{z \in Q} M_{\mathcal{C}}(v \chi_{\mathbb{R}^n \setminus 2Q})(z), \quad \text{a.e } y \in Q. \quad (5.7)$$

Let $y \in Q$ and R be any cube such that $y \in R$. If $R \setminus 2Q = \emptyset$ then $\|v \chi_{\mathbb{R}^n \setminus 2Q}\|_{\mathcal{C}, R} = 0$. Otherwise, we have that $\ell(R) > \ell(Q)/2$ which implies that $Q \subset 5R$. Then,

$$\|v \chi_{\mathbb{R}^n \setminus 2Q}\|_{\mathcal{C}, R} \lesssim \|v \chi_{\mathbb{R}^n \setminus 2Q}\|_{\mathcal{C}, 5R} \leq \operatorname{ess\,inf}_{z \in Q} M_{\mathcal{C}}(v \chi_{\mathbb{R}^n \setminus 2Q})(z),$$

and taking the supremum over all the cubes $R \ni y$ we conclude the desired estimate. Next, we use (5.7) to obtain (5.6):

$$\begin{aligned} \int_{\cup_j Q_j} g(x) M_{\bar{\mathcal{A}}} \tilde{u}(x) dx &= \sum_j \int_{Q_j} g(x) M_{\bar{\mathcal{A}}} \tilde{u}(x) dx = \sum_j f_{Q_j} \int_{Q_j} M_{\bar{\mathcal{A}}}(\tilde{u} \chi_{\mathbb{R}^n \setminus 2Q_j})(x) dx \\ &\lesssim \sum_j \int_{Q_j} f(x) dx \operatorname{ess\,inf}_{z \in Q_j} M_{\bar{\mathcal{A}}}(\tilde{u} \chi_{\mathbb{R}^n \setminus 2Q_j})(z) \leq \sum_j \int_{Q_j} f(x) M_{\bar{\mathcal{A}}}(\tilde{u} \chi_{\mathbb{R}^n \setminus 2Q_j})(x) dx \\ &= \int_{\cup_j Q_j} f(x) M_{\bar{\mathcal{A}}} \tilde{u}(x) dx. \end{aligned}$$

This completes the proof of (a).

To show (b), we only have to estimate *III*. The argument is very similar, the main change consists of proving (5.5) with \mathcal{D}_p in place of $\overline{\mathcal{A}}$. Once we have that, the argument presented above adapts trivially to the present situation and so the desired estimate for *III* follows. Note that our hypotheses guarantee that we can apply Theorem 2.4 to the adjoint of T —let us observe that $T^* = \tilde{T}$ where \tilde{T} is the singular operator with kernel $\tilde{K}(x) = K(-x) \in H_{\mathcal{A}}$ —and then Theorem 2.6 yields

$$\begin{aligned} III &= u\{x \in \mathbb{R}^n \setminus \tilde{\Omega} : |Tg(x)| > \lambda/2\} \leq \frac{2^p}{\lambda^p} \int_{\mathbb{R}^n} |Tg(x)|^p \tilde{u}(x) dx \\ &\lesssim \frac{1}{\lambda^p} \int_{\mathbb{R}^n} g(x)^p M_{\mathcal{D}_p} \tilde{u}(x) dx. \end{aligned} \quad (5.8)$$

As just mentioned, the ideas used before applied straightforward and the desired estimate follows at once.

Finally, we show (c). Given $\varepsilon > 0$ we pick $p > 1$ and $\delta > 0$ so that $p - 1 + \delta = \varepsilon$ (note that p is taken very close to 1 and δ very small). Then, by Remark 2.7, (2.3) holds for the pair of weights $(u, M_{L(\log L)^\varepsilon} u)$. Then, the previous case *mutatis mutandis* leads us to the desired estimate.

Remark 5.1. There is another argument to derive (c): Given $\varepsilon > 0$ we pick $p > 1$ and $\delta > 0$, so that $p - 1 + 2\delta = \varepsilon$. Let $\overline{\mathcal{A}}(t) = t(1 + \log^+ t)^{\delta/p}$. Note that $H_\infty \subset H_{\mathcal{A}}$ and so $K \in H_{\mathcal{A}}$. We take $\mathcal{D}(t) = t^p(1 + \log^+ t)^{p-1+2\delta}$ and $\mathcal{E}(t) \approx t^{p'}/(1 + \log^+ t)^{1+\delta(p'-1)} \in B_{p'}$. Then, we can apply (b) to obtain the desired estimate for the pair of weights $(u, M_{\mathcal{D}_p} u)$. To conclude we observe that $\mathcal{D}_p(t) = \mathcal{D}(t^{1/p}) = t(1 + \log^+ t)^\varepsilon$.

□

Proof of Theorem 3.8. The argument follows the scheme of the proof Theorem 3.1, which corresponds to the case $k = 0$, and we only give the main changes. We proceed by induction to obtain (a). The proof of (b) follows as in Theorem 3.1 from (a.2) by a suitable choice of \mathcal{A} and \mathcal{B} (see Remark 5.1).

We assume that the cases $m = 0, 1, \dots, k-1$ are proved and we show the desired estimate for T_b^k . Thus, we fix a weight $u \in L_c^\infty$ and $0 \leq f \in L_c^\infty$. By homogeneity we can also assume that $\|b\|_{\text{BMO}} = 1$.

We recall some properties of BMO to be used later. Given $b \in \text{BMO}$, a cube Q , $j \geq 0$ and $q > 0$, by John-Nirenberg's theorem we have

$$\|(b - b_Q)^j\|_{L^q, Q} \leq \|(b - b_Q)^j\|_{\tilde{\mathcal{C}}_k, Q} = \|b - b_Q\|_{\text{exp } L, Q}^j \leq C \|b\|_{\text{BMO}}^j. \quad (5.9)$$

On the other hand, for every $l \geq 1$ and $b \in \text{BMO}$, we have

$$|b_Q - b_{2^l Q}| \leq \sum_{m=1}^l |b_{2^{m-1} Q} - b_{2^m Q}| \leq 2^n \sum_{m=1}^l \|b - b_{2^m Q}\|_{L^{1, 2^m} Q} \leq 2^n l \|b\|_{\text{BMO}}. \quad (5.10)$$

We perform the Calderón-Zygmund decomposition of f at level λ . Let $g, h = \sum_j h_j$, $Q_j, \tilde{Q}_j, \tilde{\Omega}$ and \tilde{u} be as in the proof of Theorem 3.1. Then,

$$\begin{aligned} u\{x \in \mathbb{R}^n : |T_b^k f(x)| > \lambda\} &\leq u(\tilde{\Omega}) + u\{x \in \mathbb{R}^n \setminus \tilde{\Omega} : |T_b^k h(x)| > \lambda/2\} \\ &\quad + u\{x \in \mathbb{R}^n \setminus \tilde{\Omega} : |T_b^k g(x)| > \lambda/2\} \end{aligned}$$

$$= I + II + III,$$

and we estimate each term separately. For I we obtain the first estimate in (5.2) exactly as before. Then,

$$I \lesssim \frac{1}{\lambda} \int_{\mathbb{R}^n} f(x) Mu(x) dx \leq \int_{\mathbb{R}^n} \mathcal{C}_k \left(\frac{|f(x)|}{\lambda} \right) Mu(x) dx$$

and we observe that Mu is pointwise controlled by either $M_{\overline{\mathcal{A}}}u$, $M_{\mathcal{D}_p}u$ or $M_{L(\log L)^{k+\varepsilon}}u$. So the desired estimate follows in each of the cases.

Next, we estimate II by using the induction hypothesis and the conditions assumed on the kernel. As in [33] we can write

$$\begin{aligned} T_b^k h(x) &= \sum_j T_b^k h_j(x) = \sum_{m=0}^{k-1} C_{k,m} T_b^m \left(\sum_j (b - b_{Q_j})^{k-m} h_j \right)(x) \\ &\quad + \sum_j (b(x) - b_{Q_j})^k T h_j(x) = F_1(x) + F_2(x), \end{aligned} \quad (5.11)$$

and we estimate each function in turn.

For F_1 we would like to use the induction hypothesis. We start with (a.1). If $0 \leq m \leq k-1$ then $H_{A,k} \subset H_{A,m}$ and so $K \in H_{\mathcal{B}} \cap H_{A,m}$. Also, as $\overline{\mathcal{C}}_k(t) \leq \overline{\mathcal{C}}_m(t)$ we have

$$\overline{\mathcal{A}}^{-1}(t) \mathcal{B}^{-1}(t) \overline{\mathcal{C}}_m^{-1}(t) \leq \overline{\mathcal{A}}^{-1}(t) \mathcal{B}^{-1}(t) \overline{\mathcal{C}}_k^{-1}(t) \leq t.$$

Thus the hypotheses on (a) are satisfied for every $0 \leq m \leq k-1$ and therefore

$$\begin{aligned} u\{x \in \mathbb{R}^n \setminus \tilde{\Omega} : |F_1(x)| > \lambda/4\} &\leq \sum_{m=0}^{k-1} \tilde{u}\{x : |T_b^m \left(\sum_j (b - b_{Q_j})^{k-m} h_j \right)(x)| > \lambda/C\} \\ &\lesssim \sum_{m=0}^{k-1} \int_{\mathbb{R}^n} \mathcal{C}_m \left(\frac{|\sum_j (b - b_{Q_j})^{k-m} h_j|}{\lambda} \right) M_{\overline{\mathcal{A}}}\tilde{u} dx \\ &\lesssim \sum_{m=0}^{k-1} \sum_j \int_{Q_j} \mathcal{C}_m \left(\frac{|b - b_{Q_j}|^{k-m} |h_j|}{\lambda} \right) M_{\overline{\mathcal{A}}}\tilde{u} dx \\ &\lesssim \sum_{m=0}^{k-1} \sum_j \operatorname{ess\,inf}_{Q_j} M_{\overline{\mathcal{A}}}\tilde{u} \int_{Q_j} \mathcal{C}_m \left(\frac{|b - b_{Q_j}|^{k-m} |h_j|}{\lambda} \right) dx, \end{aligned}$$

where in the last estimate we have used (5.7). Let us observe that $\mathcal{C}_k^{-1}(t) \overline{\mathcal{C}}_{k-m}^{-1}(t) \lesssim \mathcal{C}_m^{-1}(t)$. Then, Young's inequality implies

$$\begin{aligned} \int_{Q_j} \mathcal{C}_m \left(\frac{|b - b_{Q_j}|^{k-m} |h_j|}{\lambda} \right) dx &\lesssim \int_{Q_j} \mathcal{C}_k \left(\frac{|h_j|}{c\lambda} \right) dx + \int_{Q_j} \overline{\mathcal{C}}_{k-m}(c|b - b_{Q_j}|^{k-m}) dx \\ &= \int_{Q_j} \mathcal{C}_k \left(\frac{|h_j|}{\lambda} \right) dx + \int_{Q_j} e^{c|b - b_{Q_j}|} dx \lesssim \int_{Q_j} \mathcal{C}_k \left(\frac{|h_j|}{\lambda} \right) dx + |Q_j| \end{aligned} \quad (5.12)$$

as $\|b\|_{\text{BMO}} = 1$ implies, by John-Nirenberg's theorem, that $\|b - b_{Q_j}\|_{\exp L, Q_j} \leq c$. Besides, using that $\mathcal{C}_k(t)^\delta$, $0 < \delta < 1$, is concave and so subadditive it follows that \mathcal{C}_k

is quasi-subadditive —that is, $\mathcal{C}_k(t_1 + t_2) \lesssim \mathcal{C}_k(t_1) + \mathcal{C}_k(t_2)$ —. Therefore, by Jensen's inequality for \mathcal{C}_k

$$\int_{Q_j} \mathcal{C}_k \left(\frac{|h_j|}{\lambda} \right) dx \leq \int_{Q_j} \mathcal{C}_k \left(\frac{f}{\lambda} \right) dx + |Q_j| \mathcal{C}_k \left(\frac{f_{Q_j}}{\lambda} \right) \leq 2 \int_{Q_j} \mathcal{C}_k \left(\frac{f}{\lambda} \right) dx.$$

Also, (5.1) implies

$$|Q_j| \leq \frac{1}{\lambda} \int_{Q_j} f dx \leq \int_{Q_j} \mathcal{C}_k \left(\frac{f}{\lambda} \right) dx.$$

Plugging these estimates into (5.12) we obtain

$$\begin{aligned} u\{x \in \mathbb{R}^n \setminus \tilde{\Omega} : |F_1(x)| > \lambda/4\} &\lesssim \sum_{m=0}^{k-1} \sum_j \operatorname{ess\,inf}_{Q_j} M_{\overline{\mathcal{A}}}\tilde{u} \int_{Q_j} \mathcal{C}_k \left(\frac{f}{\lambda} \right) dx \\ &\lesssim \sum_j \int_{Q_j} \mathcal{C}_k \left(\frac{f}{\lambda} \right) M_{\overline{\mathcal{A}}}\tilde{u} dx \leq \int_{\mathbb{R}^n} \mathcal{C}_k \left(\frac{f}{\lambda} \right) M_{\overline{\mathcal{A}}}u dx. \end{aligned}$$

This gives the desired estimate for F_1 in case (a.1). Notice that the same computations hold in case (a.2) replacing everywhere $M_{\overline{\mathcal{A}}}$ by $M_{\mathcal{D}_p}$.

Next, we estimate F_2 :

$$\begin{aligned} u\{x \in \mathbb{R}^n \setminus \tilde{\Omega} : |F_2(x)| > \lambda/4\} &\leq \frac{4}{\lambda} \sum_j \int_{\mathbb{R}^n \setminus \tilde{\Omega}} |b(x) - b_{Q_j}|^k |Th_j(x)| u(x) dx \\ &\leq \frac{4}{\lambda} \sum_j \int_{\mathbb{R}^n \setminus \tilde{\Omega}} |b(x) - b_{Q_j}|^k \int_{Q_j} |K(x-y) - K(x-x_{Q_j})| |h_j(y)| dy u(x) dx \\ &\leq \frac{4}{\lambda} \sum_j \int_{Q_j} |h_j(y)| \int_{\mathbb{R}^n \setminus \tilde{Q}_j} |K(x-y) - K(x-x_{Q_j})| |b(x) - b_{Q_j}|^k u(x) dx dy. \end{aligned}$$

We claim that for every cube Q (whose center is x_Q) and for every $y \in Q$ we have

$$\int_{\mathbb{R}^n \setminus 2Q} |K(x-y) - K(x-x_Q)| |b(x) - b_Q|^k u(x) dx \lesssim \operatorname{ess\,inf}_Q M_{\overline{\mathcal{A}}}u. \quad (5.13)$$

This estimate applied to each Q_j implies

$$\begin{aligned} u\{x \in \mathbb{R}^n \setminus \tilde{\Omega} : |F_2(x)| > \lambda/4\} &\lesssim \frac{1}{\lambda} \sum_j \operatorname{ess\,inf}_{Q_j} M_{\overline{\mathcal{A}}}u \int_{Q_j} |h_j(y)| dy \\ &\lesssim \frac{1}{\lambda} \sum_j \operatorname{ess\,inf}_{Q_j} M_{\overline{\mathcal{A}}}u \int_{Q_j} f(y) dy \leq \frac{1}{\lambda} \sum_j \int_{Q_j} f(y) M_{\overline{\mathcal{A}}}u(y) dy \\ &\leq \frac{1}{\lambda} \int_{\mathbb{R}^n} f(y) M_{\overline{\mathcal{A}}}u(y) dy \leq \int_{\mathbb{R}^n} \mathcal{C}_k \left(\frac{|f(x)|}{\lambda} \right) M_{\overline{\mathcal{A}}}u(y) dy. \end{aligned}$$

Note that this leads to the desired estimate in (a.1) and also in (a.2) (we observed in Remark 3.3 that $M_{\overline{\mathcal{A}}}u \lesssim M_{\mathcal{D}_p}u$). Collecting the obtained inequalities for F_1 and F_2 we complete the estimate of II .

We show (5.13). Let Q be a cube with center x_Q and sidelength $2r$. Using (5.10); the generalized Hölder's inequality for \mathcal{A} and $\overline{\mathcal{A}}$, and also for $\overline{\mathcal{A}}$, \mathcal{B} and $\overline{\mathcal{C}}_k$; and (5.9)

$$\begin{aligned}
& \int_{\mathbb{R}^n \setminus 2Q} |K(x-y) - K(x-x_Q)| |b(x) - b_Q|^k u(x) dx \\
& \lesssim \sum_{l=1}^{\infty} \int_{|x-x_Q| \sim 2^l r} |K(x-y) - K(x-x_Q)| |b(x) - b_{2^{l+1}Q}|^k u(x) dx \\
& \quad + \sum_{l=1}^{\infty} l^k \int_{|x-x_Q| \sim 2^l r} |K(x-y) - K(x-x_Q)| u(x) dx \\
& \lesssim \sum_{l=1}^{\infty} (2^l r)^n \|K(\cdot - y) - K(\cdot - x_Q)\|_{\mathcal{B}, |x-x_Q| \sim 2^l r} \| (b - b_{2^{l+1}Q})^k \|_{\overline{\mathcal{C}}_k, 2^{l+1}Q} \|u\|_{\overline{\mathcal{A}}, 2^{l+1}Q} \\
& \quad + \sum_{l=1}^{\infty} (2^l r)^n l^k \|K(\cdot - y) - K(\cdot - x_Q)\|_{\mathcal{A}, |x-x_Q| \sim 2^l r} \|u\|_{\overline{\mathcal{A}}, 2^{l+1}Q} \\
& \lesssim \operatorname{ess\,inf}_Q M_{\overline{\mathcal{A}}} u,
\end{aligned}$$

where we have used that $K \in H_{\mathcal{B}} \cap H_{\mathcal{A},k}$.

To complete the proof we need to estimate *III*. The proof is almost identical to that of Theorem 3.1. For the case (a.1), in (5.4) we apply Theorem 3.7 in place of Theorem 2.4. Once we have that estimate, the proof follows the same computations once we check that $|T_b^k g|^p M_r \tilde{u} \in L^1(\mathbb{R}^n)$ (we show this below). For the case (a.2) we need to show that T_b^k satisfies the corresponding estimate in (5.8). But this follows from Theorem 2.6 as we can apply Theorem 3.7 to the adjoint of T_b^k —note that $(T_b^k)^* = (T^*)^k_{-b}$ and T^* is a singular integral operator with kernel $\tilde{K}(x) = K(-x) \in H_{\mathcal{B}} \cap H_{\mathcal{A},k}$.

As just mentioned we only need to check that $|T_b^k g|^p M_r \tilde{u} \in L^1(\mathbb{R}^n)$. As $u \in L^\infty$ it suffices to see that $T_b^k g \in L^p(\mathbb{R}^n)$ for p large enough. This is trivial if one assumes that $b \in L^\infty$ as our assumption on K implies that $K \in H_1$ and thus T is bounded on $L^p(\mathbb{R}^n)$ for every $1 < p < \infty$:

$$\begin{aligned}
\|T_b^k g\|_{L^p(\mathbb{R}^n)} &= \left\| \sum_{m=0}^k C_{m,k} b^{k-m} T(b^m g) \right\|_{L^p(\mathbb{R}^n)} \lesssim \|b\|_{L^\infty}^k \|g\|_{L^p(\mathbb{R}^n)} \\
&\leq \|b\|_{L^\infty}^k \lambda^{(p-1)/p} \|f\|_{L^1(\mathbb{R}^n)}^{1/p} < \infty.
\end{aligned}$$

Thus, we obtain (3.6) with $Su = M_{\overline{\mathcal{A}}} u$ under the additional assumption that $b \in L^\infty$. We pass to an arbitrary $b \in \text{BMO}$: for any $N > 0$ we define $b_N(x) = b(x)$ if $-N \leq b(x) \leq N$, $b_N(x) = N$ if $b(x) > N$ and $b_N(x) = -N$ if $b(x) < -N$. It is not hard to prove that $|b_N(x) - b_N(y)| \leq |b(x) - b(y)|$ and hence $\|b_N\|_{\text{BMO}} \leq 2 \|b\|_{\text{BMO}}$. Therefore, as $b_N \in L^\infty$ we can use (3.6) with b_N in place of b and so

$$\begin{aligned}
u\{x \in \mathbb{R}^n : |T_{b_N}^k f(x)| > \lambda\} &\leq C \int_{\mathbb{R}^n} \mathcal{C}_k \left(\frac{\|b_N\|_{\text{BMO}}^k |f(x)|}{\lambda} \right) M_{\overline{\mathcal{A}}} u(x) dx \\
&\leq C \int_{\mathbb{R}^n} \mathcal{C}_k \left(\frac{\|b\|_{\text{BMO}}^k |f(x)|}{\lambda} \right) M_{\overline{\mathcal{A}}} u(x) dx \quad (5.14)
\end{aligned}$$

where C does not depend on N . Since $f \in L_c^\infty$ it follows that for $0 \leq m \leq k$, $(b_N)^m f \rightarrow b^m f$ as $N \rightarrow \infty$ in L^q for $q > 1$. The fact that T is bounded on L^q implies $T((b_N)^m f) \rightarrow T(b^m f)$ as $N \rightarrow \infty$ in L^q . Passing to a subsequence the convergence is almost everywhere and so using that

$$T_{b_N}^k f(x) = \sum_{m=0}^k C_{m,k} b_N(x)^{k-m} T(b_N^m f)(x)$$

it follows that $T_{b_{N_j}}^k f(x) \rightarrow T_b^k f(x)$ for a.e. $x \in \mathbb{R}^n$ as $j \rightarrow \infty$. Then, we clearly have that $\chi_{\{T_b^k f > \lambda\}}(x) \leq \liminf_{j \rightarrow \infty} \chi_{\{T_{b_{N_j}}^k f > \lambda\}}(x)$ a.e. Thus, Fatou's lemma and (5.14) drive us to the desired estimate for T_b^k . This completes the proof of (a).

To obtain (b), we proceed as in Remark 5.1. Given $\varepsilon > 0$ we pick $p > 1$ and $\delta > 0$ so that $(k+1)p - 1 + 2\delta = k + \varepsilon$. Let $\mathcal{A}(t) = \exp(t^{\frac{1}{k+\delta/p}}) - 1$ and $\mathcal{B}(t) = \exp(t^{p/\delta}) - 1$. Note that we have $\overline{\mathcal{A}}^{-1}(t) \mathcal{B}^{-1}(t) \overline{\mathcal{C}}_k^{-1}(t) \lesssim t$. Also, $H_\infty \subset H_{\mathcal{B}}$ and $H_{e^{t^{1/k}}, k} \subset H_{\mathcal{A}, k}$ (as $\mathcal{A}(t) \lesssim e^{t^{1/k}} - 1$ for $t \geq 1$). Then $K \in H_{\mathcal{B}} \cap H_{\mathcal{A}, k}$. We apply (a.2) with $\mathcal{D}(t) = t^p (1 + \log^+ t)^{(k+1)p-1+2\delta}$ and $\mathcal{E}(t) \approx t^{p'}/(1 + \log^+ t)^{1+\delta(p'-1)} \in B_{p'}$ (note that we have $\mathcal{D}^{-1}(t) \mathcal{E}^{-1}(t) \leq \overline{\mathcal{A}}^{-1}(t)$). Then we obtain the desired estimate for the pair of weights $(u, M_{\mathcal{D}_p} u)$. To conclude we observe that $\mathcal{D}_p(t) = \mathcal{D}(t^{1/p}) = t(1 + \log^+ t)^{k+\varepsilon}$. \square

Remark 5.2. The proof for the multilinear commutators follows the same scheme, we just give some of the changes, leaving details to the reader.

To estimate *II* we use ideas from [36] and replace (5.11) by

$$\begin{aligned} |T_{\vec{b}} h(x)| &\lesssim \sum_{\sigma_1, \sigma_2} |T_{\vec{b}_{\sigma_2}} \left(\sum_j \pi_{\sigma_1}(\vec{b} - \vec{\lambda}^j) h_j \right)(x)| + \sum_j |\pi_{\{1, \dots, k\}}(\vec{b} - \vec{\lambda}^j)| |Th_j(x)| \\ &= F_1(x) + F_2(x), \end{aligned}$$

where the first sum runs over all partitions σ_1, σ_2 of $\{1, \dots, k\}$ with $\sigma_1 \neq \emptyset$; $T_{\vec{b}_{\sigma_2}}$ is the multilinear commutator associated with the vector $\vec{b}_{\sigma_2} = (b_{\sigma_2(l)})_l$; $\pi_{\sigma_1}(\vec{v}) = \prod_l v_{\sigma_1(l)}$ and $\pi_{\{1, \dots, k\}}(\vec{v}) = \prod_{l=1}^k v_l$; and $\vec{\lambda}^j = ((b_1)_{Q_j}, \dots, (b_k)_{Q_j})$. With this in hand we estimate F_1 using the induction hypothesis as $\#\sigma_2 \leq k - 1$, and we estimate F_2 using that $K \in H_{\mathcal{B}, k}$ (see [23]).

The estimate for *III* is obtained by using [23, Theorem 7.1] and observing that $(T_{\vec{b}})^* = (T^*)_{-\vec{b}}$ and T^* is a singular integral operator with kernel $\tilde{K}(x) = K(-x) \in H_{\mathcal{B}, k}$.

6. PROOFS IN THE ONE-SIDED CASE

Proof of Theorem 3.12, Part (i). The proof follows the same pattern as the proof in Theorem 3.1. We will only highlight some of the details. Again, we can assume that u is bounded and has compact support, also $0 \leq f \in L_c^\infty(\mathbb{R})$. Let

$$\Omega = \{x \in \mathbb{R} : M^+ f(x) > \lambda\} = \bigcup_j I_j = \bigcup_j (a_j, b_j)$$

where $I_j = (a_j, b_j)$ are the connected component of Ω and they satisfy (see[30])

$$\frac{1}{|I_j|} \int_{I_j} f(y) dy = \lambda.$$

Note that if $x \notin \Omega$, then for all $h \geq 0$

$$\frac{1}{h} \int_x^{x+h} f(y) dy \leq \lambda.$$

Therefore $f(x) \leq \lambda$ for a.e $x \in \mathbb{R} \setminus \Omega$. Let $\widehat{I}_j^- = (c_j, a_j)$ with c_j chosen so that $|\widehat{I}_j^-| = 2|I_j|$ and set

$$\tilde{\Omega} = \bigcup_j (\widehat{I}_j^- \cup I_j) = \bigcup_j \tilde{I}_j.$$

We write $\tilde{u} = u \chi_{\mathbb{R} \setminus \tilde{\Omega}}$ and decompose f as $f = g + h$, where

$$g = f \chi_{\mathbb{R} \setminus \Omega} + \sum_{j=1}^{\infty} f_{I_j} \chi_{I_j}, \quad h = \sum_{j=1}^{\infty} h_j = \sum_{j=1}^{\infty} (f - f_{I_j}) \chi_{I_j}.$$

Observe that $0 \leq g(x) \leq \lambda$ for a.e x and also that h_j has vanishing integral. Then

$$\begin{aligned} u\{x \in \mathbb{R} : |T^+ f(x)| > \lambda\} &\leq u(\tilde{\Omega}) + u\{x \in \mathbb{R} \setminus \tilde{\Omega} : |T^+ h(x)| > \lambda/2\} \\ &\quad + u\{x \in \mathbb{R} \setminus \tilde{\Omega} : |T^+ g(x)| > \lambda/2\} \\ &= I + II + III. \end{aligned}$$

Now we proceed in the same way as in the proof of Theorem 3.1. We estimate I :

$$I = u(\tilde{\Omega}) = u(\cup_j \tilde{I}_j) \leq \sum_j (u(\widehat{I}_j^-) + u(I_j)).$$

For each j we have

$$u(\widehat{I}_j^-) = \frac{u(\widehat{I}_j^-)}{|I_j|} |I_j| = \frac{u(\widehat{I}_j^-)}{|I_j|} \frac{1}{\lambda} \int_{I_j} |f(x)| dx \leq \frac{3}{\lambda} \int_{I_j} |f(x)| M^- u(x) dx,$$

and then

$$\sum_j u(\widehat{I}_j^-) \lesssim \frac{1}{\lambda} \int_{\Omega} |f(x)| M^- u(x) dx \leq \frac{1}{\lambda} \int_{\mathbb{R}} |f(x)| M^- u(x) dx.$$

On the other hand, using that M^+ is of weak-type (1,1) with respect to the pair of weights $(u, M^- u) \in A_1^+$ (see [27]),

$$\sum_j u(I_j) = u(\Omega) \leq \frac{C}{\lambda} \int_{\mathbb{R}} f(x) M^- u(x) dx,$$

and therefore

$$I \leq \frac{C}{\lambda} \int_{\mathbb{R}} f(x) M^- u(x) dx.$$

Observe that, as before, $M^- u$ is controlled by $M_{\mathcal{A}}^- u$ in (a), by $M_{\mathcal{D}_p}^- u$ in (b) and by $M_{L(\log L)^\varepsilon}^- u$ in (c).

We turn to *II*. Let $r_j = |I_j| = |\widehat{I}_j^-|/2$. We use that h_j is supported in I_j and has vanishing integral, also that K is supported in $(-\infty, 0)$:

$$\begin{aligned} II &\leq \frac{2}{\lambda} \sum_j \int_{\mathbb{R} \setminus \widehat{\Omega}} |Th_j(x)| u(x) dx \\ &\leq \frac{2}{\lambda} \sum_j \int_{I_j} |h_j(y)| \int_{\mathbb{R} \setminus \widehat{I}_j} |K(x-y) - K(x-a_j)| u(x) dx dy \\ &= \frac{2}{\lambda} \sum_j \int_{I_j} |h_j(y)| \int_{-\infty}^{c_j} |K(x-y) - K(x-a_j)| u(x) dx dy \end{aligned}$$

Then it suffices to obtain that for every $y \in I_j$,

$$\int_{-\infty}^{c_j} |K(x-y) - K(x-a_j)| u(x) dx \lesssim \operatorname{ess\,inf}_{I_j} M_{\overline{\mathcal{A}}} u, \quad (6.1)$$

which readily leads to the desired estimate

$$\begin{aligned} II &\lesssim \frac{1}{\lambda} \sum_j \operatorname{ess\,inf}_{I_j} M_{\overline{\mathcal{A}}} u \int_{I_j} |h_j(y)| dy \lesssim \frac{1}{\lambda} \sum_j \operatorname{ess\,inf}_{I_j} M_{\overline{\mathcal{A}}} u \int_{I_j} f(y) dy \\ &\leq \frac{1}{\lambda} \sum_j \int_{I_j} f(y) M_{\overline{\mathcal{A}}} u(y) dy \leq \frac{1}{\lambda} \int_{\mathbb{R}} f(y) M_{\overline{\mathcal{A}}} u(y) dy \end{aligned}$$

We show (6.1). Let $y, z \in I_j$, using generalized Hölder's inequality, for \mathcal{A} and $\overline{\mathcal{A}}$, and that $K \in H_{\mathcal{A}}$,

$$\begin{aligned} \int_{-\infty}^{c_j} |K(x-y) - K(x-a_j)| u(x) dx &= \sum_{k=1}^{\infty} \int_{a_j - 2^{k+1} r_j}^{a_j - 2^k r_j} |K(x-y) - K(x-a_j)| u(x) dx \\ &\lesssim \sum_{k=1}^{\infty} 2^k r_j \|K(\cdot - y) - K(\cdot - a_j)\|_{\mathcal{A}, |x-a_j| \sim 2^k r_j} \|u \chi_{(a_j - 2^{k+1} r_j, a_j - 2^k r_j)}\|_{\overline{\mathcal{A}}, |x-a_j| \sim 2^k r_j} \\ &\leq \sum_{k=1}^{\infty} 2^k r_j \|K(\cdot - y) - K(\cdot - a_j)\|_{\mathcal{A}, |x-a_j| \sim 2^k r_j} \|u\|_{\overline{\mathcal{A}}, (a_j - 2^{k+1} r_j, z)} \lesssim M_{\overline{\mathcal{A}}}^- u(z). \end{aligned}$$

To estimate *III* we first claim that for any Young function \mathcal{C} , any weight v with $M_{\overline{\mathcal{C}}}^- v < \infty$ a.e., and any interval $I = (a, b)$ we have

$$M_{\overline{\mathcal{C}}}^-(v \chi_{\mathbb{R} \setminus \widehat{I}^- \cup I})(y) \approx \operatorname{ess\,inf}_{z \in I} M_{\overline{\mathcal{C}}}^-(v \chi_{\mathbb{R} \setminus \widehat{I}^- \cup I})(z), \quad \text{a.e } y \in I, \quad (6.2)$$

where $\widehat{I}^- = (c, a)$ with c so that $|\widehat{I}^-| = 2|I|$. Assuming the proofs of the three cases (a), (b) and (c) adapts readily to the one-sided setting. For (a) one uses that $M_r^- \tilde{u} \in A_1^+ \subset A_{\infty}^+$ and therefore M^+ is bounded on $L^q(M_r^- \tilde{u})$ for every $1 < q < \infty$. For (b) we apply (i) in Theorem 3.11 to the one-sided operator $T^- = (T^+)^*$ (whose kernel $\tilde{K}(x) = K(-x)$ is supported on $(0, \infty)$) and then (ii) of Theorem 3.11 to conclude (5.8) with $M_{\mathcal{D}_p}^- \tilde{u}$ in the right-hand side. In case (c) we only need to adapt Remark 5.1 to this setting.

To complete the proof of (i) we show (6.2). Fix $y, z \in I = (a, b)$ and write $\widehat{I}^- = (c, a)$, where we recall that $|\widehat{I}^-| = 2|I|$. Observe that if $c \leq t < y$ then $(t, y) \subset (c, b) =$

$\widehat{I}^- \cup I$. Thus,

$$M_{\mathcal{C}}^-(v \chi_{\mathbb{R} \setminus \widehat{I}^- \cup I})(y) = \sup_{t < y} \|v \chi_{\mathbb{R} \setminus \widehat{I}^- \cup I}\|_{\mathcal{C},(t,y)} = \sup_{t < c} \|v \chi_{\mathbb{R} \setminus \widehat{I}^- \cup I}\|_{\mathcal{C},(t,y)}.$$

Given $t < c$ and $\lambda > 0$ it follows

$$\begin{aligned} \frac{1}{y-t} \int_t^y \mathcal{C} \left(\frac{v(x) \chi_{\mathbb{R} \setminus \widehat{I}^- \cup I}(x)}{\lambda} \right) dx &= \frac{1}{y-t} \int_t^c \mathcal{C} \left(\frac{v(x) \chi_{\mathbb{R} \setminus \widehat{I}^- \cup I}(x)}{\lambda} \right) dx \\ &= \frac{z-t}{y-t} \frac{1}{z-t} \int_t^c \mathcal{C} \left(\frac{v(x) \chi_{\mathbb{R} \setminus \widehat{I}^- \cup I}(x)}{\lambda} \right) dx \leq \frac{3}{2} \frac{1}{z-t} \int_t^z \mathcal{C} \left(\frac{v(x) \chi_{\mathbb{R} \setminus \widehat{I}^- \cup I}(x)}{\lambda} \right) dx \end{aligned}$$

and therefore $\|v \chi_{\mathbb{R} \setminus \widehat{I}^- \cup I}\|_{\mathcal{C},(t,y)} \leq 3/2 \|v \chi_{\mathbb{R} \setminus \widehat{I}^- \cup I}\|_{\mathcal{C},(t,z)}$ which in turns gives the desired estimate. \square

Using the previous ideas the proof of part (ii) in Theorem 3.12 can be obtained by adapting the proof of Theorem 3.8. Further details are left to reader.

Proof of Theorem 3.14. By homogeneity it suffices to consider the case $\lambda = 1$. Let $1 < p < \infty$ and let $\Omega = \{x : |Tf(x)| > 1\}$. Then, by duality, there exists $G \in L^{p'}(u)$ with $\|G\|_{L^{p'}(u)} = 1$ such that

$$u(\Omega)^{1/p} = \|\chi_{\Omega}\|_{L^p(u)} = \int_{\Omega} G(x) u(x) dx.$$

Then, using the hypotheses and Hölder's inequality

$$u(\Omega)^{1/p} \lesssim \int_{\mathbb{R}^n} |f(x)| M_{\mathcal{F}}^-(Gu)(x) dx \leq \|f\|_{L^p(v)} \|M_{\mathcal{F}}^-(Gu)\|_{L^{p'}(v^{1-p'})}.$$

Our hypotheses guarantee that we can apply Theorem 3.15 (indeed the corresponding version for $M_{\mathcal{F}}^-$) to obtain that $M_{\mathcal{F}}^-$ maps $L^{p'}(u^{1-p'})$ into $L^{p'}(v^{1-p'})$. Therefore,

$$u(\Omega)^{1/p} \lesssim \|f\|_{L^p(v)} \|Gu\|_{L^{p'}(u^{1-p'})} = \|f\|_{L^p(v)} \|G\|_{L^{p'}(u)} = \|f\|_{L^p(v)}.$$

\square

Remark 6.1. As observed before, this proof can be easily adapted to yield Theorem 3.5: one only needs to see that (3.3) guarantee that $M_{\mathcal{F}}$ is bounded from $L^{p'}(u^{1-p'})$ into $L^{p'}(v^{1-p'})$ (see [12]). For further results and a deep treatment of extrapolation results of this kind the reader is referred to [11].

Proof of Theorem 3.15. We use ideas from [38].

First we observe that it suffices to assume that u is bounded and compactly supported (otherwise we work with $u_R = u \chi_{\{|x| \leq R, u(x) \leq R\}}$ and take $R \rightarrow \infty$).

Fix f continuous with compact support and for $k \in \mathbb{Z}$ we set $\Omega_k = \{x \in \mathbb{R} : 2^k < M_{\mathcal{A}}^+ f(x) < 2^{k+2}\}$. For any $x \in \Omega_k$ there exists $c_x > x$ such that $2^k < \|f\|_{\mathcal{A},(x,c_x)} < 2^{k+2}$. Using the continuity of the integral it is easy to show that there exists $\delta_x \in (x, c_x)$ (that

can be taken sufficiently close to x verifying $\delta_x < (c_x + x)/2$ such that $[x, \delta_x) \subset \Omega_k$ and $2^k < \|f\|_{\mathcal{A},(\delta_x,c_x)} < 2^{k+2}$. We write $I_{x,k}^- = [x, \delta_x)$ and $I_{x,k}^+ = (\delta_x, c_x)$ and therefore

$$\Omega_k = \bigcup_{x \in \Omega_k} I_{x,k}^- \quad \text{and} \quad 2^k < \|f\|_{\mathcal{A},I_{x,k}^+} < 2^{k+2}. \quad (6.3)$$

As in [5] (see also [38, Lemma 2]) there exists a finite subcollection of pairwise disjoint intervals $\{I_{j,k}^-\}_{j \in J}$ such that

$$u(\Omega_k) \leq 3 \sum_{j \in J} u(I_{j,k}^-).$$

This and (6.3) yield

$$\begin{aligned} \int_{\mathbb{R}} M_{\mathcal{A}}^+ f(x)^p u(x) dx &\leq \sum_{k \in \mathbb{Z}} \int_{\Omega_k} (M_{\mathcal{A}}^+ f)^p u dx \lesssim \sum_{k \in \mathbb{Z}} 2^{kp} u(\Omega_k) \lesssim \sum_{k \in \mathbb{Z}} 2^{kp} \sum_{j \in J} u(I_{j,k}^-) \\ &\leq \sum_{k \in \mathbb{Z}} \sum_{j \in J} \|f\|_{\mathcal{A},I_{j,k}^+}^p u(I_{j,k}^-) = \sum_{k \in \mathbb{Z}} \sum_{j \in J} \|f v^{1/p} v^{-1/p}\|_{\mathcal{A},I_{j,k}^+}^p u(I_{j,k}^-) \\ &\lesssim \sum_{k \in \mathbb{Z}} \sum_{j \in J} \|f v^{1/p}\|_{\mathcal{C},I_{j,k}^+}^p \|v^{-1/p}\|_{\mathcal{B},I_{j,k}^+}^p \|u^{1/p}\|_{L^p, I_{j,k}^-}^p |I_{j,k}^-|. \end{aligned}$$

Note that

$$\|f v^{1/p}\|_{\mathcal{C},I_{j,k}^+} \leq 2 \|f v^{1/p}\|_{\mathcal{C},I_{j,k}^- \cup I_{j,k}^+} \leq 2 M_{\mathcal{C}}(f v^{1/p})(x), \quad x \in I_{j,k}^-.$$

This, (3.7), and the fact that the intervals $I_{j,k}^-$ are pairwise disjoint and contain in Ω_k imply

$$\begin{aligned} \int_{\mathbb{R}} M_{\mathcal{A}}^+ f(x)^p u(x) dx &\lesssim \sum_{k \in \mathbb{Z}} \sum_{j \in J} \|f v^{1/p}\|_{\mathcal{C},I_{j,k}^+}^p |I_{j,k}^-| \lesssim \sum_{k \in \mathbb{Z}} \sum_{j \in J} \int_{I_{j,k}^-} M_{\mathcal{C}}(f v^{1/p})(x)^p dx \\ &\leq \sum_{k \in \mathbb{Z}} \int_{\Omega_k} M_{\mathcal{C}}(f v^{1/p})(x)^p dx \leq 2 \int_{\mathbb{R}} M_{\mathcal{C}}(f v^{1/p})(x)^p dx \\ &\lesssim 2 \int_{\mathbb{R}} |f(x)|^p v(x) dx, \end{aligned}$$

where in the last estimate we have used that $\mathcal{C} \in B_p$ and consequently $M_{\mathcal{C}}$ is bounded on $L^p(\mathbb{R})$ (see [34]). This completes the proof. \square

REFERENCES

- [1] H. Aimar, L. Forzani and F. J. Martín-Reyes, *On weighted inequalities for one-sided singular integrals*, Proc. Amer. Math. Soc. **125** (1997), 2057–2064.
- [2] J. Alvarez, R. Bagby, D. Kurtz and C. Pérez, *Weighted estimates for commutators of linear operators*, Studia Math. **104** (1993), 195–209.
- [3] C. Bennett and R. Sharpley, *Interpolation of Operators*, Academic Press, New York 1998.
- [4] A.L. Bernardis, M. Lorente, F.J. Martín-Reyes, M.T. Martínez, A. de la Torre and J.L. Torrea, *Differential transforms in weighted spaces*, J. Fourier Anal. Appl. **12** (2006), no. 1, 83–103.
- [5] J. Bliedtner and P. Loeb, *A reduction technique for limit theorems in analysis and probability theory*, Ark. Mat. **30** (1992), 25–43.
- [6] A.P. Calderón and A. Zygmund, *On the existence of certain singular integrals*, Acta Math. **88** (1952), 85–139.

- [7] R. Coifman, *Distribution function inequalities for singular integrals*, Proc. Acad. Sci. U.S.A. **69** (1972), 2838–2839.
- [8] R. Coifman, R. Rochberg and G. Weiss, *Factorization theorems for Hardy spaces in several variables*, Ann. of Math. **103** (1976), 611–635.
- [9] A. Cordoba and C. Fefferman, *A weighted norm inequality for singular integrals*, Studia Math. **57** (1976), no. 1, 97–101.
- [10] D. Cruz-Uribe, SFO, J.M. Martell and C. Pérez, *Extensions of Rubio de Francia’s extrapolation theorem*, Proceedings of the 7th International Conference on Harmonic Analysis and Partial Differential Equations (El Escorial 2004), Collect. Math. 2006, 195–231.
- [11] D. Cruz-Uribe, SFO, J.M. Martell and C. Pérez, *Weights, extrapolation and the theory of Rubio de Francia*, in preparation.
- [12] D. Cruz-Uribe, SFO and C. Pérez, *Sharp Two-Weight, Weak-Type Norm Inequalities for Singular Integral Operators*, Math. Res. Letters, **6**, (1999), 417–427.
- [13] D. Cruz-Uribe, SFO and C. Pérez, *Two weight extrapolation via the maximal operator*, J. Funct. Anal. **174** (2000), no. 1, 1–17.
- [14] G.P. Curbera, J. García-Cuerva, J.M. Martell and C. Pérez, *Extrapolation with Weights, Rearrangement Invariant Function Spaces, Modular inequalities and applications to Singular Integrals*, Adv. Math. **203** (2006), no. 1, 256–318.
- [15] J. Duoandikoetxea, *Fourier Analysis*, Grad. Stud. Math. **29**, American Math. Soc., Providence, 2000.
- [16] J. Garcia-Cuerva, E. Harboure, C. Segovia and J. L. Torrea, *Weighted norm inequalities for commutators of strongly singular integrals*, Indiana Univ. Math. J. **40** (1991), 1398–1420.
- [17] J. García-Cuerva and J.L. Rubio de Francia, *Weighted Norm Inequalities and Related Topics*, North-Holland Math. Stud. **116**, North-Holland, 1985.
- [18] S. Hofmann, *Singular integrals with power weights*, Proc. Amer. Math. Soc. **110** (1990), no. 2, 343–353.
- [19] L. Hörmander, *Estimates for translation invariant operators in L^p spaces*, Acta Math. **104** (1960), 93–140.
- [20] R.L. Jones and J. Roseblatt, *Differential and ergodic transform*, Math. Ann. **323** (2002), 525–546.
- [21] M. A. Krasnosel’skiĭ and Ya. B. Rutickiĭ, *Convex Functions and Orlicz Spaces*, P. Noordhoff, Groningen, 1961.
- [22] D.S. Kurtz and R.L. Wheeden, *Results on weighted norm inequalities for multipliers*, Trans. Amer. Math. Soc. **255** (1979), 343–362.
- [23] M. Lorente, J.M. Martell, M.S. Riveros and A. de la Torre *Generalized Hörmander’s condition, commutators and weights*, J. Math. Anal. Appl. (2008), doi:10.1016/j.jmaa.2008.01.003.
- [24] M. Lorente, M.S. Riveros and A. de la Torre, *Weighted estimates for singular integral operators satisfying Hörmander’s conditions of Young type*, J. Fourier Anal. Apl. **11** (2005), no. 5, 497–509.
- [25] M. Lorente and M.S. Riveros, *Weighted inequalities for commutators of one-sided singular integrals*, Comment. Math. Univ. Carolin. **43** (2002), no. 1, 83–101.
- [26] J.M. Martell, C. Pérez, and R. Trujillo-González, *Lack of natural weighted estimates for some singular integral operators*, Trans. Amer. Math. Soc. **357**, (2005), 385–396
- [27] F.J. Martín-Reyes, P. Ortega and A. de la Torre, *Weighted inequalities for one-sided maximal functions*, Trans. Amer. Math. Soc. **319** (1990), no. 2, 517–534.
- [28] F.J. Martín-Reyes, L. Pick and A. de la Torre, *A_∞^+ condition*, Canad. J. Math. **45** (1993), 1231–1244.
- [29] F.J. Martín-Reyes and A. de la Torre, *One Sided BMO Spaces*, J. London Math. Soc. **2** (49) (1994), no. 3, 529–542.

- [30] B. Muckenhoupt, *Weighted norm inequalities for the Hardy maximal function*, Trans. Amer. Math. Soc. **165**, (1972) 207–226.
- [31] R. O’Neil, *Fractional integration in Orlicz spaces*, Trans. Amer. Math. Soc. **115**, (1963) 300–328.
- [32] C. Pérez, *Weighted norm inequalities for singular integral operators*, J. London Math. Soc. (2) **49** (1994), no. 2, 296–308.
- [33] C. Pérez, *Endpoint estimates for commutators of singular integral operators*, J. Funct. Anal. **128** (1995), 163–185.
- [34] C. Pérez, *On sufficient conditions for the boundedness of the Hardy-Littlewood maximal operator between weighted L^p -spaces with different weights*, Proc. London Math. Soc. (3) **71** (1995), no. 1, 135–157.
- [35] C. Pérez and G. Pradolini, *Sharp weighted endpoint estimates for commutators of singular integrals*, Michigan Math. J. **49** (2001), no. 1, 23–37.
- [36] C. Pérez and R. Trujillo-Gonzalez, *Sharp weighted estimates for multilinear commutators*, J. London Math. Soc. (2) **65** (2002), 672–692.
- [37] M. Rao and Z.D. Ren, *Theory of Orlicz spaces*, Monographs and Textbooks in Pure and Applied Mathematics **146**, Marcel Dekker, Inc., New York, 1991.
- [38] M. S. Riveros, L. de Rosa and A. de la Torre, *Sufficient Conditions for one-sided Operators*, J. Fourier Anal. Appl. **6**, (2000) 607–621.
- [39] J.L. Rubio de Francia, F.J. Ruiz and J. L. Torrea, *Calderón-Zygmund theory for vector-valued functions*, Adv. in Math. **62** (1986), 7–48.
- [40] C. Segovia and J.L. Torrea, *Higher order commutators for vector-valued Calderón-Zygmund operators*, Trans. Amer. Math. Soc. **336** (1993), 537–556.
- [41] E. Sawyer, *Weighted inequalities for the one-sided Hardy-Littlewood maximal functions*, Trans. Amer. Math. Soc. **297** (1986), 53–61.
- [42] A. de la Torre and J.L. Torrea, *One-sided discrete square function*, Studia Math. **156** (2003), no. 3, 243–260.
- [43] D.K. Watson, *Weighted estimates for singular integrals via Fourier transform estimates*, Duke Math. J. **60** (1990), no. 2, 389–399.

MARÍA LORENTE, DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, FACULTAD DE CIENCIAS, UNIVERSIDAD DE MÁLAGA, CAMPUS DE TEATINOS, S/N, 29071 MÁLAGA, SPAIN
E-mail address: lorente@anamat.cie.uma.es

JOSÉ MARÍA MARTELL, INSTITUTO DE CIENCIAS MATEMÁTICAS CSIC-UAM-UC3M-UCM, CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS, C/ SERRANO 121, E-28006 MADRID, SPAIN
E-mail address: chema.martell@uam.es

CARLOS PÉREZ, DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, FACULTAD DE MATEMÁTICA, UNIVERSIDAD DE SEVILLA, 41080, SEVILLA, SPAIN
E-mail address: carlosperez@us.es

MARÍA SILVINA RIVEROS, FAMAF, UNIVERSIDAD NACIONAL DE CÓRDOBA, CIEM (CONICET), 5000 CÓRDOBA, ARGENTINA
E-mail address: sriveros@mate.uncor.edu