

A NOTE ON THE OFF-DIAGONAL MUCKENHOUP-T-WHEEDEN CONJECTURE

DAVID CRUZ-URIBE, SFO, JOSÉ MARÍA MARTELL, AND CARLOS PÉREZ

ABSTRACT. We obtain the off-diagonal Muckenhoupt-Wheeden conjecture for Calderón-Zygmund operators. Namely, given $1 < p < q < \infty$ and a pair of weights (u, v) , if the Hardy-Littlewood maximal function satisfies the following two weight inequalities:

$$M : L^p(v) \rightarrow L^q(u) \quad \text{and} \quad M : L^{q'}(u^{1-q'}) \rightarrow L^{p'}(v^{1-p'}),$$

then any Calderón-Zygmund operator T and its associated truncated maximal operator T_* are bounded from $L^p(v)$ to $L^q(u)$. Additionally, assuming only the second estimate for M then T and T_* map continuously $L^p(v)$ into $L^{q,\infty}(u)$. We also consider the case of generalized Haar shift operators and show that their off-diagonal two weight estimates are governed by the corresponding estimates for the dyadic Hardy-Littlewood maximal function.

1. INTRODUCTION AND MAIN RESULTS

In the 1970s, Muckenhoupt and Wheeden conjectured that given $p, 1 < p < \infty$, a sufficient condition for the Hilbert transform to satisfy the two weight norm inequality

$$H : L^p(v) \rightarrow L^p(u)$$

is that the Hardy-Littlewood maximal operator satisfy the pair of norm inequalities

$$\begin{aligned} M &: L^p(v) \rightarrow L^p(u), \\ M &: L^{p'}(u^{1-p'}) \rightarrow L^{p'}(v^{1-p'}). \end{aligned}$$

Moreover, they conjectured that the Hilbert transform satisfies the weak-type inequality

$$H : L^p(v) \rightarrow L^{p,\infty}(u)$$

provided that the maximal operator satisfies the second “dual” inequality. Both of these conjectures readily extend to all Calderón-Zygmund operators (see the definition below). Very recently, both conjectures were disproved:

Date: March 22, 2012.

1991 Mathematics Subject Classification. 42B20, 42B25.

Key words and phrases. Haar shift operators, Calderón-Zygmund operators, two-weight inequalities, testing conditions.

The first author was supported by a grant from the Faculty Research Committee and the Stewart-Dorwart Faculty Development Fund at Trinity College; the first and third authors are supported by grant MTM2009-08934 from the Spanish Ministry of Science and Innovation; the second author is supported by grant Grant MTM2010-16518 from the same institution.

the strong-type inequality by Reguera and Scurry [11] and the weak-type inequality by the first author, Reznikov and Volberg [5].

Remark 1.1. A special case of these conjectures, involving the A_p bump conditions, has been considered by several authors: see [1, 2, 3, 4, 5, 9].

In this note we prove the somewhat surprising fact that the Muckenhoupt-Wheeden conjectures are true for off-diagonal inequalities. Our main result is Theorem 1.2 below. We also prove an analogous result for the Haar shift operators (the so-called dyadic Calderón-Zygmund operators) with the Hardy-Littlewood maximal operator replaced by the dyadic maximal operator: see Theorem 1.3 below.

To state our results we first give some preliminary definitions. By weights we will always mean non-negative, measurable functions. Given a pair of weights (u, v) , hereafter we will assume that $u > 0$ on a set of positive measure and $u < \infty$ a.e., and $v > 0$ a.e. and $v < \infty$ on a set of positive measure. We will also use the standard notation $0 \cdot \infty = 0$.

Calderón-Zygmund operators. A Calderón-Zygmund operator T is a linear operator that is bounded on $L^2(\mathbb{R}^n)$ and

$$Tf(x) = \int_{\mathbb{R}^n} K(x, y)f(y)dy, \quad f \in L_c^\infty(\mathbb{R}^n), \quad x \notin \text{supp } f,$$

where the kernel K satisfies the size and smoothness estimates

$$|K(x, y)| \leq \frac{C}{|x - y|^n}, \quad x \neq y,$$

and

$$|K(x, y) - K(x', y)| + |K(y, x) - K(y, x')| \leq C \frac{|x - x'|^\delta}{|x - y|^{n+\delta}},$$

for all $|x - y| > 2|x - x'|$.

Associated with T is the truncated maximal operator

$$T_\star f(x) = \sup_{0 < \epsilon < \epsilon' < \infty} \left| \int_{\epsilon < |x-y| < \epsilon'} K(x, y)f(y)dy \right|.$$

Let M denote the Hardy-Littlewood maximal operator, that is,

$$Mf(x) = \sup_{Q \ni x} \int_Q |f(y)|dy = \sup_{Q \ni x} \frac{1}{|Q|} \int_Q |f(y)|dy.$$

where the supremum is taken over all cubes in \mathbb{R}^n with sides parallel to the coordinate axes.

Theorem 1.2. *Given a Calderón-Zygmund operator T , let $1 < p < q < \infty$ and let (u, v) be a pair of weights. If the maximal operator satisfies*

$$(1.1) \quad M : L^p(v) \rightarrow L^q(u) \quad \text{and} \quad M : L^{q'}(u^{1-q'}) \rightarrow L^{p'}(v^{1-p'}),$$

then

$$(1.2) \quad \|Tf\|_{L^q(u)} \leq C\|f\|_{L^p(v)} \quad \text{and} \quad \|T_\star f\|_{L^q(u)} \leq C\|f\|_{L^p(v)}.$$

Analogously, if the maximal operator satisfies

$$(1.3) \quad M : L^{q'}(u^{1-q'}) \rightarrow L^{p'}(v^{1-p'}),$$

then

$$(1.4) \quad \|Tf\|_{L^{q,\infty}(u)} \leq C\|f\|_{L^p(v)} \quad \text{and} \quad \|T_*f\|_{L^{q,\infty}(u)} \leq C\|f\|_{L^p(v)}.$$

If the pairs of weights (u, v) satisfy any of the conditions in (1.1), then the weights u and $v^{1-p'}$ are locally integrable. This is a consequence of a characterization of the two weight norm inequalities for the maximal operator due to Sawyer [12]. He proved that the $L^p - L^q$ inequality holds if and only if for every cube Q ,

$$\left(\int_Q M(v^{1-p'} \chi_Q)(x)^q u(x) dx \right)^{1/q} \leq C \left(\int_Q v(x)^{1-p'} dx \right)^{1/p} < \infty,$$

and the $L^{q'} - L^{p'}$ inequality holds if and only if

$$\left(\int_Q M(u \chi_Q)(x)^{p'} v(x)^{1-p'} dx \right)^{1/p'} \leq C \left(\int_Q u(x) dx \right)^{1/q'} < \infty.$$

It is straightforward to construct pairs of weights that satisfy these conditions. For instance, in \mathbb{R} both of these conditions follow easily for every $1 < p \leq q < \infty$ and the pair of weights (u, v) with $u = \chi_{[0,1]}$ and $v^{-1} = \chi_{[2,3]}$ (i.e., $v = 1$ in $[2, 3]$ and $v = \infty$ elsewhere). Indeed, we only need to check Sawyer's inequalities for cubes Q that intersect both $[0, 1]$ and $[2, 3]$, in which case we have $M(\chi_{[2,3] \cap Q})(x) \leq |[2, 3] \cap Q|$ for every $x \in [0, 1] \cap Q$, and $M(\chi_{[0,1] \cap Q})(x) \leq |[0, 1] \cap Q|$ for every $x \in [2, 3] \cap Q$. These readily imply the desired estimates.

Dyadic Calderón-Zygmund operators. A generalized dyadic grid \mathcal{D} in \mathbb{R}^n is a set of generalized dyadic cubes with the following properties: if $Q \in \mathcal{D}$ then $\ell(Q) = 2^k$, $k \in \mathbb{Z}$; if $Q, R \in \mathcal{D}$ and $Q \cap R \neq \emptyset$ then $Q \subset R$ or $R \subset Q$; the cubes in \mathcal{D} with $\ell(Q) = 2^{-k}$ form a disjoint partition of \mathbb{R}^n (see [9] and [10] for more details).

We say that g_Q is a generalized Haar function associated with $Q \in \mathcal{D}$ if

- (a) $\text{supp}(g_Q) \subset Q$;
- (b) if $Q' \in \mathcal{D}$ and $Q' \subsetneq Q$, then g_Q is constant on Q' ;
- (c) $\|g_Q\|_\infty \leq 1$.

Given a dyadic grid \mathcal{D} and a pair $(m, k) \in \mathbb{Z}_+^2$, a linear operator \mathcal{S} is a generalized Haar shift operator (that is, a dyadic Calderón-Zygmund operator) of complexity type (m, k) if it is bounded on $L^2(\mathbb{R}^n)$ and

$$\mathcal{S}f(x) = \sum_{Q \in \mathcal{D}} \mathcal{S}_Q f(x) = \sum_{Q \in \mathcal{D}} \sum_{\substack{Q' \in \mathcal{D}_m(Q) \\ Q'' \in \mathcal{D}_k(Q)}} \frac{\langle f, g_{Q''} \rangle}{|Q|} g_{Q'}(x),$$

where $\mathcal{D}_j(Q)$ stands for the dyadic subcubes of Q with side length $2^{-j}\ell(Q)$, $g_{Q'}^{Q''}$ is a generalized a Haar function associated with Q' and $g_{Q''}^{Q'}$ is a generalized a Haar function associated with Q'' . We say that the complexity of \mathcal{S} is $\kappa = \max(m, k)$. We also define the truncated Haar shift operator

$$\mathcal{S}_\star f(x) = \sup_{0 < \epsilon < \epsilon' < \infty} |\mathcal{S}_{\epsilon, \epsilon'} f(x)| = \sup_{0 < \epsilon < \epsilon' < \infty} \left| \sum_{\substack{Q \in \mathcal{D} \\ \epsilon \leq \ell(Q) \leq \epsilon'}} \mathcal{S}_Q f(x) \right|.$$

An important example of a Haar shift operator on the real line is the Haar shift (also known as the dyadic Hilbert transform) H^d , defined by

$$H^d f(x) = \sum_{I \in \Delta} \langle f, h_I \rangle (h_{I_-}(x) - h_{I_+}(x)),$$

where, given a dyadic interval I , I_+ and I_- are its right and left halves, and

$$h_I(x) = |I|^{-1/2} (\chi_{I_-}(x) - \chi_{I_+}(x)).$$

After renormalizing, h_I is a Haar function on I and one can write H^d as a generalized Haar shift operator of complexity 1. These operators have played a very important role in the proof of the A_2 conjecture: see [4, 6, 7] and the references they contain for more information.

Associated with the dyadic grid \mathcal{D} is the dyadic maximal function

$$M_{\mathcal{D}} f(x) = \sup_{x \in Q \in \mathcal{D}} \int_Q |f(y)| dy.$$

Note that $M_{\mathcal{D}}$ is dominated pointwise by the Hardy-Littlewood maximal operator.

We can now state our result for dyadic Calderón-Zygmund operators.

Theorem 1.3. *Let \mathcal{S} be a generalized Haar shift operator of complexity κ . Given $1 < p < q < \infty$ and a pair of weights (u, v) , if the dyadic maximal operator satisfies*

$$(1.5) \quad M_{\mathcal{D}} : L^p(v) \rightarrow L^q(u) \quad \text{and} \quad M_{\mathcal{D}} : L^{q'}(u^{1-q'}) \rightarrow L^{p'}(v^{1-p'}),$$

then

$$(1.6) \quad \|\mathcal{S}f\|_{L^q(u)} \leq C\kappa^2 \|f\|_{L^p(v)} \quad \text{and} \quad \|\mathcal{S}_\star f\|_{L^q(u)} \leq C\kappa^2 \|f\|_{L^p(v)}.$$

Analogously, if the dyadic maximal operator satisfies

$$(1.7) \quad M_{\mathcal{D}} : L^{q'}(u^{1-q'}) \rightarrow L^{p'}(v^{1-p'})$$

then

$$(1.8) \quad \|\mathcal{S}f\|_{L^{q,\infty}(u)} \leq C\kappa^2 \|f\|_{L^p(v)} \quad \text{and} \quad \|\mathcal{S}_\star f\|_{L^{q,\infty}(u)} \leq C\kappa^2 \|f\|_{L^p(v)}.$$

2. PROOFS OF THE MAIN RESULTS

Proof of Theorem 1.2. We will prove our estimates for T_* ; the ones for T are completely analogous.

Given a dyadic grid \mathcal{D} we say that $\{Q_j^k\}_{j,k}$ is a *sparse family* of dyadic cubes if for any k the cubes $\{Q_j^k\}_j$ are pairwise disjoint; if $\Omega_k := \cup_j Q_j^k$, then $\Omega_{k+1} \subset \Omega_k$; and $|\Omega_{k+1} \cap Q_{j,k}| \leq \frac{1}{2}|Q_j^k|$. Given \mathcal{D} and a sparse family $\mathcal{S} = \{Q_j^k\}_{j,k} \subset \mathcal{D}$, define the positive dyadic operator \mathcal{A} by

$$\mathcal{A}f(x) = \mathcal{A}_{\mathcal{D},\mathcal{S}}f(x) = \sum_{j,k} f_{Q_j^k} \chi_{Q_j^k}(x)$$

where $f_Q = \int_Q f(y)dy$.

For our proof we will use the main result in [9, 10]. Given a Banach function space X and a non-negative function f ,

$$(2.1) \quad \|T_*f\|_X \leq C(T, n) \sup_{\mathcal{D},\mathcal{S}} \|\mathcal{A}_{\mathcal{D},\mathcal{S}}f\|_X,$$

where the supremum is taken over all dyadic grids \mathcal{D} and sparse families $\mathcal{S} \subset \mathcal{D}$. To prove Theorem 1.2 we apply this result with $X = L^q(u)$ or $X = L^{q,\infty}(u)$; it will then suffice to show that our assumptions on M guarantee that $\mathcal{A}_{\mathcal{D},\mathcal{S}}$ satisfies the corresponding two weight inequalities.

To prove this fact we will use a result by Lacey, Sawyer and Uriarte-Tuero [8]. Given a sequence of non-negative constants $\alpha = \{\alpha_Q\}_{Q \in \mathcal{D}}$, define the positive operator

$$T_\alpha f(x) = \sum_{Q \in \mathcal{D}} \alpha_Q f_Q \chi_Q(x).$$

Further, given $R \in \mathcal{D}$ we define the ‘‘outer truncated’’ operator

$$T_\alpha^R f(x) = \sum_{\substack{Q \in \mathcal{D} \\ Q \supset R}} \alpha_Q f_Q \chi_Q(x).$$

In [8] it was shown that for all $1 < p < q < \infty$, $T_\alpha : L^p(v) \rightarrow L^q(u)$ if and only if there exist constants C_1 and C_2 such that for every $R \in \mathcal{D}$

$$(2.2) \quad \left(\int_{\mathbb{R}^n} T_\alpha^R(v^{1-p'} \chi_R)(x)^q u(x) dx \right)^{\frac{1}{q}} \leq C_1 \left(\int_R v(x)^{1-p'} dx \right)^{\frac{1}{p}},$$

and

$$(2.3) \quad \left(\int_{\mathbb{R}^n} T_\alpha^R(u \chi_R)(x)^{p'} v(x)^{1-p'} dx \right)^{\frac{1}{p'}} \leq C_2 \left(\int_R u(x) dx \right)^{\frac{1}{q}}.$$

Furthermore, for $1 < p < q < \infty$, $T_\alpha : L^p(v) \rightarrow L^{q,\infty}(u)$ holds if and only if there exists a constant C_2 such that for every $R \in \mathcal{D}$, (2.3) holds.

We can apply these results to the operator $\mathcal{A} = \mathcal{A}_{\mathcal{D},\mathcal{S}}$ where \mathcal{D} and \mathcal{S} are fixed, since $\mathcal{A} = T_\alpha$ with $\alpha_Q = 1$ if $Q \in \mathcal{S}$ and $\alpha_Q = 0$ otherwise. Fix $R \in \mathcal{D}$; to estimate \mathcal{A}^R , take the increasing family of cubes $R = R_0 \subsetneq R_1 \subsetneq R_2 \subsetneq \dots$ with $R_k \in \mathcal{D}$ and $\ell(R_k) = 2^k \ell(R)$. Define $R_{-1} = \emptyset$. Note that

$\text{supp } \mathcal{A}^R \subset \cup_{k \geq 0} R_k$. Then for every non-negative function f and for every $x \in R_k \setminus R_{k-1}$ with $k \geq 0$ we have that

$$\begin{aligned} 0 \leq \mathcal{A}^R(f\chi_R)(x) &\leq \sum_{j=0}^{\infty} (f\chi_R)_{R_j} \chi_{R_j}(x) = f_R \sum_{j=k}^{\infty} 2^{-jn} \\ &\lesssim f_R 2^{-kn} = (f\chi_R)_{R_k} \leq M_{\mathcal{D}}(f\chi_R)(x). \end{aligned}$$

Consequently, for every $x \in \mathbb{R}^n$,

$$(2.4) \quad 0 \leq \mathcal{A}^R(f\chi_R)(x) \lesssim M_{\mathcal{D}}(f\chi_R)(x) \leq M(f\chi_R)(x).$$

Inequality (2.4) together with our hypothesis (1.1) implies (2.2) and (2.3). Therefore, we have that $\mathcal{A} : L^p(v) \rightarrow L^q(u)$ with constants depending on the dimension, p , q and the implicit constants in (1.1). Therefore, by Lerner's estimate (2.1) we get $T_{\star} : L^p(v) \rightarrow L^q(u)$ as desired.

For the weak-type estimates we proceed in the same manner, using the fact that (1.3) yields (2.3) and therefore $\mathcal{A} : L^p(v) \rightarrow L^{q,\infty}(u)$. This in turn implies, by Lerner's estimate (2.1) applied to $X = L^{q,\infty}(u)$, that $T_{\star} : L^p(v) \rightarrow L^{q,\infty}(u)$.

Proof of Theorem 1.3. Fix \mathcal{D} and a generalized Haar shift operator of complexity κ . As before we can work with \mathcal{S}_{\star} . We can repeat the previous argument except that we want to keep the fixed dyadic structure \mathcal{D} . A careful examination of [9, Section 5] shows that, given a Banach function space X , we have

$$(2.5) \quad \|\mathcal{S}_{\star}f\|_X \leq C_n \kappa^2 \sup_{\mathcal{J}} \|\mathcal{A}_{\mathcal{D},\mathcal{J}}f\|_X, \quad f \geq 0,$$

where the supremum is taken over all sparse families $\mathcal{J} \subset \mathcal{D}$. We emphasize that in [9, Section 5] there is an additional supremum over the dyadic grids \mathcal{D} . This is because at some places the dyadic maximal operator is majorized by the regular Hardy-Littlewood maximal operator and the latter is in turn controlled by a sum of $\mathcal{A}_{\mathcal{D}_{\alpha},\mathcal{J}_{\alpha}}$ for 2^n dyadic grids \mathcal{D}_{α} . However, keeping $M_{\mathcal{D}}$ one can easily show that (2.5) holds. Details are left to the interested reader.

Given (2.5), we fix a sparse family $\mathcal{J} \subset \mathcal{D}$ and write $\mathcal{A} = \mathcal{A}_{\mathcal{D},\mathcal{J}}$. Arguing exactly as before we obtain (2.4). Thus, (1.5) implies (2.2) and (2.3) and therefore the result from [8] yields $\mathcal{A} : L^p(v) \rightarrow L^q(u)$ with constants depending on the dimension, p , q and the implicit constants in (1.5). Combining this with Lerner's estimate (2.5) applied to $X = L^q(u)$ we conclude as desired that $\mathcal{S}_{\star} : L^p(v) \rightarrow L^q(u)$. We get the weak-type estimate by adapting the above proof in exactly the same way.

REFERENCES

- [1] D. Cruz-Uribe, J. M. Martell, and C. Pérez. Sharp two-weight inequalities for singular integrals, with applications to the Hilbert transform and the Sarason conjecture. *Adv. Math.*, 216(2):647–676, 2007.
- [2] D. Cruz-Uribe, J. M. Martell, and C. Pérez. Sharp weighted estimates for approximating dyadic operators. *Electron. Res. Announc. Math. Sci.*, 17:12–19, 2010.

- [3] D. Cruz-Uribe, J.M. Martell, and C. Pérez. *Weights, extrapolation and the theory of Rubio de Francia*, volume 215 of *Operator Theory: Advances and Applications*. Birkhäuser/Springer Basel AG, Basel, 2011.
- [4] D. Cruz-Uribe, J.M. Martell, and C. Pérez. Sharp weighted estimates for classical operators. *Adv. in Math*, 229:408–441, 2012.
- [5] D. Cruz-Uribe, A. Reznikov, and A. Volberg. Logarithmic bump conditions and the two-weight boundedness of Calderón–Zygmund operators. *Preprint*, 2012. arXiv:1112.0676.
- [6] T. Hytönen. The sharp weighted bound for general Calderón–Zygmund operators. *Ann. of Math. (2)*, (to appear). arXiv:1007.4330 (2010).
- [7] T. Hytönen, C. Pérez, S. Treil, and A. Volberg. Sharp weighted estimates of the dyadic shifts and A_2 conjecture. *Preprint*, 2010. arXiv:1010.0755.
- [8] M. Lacey, E. Sawyer, and I. Uriarte-Tuero. Two weight inequalities for discrete positive operators. *Preprint*, 2010. arXiv:0911.3437.
- [9] A. Lerner. On an estimate of Calderón–Zygmund operators by dyadic positive operators. *Preprint*, 2012. arXiv:1202.1860.
- [10] A. Lerner. A simple proof of the A_2 conjecture. *Preprint*, 2012. arXiv:1202.2824.
- [11] M. Reguera and J. Scurry. On joint estimates for maximal functions and singular integrals in weighted spaces. *Preprint*, 2011. arXiv:1109.2027.
- [12] E.T. Sawyer. A characterization of a two-weight norm inequality for maximal operators. *Studia Math.*, 75(1):1–11, 1982.

DAVID CRUZ-URIBE, SFO, DEPT. OF MATHEMATICS, TRINITY COLLEGE, HARTFORD, CT 06106-3100, USA

E-mail address: david.cruzuribe@trincoll.edu

JOSÉ MARÍA MARTELL, INSTITUTO DE CIENCIAS MATEMÁTICAS CSIC-UAM-UC3M-UCM, CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS, C/ NICOLÁS CABRERA, 13-15, E-28049 MADRID, SPAIN

E-mail address: chema.martell@icmat.es

CARLOS PÉREZ, DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, FACULTAD DE MATEMÁTICAS, UNIVERSIDAD DE SEVILLA, 41012 SEVILLA, SPAIN

E-mail address: carlosperez@us.es