

Scattering of ^8He on ^{208}Pb at 22 MeV

G. Marquín-Durán*, A.M. Sánchez-Benítez*, I. Martel*, L. Acosta*.[†],
 K. Rusek**, M.A.G. Álvarez^{‡,§}, R. Berjillos*, M.J.G. Borge[¶], A. Chbihi[¶],
 C. Cruz[¶], M. Cubero[¶], J.A. Dueñas*, J.P. Fernández-García[§],
 B. Fernández-Martínez[‡], J.L. Flores^{††}, J. Gómez-Camacho^{‡,§}, N. Keeley^{‡‡},
 J.A. Labrador[‡], M. Marqués^{§§}, A.M. Moro[§], M. Mazzocco^{¶¶}, A. Pakou^{***},
 V.V. Parkar*, N. Patronis^{***}, V. Pseudo[¶], D. Pierroutsakou^{†††}, R. Raabe^{‡‡‡},
 R. Silvestri^{††††}, N. Soic^{§§§}, Ł. Standylo^{‡‡}, I. Strojek^{‡‡}, O. Tengblad[¶],
 R. Wolski^{¶¶¶} and A.H. Ziad[‡]

**Depto. de Física Aplicada, Universidad de Huelva, 21071 Huelva, Spain*

[†]*Laboratori Nazionali del Sud, INFN, Via Santa Sofia 62, 95123, Catania, Italy*

***Heavy Ion Laboratory, University of Warsaw*

[‡]*Centro Nacional de Aceleradores, 41092, Sevilla, Spain*

[§]*Depto. de Física Atómica, Molecular y Nuclear, Universidad de Sevilla, 41080 Sevilla, Spain*

[¶]*Instituto de Estructura de la Materia, CSIC, 28006 Madrid, Spain*

^{¶¶}*GANIL, CEA and IN2P3-CNRS, B.P. 5027, 14076 Caen cedex, France*

^{††}*Depto. de Ingeniería Eléctrica, Universidad de Huelva, 21071 Huelva, Spain*

^{‡‡}*National Centre for Nuclear Research, A.Soltana 7, 05-400 Otwock, Poland*

^{§§}*Laboratoire de Physique Corpusculaire, 14050 Caen cedex, France*

^{¶¶¶}*Dip. di Fisica and INFN, Università di Padova, via F. Marzolo 8, 35131 Padova, Italy*

^{***}*Department of Physics, University of Ioannina, 45110 Ioannina, Greece*

^{†††}*INFN - Sezione di Napoli, via Cintia, 80126 Napoli, Italy*

^{‡‡‡}*Instituut voor Kern-en Stralingsfysica, K.U. Leuven, Belgium*

^{§§§}*Rudjer Boskovic Institute, Bijenicka 54, 10000 Zagreb, Croatia*

^{¶¶¶¶}*Flerov Laboratory of Nuclear Reactions, JINR, Dubna 141980, Russian Federation*

Abstract. The skin nucleus ^8He is investigated by measuring the angular distribution of the elastically scattered ^8He and the $^6,4\text{He}$ fragments produced in the collision with a ^{208}Pb target at 22 MeV, just above the Coulomb barrier. The experiment was carried out at SPIRAL/GANIL in 2010. Here we present preliminary results for the elastic scattering.

Keywords: Direct reactions, Elastic scattering, ^8He

PACS: 01.30.Cc, 13.85.Dz, 24.10.Ht, 25.10.+s

INTRODUCTION

The near-barrier scattering of ^8He with heavy targets remains still not very well known. Only a few scattering data sets are available nowadays [1, 2]. The skin nucleus ^8He presents the largest N-Z asymmetry of the particle-stable nucleus and its particular features makes it an attractive field of study. This work is focused in the experimental results for the elastic scattering up to 95° , for the $^8\text{He}+^{208}\text{Pb}$ system at 22 MeV.

EXPERIMENTAL SETUP

The experiment was performed at SPIRAL/GANIL. The experimental setup consisted of a portable reaction chamber where the array of detectors was mounted, a set of collimators and a beam diagnostics system. The detection system, GLORIA (GLObal Reaction Array), consists of 12 DSSSD detectors arranged in 6 particle telescopes (40 μ m, 1mm) covering a continuous angular range between 15° and 165°.

EXPERIMENTAL RESULTS FOR THE ELASTIC SCATTERING

The experimental results obtained are presented in Fig. 1. The three first telescopes, covering up to 95°, are plotted separately in order to show the existing matching between them. The elastic cross section decreases in a smooth way losing the Coulomb-nuclear rainbow, characteristic for light stable nuclei.




FIGURE 1. Angular distribution of the elastic cross section for the system ${}^8\text{He}+{}^{208}\text{Pb}$ at 22 MeV. The two forward and the bottom telescopes are represented separately.

The experimental data is compared with an OM calculation in which the optical potential derived from the elastic scattering of ${}^8\text{Li}+{}^{208}\text{Pb}$ [3] is used.

CONCLUSIONS

The elastic scattering of ${}^8\text{He}+{}^{208}\text{Pb}$ at 22 MeV has been measured at SPIRAL-GANIL using GLORIA, a new detector developed at the University of Huelva. The angular distribution of the elastic channel decreases in a smooth way losing the Coulomb-nuclear rainbow.

REFERENCES

1. A. Lemasson *et al.*, *Phys. Rev.* **C82**, 044617 (2010).
2. A. Lemasson *et al.*, *Phys. Lett.* **B697**, 454 (2011).
3. J.J. Kolata *et al.*, *Phys. Rev.* **C65**, 054616 (2002).