



# Complete Genome Sequence of an Aquaculture-Associated Phage, FL-1, Infecting *Flavobacterium* spp.

Elina Laanto,<sup>a</sup> Janne J. Ravantti,<sup>b</sup> Lotta-Riina Sundberg<sup>a</sup>

Department of Biological and Environmental Science and Nanoscience Center, Centre of Excellence in Biological Interactions, University of Jyväskylä, Jyväskylä, Finland<sup>a</sup>; Department of Biosciences and Institute of Biotechnology, University of Helsinki, Helsinki, Finland<sup>b</sup>

**ABSTRACT** FL-1, a myophage of *Flavobacterium*, was found to have a 53-kb genome with 87 putative coding sequences.

*Flavobacterium* spp. are abundant and widespread in freshwater environments (1, 2). Despite the prevalence of the members in this genus, the phages infecting *Flavobacterium* spp. are less known. So far, most of the phages studied in detail infect fish pathogenic members of the genus, such as *F. psychrophilum* (3).

The phage FL-1 and its *Flavobacterium* sp. host strain B183 were previously isolated from a water sample from a fish farm in Central Finland. Transmission electron microscopy revealed FL-1 to be a member of the family *Myoviridae* (4). In addition to the isolation host, FL-1 also infects multiple *Flavobacterium* sp. isolates and strains of the fish pathogen *F. columnare*, the causative agent of columnaris disease in fish (5).

DNA was extracted from FL-1 phage lysate, as described by Santos (6) with slight modifications as described earlier (7). DNA was sequenced on two platforms—Ion Torrent PMG with a 100-bp kit and commercially with Roche 454 (LGC Genomics)—and the data were combined (neither method resulted in a whole-genome sequence). Analyses were done using GS De Novo Assembler version 2.9 (Roche 454 Life Sciences), which uses an overlap layout consensus methodology. The average coverage was 11×, and the inferred read error rate was 0.94%. The assembly resulted in three scaffolds that were combined using Sanger sequencing. Glimmer and GeneMark were used for predicting the open reading frames (ORFs) using Geneious version 7.1 (Biomatters Ltd.), with possible functions predicted using BLASTp (8). tRNAscan-SE (9) was used to search for putative genes coding for tRNAs, but none were detected. The genome of FL-1 comprises 53,088 bp with a G+C content of 32.4%. Of the 87 predicted ORFs (ranging from 120 to 2,043 bp in length), only two were leftward oriented. Start codon usage was 94% for ATG, 3% for TTG as an initiation codon, and 2% for GTG. Putative functions were assigned for only six of the predicted coding sequences (CDSs), including a terminase, a portal protein, a phage lysin, and two tail proteins. The remaining 81 ORFs were assigned as hypothetical proteins. Primer walking confirmed the ends of the genome. Interestingly, only five of the CDSs had a recognizable Shine-Dalgarno sequence. However, the ribosomal binding site consensus sequence, TAAAA, has been proposed for environmental *F. hibernum* (10). Indeed, this sequence was also found in the upstream (20 bp) sequence for 34 of the CDSs and a TAAA sequence for 24 of the CDSs.

BLASTp analysis of the FL-1 predicted ORFs found several hits to the *Cellulophaga* phage phiSM and *F. columnare* phage FCL-2 genomes (21 and 30, respectively). An alignment of these three genomes using Mauve (11) revealed FL-1's relationship to both FCL-2 and phiSM, as all shared a conserved region that includes the putative structural proteins, indicating shared synteny.

Received 15 January 2017 Accepted 18 April 2017 Published 8 June 2017

**Citation** Laanto E, Ravantti JJ, Sundberg L-R. 2017. Complete genome sequence of an aquaculture-associated phage, FL-1, infecting *Flavobacterium* spp. Genome Announc 5: e00014-17. <https://doi.org/10.1128/genomeA.00014-17>.

**Copyright** © 2017 Laanto et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Elina Laanto, [elina.laanto@jyu.fi](mailto:elina.laanto@jyu.fi).

**Accession number(s).** The complete genome of *Flavobacterium* sp. phage FL-1 was deposited in GenBank under the accession number [KY421186](https://doi.org/10.1111/j.1462-2920.2004.00657.x).

## ACKNOWLEDGMENTS

This work was supported by the Centre of Excellence Program of the Academy of Finland; by the CoE in Biological Interactions 2012–2017 (grant no. 252411); by the Academy of Finland (grant no. 266879 and 304615); and by the Jane and Aatos Erkko Foundation. We thank Pilvi Ruotsalainen for skillful technical assistance.

## REFERENCES

- Eiler A, Bertilsson S. 2004. Composition of freshwater bacterial communities associated with cyanobacterial blooms in four Swedish lakes. *Environ Microbiol* 6:1228–1243. <https://doi.org/10.1111/j.1462-2920.2004.00657.x>.
- Eiler A, Bertilsson S. 2007. *Flavobacteria* blooms in four eutrophic lakes: linking population dynamics of freshwater bacterioplankton to resource availability. *Appl Environ Microbiol* 73:3511–3518. <https://doi.org/10.1128/AEM.02534-06>.
- Castillo D, Middelboe M. 2016. Genomic diversity of bacteriophages infecting the fish pathogen *Flavobacterium psychrophilum*. *FEMS Microbiol Lett* 363:272. <https://doi.org/10.1093/femsle/fnw272>.
- Laanto E, Sundberg LR, Bamford JKH. 2011. Phage specificity of the freshwater fish pathogen *Flavobacterium columnare*. *Appl Environ Microbiol* 77:7868–7872. <https://doi.org/10.1128/AEM.05574-11>.
- Declercq AM, Haesebrouck F, Van den Broeck W, Bossier P, Decostere A. 2013. Columnaris disease in fish: a review with emphasis on bacterium-host interactions. *Vet Res* 44:27. <https://doi.org/10.1186/1297-9716-44-27>.
- Santos MA. 1991. An improved method for the small scale preparation of bacteriophage DNA based on phage precipitation by zinc chloride. *Nucleic Acids Res* 19:5442–5442. <https://doi.org/10.1093/nar/19.19.5442>.
- Laanto E, Bamford JKH, Ravantti JJ, Sundberg LR. 2015. The use of phage FCL-2 as an alternative to chemotherapy against columnaris disease in aquaculture. *Front Microbiol* 6:829. <https://doi.org/10.3389/fmicb.2015.00829>.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* 215:403–410. [https://doi.org/10.1016/S0022-2836\(05\)80360-2](https://doi.org/10.1016/S0022-2836(05)80360-2).
- Schattner P, Brooks AN, Lowe TM. 2005. The tRNAscan-SE, snoscan and snoGPS web servers for the detection of tRNAs and snoRNAs. *Nucleic Acids Res* 33:W686–W689. <https://doi.org/10.1093/nar/gki366>.
- Chen S, Bagdasarian M, Kaufman MG, Walker ED. 2007. Characterization of strong promoters from an environmental *Flavobacterium hibernum* strain by using a green fluorescent protein-based reporter system. *Appl Environ Microbiol* 73:1089–1100. <https://doi.org/10.1128/AEM.01577-06>.
- Darling ACE, Mau B, Blattner FR, Perna NT. 2004. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res* 14:1394–1403. <https://doi.org/10.1101/gr.2289704>.