

Preface to the Special Issue “Integrated Monitoring in the Valkea-Kotinen Catchment during 1990–2009: Abiotic and Biotic Responses to Changes in Air Pollution and Climate”

Martti Rask¹, Lauri Arvola², Martin Forsius³ and Jussi Vuorenmaa³

¹ Finnish Game and Fisheries Research Institute, Evo Fisheries Research Station, Rahtijärventie 291, FI-16970 Evo, Finland

² University of Helsinki, Lammi Biological Station, FI-16900 Lammi, Finland

³ Finnish Environment Institute, P.O. Box 140, FI-00251 Helsinki, Finland

Increased emissions of air pollutants and greenhouse gases into the atmosphere have caused severe environmental problems at local and global scales. The long-range transport of sulphur and nitrogen oxides resulted in widespread acidification of acid-sensitive aquatic ecosystems in Europe and North America in the late 1900s (e.g. Rodhe *et al.* 1995). Airborne contamination of trace metals and persistent organic pollutants also reached the most remote regions of the world (e.g. AMAP 1998). At the same time, emissions of greenhouse gases into the atmosphere are causing global warming, and consequent climate change is considered to be among the most severe threats to the ecosystems (e.g. Rosenzweig *et al.* 2007).

Detrimental effects of transboundary air pollution led to international agreements on reducing emissions of SO₂ and NO_x in Europe and North America. International negotiations on emission reductions have been conducted under the Convention on Long-Range Transboundary Air Pollution (CLRTAP), signed in 1979 under the UN Economic Commission of Europe (UNECE 1996). Since the 1980s, environmental regulations have led to decreased emissions of air pollutants in Europe and North America. The effects of emission reductions have been monitored by the International Cooperative Programmes (ICP) of the CLRTAP convention (Bull *et al.* 2001).

The International Cooperative Programme on Integrated Monitoring of Air Pollution Effects on Ecosystems (UNECE ICP IM) contains today 44 research catchments in 16 countries. In Finland, the Integrated Monitoring Programme (IM) was initiated in 1987 (Bergström 1998). Finnish Environment Institute is the leading partner in the monitoring, accompanied by other governmental research institutes (Finnish Forest Research Institute, Finnish Game and Fisheries Research Institute, Finnish Meteorological Institute and Geological Survey of Finland) and universities (Helsinki, Eastern Finland and Oulu).

The Valkea-Kotinen Integrated Monitoring catchment is one of the three active IM sites in Finland and is also part of the Finnish Long-Term Socio-Ecological Research network (FinLTSER). As a consequence of the comprehensive long-term environmental monitoring together with diverse terrestrial and aquatic ecological research (e.g. Kurka and Starr 1997, Starr and Ukonmaanaho 2004, Jones *et al.* 1999, Vähätalo *et al.* 2003, Huotari *et al.* 2009, Peltonmaa *et al.* 2013a, 2013b), the Valkea-Kotinen site has grown into a major Finnish research infrastructure and data source for environmental modelling (Forsius *et al.* 1998, Futter *et al.* 2009, Saloranta *et al.* 2009, Holmberg *et al.* 2014).

This special issue of *Boreal Environment Research* summarizes the main results of the Integrated Monitoring of the Valkea-Kotinen

catchment over a 20-year period (1990–2009). The seven papers included provide findings on the climate variability and trends (Jylhä *et al.* 2014), development in bulk deposition and atmospheric concentration of acidifying compounds and trace elements (Ruoho-Airola *et al.* 2014), long-term water quality trends in Lake Valkea-Kotinen (Vuorenmaa *et al.* 2014), and modelling of climate change effects on the hydrology and carbon processes of Valkea-Kotinen catchment (Holmberg *et al.* 2014). The remaining three papers present long-term biological responses to environmental changes in Lake Valkea-Kotinen, including plankton metabolism and sedimentation (Arvola *et al.* 2014), zooplankton community patterns (Lehtovaara *et al.* 2014), and perch (*Perca fluviatilis*) population dynamics (Rask *et al.* 2014).

In summary, the results clearly demonstrate the complexity of boreal nature and interactions within and between the aquatic and terrestrial ecosystems, and atmosphere. Further, these studies strongly emphasize the importance of integrated long-term monitoring of physical, chemical and biological variables for detecting the variety of impacts of changing environmental conditions on ecosystems.

References

- AMAP 1998. *AMAP assessment report: Arctic pollution issues*. Arctic Monitoring and Assessment Programme (AMAP), Oslo, Norway.
- *Arvola L., Salonen K., Keskitalo J., Tulonen T., Järvinen M. & Huotari J. 2014. Plankton metabolism and sedimentation in a small boreal lake — a long-term perspective. *Boreal Env. Res.* 19 (suppl. A): 83–96.
- Bergström I. 1998. The Integrated Monitoring Programme in Finland. *Boreal Env. Res.* 3: 201–203.
- Bull K.R., Achermann B., Bashkin V., Christ R., Fenech G., Forsius M., Gregor H.-D., Guardans R., Haußmann T., Hayes F., Hettelingh J.-P., Johannessen T., Krzyzanowski M., Kucera V., Kvaeven B., Lorenz M., Lundin L., Mills G., Posch M., Skjelkvåle B.L. & Ulstein J.M. 2001. Coordinated effects monitoring and modelling for developing and supporting international air pollution control agreements. *Water Air and Soil Pollution* 130: 119–130.
- Forsius M., Alveteg M., Jenkins A., Johansson M., Kleemola S., Lükeville A., Posch M., Sverdrup H. & Walse C. 1998. MAGIC, SAFE and SMART model applications at Integrated Monitoring sites: Effects of emission reduction scenarios. *Water Air and Soil Pollution* 105: 21–30.
- Futter M.N., Forsius M., Holmberg M. & Starr M. 2009. A long-term simulation of the effects of acidic deposition and climate change on surface water dissolved organic carbon concentrations in a boreal catchment. *Hydrology Research* 40: 291–305.
- *Holmberg M., Futter M.N., Kotamäki N., Fronzek S., Forsius M., Kiuru P., Pirttioja N., Rasmus K., Starr M. & Vuorenmaa J. 2014. Effects of a changing climate on the hydrology of a boreal catchment and lake DOC — probabilistic assessment of a dynamic model chain. *Boreal Env. Res.* 19 (suppl. A): 66–82.
- Huotari J., Ojala A., Peltomaa E., Pumpanen J., Hari P. & Vesala T. 2009. Temporal variations in surface water CO₂ concentration in a boreal humic lake based on high-frequency measurements. *Boreal Env. Res.* 14: 48–60.
- *Jylhä K., Laapas M., Ruosteenoja K., Arvola L., Drebs A., Kersalo J., Saku S., Gregow H., Hannula H.-R. & Pirinen P. 2014. Climate variability and trends in the Valkea-Kotinen region, southern Finland: comparisons between the past, current and projected climates. *Boreal Env. Res.* 19 (suppl. A): 4–30.
- Kurka A.-M., Starr M., Heikinheimo M. & Salkinoja-Salonen M. 2000. Decomposition of cellulose strips in relation to climate, litterfall, nitrogen, phosphorus and C/N ratio in natural boreal forests. *Plant and Soil* 219: 91–101.
- *Lehtovaara A., Arvola L., Keskitalo J., Olin M., Rask M., Salonen K., Sarvala J., Tulonen T. & Vuorenmaa J. 2014. Responses of zooplankton to long-term environmental changes in a small boreal lake. *Boreal Env. Res.* 19 (suppl. A): 97–111.
- Peltomaa E., Zingel P. & Ojala A. 2013. Weak response of the microbial food web of a boreal humic lake to hypolimnetic anoxia. *Aquat. Microb. Ecol.* 68: 91–105.
- Peltomaa E., Ojala A., Holopainen A.-L. & Salonen K. 2013. Changes in phytoplankton in a boreal lake during a 14-year period. *Boreal Env. Res.* 18: 387–400.
- *Rask M., Sairanen S., Vesala S., Arvola L., Estlander S. & Olin M. 2014. Population dynamics and growth of perch in a small, humic lake over a twenty year period — importance of abiotic and biotic factors. *Boreal Env. Res.* 19 (suppl. A): 112–123.
- Rodhe H., Grennfelt P., Wisniewski J., Ågren C., Bengtsson G., Johansson K., Kauppi P., Kucera V., Rasmussen L., Rosseland B., Schotte L. & Sellden G. 1995. Acid Reign '95? — Conference summary statement. *Water Air Soil Pollution* 85: 1–14.
- Rosenzweig C., Casassa G., Karoly D.J., Imeson A., Liu C., Menzel A., Rawlins S., Root T.L., Seguin B. & Tryjanowski P. 2007. Assessment of observed changes and responses in natural and managed systems. In: Parry M.L., Canziani O.F., Palutikof J.P., van der Linden P.J. & Hanson C.E. (eds.), *Climate change 2007: Impacts, adaptation and vulnerability*, Contribution of Working Group II to the Fourth Assessment, Report of the Intergovernmental Panel on Climate Change, Cambridge Univ. Press, Cambridge, pp. 79–131.
- *Ruoho-Airola T., Hatakka T., Kyllönen K., Makkonen U. & Porvari P. 2014. Temporal trends in the bulk deposition and atmospheric concentration of acidifying compounds and trace elements in the Finnish Integrated Monitoring catchment Valkea-Kotinen during 1988–2011. *Boreal*

Env. Res. 19 (suppl. A): 31–46.

Saloranta T.M., Forsius M., Järvinen M. & Arvola L. 2009. Impacts of projected climate change on the thermodynamics of a shallow and a deep lake in Finland: model simulations and Bayesian uncertainty analysis. *Hydrology Research* 40: 234–248.

Starr M. & Ukonmaanaho L. 2004. Levels and characteristics of TOC in throughfall, forest floor leachate and soil solution in undisturbed boreal forest ecosystems. *Water Air Soil Pollution, Focus* 4: 715–729.

UNECE 1996. *1979 Convention on long-range transbound-*

ary air pollution and its protocols. United Nations Economic Commission for Europe, New York and Geneva.

*Vuorenmaa J., Salonen K., Arvola L., Mannio J., Rask M. & Horppila P. 2014. Water quality of a small headwater lake reflects long-term variations in deposition, climate and in-lake processes. *Boreal Env. Res.* 19 (suppl. A): 47–65.

Vähätalo A.V., Salonen K., Münster U., Järvinen M. & Wetzel R.G. 2003. Photochemical transformation of allochthonous organic matter provides bioavailable nutrients in a humic lake. *Arch. Hydrobiol.* 156: 287–314.