

Trematominae and Artedidraconinae: contrasted mitogenome evolution for two Antarctic radiations

Dettai Agnes¹, Juliette Auvinet², Bruno Frederich³, Henrik Christiansen³, Najda Aboubakari¹, Céline Bonillo⁴ and Dominique Higuët²

¹ Sorbonne Universités, UMR 7205 ISYEB Institut de Systématique, Evolution, Biodiversité, Muséum National d'Histoire Naturelle, 57 rue Cuvier, 75231 Paris cedex 05, France

E-mail: agnes.dettai@mnhn.fr

² Sorbonne Universités, UMR 7138 Systématique Adaptation Evolution, Université Pierre et Marie Curie, 7-9 quai Staint-Bernard, 75005 Paris France.

³ KU Leuven, Laboratory of Biodiversity and Evolutionary Genomics (LBEG), Ch. Deberiotstraat 32, B-3000 Leuven, Belgium.

⁴ Sorbonne Universités, UMS2700 OMSI Outils et Methodes de la Systématique Intégrative, Muséum National d'Histoire Naturelle, 57 rue Cuvier, 75231 Paris cedex 05, France

Cellular respiration has been widely studied in Antarctic teleost fishes because of their peculiar adaptations to an extreme environment. In parallel mitochondrial sequence markers have become highly popular for molecular systematics. However, there are few whole mitochondrial genome sequences published, and none available for some of the subfamilies. Here, we present two large mitogenome datasets including most species and multiple sequences for many species of two subfamilies, Trematominae and Artedidraconinae (Duhamel et al. 2014). These include two highly diverse but very different adaptative radiations, with contrasting divergence dates, morphological polymorphism, and habitat dominance. The sampling is based on a well identified, extensive collection resulting from the 2008 CEAMARC survey and the subsequent REVOLTA surveys in Terre Adélie (IPEV), already DNA barcoded and sequenced in previous studies. The mitogenome sequences for these two subfamilies differ in composition, gene order, and relative divergence of mitochondrial markers, with strong, taxon-specific biases like very high C contents in some regions. The gene order change provides a synapomorphy for the subfamily Trematominae and an interesting development in teleost mitogenomes. The complete Artedidraconinae mitogenomes provide a much higher amount of variable sites (approx*30), while previous sequence datasets were plagued by low informativeness (Lecointre et al. 2011). As already established on single mitochondrial genes, intraspecific variability is lower than interspecific variability within each subfamily, however interspecific variability in Artedidraconinae is lower or similar to intraspecific variability in Trematominae. This expanded dataset confirms the unusual evolution of the mitochondrial coded sequences involved in the cellular respiration in Antarctic Nototheniidae, as well as the usefulness of complete mitochondrial genomes for their systematics. The two level multiplexing (Timmermans et al. 2010) and next generation sequencing of long PCR amplicons (following Hinsinger et al. 2015) is efficient to obtain large mitogenomic datasets representative of both inter- and intraspecific variability, key to the understanding of mitochondrial evolution and a step closer to resolving the relationships among these taxa.

References

- Duhamel, G., Hulley, P.-A., Causse, R., Koubbi, P., Vacchi, M., Pruvost, P., Vigetta, S., Irisson, J.-O., Mormède, S., Belchier, M., Dettai, A., Detrich, H.W., Gutt, J., Jones, C.D., Kock, K.-H., Lopez Abellan, L. J., Van de Putte, A. 2014. Chapter 7: Biogeographic Patterns of Fish. In: *Biogeographic Atlas of the Southern Ocean*. De Broyer, C, Koubbi, P., Griffiths, H., Raymond, B., d'Udekem d'Acoz, C., Van de Putte, A., Danis, B., David, S., Grant, S., Gutt, J., Held, C., Hosie, G., Huettmann, F., Post, A., Ropert-Coudert, Y. (eds.), Cambridge, Scientific Committee on Antarctic Research, 328-362.
- Hinsinger D.D., Debruyne R., Thomas M., Denys G.P.J., Mennesson M., Utge J., Dettai A. 2015. Fishing for barcodes in the Torrent: from COI to complete mitogenomes on NGS platforms. *DNA Barcodes* 3: 170-186.
- Lecointre, G., Gallut, C., Bonillo, C., Couloux, A., Ozouf-Costaz, C., Dettai, A. 2011. The Antarctic Fish Genus *Artedidracon* is paraphyletic (Teleostei, Notothenioidae, Artedidraconidae). *Polar Biol.* 34(8): 1135-1145 DOI: 10.1007/s00300-011-0974-5.
- Timmermans M.J., Dodsworth S, Culverwell CL, Bocak L, Ahrens D, Littlewood DT, Pons J, Vogler AP 2010. Why barcode? High-throughput multiplex sequencing of mitochondrial genomes for molecular systematics. *Nucleic Acids Res* 38(21): e197. doi: 10.1093/nar/gkq807.