

SCIENTIFIC REPORTS

OPEN

The association of bacterial C₉-based TTX-like compounds with *Prorocentrum minimum* opens new uncertainties about shellfish seafood safety

Received: 30 September 2016

Accepted: 12 December 2016

Published: 20 January 2017

Inés Rodríguez¹, Amparo Alfonso¹, Eva Alonso¹, Juan A. Rubiolo¹, María Roel¹, Aristidis Vlamis^{1,2}, Panagiota Katikou², Stephen A. Jackson³, Margassery Lekha Menon³, Alan Dobson³ & Luis M. Botana¹

In 2012, Tetrodotoxin (TTX) was identified in mussels and linked to the presence of *Prorocentrum minimum* (*P. minimum*) in Greece. The connexion between TTX and *P. minimum* was further studied in this paper. First, the presence of TTX-producer bacteria, *Vibrio* and *Pseudomonas* spp, was confirmed in Greek mussels. In addition these samples showed high activity as inhibitors of sodium currents (I_{Na}). *P. minimum* was before associated with neurotoxic symptoms, however, the nature and structure of toxins produced by this dinoflagellate remains unknown. Three *P. minimum* strains, ccmp1529, ccmp2811 and ccmp2956, growing in different conditions of temperature, salinity and light were used to study the production of toxic compounds. Electrophysiological assays showed no effect of ccmp2811 strain on I_{Na} , while ccmp1529 and ccmp2956 strains were able to significantly reduce I_{Na} in the same way as TTX. In these samples two new compounds, m/z 265 and m/z 308, were identified and characterized by liquid chromatography tandem high-resolution mass spectrometry. Besides, two TTX-related bacteria, *Roseobacter* and *Vibrio* sp, were observed. These results show for the first time that *P. minimum* produce TTX-like compounds with a similar ion pattern and C₉-base to TTX analogues and with the same effect on I_{Na} .

Prorocentrum minimum (*P. minimum*) is a widely distributed neritic bloom-forming dinoflagellate first described by Pavillard in 1916. Since then, more than ten taxonomy synonyms have been used to describe the same micro-organism. This controversy and confusion with names was related to the geographic area where it was isolated, including tropical, subtropical and temperate climates¹. Blooms can occur in a wide range of environmental conditions since *P. minimum* has been described as eurythermal and euryhaline².

P. minimum is potentially harmful to humans through consumption of toxic seafood. Over the years several human intoxications have been associated with the presence of *P. minimum* blooms. This dinoflagellate was considered responsible for more than 200 deaths in Japan, in 1942 and 1943, with symptoms such as liver injury, haemorrhage, unconsciousness, coma, and death after 24–48 hours. The toxin described in those episodes was a hydrophilic compound named venerupin and the syndrome was called Venerupin Shellfish Poisoning (VSP)^{3–5}. At that time, *P. minimum* was considered the source of venerupin although no definitive links between dinoflagellate and toxin were reported¹. Later on *P. minimum* was involved in several episodes of human poisoning after shellfish consumption in Portugal (Obidos Lagoon). The symptoms were considered to be characteristic of paralytic shellfish poisoning (PSP)⁶. The definitive link between *P. minimum* and neurotoxic effects was established with the analysis of several axenic clones of *P. minimum* from French coastal samples. These dinoflagellates produced a water-soluble neurotoxic compound, different from PSP, which was able to rapidly kill mice after

¹Departamento de Farmacología. Facultade de Veterinaria, Universidade de Santiago de Compostela, Lugo, Spain.

²National Reference Laboratory on Marine Toxins, Veterinary Center of Thessaloniki, Ministry of Rural Development and Food, 3A Limnou street, GR-54627 Thessaloniki, Greece. ³School of Microbiology, University College Cork, Cork, Ireland. Correspondence and requests for materials should be addressed to L.M.B. (email: luis.botana@usc.es)




Figure 1. (A) Rarefaction curves for white tissue (WT) and TTX contaminated samples (1770/2012 and 1774/2012) samples showing the OTU based (97% 16S rRNA gene sequence similarity) number of observed species as a function of number of sequences per sample. (B) Bootstrapped tree for hierarchical samples clustering.

neurotoxic symptoms^{7,8}. This unknown compound was able to block sodium and calcium channels and was accumulated by shellfish⁷. However, since some of the toxic episodes related to *P. minimum* were also associated with the presence of other toxic microalgae such as *Dinophysis* spp., its toxicity was discussed¹. This coexistence does not eliminate the potential risk of *P. minimum* blooms for human health, being evident that the reported toxicity of this dinoflagellate was clone-related and also related to environmental conditions¹. In this sense, although most *P. minimum* clones were reported as non-toxic, and rare human toxicity episodes have been attributed to this specie, it should be considered for risk assessment.

Tetrodotoxin (TTX) is one of the most important neurotoxins known to block sodium channels and thus it inhibits the propagation of action potentials in muscle and nerve cells. The minimum lethal dose for TTX is 8 µg/Kg and the median lethal dose is 10 µg/Kg⁹. Symptoms can appear between 10 and 45 minutes after exposure, although depending of the amount of the toxin ingested some manifestations could appear 6 hours later. This compound blocks site 1 of voltage dependent sodium channels¹⁰ and it has interesting pharmacological applications in the study of excitable membranes as well as therapeutic uses in treating migraines, addictions, or as an anaesthetic agent for pain¹¹.

TTX has been described in different aquatic animals such as fish, arthropods, echinoderms, molluscs, worms, newts, frogs or toads as well as several bacteria species and dinoflagellates^{11,12}. However, the origin of TTX is still unclear, although the presence of this toxin in such a diverse group of animals suggests that bacteria might be as the primary source¹³. In this sense, TTX can be produced without bacteria, but the presence of microorganisms could, in some way, simplify TTX biosynthesis¹¹.

To date, few appearances of TTX in bivalve molluscs have been reported^{14,15}. Recently, TTX has been detected in mussels and Pacific oyster from the English coast (2013–2014)¹⁶. In addition in 2012, in Greece, during routine controls of shellfish, an unexplained toxicity associated with neurological symptoms was observed in a series of mouse bioassays (MBA). This atypical toxicity coincided with the absence of known dinoflagellates and other toxins, while only *P. minimum* was present in seawater. After mass-spectrometry analysis, both TTX and TTX analogues were confirmed¹⁷. Therefore a potential TTX presence-dinoflagellate link was hypothesized. In this context the aim of this work was to further study the relationship between TTX and *P. minimum* blooms in order to identify and characterize the compounds produced by this dinoflagellate.

Results

To further study the occurrence and origin of TTXs in mussels collected in the unusual toxic episode in Greece, it was evaluated the presence of bacteria as TTX producers as well as the effect on Na_v channels.

Firstly, the presence of bacteria was checked in contaminated mussel-samples, 1770/2012 and 1774/2012, from the Greek episode. Levels of 222.9 µg/Kg and 206.3 µg/Kg of TTXs respectively were reported when these samples were previously analysed by mass spectrometry¹⁷. Sequence reads were clustered at 97% identity resulting in 297 unique Operational Taxonomic Units (OTUs). A rarefaction plot of observed bacterial species in control (white non-contaminated tissue) and TTX-mussel samples showed higher species diversity in the uncontaminated material (Fig. 1A). Although high species diversity was achieved, uncontaminated sample was under represented when compared to TTX-mussel samples. Jackknifed hierarchical clustering by UPGMA (unweighted pair group method with arithmetic mean) showed a high support for the deputed samples clustering separate from the non-deputed one, while a lower support was observed for contaminated samples (Fig. 1B). In this way, the identified OTUs corresponded to 9 bacterial phyla (Supplementary Table 2 and Sub-Table 2). Higher percentages of proteobacterial OTUs, the phylum containing *Pseudomonas*, *Pseudoalteromonas*, and *Vibrio*, two known TTX producers, were detected in TTX-contaminated mussels. However these genera were not represented in the Proteobacteria of non-contaminated sample. To better identify *Vibrio* and *Pseudomonas* species, OTUs were re-picked with 100% homology. When compared to the GreenGenes 16S RNA database¹⁸, several potential TTX




Figure 2. Effect of mussel sample 1786/2012 over I_{Na} current activity. Automated patch clamp experiments were carried out in hNav1.6 transfected cells. I_{Na} magnitude is expressed as percentage of basal current. (A) Dose-response inhibition of serial dilutions of 1786/2012 mussel sample over I_{Na} . 0.4, 2 and 4 μ l of extract were added to 200 μ l of extracellular solution. Significant differences (*) $p < 0.05$. (B) Dose-response inhibition of TTX standard over I_{Na} with an IC_{50} of 0.43 to 1.2 nM. A discontinuous line marks the 50% of sodium current (A and B) and the corresponding TTX standard concentration (B).

producers were identified in both TTX-contaminated samples (Supplementary Table 3). The distribution of these bacteria showed that in both samples *Vibrio parahaemolyticus* and *Vibrio alginolyticus* were the dominant species, while *Pseudomonas* appeared at lower levels as well as other *Vibrio* species.

Next, to further support the existence of TTX in mussel extracts, the effect over Na_v channels, the main cellular target of TTX, was checked by patch clamp measurements in stable transfected hNav1.6 HEK cells^{19,20}. Aliquots of 2 g of DG of sample 1786/2012 collected in Greece were extracted as previously described¹⁷. As Fig. 2 shows, serial dilutions of the TTX-positive mussel extract 1786/2012 were tested (0.4, 2 and 4 μ l) and a dose response effect was reported. At the highest concentration tested, an inhibition of $54.65 \pm 14.08\%$ versus control current was observed (Fig. 2A). In the same cells, the control of TTX standard elicited a dose response sodium current (I_{Na}) inhibition, IC_{50} 0.72 nM (95% confidence interval 0.43 to 1.2 nM) (Fig. 2B).

Although the source of TTXs in contaminated mussels seemed to be related to bacteria, the dinoflagellate *P. minimum* was present in water in high abundance when toxic mussels were harvested, therefore the presence of toxins and bacteria in this microalgae was studied. Three strains of *P. minimum* from different collection sites, ccmp1529 from Ecuador, temperature range 18 °C to 22 °C, ccmp2811 from Sarasota, Florida (USA), temperature range 18 °C to 22 °C, and ccmp2956 from Johor Strait, between Singapore and Malaysia, temperature range 22 °C to 28 °C, were cultured in L1 medium in different growth conditions, varying light intensity (785, 840 and 865), salinity (ranging from 31‰ to 37‰) and temperature (19 °C or 24 °C), with each condition being carried out in triplicate. After 15 days of growth, cells were harvested and the extracts obtained dissolved in acetic acid 0.03 M and analysed for the presence of active compounds with similar characteristics to TTXs. The same rate of growth was obtained for ccmp1529 strain either at 19 °C and 24 °C, slightly less growth was obtained when the ccmp2956 strain was grown at 19 °C compared to 24 °C and no growth was observed when the ccmp2811 strain was cultured at 24 °C. Therefore ccmp1529 and ccmp2956 strains were analysed both at 19 °C and 24 °C, while only cultures obtained at 19 °C were analysed in the case of ccmp2811. No significant effect, in terms of growth, was observed when light was modified and only ccmp2956 strain was able to grow at 37‰ of salinity. Then, the effect of dinoflagellate extracts on I_{Na} was checked. Electrophysiological recordings showed that while ccmp1529 extracts did not produce any effect on I_{Na} when dinoflagellates were grown at 19 °C, the addition of serial dilutions of 24 °C extract culture reveal a dose dependent current reduction with a maximal inhibition of $49.75 \pm 8.04\%$ versus control current (Fig. 3A). This effect was not dependent on light intensity. Likewise, the ccmp2956 strain cultured at 19 °C had no effect on I_{Na} but when it was grown at 24 °C and 37‰ salinity, a maximal current inhibition of $47.65 \pm 10.6\%$ versus control was observed (Fig. 3B). When salinity was 31–33‰, I_{Na} inhibition was dependent on light intensity, since no effect was observed with dinoflagellates cultured at 18 W/840, while $39.69 \pm 12.03\%$ inhibition was obtained at 18 W/865. ccmp2811 extracts from cultures grown at 19 °C did not produce a significant I_{Na} inhibition when consecutive dilutions were added (Fig. 3C). For all these assays TTX standard was used as control of I_{Na} inhibition (Fig. 3D). Therefore, these results indicate the presence of some compound produced by *P. minimum* with a similar effect than TTX on I_{Na} . The profile of compounds present in cultures grown at 19 °C or 24 °C seems to be different, at least with regard to potency over I_{Na} inhibition.

Next, we proceeded to the identification of compounds produced by *P. minimum* cultures using mass spectrometry technology. The first screening of culture extracts was done by Multiple Reaction Monitoring (MRM) mode, searching for the most common TTX analogues described in the literature (Supplementary Table 1). In this way, 5 compounds with same product ions, but different retention times than TTX standards were obtained, m/z 320 > 302, m/z 304 > 162, m/z 302 > 162, m/z 290 > 272, m/z 272 > 254. Then, by operating in the Product Ion Scan mode, the fragmentation pathway of these five compounds was checked following the pattern fragmentation observed in TTX, 4,9-anhydroTTX and 11-deoxyTTX standards (Supplementary Figure 1B,C and D). That is, for TTX (m/z 320) the ions m/z 302 [$M + H - H_2O$]⁺, m/z 162 and m/z 178 (2-aminohydroxyquinazoline and 2-aminodihydroxyquinazoline, respectively). For 4,9-anhydroTTX (m/z 302) the ions m/z 284 [$M + H - H_2O$]⁺,




Figure 3. Effect of *P. minimum* extracts growing in different conditions over I_{Na} activity. Automated patch clamp experiments were carried out in hNav1.6 transfected cells. I_{Na} magnitude is expressed as percentage of basal current. **(A)** Effect of extracts from *ccmp1529* strain cultured at 19 and 24 °C and different light exposures over I_{Na} . **(B)** Effect of extracts from *ccmp2956* strain cultured at 19 and 24 °C, two light exposures and two different salinities over I_{Na} . **(C)** Effect of extracts from *ccmp2811* strain cultured at 19 °C over I_{Na} . **(D)** TTX dose-response inhibition of I_{Na} . **(E and F)** Representative current traces of I_{Na} in the presence of an extract from *ccmp1529* and TTX. Results are mean \pm SEM of 3 experiments, each performed in duplicate. Significant differences * $p < 0.05$, ** $p < 0.01$.

m/z 162 and m/z 178. And in the case of 11-deoxyTTX (m/z 304) the ions m/z 286 $[M + H - H_2O]^+$, m/z 162 and m/z 176²¹. The five compounds initially observed in *P. minimum* cultures did not follow this fragmentation pathway (data not show), therefore these five compounds could not be identified as TTX analogues. However, since *P. minimum* extracts had some TTX activity, mass spectrometry operating procedure was changed from Product to Precursor Ion Scan Mode. Thus the search of compounds was addressed from the common product ion for all TTX analogues, m/z 162, to precursor ions. In this way, operating in the Precursor Ion Scan mode, the extract from *ccmp2956* strain (24 °C and 37‰) with high inhibitory effect on I_{Na} , was checked. In this sample two precursor ions, m/z 265 and m/z 308, were observed (Fig. 4). Chromatograms obtained by Ultra performance Liquid Chromatography tandem mass spectrometry (UPLC-MS/MS) show two peaks with retention times of 3.8 min and 8 min respectively (Fig. 4A and C). The Product Ion Scan of these peaks shows for the parent peak m/z 265 $[M + H]^+$, the ions m/z 247 corresponding to $[M + H - H_2O]^+$, m/z 178.9 and m/z 162, Fig. 4B, and in the case of m/z 308 $[M + H]^+$ the ions m/z 290 corresponding to $[M + H - H_2O]^+$, m/z 162 and m/z 179.8 (Fig. 4D).




Figure 4. Product Ion Scan of TTX analogues m/z 265 and m/z 308 in ccmp2956 strain extract. (A) Product Ion Scan chromatogram obtained in positive mode of compound m/z 265. (B) Product Ion Scan chromatogram obtained in positive mode of compound m/z 308. (C) Mass spectrum obtained in Product Ion Scan of compound m/z 265. (D) Mass spectrum obtained in Product Ion Scan of compound m/z 308.

Compound	MS ⁿ	Elemental composition	Theoretical value (m/z)	[M + H] ⁺ (m/z)	Error (mDa)	Error (ppm)
TTX	MS ¹	C ₁₁ H ₁₇ N ₃ O ₈	320.1088	320.1099	1.1	3.44
	MS ²	C ₁₁ H ₁₅ N ₃ O ₇	302.0983	302.0984	0.1	0.33
	MS ³	C ₁₁ H ₁₃ N ₃ O ₆	284.0877	284.0923	4.6	16.19
		C ₁₁ H ₁₁ N ₃ O ₅	266.0771	266.0828	5.7	21.42
		C ₁₀ H ₁₃ N ₃ O ₅	256.0928	256.0957	2.9	11.32
		C ₈ H ₇ N ₃ O ₂	178.0611	178.0621	1.0	5.62
		C ₈ H ₇ N ₃ O	162.0662	162.0683	2.1	12.96
m/z 265	MS ¹	C ₉ H ₂₀ N ₄ O ₅	265.1506	265.1543	3.7	13.95
	MS ²	C ₉ H ₁₈ N ₄ O ₄	247.1401	247.1434	3.3	13.35
		C ₉ H ₁₀ N ₄ O	191.0927	191.1192	26.5	138.67
		C ₉ H ₁₄ N ₄	179.1291	179.1199	-9.2	-51.36
		C ₉ H ₁₁ N ₃	162.1026	162.0927	-9.9	-61.08
m/z 308	MS ¹	C ₁₁ H ₂₁ N ₃ O ₇	308.1452	308.1183	-26.9	-87.30
	MS ²	C ₁₁ H ₁₉ N ₃ O ₆	290.1347	290.1317	-3.0	-10.34
		C ₉ H ₁₃ N ₃ O	180.1131	180.0999	-13.2	-73.29
		C ₉ H ₁₁ N ₃	162.1026	162.0842	-18.4	-113.52

Table 1. MSⁿ Data for TTX standard and compounds m/z 308 and m/z 265.

To characterize and predict the molecular formula of these two new compounds, m/z 308 and m/z 265, and their product-ions, a high-resolution mass spectrometry Ion Trap-Time of Flight (IT-TOF) was used. This technology allows us to predict the formula using *ad oc* predictor software based on the accurate mass data recorded. TTX standard was employed as control to predict the molecular formula of precursor and ion products. In this way, elemental composition, accurately measured mass/charge ratio, theoretical value m/z , and mass errors expressed in ppm and mDa for TTX, m/z 308 and m/z 265 were obtained after MSⁿ spectra. These data are summarized in Table 1 and shown in Figs 5, 6 and 7. As Fig. 5 shows, TTX (m/z 320.1099, [C₁₁H₁₇N₃O₈ + H]⁺) ion was observed in MS¹ spectrum, then in MS² spectrum ions [M + H - H₂O]⁺ at m/z 302.0984 and [M + H - 2H₂O]⁺ at m/z 284.0899 were obtained. The other characteristic ions, [M + H - C₃H₁₀O₇]⁺ at m/z 162.0683 and [M + H - C₃H₁₀O₆]⁺ at m/z 178.0621 and the losses 2 and 3 H₂O molecules at m/z 284.0923 and m/z 266.0828 respectively and [M + H - CH₄O₃]⁺ at m/z 256.0957 were observed at MS³ spectrum. The errors between these measured ion products and the theoretical values range from 0.1–5.7 mDa, Table 1, indicating a good accuracy




Figure 5. Accurate MS^{1-3} spectra of TTX standard in positive mode (m/z 320 and molecular formula $C_{11}H_{17}N_3O_8$). (A) MS^1 range m/z 150–50. (B) MS^2 of m/z 320. (C) MS^3 of m/z 302 with the predicted structures of the fragments 256, 178 and 162.

between the predicted molecular formula either for TTX or its ion products. Similarly, m/z 265 and m/z 308 compounds were analysed and their MS^n product-ions spectra are shown in Figs 6 and 7. The ion products of 265-compound (m/z 265.1543, $[C_9H_{20}N_4O_5 + H]^+$) observed in MS^2 spectrum were: $[M + H - H_2O]^+$ ion at m/z 247.1434, $[M + H - H_{10}O_4]^+$ ion at m/z 191.1192, $[M + H - H_6O_5]^+$ ion at m/z 179.1199, and $[M + H - H_9NO_5]^+$ ion at m/z 162.0927 (Fig. 6). For the 308-compound (m/z 308.1183, $[C_{11}H_{21}N_3O_7 + H]^+$), the ion products observed in MS^2 spectrum were: $[M + H - H_2O]^+$ ion at m/z 290.1317, $[M + H - C_2H_8O_6]^+$ ion at m/z 180.0999 and $[M + H - C_2H_{10}O_7]^+$ ion at m/z 162.0842 (Fig. 7). Since these compounds are present in low concentration in *P. minimum* extracts only MS^2 spectra were obtained. The errors between these measured product-ions and the theoretical values calculated are shown in Table 1. In this case the higher range observed was probably related to the low concentration. In addition the predicted structure proposed for each compound based on the high resolution masses of fragment ions are shown in Figs 5, 6 and 7.

Next, the presence of this characterized compounds, m/z 308 and m/z 265, was checked in all *P. minimum* extracts, using $[M + H - H_2O]^+$ ions for identification and m/z 162 ions for quantification. Since no standards of these compounds were available, results were obtained as peak area and also as ng of TTX equivalent/mL (see Supplementary Table 4). Both compounds were present in all *P. minimum* extracts. While the amount of compound 308 did not show a direct relation with the effect on I_{Na} , the extracts with higher amount of compound 265 produced higher I_{Na} inhibition.

Finally, the presence of TTX-producing bacteria was checked in *P. minimum* cultures. For this experiment, ccmp1529 strain was cultured both at 19 and 24 °C, ccmp2956 strain was cultured at 19 °C, and ccmp2811 cultures were obtained at 24 °C all 31–33‰ salinity. Firstly, we checked for the presence of bacteria described in contaminated mussels, *Vibrio* and *Pseudomonas* spp. *P. minimum* cultures were analysed by PCR using specific primers for *Vibrio alginolyticus*, *Vibrio parahaemolyticus*, *Vibrio vulnificus*, and *Vibrio cholerae*. After PCR amplification, *V. alginolyticus* was observed in ccmp2811 and ccmp1529 cultures, while none were detected in sample ccmp2956 (Supplementary Figure 2). When the same assay was performed using specific primers for *Pseudomonas*, no positive amplification product was detected (results not shown). In addition, *P. minimum* cultures were analysed for




Figure 6. Accurate MS¹⁻² spectra of compound *m/z* 265 and molecular formula C₉H₂₀N₄O₅. (A) MS¹ range *m/z* 150–500. (B) MS² of *m/z* 265 with the proposed structures of fragments 191, 179 and 162. (C) Proposed structure for *m/z* 265 (NH₂ group can be in different positions).

the presence of other marine bacteria, resulting in the identification of several bacteria from α -*Proteobacterium* class, including *Roseobacter* and *Flavobacteria* (Supplementary Figure 3 and Table 5).

Discussion

The presence of TTX in mussels is an important issue from both a toxicological and human-health risk standpoint. Although this compound is a regulated toxin in Europe and around the world, it is not included in regular monitoring programs because the consumption of species that usually accumulate the toxin is prohibited. Before 2014 only one highly toxic TTX-accumulation in scallops from Japan had been reported¹⁴. However, from 2014 to date three reports have identified the presence of TTX in shellfish and it was also detected in several edible gastropods^{15–17,22,23}. Therefore, the presence of TTXs in shellfish as well as the source of these toxins should be taken into account, since TTX-contaminated samples will not be detected by official analytical methods, such as LC-MS/MS or HPLC with fluorescence detection, because TTX is not required to be included in these methods. In addition, the MBA, that could detect all toxins, is no longer used in Europe as an official assay. In this paper the presence of TTX in contaminated samples collected in Greece was confirmed through the effect on I_{Na}. In addition, *V. parahaemolyticus* and *V. alginolyticus* species, two known TTX-producers, were identified in these samples¹¹. Besides this, several species from *Pseudomonas* genus were also observed. This genus had previously been reported in TTX production, although the species were not specified^{24,25}. From our data, both *Vibrio* and *Pseudomonas* sp. could be detected in TTX-contaminated mussel samples. In the case of *Pseudomonas*, several species are confirmed by 99% sequence homology and for the first time they can be clearly linked to TTX and contaminated mussels¹⁷.

Therefore, the connection between shellfish, bacteria and TTX in the toxic episode earlier described in Greece was clear, however, a *P. minimum* bloom was occurring at the same time and it could also be connected. Since the toxin production by this dinoflagellate has not to date been clarified, we studied its growth and toxicity in different conditions. In this way, depending on the salinity, temperature and light, *P. minimum* strains produce some compounds with an inhibitory effect on I_{Na}, the same as TTX. The inhibitory effect on transfected hNa_v1.6 HEK cells was higher when *P. minimum* strains from Ecuador and Malaysia were grown at 24 °C, while the Florida strain was not able to grow at this temperature. In the Malaysia strain the effect was also dependent of light and salinity conditions. These results confirm previous data where a water-soluble component produced by *P. minimum* was able to block sodium channels⁷. The variations of *P. minimum* toxicity were previously linked to the clone studied and to environmental circumstances^{1,8}. Our results also confirm these hypotheses, since depending on the strain and on the growth conditions the effect on I_{Na} is different, as well as the toxin production.

From electrophysiological experiments with *P. minimum* extracts, the presence of some active compound produced by dinoflagellates was evident. To determine its structure, UPLC-MS/MS technology with Electrospray




Figure 7. Accurate MS¹⁻² spectra of compound *m/z* 308 and molecular formula C₁₁H₂₁N₃O₇. (A) MS¹ range *m/z* 150–500. (B) MS² of *m/z* 308 with the proposed structures of fragments 180 and 162. (C) Proposed structure for *m/z* 308.

Ionization (ESI) source was employed to follow the fragmentation pathway of TTX and analogues earlier described^{21,26}. In this case a column for Hydrophilic Interaction Chromatography (HILIC) with polar analytes was used and 5 compounds with apparently the same product ions but different retention times to TTX and analogues were obtained^{22,26,27}. However, these compounds did not follow the characteristic fragmentation pathway of TTXs. TTX and analogues follow the same pattern, first dehydrated ions by losing a molecule of water [M + H - H₂O], and sometimes a second one [M + H - 2H₂O], and then the *m/z* 162 and *m/z* 178 or 176 ion products. These conditions are deemed necessary to affirm that the compounds are TTX analogues^{27,28}. Therefore, *P. minimum* extracts were checked for ion precursors of the common ion product for all TTX analogues, *m/z* 162, and two peaks *m/z* 265 and 308 were obtained. To determine its structure, high-resolution UPLC-IT-TOF technology was performed using TTX as model. Based on the predicted elemental composition and the structure of a parent compound, the structures of product ions can be obtained with a high degree of confidence^{21,29–32}. As mentioned, it is well known that TTX can easily lose a molecule of water (–18 Da), yielding the abundant ion *m/z* 302. Taking into account the structure for the TTX, the loss of water can occur in different carbons but it is considered that the easiest loss is in C⁴. This fragmentation was confirmed with a MS³ spectrum. In this spectrum, the most intense product ions were *m/z* 284.0877, 266.0828, 256.0957, 178.0621 and 162.0683. The product ions *m/z* 284.0877, [C₁₁H₁₃N₃O₆ + H]⁺, and 266.0828, [C₁₁H₁₁N₃O₅ + H]⁺, were formed by elimination of one (–18 Da) and two molecules of water (–36 Da), respectively. The next ion product, *m/z* 256.0957, [C₁₀H₁₃N₃O₅ + H]⁺, was formed by the elimination of CO (–28 Da) from *m/z* 284. Finally, the two last ions, *m/z* 178.0621, [C₈H₇N₃O₂ + H]⁺, and 162.0683, [C₈H₈N₃O + H]⁺, were formed by the elimination of C₂H₆O₃ (–78 Da) and C₂H₆O₄ (–94 Da), respectively, from *m/z* 284. According to the ESI-MSⁿ results for *m/z* 265 and 308, these two compounds had similar fragmentation behaviour. Due to the small amount of compounds present in samples comparing to the TTX amount used for these experiments (2000 ng/mL), only MS² could be performed and the signal intensity obtained in these spectra was low. Compound *m/z* 265, [C₉H₂₀N₄O₅ + H]⁺, loses a molecule of water (–18 Da) giving rise to ion *m/z* 247.1434, [C₉H₁₈N₄O₄ + H]⁺. The most intense product ions in MS² spectra for this compound were *m/z* 191.1192, 179.1199 and 162.0927. The product ion *m/z* 191.1192, [C₉H₁₀N₄O + H]⁺, was formed by elimination of H₁₀O₄ (–74 Da) and *m/z* 179.1199, [C₉H₁₄N₄ + H]⁺, and *m/z* 162.0927, [C₉H₁₁N₃ + H]⁺ by elimination of 86 and 104 Da respectively. Compound *m/z* 308 [C₁₁H₂₁N₃O₇ + H]⁺ loses a molecule of water, *m/z* 290.1317, [C₁₁H₁₉N₃O₆ + H]⁺. Besides, *m/z* 180.0999, [C₉H₁₃N₃O + H]⁺, and *m/z*

162.0842, $[C_9H_{12}N_3 + H]^+$, are also obtained by elimination of $C_2H_8O_6$ and of $C_2H_{10}O_7$ respectively. The ion m/z 162.0662 identified, as product ion of TTX is different from m/z 162.1026 identified as product ion m/z 265 and m/z 308. In the first case it is a C8 molecule while in the second it is a C9. These structures and molecular formula were also proposed as fragment ions of 5,6,11-trideoxyTTX²¹. In addition, a similar molecule $C_9H_9N_3O_2$, molecular weight 191.1, is produced after strong base treatment of TTX¹⁵. When the errors between theoretical and experimental data are compared some differences are observed. In the case of TTX, both precursor and product ions have a narrow mass difference (0.1–5.7 mDa). In the case of compound m/z 265, mass differences for precursor and 3 of 4 product ions are within the range (3.7–9.9 mDa). While in the case of compound m/z 308, due to the low amount, the mass differences are wider (3–26.9 mDa). Therefore the increase in error should be related to the decrease in the amount of compound used. In addition the sample were not purified, and some matrix effect is also present. However, the errors observed do not invalidate these results since the fragmentation pattern of the TTX analogues observed show the loses of water (–18 Da), m/z 162 and m/z 179–180. Therefore, m/z 265 and 308 compounds could be precursors, maybe important molecules for later TTXs synthesis. In this regard, bacteria associated with TTX production, *Roseobacter* genus and *V. alginolyticus* specie, have also previously been identified in *P. minimum* cultures¹¹. So far, the biosynthetic pathway in organisms that produce TTX is not known and some TTX analogues have been proposed as intermediates of TTX in microorganisms²¹. Therefore, compounds m/z 308 and m/z 265 identified in *P. minimum* cultures are TTX analogues produced in or by the dinoflagellate, with the same activity on I_{Na} as TTX. These molecules could be TTX precursors in mussels.

In terms of activity it seems that the effect on Na_v channels is directly related to m/z 265 presence. However, the quantification of both molecules was done with TTX standard and can be not accurate. In addition, as happens with other TTX analogues, the effect on I_{Na} can be different²². In this sense, it has been described that the hydroxyl group at C6 and a CH_2OH group in the same carbon, are necessary for TTX binding to Na_v channels³³. These groups are also present in compounds m/z 308 and m/z 265. It is important to emphasize that the presence and above all the position of CH_2OH group is important for TTX- Na_v channels binding, and the compounds reported in this work have the same group. Thus, although when compared to TTX structure, some groups are missing, these molecules show inhibition of I_{Na} since the C9 base of TTX is maintained. This structure is common to all TTX active analogues²¹. In this sense the C9 base of TTX is often used to determine the total amount of TTX analogues in samples^{15,16}. However, the affinity and toxicity of analogues should be different^{22,34}.

A symbiosis between bacteria and dinoflagellates has been often proposed, however little is known about this phenomenon. Sometimes the bacteria induce death of the algae, while in other situations it provides essential vitamins for microalgae. This is the case for *Roseobacter* sp and *P. minimum* cultures, which the symbiotic association increases the production of B1 and B12 vitamins essential for the dinoflagellate^{35,36}. *Roseobacter* sp have been associated with the production of TTX in the copepod *Pseudocaligus fugu*³⁷. In this case, m/z 320 and m/z 302 ions were identified as TTX and anhydroTTX in *Roseobacter* sp. cultures. In our experiments both ions (m/z 320 and m/z 302) and other fragment ions were also detected, however since they did not follow the TTX characteristic fragmentation pathway they were not identified as TTXs. As it was reported before, TTX and analogues have been detected in many different animals and environments, in addition considerable differences were found in the analogues identified in pufferfish (TTX derivatives) and amphibians (chiquiritoxin)^{21,38}. It has been suggested that this wide range of compounds identified is probably due to the different biosynthesis or metabolism of TTX between animals³⁹. Therefore, our results point to compounds 307 ($C_{11}H_{21}N_3O_7$) and 264 ($C_9H_{20}N_4O_5$) as other TTX analogues produced by *P. minimum* associated with bacteria (*Roseobacter* and *Vibrio* sp). This symbiotic production may explain why the strain is not always toxic. As far as we know, this is the first time that the toxic compounds produce by *P. minimum* have been identified and related to symbiotic bacteria. The structures proposed for the C9 base in these two compounds are shown in Figs 6C and 7C. In the case of m/z 265, the position of the NH_2 group is not defined. For compound m/z 308 two structures could be proposed because the double bond can be located in several positions. To finally clarify this, the structural NMR determination and crystallographic analysis should be performed, although this would require a massive scaling of cultures to obtain enough pure compound. A consequence of TTX being produced by bacteria in algae is that this would potentially be highly influenced by environmental conditions, hence opening a justification as to why climate change provide a link to the recent increase of TTX presence in previously unreported areas.

Methods

Chemicals and Solutions. TTX certified standard was purchased from Laboratorios CIFGA S.A. (Lugo, Spain). 0.5 mL of solution contains $80.7 \pm 6.6 \mu\text{mol}$ TTX/Kg and $9.9 \pm 1.1 \mu\text{mol}$ 4,9-anhTTX/Kg.

Acetonitrile and methanol were obtained from Panreac (Barcelona, Spain). All solvents were HPLC or analytical grade and the water was obtained from a water purification system (Milli-Q, Millipore, Spain). Formic acid was purchased from Merck (Darmstadt, Germany). Ammonium formate was from Fluka (Sigma-Aldrich, Spain). Other reagents were from Sigma (Sigma-Aldrich, Spain).

Plastic tissue cultures dishes were purchased from Falcon (Madrid, Spain). Fetal calf serum, Dulbecco's modified Eagle medium/F12 nutrient mixture (DMED/F12), Glutamax, Minimum essential medium, non essential amino acids (MEM NEAA) and G418 were purchased from Gibco (Glasgow, UK). Detachin™ was purchased from Genlantis (USA).

Mussel samples treatment. Aliquots of 2 g of DG (digestive glands) of sample 1786/2012 collected in Greece were extracted as previously described¹⁷. The extract obtained was used for patch clamp recordings.

For DNA extraction DG of samples 1770/2012 and 1774/2012¹⁷, and one white sample (WT) (not toxic) were separated from mussels and stored at -80°C until processed. At least 5 DG from each sample were mechanically homogenized, and DNA was extracted using the DNA extraction kit NucleoSpin Tissue DNA, RNA and protein purification kit (Macherey Nagel, forensic quality certified free of detectable DNA). The bacterial DNA extraction

protocol was used, following the manufacturer instructions. After purification DNA concentration was determined with a NanoDrop (Thermo Scientific). To study bacteria presence, 16s DNA was amplified, sequenced and analysed (See Supplementary Information).

Automated Patch clamp electrophysiological recordings. For Na current measurements HEK-293 cells stably transfected with the Na channel hNa_v1.6 were used^{20,40–42}. This cell line was kindly provided by Dr Andrew Powell (GlaxoSmithKline R&D, Stevenage, UK). Cells were cultured in DMEM/F12 medium supplemented with 10% of fetal bovine serum, Glutamax and MEM NEAA (1% w/v). G418 was freshly added at a final concentration of 0.4 mg/ml in each cell passage. Cells were incubated in a humidified 5% CO₂/95% air atmosphere at 37 °C until 80% of confluence. Then, cells were incubated at 30 °C for 24–48 h before electrophysiological measurements. Cells were split twice per week.

All the measurements were recorded in whole-cell patch clamp configuration using an IonFlux 16 system (Fluxion, California, USA) and the corresponding Ionflux 16 software for cell capture, seal formation, whole cell obtaining, data acquisition and analysis. After incubation at 30 °C for 24–48 h, cells were washed twice with Ca²⁺ and Mg²⁺ free phosphate buffered saline (PBS) and harvested with 5 ml of Detachin™ solution. Detached cells were resuspended in extracellular solution containing (mM): 2 CaCl₂, 1 MgCl₂, 100 Hepes, 4 KCl, 145 NaCl, 10 TEA-Cl and 10 Glucose. pH 7.4 and 320 mOsm. Electrophysiological recordings were carried out at room temperature (±22 °C) in a 96-well IonFlux microfluidic plate⁴³.

I_{Na} (Na current) was evoked by depolarizing to –10 mV for 50 ms after a 100 ms step to –120 mV from –90 mV holding potential (V_h). The intracellular solution composition for I_{Na} recordings was (in mM): 100 CsF, 45 CsCl, 10 Hepes, 5 NaCl, 5 EGTA corrected to pH 7.1 using CsOH. The contaminating effects of resistance and capacitance currents were compensated electronically by the software. Leak resistance is measured by introducing a short 20 mV pulse at the beginning of each sweep and measuring the current difference⁴³. A sampling frequency of 10 kHz was used.

P. minimum cultures. Three strains of *P. minimum*, from the National Center for Marine Algae and Microbiota, Bigelow (Maine, USA) ccmp1529, ccmp2811 and ccmp2956 were incubated in L1 medium with salinity of 31–33 and 37^{0/00}. The salinity was adjusted to these proportions by the addition of freshwater removing chlorine by aeration. The three strains were incubated at 19 °C with 16:8 h light-dark photoperiod or 24 °C with 14:10 h light-dark photoperiod and different light properties (18 W/765, 18 W/840 and 18 W/865). ccmp2811 only properly grew at 19 °C. The cells were counted using an Utermöhl camera. After 15 days, the cells were harvested by filtration and extracted with methanol. The extracts were always adjusted to 7 × 10⁷ cells/mL. Methanol extracts were vacuum dried and dissolved in acetic acid 0.03 M for further electrophysiological or spectrometry purposes. For bacteria identification, *P. minimum* cultures were harvested after 15 days growth, filtered and frozen at –80 °C until use (see below).

L1 medium properties: To 1 L of sterilized seawater was added: 0.075 g NaNO₃, 0.00565 g, NaH₂PO₄ · 2H₂O, 1.0 mL of trace elements stock solution (1) and 1.0 mL of vitamin mix stock solution (2). (1) was made by adding (per 1 L): FeCl₃ · 6H₂O (3.15 g), Na₂EDTA · 2H₂O (4.36 g), CuSO₄ · 5H₂O (1 × 10^{–8} M), Na₂MoO₄ · 2H₂O (9 × 10^{–8} M), ZnSO₄ · 7H₂O (8 × 10^{–8} M), CoCl₂ · 6H₂O (5 × 10^{–8} M), MnCl₂ · 4H₂O (9 × 10^{–7} M), H₂SeO₃ (1 × 10^{–8} M), Na₃VO₄ (1 × 10^{–8} M), K₂CrO₄ (1 × 10^{–9} M). (2) was made by adding (per 1 L): Cyanocobalamin (0.0005 g), Thiamine HCl (0.1 g), Biotin (0.0005 g).

UPLC Conditions. Chromatographic separation was carried out using both a 1290Infinity ultra-high-performance liquid chromatography system coupled to a 6460 Triple Quadrupole mass spectrometer (Agilent Technologies, Waldbronn, Germany) and from Shimadzu (Kyoto, Japan), two pumps (LC-30AD), autoinjector (SIL-10AC) with refrigerated rack, degasser (DGU-20A), column oven (CTO-10AS) and a system controller (SCL-10AVP) coupled to IT-TOF. The toxins were separated using a column ACQUITY UPLC BEH Amide (2.1 × 100 mm, 1.7 μm, Waters) at 35 °C. Mobile phase A was 100% water with 10 mM formic acid and 10 mM ammonium formate. Mobile phase B was acetonitrile-water (95:5), containing 5 mM formic acid and 2 mM ammonium formate. The gradient program with a flow rate of 0.4 mL/min was started with 100% B and then a linear gradient to 65% B in 7 minutes. After an isocratic hold time linear of 2 minutes at 65% B and return to the starting conditions of 100% B in 0.5 minutes. Finally, 100% B was kept for 3.5 minutes before the next injection. The injection volume was 5 μL.

MS detection. MS detection was performed using an Agilent G6460C Triple Quadrupole mass spectrometer equipped with an Agilent Jet Stream ESI source (Agilent Technologies, Waldbronn, Germany) and an IT-TOF-MS system with an electrospray ionization (ESI) interface (Shimadzu, Kyoto, Japan). The nitrogen generator is a Nitrocraft NCLC/MC from Air Liquid (Spain).

Agilent source conditions were optimized to achieve the best sensitivity for all compounds: drying gas temperature of 250 °C and flow of 5 L/min, nebulizer gas pressure of 55 psi (Nitrocraft NCLC/MS from Air Liquid), sheath gas temperature of 400 °C and flow of 12 L/min. The capillary voltage was set to 3000 V in positive mode with a nozzle voltage of 0 V. The fragmentor was 152 and the cell accelerator voltage was 2 for each toxin in this method.

IT-TOF-MS source conditions were nebulizing gas flow, 1.5 L/min, heat block temperature and CDL temperature, 200 °C and detector voltage, 1.65 kV.

MS/MS analysis. Initial simple analysis was done in MRM mode. The mass spectrometer was operated in positive mode and the collision energy optimized using MassHunter Optimizer software (Supplementary Table 1). Two ions product were analysed per compound, one for quantification and another for confirmation.

Mass spectrum analysis. To confirm and identify TTX analogues the fragmentation pathway of each molecule was used. The mass spectrometer was operated in the Product Ion Scan positive mode for each TTX analogue with scan range from 50 to 350 m/z , scan time 1120 msec and collision energy from 20 to 50 eV. Three characteristic ions were always formed $[M + H - H_2O]^+$, m/z 162 and m/z 178 (or 176) and sometimes also $[M + H - 2H_2O]^+$ ^{21,26,27}. To confirm the presence of TTX analogues, the Precursor Ion Scan of m/z 162 product ion was checked with scan range from 50 to 450 m/z , scan time 40 msec and collision energy from 20 to 50 eV.

IT-TOF-MS analysis. The UPLC from Shimadzu (Kyoto, Japan) mentioned above was connected with an Ion Trap-Time of Flight (IT-TOF) mass spectrometer from Shimadzu. The molecules were analysed using an ion accumulation time of 10 msec and the collision energy for MS^n was adjusted to 50% in the analysis and the isolation width of precursor ions was 3.0Th. For full-scan MS analyses, the spectra were recorded in the range of m/z 150–500. Data-dependent acquisition was set such that the most abundant ions in full-scan MS would trigger tandem mass spectrometry (MS^n , $n = 2-3$). Data were acquired and processed by LC/MS solution software including a formula predictor to calculate elemental compositions.

Formula assignments. Accurate masses of fragment ions were processed using the Lab Solution software supplied with the instrument. Any mass corresponding to the formula predictor also calculated particular compositions. To assign the elemental composition of fragment ions, the error ranges were set less than 1 Da as a limit to the calculation of possible elemental compositions using the formula predictor. The other conditions for calculating elemental compositions that were taken into account were the upper limits on the number of C, H, O, N atoms, C/H ratios and the range of double-bond-equivalent (DBE).

References

- Heil, C. A., Glibert, P. M. & Fan, C. Prorocentrum minimum (Pavillard) Schiller: A review of a harmful algal bloom species of growing worldwide importance. *Harmful Algae* **4**, 449–470 (2005).
- Tango, P. J. *et al.* Impacts and potential effects due to Prorocentrum minimum blooms in Chesapeake Bay. *Harmful Algae* **4**, 525–531 (2005).
- Akiba, T. & Hattori, Y. Food poisoning caused by eating asari and oyster-toxic substance, venerupin. *Jpn. J. Exp. Med.* **20**, 271–284 (1949).
- Nakazima, M. Studies on the source of shellfish poison in Lake Hamana. I Relation of the abundance of a species of dinoflagellate, Prorocentrum sp. to shellfish toxicity. *Bull. Jpn. Soc. Sci. Fish.* **31**, 198–203 (1965).
- Nakazima, M. Studies on the source of shellfish poison in Lake Hamana. II. Shellfish toxicity during the 'red-tide'. *Bull. Jpn. Soc. Sci. Fish.* **31** (1965).
- Silva, E. S. & Sousa, I. Experimental Work on the Dinoflagellate Toxin Production. *Arq. Inst. Nac. Saude* **6**, 381–387 (1981).
- Denardou-Queneherve, A. *et al.* Toxicity of French strains of the dinoflagellate Prorocentrum minimum experimental and natural contaminations of mussels. *Toxicon: official journal of the International Society on Toxinology* **37**, 1711–1719 (1999).
- Grzebyk, D., Denardou, A., Berland, B. & Pouchus, Y. F. Evidence of a new toxin in the red-tide dinoflagellate Prorocentrum minimum. *J. Plankton Res.* **19**, 1111–1124 (1997).
- Noguchi, T. *et al.* Occurrence of tetrodotoxin and anhydrotetrodotoxin in *Vibrio* sp. isolated from the intestine of a xanthid crab, *Atergatis floridus*. *J. Biochem.* **99**, 311–314 (1986).
- Messner, D. J. & Catterall, W. A. The sodium channel from rat brain. Role of the beta 1 and beta 2 subunits in saxitoxin binding. *The Journal of biological chemistry* **261**, 211–215 (1986).
- Jal, S. & Khora, S. S. An overview on the origin and production of tetrodotoxin, a potent neurotoxin. *Journal of applied microbiology* **119**, 907–916, doi: 10.1111/jam.12896 (2015).
- Prathepa, V. & Vasconcelos, V. Microbial diversity associated with tetrodotoxin production in marine organisms. *Environmental toxicology and pharmacology* **36**, 1046–1054, doi: 10.1016/j.etap.2013.08.013 (2013).
- Lago, J., Rodriguez, L. P., Blanco, L., Vieites, J. M. & Cabado, A. G. Tetrodotoxin, an Extremely Potent Marine Neurotoxin: Distribution, Toxicity, Origin and Therapeutical Uses. *Marine drugs* **13**, 6384–6406, doi: 10.3390/md13106384 (2015).
- Kodama, M., Sato, S., Sakamoto, S. & Ogata, T. Occurrence of tetrodotoxin in Alexandrium tamarense, a causative dinoflagellate of paralytic shellfish poisoning. *Toxicon: official journal of the International Society on Toxinology* **34**, 1101–1105 (1996).
- McNabb, P. S. *et al.* First detection of tetrodotoxin in the bivalve Paphies australis by liquid chromatography coupled to triple quadrupole mass spectrometry with and without precolumn reaction. *Journal of AOAC International* **97**, 325–333 (2014).
- Turner, A. D., Powell, A., Schofield, A., Lees, D. N. & Baker-Austin, C. Detection of the pufferfish toxin tetrodotoxin in European bivalves, England, 2013 to 2014. *Euro surveillance: bulletin Europeen sur les maladies transmissibles = European communicable disease bulletin* **20** (2015).
- Vlami, A. *et al.* First Detection of Tetrodotoxin in Greek Shellfish by UPLC-MS/MS Potentially Linked to the Presence of the Dinoflagellate Prorocentrum minimum. *Toxins* **7**, 1779–1807, doi: 10.3390/toxins7051779 (2015).
- Edgar, R. C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**, 2460–2461, doi: 10.1093/bioinformatics/btq461 (2010).
- Alonso, E., Alfonso, A., Vieytes, M. R. & Botana, L. M. Evaluation of toxicity equivalent factors of paralytic shellfish poisoning toxins in seven human sodium channels types by an automated high throughput electrophysiology system. *Archives of toxicology* **90**, 479–488, doi: 10.1007/s00204-014-1444-y (2016).
- Caldwell, J. H., Schaller, K. L., Lasher, R. S., Peles, E. & Levinson, S. R. Sodium channel Na(v)1.6 is localized at nodes of ranvier, dendrites, and synapses. *Proceedings of the National Academy of Sciences of the United States of America* **97**, 5616–5620, doi: 10.1073/pnas.090034797 (2000).
- Yotsu-Yamashita, M. *et al.* First identification of 5,11-dideoxytetrodotoxin in marine animals, and characterization of major fragment ions of tetrodotoxin and its analogs by high resolution ESI-MS/MS. *Marine drugs* **11**, 2799–2813, doi: 10.3390/md11082799 (2013).
- Rodriguez, P. *et al.* First toxicity report of tetrodotoxin and 5,6,11-trideoxyTTX in the trumpet shell Charonia lampas lampas in Europe. *Anal Chem* **80**, 5622–5629 (2008).
- Silva, M. *et al.* New gastropod vectors and tetrodotoxin potential expansion in temperate waters of the Atlantic Ocean. *Marine drugs* **10**, 712–726, doi: 10.3390/md10040712 (2012).
- Cheng, C. A. *et al.* Microflora and tetrodotoxin-producing bacteria in a gastropod, Niotta clathrata. *Food and chemical toxicology: an international journal published for the British Industrial Biological Research Association* **33**, 929–934 (1995).
- Yotsu, M. *et al.* Production of tetrodotoxin and its derivatives by Pseudomonas sp. isolated from the skin of a pufferfish. *Toxicon: official journal of the International Society on Toxinology* **25**, 225–228 (1987).

26. Rodriguez, P. *et al.* Liquid chromatography–mass spectrometry method to detect Tetrodotoxin and Its analogues in the puffer fish *Lagocephalus sceleratus* (Gmelin, 1789) from European waters. *Food Chemistry* **132**, 1103–1111 (2012).
27. Shoji, Y., Yotsu-Yamashita, M., Miyazawa, T. & Yasumoto, T. Electrospray ionization mass spectrometry of tetrodotoxin and its analogs: liquid chromatography/mass spectrometry, tandem mass spectrometry, and liquid chromatography/tandem mass spectrometry. *Analytical biochemistry* **290**, 10–17, doi: 10.1006/abio.2000.4953 (2001).
28. Nakagawa, T., Jang, J. & Yotsu-Yamashita, M. Hydrophilic interaction liquid chromatography-electrospray ionization mass spectrometry of tetrodotoxin and its analogs. *Analytical biochemistry* **352**, 142–144, doi: 10.1016/j.ab.2006.02.010 (2006).
29. Hopfgartner, G., Chernushevich, I. V., Covey, T., Plomley, J. B. & Bonner, R. Exact Mass Measurement of Product Ions for the Structural Elucidation of Drug Metabolites with a Tandem Quadrupole Orthogonal- Acceleration Time-of-Flight Mass Spectrometer. *American Society for Mass Spectrometry* **10**, 1305–1314 (1999).
30. Lin, Z. *et al.* Identification of antioxidants in *Fructus aurantii* and its quality evaluation using a new on-line combination of analytical techniques. *Food Chem* **134**, 1181–1191, doi: 10.1016/j.foodchem.2012.02.161 (2012).
31. Schymanski, E. L., Bataineh, M., Goss, K.-U. & Brack, W. Integrated analytical and computer tools for structure elucidation in effect-directed analysis. *Trends in Analytical Chemistry* **28**, 550–561 (2009).
32. Rockwood, A. L., Kushnira, M. M. & Nelsona, G. J. Dissociation of individual isotopic peaks: predicting isotopic distributions of product ions in MSn. *Journal of the American Society for Mass Spectrometry* **14**, 311–322 (2003).
33. Yotsu-Yamashita, M., Sugimoto, A., Takai, A. & Yasumoto, T. Effects of specific modifications of several hydroxyls of tetrodotoxin on its affinity to rat brain membrane. *The Journal of pharmacology and experimental therapeutics* **289**, 1688–1696 (1999).
34. Jang, J. & Yotsu-Yamashita, M. Distribution of tetrodotoxin, saxitoxin, and their analogs among tissues of the puffer fish *Fugu pardalis*. *Toxicon: official journal of the International Society on Toxinology* **48**, 980–987, doi: 10.1016/j.toxicon.2006.07.034 (2006).
35. Wagner-Dobler, I. *et al.* The complete genome sequence of the algal symbiont *Dinoroseobacter shibae*: a hitchhiker's guide to life in the sea. *The ISME journal* **4**, 61–77, doi: 10.1038/ismej.2009.94 (2010).
36. Wang, H., Tomasch, J., Jarek, M. & Wagner-Döbler, I. A dual-species co-cultivation system to study the interactions between Roseobacters and dinoflagellates. *Front. Microbiol.* **5**, 1–11 (2014).
37. Maran, B. A. *et al.* Isolation and characterization of bacteria from the copepod *Pseudocaligus fugu* ectoparasitic on the panther puffer *Takifugu pardalis* with the emphasis on TTX. *Toxicon: official journal of the International Society on Toxinology* **50**, 779–790, doi: 10.1016/j.toxicon.2007.06.021 (2007).
38. Adachi, M. *et al.* Total synthesis of chiriquitoxin, an analogue of tetrodotoxin isolated from the skin of a dart frog. *Chemistry* **20**, 1247–1251, doi: 10.1002/chem.201304110 (2014).
39. Kudo, Y., Yasumoto, T., Konoki, K., Cho, Y. & Yotsu-Yamashita, M. Isolation and structural determination of the first 8-epi-type tetrodotoxin analogs from the newt, *Cynops ensicauda popei*, and comparison of tetrodotoxin analogs profiles of this newt and the puffer fish, *Fugu poecilonotus*. *Marine drugs* **10**, 655–667, doi: 10.3390/md10030655 (2012).
40. Plummer, N. W. *et al.* Exon organisation, coding sequence, physical mapping and polymorphic intragenic markers for the human neuronal sodium channel gene SCN8A. *Genomics* **54**, 287–296 (1998).
41. Burbidge, S. A. *et al.* Molecular cloning, distribution and functional analysis of the NA(V)1.6. Voltage-gated sodium channel from human brain. *Brain research. Molecular brain research* **103**, 80–90 (2002).
42. Chatelier, A., Zhao, J., Bois, P. & Chahine, M. Biophysical characterisation of the persistent sodium current of the Nav1.6 neuronal sodium channel: a single-channel analysis. *Pflugers Archiv: European journal of physiology* **460**, 77–86, doi: 10.1007/s00424-010-0801-9 (2010).
43. Spencer, C. I. *et al.* Ion channel pharmacology under flow: automation via well-plate microfluidics. *Assay Drug Dev Technol* **10**, 313–324 (2012).

Acknowledgements

Inés Rodríguez is supported by a fellowship from Subprograma de Formación de Personal Investigador MINECO (AGL2012-40185-CO2-01), Spain. The research leading to these results has received funding from the following FEDER cofunded-grants. From CDTI and Technological Funds, supported by Ministerio de Economía y Competitividad, AGL2012-40185-CO2-01, AGL2014-58210-R, and Consellería de Cultura, Educación e Ordenación Universitaria, GRC2013-016. From CDTI under ISIP Programme, Spain, IDI-20130304 APTAFOOD. From the European Union's Seventh Framework Programme managed by REA – Research Executive Agency (FP7/2007-2013) under grant agreement 312184 PHARMASEA.

Author Contributions

A.A. and L.B. conceived, designed the experiments and manuscript elaboration; I.R. UPLC-MS/MS analysis, IT-TOF analysis and formula assignments; E.A. automated patch clamp electrophysiological analysis; A.V. and P.K. mussel samples analysis and J.R., M.R., S.J., M.L.M. and A.D. bacteria analysis. All authors read and approved the final manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Rodríguez, I. *et al.* The association of bacterial C₉-based TTX-like compounds with *Prorocentrum minimum* opens new uncertainties about shellfish seafood safety. *Sci. Rep.* **7**, 40880; doi: 10.1038/srep40880 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017