

ACCEPTED MANUSCRIPT

# Fusion product losses due to fishbone instabilities in deuterium JET plasmas

To cite this article before publication: Vasily Kiptily *et al* 2017 *Nucl. Fusion* in press <https://doi.org/10.1088/1741-4326/aa9340>

## Manuscript version: Accepted Manuscript

Accepted Manuscript is “the version of the article accepted for publication including all changes made as a result of the peer review process, and which may also include the addition to the article by IOP Publishing of a header, an article ID, a cover sheet and/or an ‘Accepted Manuscript’ watermark, but excluding any other editing, typesetting or other changes made by IOP Publishing and/or its licensors”

This Accepted Manuscript is © 2017 Culham Centre for Fusion Energy.

During the embargo period (the 12 month period from the publication of the Version of Record of this article), the Accepted Manuscript is fully protected by copyright and cannot be reused or reposted elsewhere.

As the Version of Record of this article is going to be / has been published on a subscription basis, this Accepted Manuscript is available for reuse under a CC BY-NC-ND 3.0 licence after the 12 month embargo period.

After the embargo period, everyone is permitted to use copy and redistribute this article for non-commercial purposes only, provided that they adhere to all the terms of the licence <https://creativecommons.org/licenses/by-nc-nd/3.0>

Although reasonable endeavours have been taken to obtain all necessary permissions from third parties to include their copyrighted content within this article, their full citation and copyright line may not be present in this Accepted Manuscript version. Before using any content from this article, please refer to the Version of Record on IOPscience once published for full citation and copyright details, as permissions will likely be required. All third party content is fully copyright protected, unless specifically stated otherwise in the figure caption in the Version of Record.

View the [article online](#) for updates and enhancements.

## Fusion product losses due to fishbone instabilities in deuterium JET plasmas

V G Kiptily<sup>1</sup>, M Fitzgerald<sup>1</sup>, V Goloborodko<sup>2</sup>, S E Sharapov<sup>1</sup>, C D Challis<sup>1</sup>, D Frigione<sup>3</sup>, J Graves<sup>4</sup>, M J Mantinen<sup>5,6</sup>, P Beaumont<sup>1</sup>, M Garcia-Munoz<sup>7</sup>, C Perez von Thun<sup>8,9</sup>, J F R Rodriguez<sup>7</sup>, D Darrow<sup>10\*</sup>, D Keeling<sup>1</sup>, D King<sup>1</sup>, K G McClements<sup>1</sup>, E R Solano<sup>11</sup>, S Schmuck<sup>1</sup>, G Sips<sup>12</sup>, G Szepesi<sup>1</sup> and JET contributors\*\*

<sup>1</sup> Culham Centre for Fusion Energy, UKAEA, Culham Science Centre, Abingdon, OX14 3DB, UK

<sup>2</sup> OEAW, Institute for Theoretical Physics, University of Innsbruck, Austria

<sup>3</sup> Unità Tecnica Fusione - ENEA C. R. Frascati - via E. Fermi 45, 00044 Frascati (Roma), Italy

<sup>4</sup> Ecole Polytechnique Fédérale de Lausanne, Swiss Plasma Center, CH-1015 Lausanne, Switzerland

<sup>5</sup> Barcelona Supercomputing Center, Barcelona, Spain

<sup>6</sup> ICREA, Pg. Lluís Companys 23, 08010 Barcelona, Spain

<sup>7</sup> FAMN Department, Faculty of Physics, University of Seville, 41012 Seville, Spain

<sup>8</sup> EUROfusion Programme Management Unit, Culham Science Centre, Culham, OX14 3DB, UK

<sup>9</sup> Forschungszentrum Jülich GmbH, Institut für Energie- und Klimaforschung - Plasmaphysik, 52425 Jülich, Germany

<sup>10</sup> Princeton Plasma Physics Laboratory, Princeton, NJ 08543, New Jersey, USA

<sup>11</sup> Laboratorio Nacional de Fusión, CIEMAT, Madrid, Spain

<sup>12</sup> European Commission, B-1049 Brussels, Belgium

\*Now deceased

\*\*See the author list of "X. Litaudon et al 2017 Nucl. Fusion 57 102001"

E-mail: [vasili.kiptily@ukaea.uk](mailto:vasili.kiptily@ukaea.uk)

During development of a high-performance hybrid scenario for future deuterium-tritium experiments on the Joint European Torus, an increased level of fast ion losses in the MeV energy range was observed during the instability of high-frequency  $n=1$  fishbones. The fishbones are excited during deuterium neutral beam injection combined with ion cyclotron heating. The frequency range of the fishbones, 10 – 25 kHz, indicates that they are driven by a resonant interaction with the NBI-produced D beam ions in the energy range  $\leq 120$  keV. The fast particle losses in a much higher energy range are measured with a fast ion loss detector, and the data show an expulsion of deuterium plasma fusion products, 1 MeV tritons and 3 MeV protons, during the fishbone bursts. An MHD mode analysis with the MISHKA code combined with the nonlinear wave-particle interaction code HAGIS shows that the loss of toroidal symmetry caused by the  $n=1$  fishbones affects strongly the confinement of non-resonant high energy fusion-born tritons and protons by perturbing their orbits and expelling them. This modelling is in a good agreement with the experimental data.

Fusion reactions as well as additional heating of fusion-grade tokamak plasmas generate large numbers of fast ions in the plasma core that can interact with the magneto-hydro-dynamic (MHD) perturbations of the plasma discharge. The MHD instabilities can eject fusion products (FP), in particular alpha-particles, which are the principal source of plasma heating in burning deuterium-tritium (D-T) plasmas. The interactions between MHD perturbations and alpha-particles could be resonant if the alpha-particles are in resonance with high-frequency waves, e.g. Toroidal Alfvén Eigenmodes (TAE) or non-resonant, when MHD perturbations are driven by thermal plasma or fast particles other than the alpha-particles. Both resonant and non-resonant interactions between alpha-particles and MHD

1  
2  
3  
4 perturbations could lead to the particle redistribution and losses [1] thus affecting the fusion plasma  
5  $Q=P_{out}/P_{in}$  and the first wall.

6  
7 The development of high fusion performance scenarios, both “baseline” sawtooth H-mode  
8 with safety factor at the magnetic axis  $q_0 \leq 1$  and the so-called “hybrid” with  $q_0 \geq 1$ , for the forthcoming  
9 high-power D-T experiments was a main priority of recent deuterium and hydrogen experimental  
0 campaigns on the Joint European Torus (JET) with ITER-like wall (ILW). In the hybrid scenario with  
1 a low magnetic shear in the plasma centre [2], the sawtooth oscillations are almost always avoided but  
2 a strong beam ion pressure drives fishbones (FB) with toroidal mode number  $n=1$ , an instability with  
3 bursting amplitude and sweeping frequency in the range  $\approx 10 - 25$  kHz.

4  
5 The fishbones occur in plasmas with high- $\beta_{poloidal}$  (ratio of plasma pressure to poloidal magnetic  
6 field pressure) and they are destabilized by energetic ions produced from perpendicular and tangential  
7 neutral beam injection [3, 4], which in the case of JET experiments had primary energies 90 keV and  
8 110 keV. As it has been shown [5, 6], trapped and circulating beam ions can resonate with fishbone  
9 perturbation, the core-localised  $m=1/n=1$  mode. In JET and other tokamaks with fusion reactivity  
0 dominated by beam-plasma fusion reactions, neutron rate drops up to 10% were observed and linked to  
1 the resonant interaction between the beam ions and fishbones [7, 8].

2  
3 In addition to the well-understood resonant interaction described in [5-8], the presence of  
4 fishbones also affects significantly the confinement of non-resonant ions in the MeV energy range.  
5 These ions are p, t and  $^3\text{He}$  produced in the deuterium plasma due to the following fusion reactions:  
6  $D+D = p$  (3MeV) + t (1MeV),  $D+D = n$  (2.5MeV) +  $^3\text{He}$  (0.82MeV). During the fishbone periods, a  
7 “burn-up” of tritons and  $^3\text{He}$  was reduced. The reduction of the burn-up was measured via 14.1-MeV  
8 neutrons produced in the  $D(t, n)^4\text{He}$  fusion reaction and 14.7-MeV protons generated in another fusion  
9 reaction  $D(^3\text{He}, p)^4\text{He}$ . The anomalously low burn-up rate during fishbones has been explained by  
0 prompt and non-prompt losses of MeV-particles [9 – 11]. However, a part of the non-prompt losses,  
1 e.g. slow tritons, could be due to the resonant mechanism of interaction with fishbones (the cross-section  
2 has a maximum at  $E_t \approx 160$  keV), as it was observed with the resonant beam particles. An early study  
3 of fusion product losses via the  $^3\text{He}$  burn-up measurements (the cross-section has a maximum at  $E_{^3\text{He}} \approx$   
4 650 keV) was made on tokamak PDX [10]. In spite of the small size of the machine (a low rate of the  
5  $D(^3\text{He}, p)^4\text{He}$  reaction), it was possible to identify the non-resonant character of the  $^3\text{He}$  losses, and draw  
6 a conclusion on the possible applicability of the loss mechanism to ignition experiments.

7  
8 A non-resonant loss mechanism due to high amplitude fishbones causing a distortion of the  
9 toroidal symmetry with the  $m=1/n=1$  perturbation, was predicted for non-resonant alphas in burning  
0 plasmas in [12]. Experimentally, such non-resonant losses of fast ions in the MeV energy range caused  
1 by the fishbones driven by the beam deuterons with energy 80 - 100 keV were observed in JET  
2 discharges with ion cyclotron range of frequencies (ICRF) heating [13]. The high energy particles  
3 ejected from the plasma during the FB oscillations were identified as H-ions accelerated during ICRH.  
4  
5  
6  
7  
8  
9  
0

The losses were enhanced by about a factor of  $\approx 10 - 20$  with respect to the MHD-quiescent levels, and the magnitude of the losses was found to increase quadratically with the FB amplitude.

In this Letter, we report on directly measured non-resonant losses of fusion protons and tritons in high performance experiments with hybrid plasmas. The fishbones were observed to trigger sawtooth-like reconnection events and neoclassical tearing modes (NTM), the latter substantially reducing the plasma performance.

JET is equipped well for the studies of confined and escaped fast ions [14]. The lost ion measurements are carried out with a scintillator probe [15], which is called “fast ion loss detector”, or FILD. FILD is located about 28 cm below the mid-plane of the torus, just outside the plasma and provides information on the lost ion pitch angle,  $\theta = \cos^{-1}(v_{\parallel}/v)$ , with 5% resolution in the range  $35^{\circ} - 85^{\circ}$  and gyro-radius between 3cm and 14cm with 15% resolution (for 12-cm gyro-radius  $\sim 2$  cm). The light emitted by the scintillator (decay time  $\sim 0.5\mu\text{s}$ ) during an ion collision, is transferred through a coherent fibre bundle to a charge-coupled device (CCD) camera and a photomultiplier tube (PMT) 4x4 array. The 128x256 pixel CCD camera can provide 20-kHz snapshots of light emission on the pitch-angle – gyro-radius grid calculated with the EfiDesign code [16]. The FILD has recently been upgraded, so the fast PMT signals are digitised by 2MHz thus allowing a loss spectrogram to be made with the MHz bandwidth. Such loss spectrograms could be then compared to magnetic spectrograms showing the MHD perturbations.

In the hybrid scenario experiments, with plasma currents in the range  $I_p = 2.0 - 2.4$  MA at central toroidal field  $B_T(0) = 2.8\text{T}$ , deuterium neutral beam injection (NBI) and H-minority heating of deuterium plasmas with ICRF at  $f = \omega/(2\pi) \approx 42.5$  MHz of dipole phasing with hydrogen concentration  $n_H/(n_H+n_D) \approx 1-3\%$  were used. Fishbones were observed in discharges with normalized poloidal beta  $\beta_N > 1.9$ , however not all of them led to the FP losses. A typical example of a discharge with fast ion losses during period of strong FB instabilities is shown in figure 1. The MHD instability in this discharge has been identified as  $n=1$  fishbones by means of a comparison of the signal phase in toroidally-separated Mirnov coils. The result of the mode analysis is presented on figure 2.

An example of losses measured with the FILD in discharge #92394 is shown in the gyro-radius ( $R_{\text{gyr}}$ ) vs the pitch-angle grid in figure 3a. This is a typical fusion product first orbit (FO) loss footprint recorded with CCD camera in the FB-free period 7.843s – 7.860s just before a fishbone. It is clearly seen that the maxima of losses are localised along the gyro-radius (red dash line), corresponding to the FP birth energies, 1-MeV for tritons and 3-MeV for protons. Prompt ICRH losses, H-ions and D-NBI ions accelerated at  $\omega = \omega_{cH} = 2\omega_{cD}$ , would be expected to occur along the red solid line, corresponding to the IC resonance layer in the plasma, but such losses appear to be rather low. The image of losses recorded during the fishbone period 7.860s – 7.877s is presented in figure 3b. This image was obtained by a subtraction of the signal recorded during the previous no fishbone time bin shown in figure 3a. It is important to note that the loss ion image obtained by the subtraction of the signal recorded after the

FB period, 7.877s – 7.893s, is indistinguishable from the one shown in figure 3b. One can see that the losses related to the FB instability are localised along the red dot line, which is a separatrix between trapped and circulating orbits in the phase space. This indicates that the fishbone perturbation causes losses by pushing core-localised fusion products with confined passing orbits to a space phase area corresponding to the unconfined trapped orbits.

The FILD data analysis shows that in the FB period the fusion tritons and proton prompt losses are localised at the pitch-angle  $\theta \sim 55^\circ$  and the gyro-radius  $R_G \sim 11$  cm that is related to a maximal signal. The gyro-radius distribution associated with the energy distribution has roughly the same shape as distributions before and after this period (although the total losses are higher); for tritons, the maximum of the losses is at  $E_t \sim 1$  MeV, for protons at  $E_p \sim 3$  MeV. However, the pitch-angle distribution of the prompt losses during the fishbone is slightly shifted relative to the FB-free distributions, from  $\theta \sim 57^\circ$  to  $\sim 55^\circ$ . Assuming the losses are of the particles with trapped orbits, it is important to note that the major radius at the bounce reflection point for these particles and the pitch-angle value on the scintillator plate are related by  $R(\theta) = R_{\text{FILD}}[1 - \cos^2(\theta)]$ , where  $R_{\text{FILD}}$  is radial position of the scintillator. As can be seen from the differential loss footprint in figure 3b, particles escaped during the FB period are mostly coming from the plasma region near  $55^\circ$  related to  $R(55^\circ) = 2.45$  m. A back-in-time FP orbit calculation starting from the FILD scintillator plate (figure 4a) shows that bounce points of these particles are close to the trapped/passing boundary in the mid-plane at  $\sim 2.45$  m. However, the orbits of H-ions accelerated by ICRH, calculated with pitch-angle  $\theta \sim 63^\circ$  are trapped and their turning points are on the IC resonance layer at  $R \approx 2.97$  m (figure 4b). These calculations give us a proof that ions escaped from the plasma during the fishbone period are mostly fusion products.

An  $n=1$  ideal internal kink mode was found with the MISHKA-1 code [17] for a JET equilibrium reconstruction obtained with EFIT [18]. The HAGIS [19] code was used to compute the fusion product drift orbits assuming the EFIT equilibrium, and the drift orbit changes were computed assuming the MISHKA-1 solution (full details of these calculations will be the subject of a following publication). The largest losses are predicted for particles near the boundary between trapped and counter passing lost orbits. The results are shown in figure 5, where  $Z_{R=R_{\text{mag}}}$  denotes the Z particle location in the upper midplane where it crosses  $R=R_{\text{mag}}$ . This is used in place of the canonical momentum  $P_\phi$ . The three constants of motion used in the plots are Z (canonical momentum),  $\Lambda$  (magnetic moment) and energy E. The plots are at fixed E corresponding to the birth energy ignoring thermal spread. One can see that the lack of particles at the parameter  $\Lambda=1$  ( $\Lambda \equiv \mu_m B(R_{\text{mag}})/E = B(R_{\text{mag}})[1 - \cos^2\theta]/B$ ) related to on-axis trapped orbits of H- and D-beam ions accelerated by ICRH is consistent with the footprint in Figure 3b. Figure 6 gives an illustration of the orbit types, where the greatest prompt fusion product losses are predicted to occur. The counter travelling fusion product orbits near the trapped/passing/lost triple-point cross the internal kink mode. So, the fishbone is producing a radial convection of these non-resonant fusion products into loss regions.









## Figures




Fig.1






Fig.4




Fig.5




Fig.6






Fig.7




Fig.8




Fig.9