

Serveur Académique Lausannois SERVAL serval.unil.ch

Publisher's version PDF

Faculty of Biology and Medicine Publication

Title: Bone Formation and the Wnt Signaling Pathway.

Authors: Gori F, Superti-Furga A, Baron R

Journal: The New England journal of medicine

Year: 2016 Nov 10

Volume: 375

Issue: 19

Pages: 1902-1903

DOI: 10.1056/NEJMc1609768

“From [Publication Title, Author(s), Title of Article, Volume No., Page No. Copyright © (notice year) Massachusetts Medical Society. Reprinted with permission.

Stéphane Gaudry, M.D.
David Hajage, M.D.
Didier Dreyfuss, M.D.

Assistance Publique–Hôpitaux de Paris
Paris, France
didier.dreyfuss@aphp.fr

Since publication of their article, the authors report no further potential conflict of interest.

1. Zarbock A, Kellum JA, Schmidt C, et al. Effect of early vs delayed initiation of renal replacement therapy on mortality in

critically ill patients with acute kidney injury: the ELAIN randomized clinical trial. *JAMA* 2016;315:2190-9.

2. Wald R, Adhikari NKJ, Smith OM, et al. Comparison of standard and accelerated initiation of renal replacement therapy in acute kidney injury. *Kidney Int* 2015;88:897-904.

3. Ridgeon EE, Young PJ, Bellomo R, et al. The fragility index in multicenter randomized controlled critical care trials. *Crit Care Med* 2016;44:1278-84.

4. Pocock SJ, Stone GW. The primary outcome is positive — is that good enough? *N Engl J Med* 2016;375:971-79.

DOI: 10.1056/NEJMc1610500

Bone Formation and the Wnt Signaling Pathway

TO THE EDITOR: Simsek Kiper et al. (June 30 issue)¹ link Pyle's disease to a deficiency of secreted frizzled-related protein 4 (sFRP4), a soluble Wnt inhibitor, and demonstrate its specific effects on cortical bone (which is unusually thin in persons with Pyle's disease) versus trabecular bone (which is increased in persons with Pyle's disease). In sFrp4-deficient mice, the inhibition of cortical bone formation, which determines bone strength at large, was shown to be mediated through bone morphogenetic protein (BMP)-dependent induction of sclerostin, a key Wnt inhibitor in bone. Given the small amount of trabecular bone in the mouse femur, are bone marrow-derived osteoblasts that are obtained from the femur indeed representative of trabecular bone? Second, did the authors assay levels of dickkopf 1 (Dkk1)? Dkk1 also inhibits Wnt and — like sclerostin — is a downstream target of BMP signaling. Therefore, Dkk1, in addition to sclerostin, may contribute to the suppression of bone formation.² Moreover, blocking sclerostin increases the expression of Dkk1 in bone, which partially limits the bone-forming effects of sclerostin inhibition.³ Thus, Wnt signaling has multiple redundant safeguards to fine-tune bone formation. Therapeutically, this complexity may require the use of bispecific antibodies against more than one Wnt inhibitor to fully restore Wnt signaling, bone formation, and ultimately bone strength.

Martina Rauner, Ph.D.
Tilman D. Rachner, M.D.
Lorenz C. Hofbauer, M.D.

TU Dresden Medical Center
Dresden, Germany
lorenz.hofbauer@uniklinikum-dresden.de

No potential conflict of interest relevant to this letter was reported.

1. Simsek Kiper PO, Saito H, Gori F, et al. Cortical-bone fragility — insights from sFRP4 deficiency in Pyle's disease. *N Engl J Med* 2016;374:2553-62.

2. Kamiya N, Kobayashi T, Mochida Y, et al. Wnt inhibitors Dkk1 and Sost are downstream targets of BMP signaling through the type IA receptor (BMPRIA) in osteoblasts. *J Bone Miner Res* 2010;25:200-10.

3. Florio M, Gunasekaran K, Stolina M, et al. A bispecific antibody targeting sclerostin and DKK-1 promotes bone mass accrual and fracture repair. *Nat Commun* 2016;7:11505.

DOI: 10.1056/NEJMc1609768

THE AUTHORS REPLY: We isolated bone marrow-derived osteoblasts from the femora and tibiae of mice that were 10 weeks old, an age at which both the femur and the tibia contain trabecular bone. It is well established that enhancement of canonical Wnt signaling is associated with increases in bone mass.¹ Our studies clearly show that sFrp4 deficiency has a marked anabolic effect on trabecular bone in humans and mice. Given that trabecular bone resides in the marrow and that the properties of sFrp4-deficient bone marrow-derived osteoblasts (e.g., the Wnt canonical pathway — but not the noncanonical pathway — is activated in these cells) are consistent with the changes observed in trabecular bone (increased bone mass), we concluded that these cells better reflect the cellular activities in the trabecular bone compartment than do calvarial cortical osteoblasts. With regard to the second point, sFrp4 deficiency was associated with increased *Dkk1* expression in both bone marrow-derived and calvarial osteoblasts (Fig. 1). This is not surprising; *Dkk1* is a well-known canonical




Figure 1. Effect of *Sfrp4* Deletion on *Dkk1* Messenger RNA (mRNA) Expression.

Dkk1 mRNA expression in calvarial osteoblasts and bone marrow–derived osteoblasts of wild-type and *Sfrp4*^{-/-} mice after 0, 14, and 28 days of osteogenic differentiation is shown. Fold differences in mRNA expression relative to that in wild-type cells were calculated as described by Livak and Schmittgen.² The bars show mean values; T bars indicate standard deviations. Three independent experiments were performed at each time point.

on the maintenance of cortical bone homeostasis than do antibodies to only sclerostin.

Francesca Gori, Ph.D.

Harvard School of Dental Medicine
Boston, MA

Andrea Superti-Furga, M.D.

Lausanne University
Lausanne, Switzerland

Roland Baron, D.D.S., Ph.D.

Harvard School of Dental Medicine
Boston, MA
roland_baron@hms.harvard.edu

Since publication of their article, the authors report no further potential conflict of interest.

1. Baron R, Kneissel M. WNT signaling in bone homeostasis and disease: from human mutations to treatments. *Nat Med* 2013; 19:179-92.

2. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 2001;25:402-8.

3. Florio M, Gunasekaran K, Stolina M, et al. A bispecific antibody targeting sclerostin and DKK-1 promotes bone mass accrual and fracture repair. *Nat Commun* 2016;7:11505.

DOI: 10.1056/NEJMc1609768

Wnt signaling target gene. We agree with Rauner and colleagues: bispecific antibodies against sclerostin and *Dkk1*³ could have a stronger effect

Fire-Related Inhalation Injury

TO THE EDITOR: In his proposed algorithm for the early management of fire-related inhalation injury, Sheridan (Aug. 4 issue)¹ excluded bronchoscopy from the recommendations listed as standards of care because of the lack of “high-level proof” of its efficacy. That stance raises concern because there are reports that favor the use of bronchoscopy in this context. The degree of inhalation injury identified on bronchoscopy has been reported as a determinant of an inflammatory response to smoke inhalation,² and bronchoscopic findings have been positively correlated with mortality.³ Our experience also supports

the use of bronchoscopy in this situation. In providing medical care for the victims of a nightclub fire in Brazil, we found that serial, flexible bronchoscopy provided us with information that was crucial to the decision-making process with regard to treatment, allowing us to identify the extent of burns in the airways and to remove large amounts of debris (Fig. 1). No complications were observed with the use of this procedure. Among 18 patients who underwent bronchoscopy at our hospital, 16 are still receiving follow-up care and have had good clinical outcomes. Thus, the limited data available on bronchoscopy can