



CCAFS site atlas

Nyando / Katuk Odeyo Kenya

CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS)

Site Atlas

Correct citation:

Sijmons K., Kiplimo J., Förch W., Thornton P.K., Radeny, M. and Kinyangi, J. (2013). CCAFS Site Atlas – Nyando / Katuk Odeyo. CCAFS site atlas series. The CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS). Copenhagen, Denmark. Available online at: www.ccafs.cgiar.org

Titles in this series aim to disseminate interim climate change, agriculture and food security research and practices and stimulate feedback from the scientific community.

This document is published by the CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS) which is a strategic partnership between CGIAR and Future Earth. CCAFS is supported by the CGIAR Fund, the Danish International Development Agency (DANIDA), the Australian Government Overseas Aid Program (AusAid), Irish Aid, Environment Canada, the Ministry of Foreign Affairs for the Netherlands, the Swiss Agency for Development and Cooperation (SDC), Instituto de Investigação Científica Tropical (IICT), UK Aid, the Government of Russia, and the European Union (EU). The Program is carried out with technical support from the International Fund for Agricultural Development (IFAD)

Contact:

CCAFS Coordinating Unit - Faculty of Science, Department of Plant and Environmental Sciences, University of Copenhagen, Rolighedsvej 21, DK-1958 Frederiksberg C, Denmark. Tel: +45 35331046; Email: ccaafs@cgiar.org

Creative Commons License



This Atlas is licensed under a Creative Commons Attribution – NonCommercial–NoDerivs 3.0 Unported License.

Articles appearing in this publication may be freely quoted and reproduced provided the source is acknowledged. No use of this publication may be made for resale or other commercial purposes.

©2013 CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS).

DISCLAIMER:

This Atlas Series has been prepared as an output for Theme 4.2 under the CCAFS program and has not been peer reviewed. Any opinions stated herein are those of the author(s) and do not necessarily reflect the policies or opinions of CCAFS, donor agencies, or partners. The geographic designation employed and the presentation of material in this publication do not imply the expression of any opinion whatsoever on the part of CCAFS concerning the legal status of any country, territory, city or area or its authorities, or concerning the delimitation of its frontiers or boundaries.

All images remain the sole property of their source and may not be used for any purpose without written permission of the source.

Contents

Chapter 1: Introduction	iv
Regional Map.....	1
Topographic Map	2
Satellite Map	3
Chapter 2: Climate and Climate Variability	
Annual Rainfall	4
Annual Temperature	5
Aridity Index	6
Chapter 3: Bio-Physical Characteristics	
Altitude.....	7
Landforms.....	8
Soil Type	9
Agro-Ecological Zones	10
Landcover	11
Landuse	12
Length of Growing Period 2000	13
Length of Growing Period 2030	14

Crop Suitability	15
Livestock Production Systems	16
Livestock Density	17
Chapter 4: Socio-Economic Factors	
Livelihood Zones	18
Human Population Density	19
Market Access	20
Poverty	21
Conservations Areas	22
References and Data Sources	23

Introduction

The CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS) seeks to promote a food-secure world through the provision of science-based efforts that support sustainable agriculture and enhance livelihoods while adapting to climate change and conserving natural resources and environmental services.

Climate change is an unprecedented threat to the food security of hundreds of millions of people who depend on small-scale agriculture for their livelihoods. Climate change affects agriculture and food security, and likewise, agriculture and natural resource management affect the climate system.

CCAFS has initially focused on three regions; East Africa (EA), West Africa (WA) and South Asia (SA) to carry out its research. The 15 CCAFS sites in these areas represent areas that are becoming both drier and wetter, and are focal locations that will generate results that can be applied and adapted to other regions worldwide. In this year, 2013, CCAFS is expanding its portfolio to additional sites in Latin America and South-East Asia.

These sites serve as the initial focus of CCAFS partnership-building and long-term research activities falling within the following CCAFS Research Themes; Adaptation to Progressive Climate Change, Adaptation through Managing Climate Risk, Pro-Poor Climate Change Mitigation and Integration for Decision Making. At all 15 CCAFS sites, baseline surveys have been conducted, including three levels of data collection and analysis at household, village and organizational levels (see: <http://ccaafs.cgiar.org/resources/baseline-surveys>).

More information on CCAFS work in all the three regions can be accessed at www.ccaafs.cgiar.org

To better understand the CCAFS sites' characteristics, a list of geospatial indicators for climate variability, bio-physical characteristics and socio-economic variables have been mapped into site atlases.

This Atlas was developed for the CCAFS site at Nyando / Katuk Odeyo in Kenya, in East Africa Region.

CCAFS Sites: East Africa



- Ethiopia: Borana (ET01)
- Kenya: Nyando (KE01)
- Kenya: Makueni (KE02)
- Uganda: Albertine Rift (UG01)
- Uganda: Kagera Basin (UG02)
- Tanzania: Usambara (TZ01)

CCAFS Country Sites

Citation: GeoMapa (2013a)


Topography Nyando

CCAFS Site KE01, Nyando / Katuk Odeyo, Kenya




Coordinates of the CCAFS Baseline Sampling frame

35.068E 0.269S
 35.068E 0.361S
 34.978E 0.361S
 34.978E 0.269S








Sampling frame size: 10km x 10km



Satellite Image Katuk Odeyo






RapidEye imagery from 05-02-2011
 at 5m ground resolution

HBS= Household Baseline Survey
 VBS= Village Baseline Survey
 OBS= Organizational Baseline Survey


-  Road
-  River
-  Settlement
-  CCAFS VBS/OBS village
-  CCAFS HBS villages



Annual Rainfall




Katuk Odeyo Mean Monthly Rainfall Distribution






Citation: Jones et al (2002)

Citation: Hijmans et al (2005)


Annual Rainfall data of current interpolations of observed data, representative of 1950 - 2000





Annual Temperature



Katuk Odeyo Mean Monthly Temperature Distribution




Citation: Jones et al (2002)



Annual Temperature represents annual temperature data of current interpolations of observed data, averaged for 1950 - 2000

Citation: Hijmans et al (2005)

Aridity Index



— International boundary
 □ Corresponds to the map on the left



- Aridity Index**
- Hyper Arid
 - Arid
 - Semi Arid
 - Dry sub-humid
 - Humid

Aridity Index indicates the level of dryness, taking evapotranspiration into account, at a given location of known rainfall



Katuk Odeyo CCAFS sampling frame

Citation: Trabucco et al (2009)

Altitude




Altitude indicates the height above sea level in meters






Citation: Jarvis et al (2008)

Landforms



- Town
- Settlement
- Road
- River
- Wetland





Corresponds to the map on the left

- Landforms ***
- | | |
|----------------------------------|--|
| ■ Badlands | ■ Piedmont plains |
| ■ Bottomlands | ■ Plateaus and high-level structural plains |
| ■ Floodplains | ■ Swamps |
| ■ Footslopes | ■ Upland/high-level plain transitional lands |
| ■ Hills and minor scarps | ■ Upper middle-level uplands |
| ■ Lacustrine plains | ■ Upper-level uplands |
| ■ Lower middle-level uplands | ■ Volcanic footridges |
| ■ Minor Valleys | ■ Undifferentiated or various rocks |
| ■ Mountains and major scarps | ■ Urban area |
| ■ Non-dissected erosional plains | |
- * Legend corresponds to left map

Landforms comprise the geomorphological units that make up the Earth's surface, largely defined by its surface form and location in the landscape

Citation: FAO Africover (2002)

Soil Type



International boundary
 Scale 1:12,500,000
 0 125 250 500 Kilometers
 Corresponds to the map on the left


Soil Type *

Acrisols	Lixisols
Alisols	Luvisols
Andosols	Nitisols
Arenosols	Phaeozems
Cambisols	Planosols
Ferralsols	Regosols
Fluvisols	Vertisols
Gleysols	Greyzems


* Legend corresponds to left map

Soil Type refers to the soil group as per the FAO classification. Soil groups are defined by their parent material and morphogenetic characteristics in terms of structural properties and texture (sand, silt and clay content), as well as organic matter content.

Scale 1:750,000
 0 7.5 15 30 45 Kilometers
 1 cm = 7.5 km




Agro-Ecological Zones






- Agro-Ecological Zones ***
- High Altitude Derived Savanna
 - Mid Altitude Derived Savanna
 - High Altitude Humid Forest
 - Mid Altitude Humid Forest
- * Legend corresponds to left map

Agro-Ecological Zones indicate the division of land areas that have similar characteristics related to land suitability, potential agricultural production and environmental impact.






Landcover



- Landcover**
- Rainfed croplands
 - Mosaic Croplands/Vegetation
 - Mosaic Vegetation/Croplands
 - Closed broadleaved deciduous forest
 - Open broadleaved deciduous forest
 - Open needleleaved deciduous or evergreen forest
 - Mosaic Forest-Shrubland/Grassland
 - Urban area
 - Closed to open shrubland
 - Closed to open broadleaved evergreen or semi-deciduous forest

Landcover shows the observed (bio)physical cover of the earth's surface, i.e. dominant vegetation, land use and man-made features.





— International boundary

Scale 1:12,500,000

0 125 250 500 Kilometers

□ Corresponds to the map on the left

- Landuse ***
- Forest protected
 - Forest with agricultural activities
 - Forest with moderate or higher livestock density
 - Shrubs unmanaged
 - Shrubs protected
 - Shrubs moderate livestock density
 - Shrubs high livestock density
 - Rainfed crops (Subsistence/Commercial)
 - Crops and moderate intensive livestock density
 - Crops and high livestock density
 - Agriculture protected
 - Urban area

* Legend corresponds to left map
 Citation: Natchtergaele et al (2010)

Landuse is a description of how people utilize the land. It involves socio-economic activity, i.e. the management and modification of the natural environment into built environment, such as agricultural fields and settlements. At any place, there may be multiple land uses, the dominant one is presented here.

Length of Growing Period 2000




— International boundary

Scale 1:12,500,000

0 125 250 500 Kilometers

□ Corresponds to the map on the left

Length of Growing Period (Days)




The Length of Growing Period (LGP) is defined as the number of days in a year during which there is available rainfed soil moisture supply for plant growth.

Length of Growing Period 2030





Scale 1:12,500,000
0 125 250 500 Kilometers



Corresponds to the map on the left



The Length of Growing Period (LGP) is defined as the number of days in a year during which there is available rainfed soil moisture supply for plant growth; here modeled for 2030




Crop Suitability



Crop Suitability refers to the land resource assessment that considers agricultural land use options with relevant agro-ecological condition to estimate expected cropping activities.

Livestock Production Systems



International boundary

Corresponds to the map on the left

- | | |
|------------------------|--------------------------|
| Mixed Rainfed | Livestock only |
| Arid / Semi-arid | Arid / semi-arid |
| Humid / sub-humid | Temperate / highland |
| Temperate / highland | Closed to open shrubland |
| Mixed Irrigated | Urban area |
| Arid / semi-arid | Other |
| Humid / sub-humid | |
| Temperate / highland | |


Livestock Production Systems as part of agricultural systems take agro-climatic conditions into account and are classified in terms of feed and livestock resources; livestock commodities produced; production technology; product use and livestock functions; area covered; geographic locations; and human populations supported.

Livestock Density





Scale 1:12,500,000
0 125 250 500 Kilometers

Corresponds to the map on the left





Livestock Density is measured in numbers of livestock, including cattle, goats and sheep, per km²



Citation: Wint et al (2007)

Livelihood Zones



Scale 1:12,500,000
 0 125 250 500 Kilometers

International boundary

Corresponds to the map on the left

Livelihood Zones *


- Lake Victoria Fishing Zone
- Western Lakeshore Marginal Mixed Farming Zone
- Western High Potential Zone

* Legend corresponds to left map


Livelihoods are complex and shaped by a variety of factors. These livelihood zone maps delineate geographic areas within which people broadly share the same livelihood patterns including access to food, income, and markets.

Scale 1:750,000
 0 7.5 15 30 45 Kilometers
 1 cm = 7.5 km

Town (circle with dot)
 Settlement (circle)
 Road (red line)
 River (blue line)
 Wetland (dotted pattern)



Human Population Density




Scale 1:12,500,000

0 125 250 500 Kilometers


International boundary

Corresponds to the map on the left

Number of persons per km²




Human Population Density is the gridded number of persons per km² in 2005.




Katuk Odeyo CCAFS sampling frame

Citation: CIESIN (2005)

Market Access



Travel time to nearest large town/city (Hours)




Travel time is a measure of accessibility determined in the time (hours) taken to the nearest urban centre, town or city of a population of 50,000 people or more (taking different means of transportation into account)

Poverty





Percentage of People living on less than 2 US\$ per day




CIESIN constructed global data sets of poverty that are based on estimates of subnational infant mortality and child malnutrition data, recognizing that both are proxies for poverty and welfare rather than direct measures.

Citation: CIESIN (2005)

Conservation Areas



Conservation Areas represent protected areas that, according to IUCN, are clearly defined geographic spaces, recognized, dedicated and managed through legal or other effective means, to achieve long-term conservation of nature with associated ecosystem services and cultural value.



Katuk Odeyo
CCAFS sampling frame

Citation: UNEP and WCMC (2012)

References and Data Sources

Regional Map

Sijmons K. 2013a. Digital Satellite Image based on, MODIS (Moderate Resolution Imaging Spectroradiometer) NASA, 2009, Ground resolution : 1 Kilometer. GTOPO30, (DEM) Global Digital Elevation Model U.S Geological Survey, Ground resolution: 1 Kilometer. Topographic Features derived from: Global GIS, U.S. Geological Survey and Google Earth. Projection: Geographic, Lat/Long, WGS84

Topographic Map

Sijmons K. 2013b. Relief representation derived from Digital Elevation Model (DEM) of SRTM (Shuttle Radar Topographic Mission) 2000, Ground resolution 90 meter and ASTER GDEM, Ground resolution 30 meter, NASA. Topographic Features digitized from Google Earth Projection: Geographic, Lat/Long, WGS84

Satellite Image

RapidEye Satellite Image, 5 meter ground resolution, Image acquisition, 17-01-2011

Annual Rainfall

Hijmans, R.J., S.E. Cameron, J.L. Parra, P.G. Jones and A. Jarvis, 2005. Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* 25: 1965-1978.

Annual Rainfall Graph

Jones P G, Thornton P K, Diaz W and Wilkens P W. 2002. MarkSim, a computer tool that generates simulated weather data for crop modeling and risk assessment. Version 1, 2002. CD-ROM and Users Manual. CIAT, AA6713, Cali, Colombia, 87 pp.

Annual Temperature

Hijmans, R.J., S.E. Cameron, J.L. Parra, P.G. Jones and A. Jarvis, 2005. Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* 25: 1965-1978.

Annual Temperature Graph

Jones P G, Thornton P K, Diaz W and Wilkens P W. 2002. MarkSim, a computer tool that generates simulated weather data for crop modeling and risk

assessment. Version 1, 2002. CD-ROM and User's Manual. CIAT, AA6713, Cali, Colombia, 87 pp.

Aridity Index

Trabucco, A., and Zomer, R.J. 2009. Global Aridity Index (Global-Aridity) and Global Potential Evapo-Transpiration (Global-PET) Geospatial Database. CGIAR Consortium for Spatial Information. Published online, available from the CGIAR-CSI GeoPortal at: <http://www.csi.cgiar.org/>

Altitude

Jarvis, A., H.I. Reuter, A. Nelson, E. Guevara, 2008, Hole-filled SRTM for the globe Version 4, available from the CGIAR-CSI SRTM 90m Database. Available at <http://srtm.csi.cgiar.org>

Landforms

FAO Africover, 2002. Available at <http://www.africover.org/system/metadata.php?metadataid=65>

Soil Type

FAO/IIASA/ISRIC/ISS-CAS/JRC. 2009. Harmonized World Soil Database (version 1.1). FAO, Rome, Italy and IIASA, Laxenburg, Austria.

Agro-Ecological Zones

Fischer, G., F. Nachtergaele, S. Prieler, H.T. van Velthuisen, L. Verelst, D. Wiberg, 2008. Global Agro-ecological Zones Assessment for Agriculture (GAEZ 2008). IIASA, Laxenburg, Austria and FAO, Rome, Italy

Landcover

Arino, O., Perez J. R., Kalgirou V., Defourny P., Achard F. 2009. GlobCover. Version 2.3 Global. Accessed at <http://ionia1.esrin.esa.int/>

Landuse

Nachtergaele F., Petri M., 2010. Mapping Land Use at global and Regional Scales for Land Degradation Assessment Analysis (LADA).Version 1.1. LADA Project FAO/UNEP GEF/. <http://www.fao.org/geonetwork/srv/en/metadata.show?id=37139&currTab=simple>

Length of Growing Period 2000

Thornton P K, Jones P G, Owiyo T, Kruska R L, Herrero M, Kristjanson P, Notenbaert A, Bekele N and Omolo A, with contributions from Orindi V, Adwerah A, Otiende B, Bhadwal S, Anantram K, Nair S and Kumar V. 2006. Mapping climate vulnerability and poverty in Africa. Report to the Department for International Development, International Livestock Research Institute, Nairobi, Kenya, 200 p.

Length of Growing Period 2030

Thornton P K, Jones P G, Owiyo T, Kruska R L, Herrero M, Kristjanson P, Notenbaert A, Bekele N and Omolo and Kumar V. 2006. Mapping climate vulnerability and poverty in Africa. Report to the Department for International Development, International Livestock Research Institute, Nairobi, Kenya, 200 p.

Crop Suitability

FAO-IIASA 2007. Mapping biophysical factors that influence agricultural production and rural vulnerability. Food and Agriculture Organization and International Institute for Applied Systems Analysis, Rome 2007.

Livestock Production Systems

FAO. 2007. Gridded livestock of the world 2007, by G.R.W. Wint and T.P. Robinson. Rome, pp 131.

Livestock Density

Wint W, Robinson T, 2007. Gridded Livestock of the World. FAO, 131 pp.

Livelihood Zones

Livelihood Zone Products. Famine Early Warning Systems Network (FEWS NET). Accessed at <http://www.fews.net/pages/livelihoods.aspx?loc=6&l=en>.

Human Population Density

Center for International Earth Science Information Network (CIESIN)/Columbia University, and Centro Internacional de Agricultura Tropical (CIAT, 2005. Gridded Population of the World, Version 3 (GPWv3): Population Density Grid, Future Estimates, Edition: 3.0. NASA Socioeconomic Data and Applications Center (SEDAC. Online_Linkage: <http://sedac.ciesin.columbia.edu/data/set/gpw-v3-population-density-future-estimates>

Market Access

Nelson, A. (2008) Estimated travel time to the nearest city of 50,000 or more people in year 2000. Global Environment Monitoring Unit - Joint Research Centre of the European Commission, Ispra Italy. Available at <http://www.tem.jrc.it/accessibility> (accessed 06/03/2004)

Poverty

Center for International Earth Science Information Network (CIESIN), Columbia University, 2005. Small Area Estimates of Poverty and Inequality (SAEPI) database. Palisades, NY: CIESIN,

University. Available at <http://www.ciesin.columbia.edu/povmap>. (Version 1.0) Available at <http://sedac.ciesin.columbia.edu/theme/poverty>

Conservation Areas

UNEP-WCMC (2012). Data Standards for the World Database on Protected Areas. UNEP-WCMC: Cambridge, UK.

The CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS) brings together the world's best researchers in agricultural science, development research, climate science and Earth System science, to identify and address the most important interactions, synergies and tradeoffs between climate change, agriculture and food security. CCAFS is a strategic partnership of CGIAR and Future Earth, led by the International Center for Tropical Agriculture (CIAT).

For more information, visit www.ccafs.cgiar.org and www.geomapa.nl



RESEARCH PROGRAM ON
**Climate Change,
Agriculture and
Food Security**

