



Tropentag, September 19-21, 2012, Göttingen -  
Kassel/Witzenhausen

“Resilience of agricultural systems against crises”

## Performance of Farmland Terraces in Maintaining Crop Productivity in Wello, Northern Highlands of Ethiopia

SHIMELES DAMENE SHIENE<sup>1</sup>, PAUL L. G. VLEK<sup>1</sup>, LULSEGED TAMENE DESTA<sup>2</sup>

<sup>1</sup>University of Bonn, Center for Development Research (ZEF), Germany

<sup>2</sup>International Center for Tropical Agriculture (CIAT), Trop. Soil Biology and Fertility (TSBF), Malawi

### Abstract

Soil-erosion-induced land degradation and crop production loss is a great challenge in the Ethiopian highlands. Consequently, the government has invested in soil and water conservation (SWC) measures where farmland terracing is among the widely implemented practices. The purpose of this study was to analyse the performance of terracing with respect to crop productivity. Yield differences across the slope of the terrain, within a terrace and across terrace age were evaluated. The study was conducted in the Maybar soil conservation research site (MSCRS) in Wello, northern Ethiopia. It is located between 10°58' and 11°02' N latitude and 39°38' and 39°40' E longitude. Yield data (grain and biomass) of seven crops such as wheat (*Triticum* spp), emmer wheat (*Triticum* spp), horse bean (*Vicia faba*), field pea (*Pisum sativum*), barley (*Hordeum* spp), maize (*Zea mays*) and teff (*Eragrostis tef*) collected between 1995 and 2009 from 40 fixed plots on three terrace positions (upper, middle and lower) were statistically analysed using a mixed linear model that employed Tukey-Kramer adjustment in SAS (Version 9.2). Generally, yields of all crops except wheat decreased significantly (on average 0.37 to 0.15 t ha<sup>-1</sup>) from the upper terrace position towards the lower position. Unlike in other studies, the differences could be due to the soil depth gradient that in turn influenced nutrient and soil-water storage. On the other hand, yields showed insignificant differences across the terrain, they showed a decreasing tendency with slope increase. This indicates that terracing reduced soil erosion and nutrient translocation. With terrace age, yields showed only slight changes. The stable yield under limited fertility improvement measures and reduced fallowing indicates that terracing reduced soil and nutrient loss through erosion. However, terracing alone does not improve soil fertility and thereby crop production. Thus, in order to significantly increase yields terracing should be supplemented by soil fertility amendments.

**Keywords:** Crop yield, farmland terrace, Maybar, northern Ethiopia, soil erosion