

# ‘NASPOT 12 O’ and ‘NASPOT 13 O’ Sweetpotato

**Robert O.M. Mwanga<sup>1</sup> and Gerald Kyalo**

*International Potato Center (CIP), Naguru Hill, Ntinda II Road, Box 22274, Kampala, Uganda*

**Gorrettie N. Ssemakula, Charles Niringiye, Benard Yada,  
Milton A. Otema, Joweria Namakula, Agnes Alajo, Benjamin Kigozi,  
and Rose N.M. Makumbi**

*National Agricultural Research Organization (NARO), National Crops Resources Research Institute (NaCRRI), Namulonge, P.O. Box 7084, Kampala, Uganda*

**Anna-Marie Ball**

*HarvestPlus, International Food Policy Research Institute, P.O. Box 28565, Kampala, Uganda*

**Wolfgang J. Grüneberg**

*International Potato Center, Apartado 1558, Lima, Peru*

**Jan W. Low**

*International Potato Center, Box 25171, Nairobi, Kenya*

**G. Craig Yencho**

*Department of Horticultural Science, North Carolina State University, Rm. 214A Kilgore Hall, Box 7609, Raleigh, NC 27695-7609*

*Additional index words.* *Ipomoea batatas*, sweetpotato virus disease, *alternaria bataticola* blight, provitamin A

Two sweetpotato [*Ipomoea batatas* L. (Lam.)] cultivars, NASPOT 12 O (Namulonge sweetpotato 12 orange-fleshed) and NASPOT 13 O (Namulonge sweetpotato 13 orange-fleshed) were approved for release by the Ugandan Plant Variety Release Committee (UPVRC) in Nov. 2013 (Ssemakula et al., 2013). This brings to 22, the number of sweetpotato cultivars officially released between 1999 and 2013 by the National Sweetpotato Program in Uganda (Mwanga et al., 2011; Ssemakula et al., 2013). The two cultivars herein described and released in 2013, have high average storage root yields, 43.1 t·ha<sup>-1</sup> (‘NASPOT 12 O’) and 27.8 t·ha<sup>-1</sup> (‘NASPOT 13 O’) on station, and 14.9 t·ha<sup>-1</sup> (‘NASPOT 12 O’) and 9.7 t·ha<sup>-1</sup> (‘NASPOT 13 O’) on farm compared with the national on-farm average for Uganda of 4.5 t·ha<sup>-1</sup> (FAOSTAT 2010; Low et al., 2009). The cultivars have acceptable root shape, high dry matter content (DMC) (>30%) with

good-to-excellent consumer qualities. The cultivars also have moderate levels of field resistance to sweetpotato virus disease (SPVD) and *alternaria bataticola* blight. The two cultivars were bred targeting development of vitamin A-rich (biofortified) orange-fleshed sweetpotato (OFSP). OFSP have been shown to be both effective for increasing maternal and child vitamin A intake and status (Hotz et al., 2012; Ruel and Alderman, 2013). The potential of the two OFSP cultivars to contribute to food and nutrition security in Uganda and the developing world is high (Low et al., 2007; Ruel, 2001), especially where high dry matter and starchy sweetpotatoes are preferred. The cultivars can be used directly if adapted in similar agroecologies in sub-Saharan Africa and globally and/or used as parents in breeding programs to develop locally adapted cultivars that meet high dry matter consumer preferences.

## Origin

Before release, ‘NASPOT 12 O’ and ‘NASPOT 13 O’ were code named SPK004/2006/1136 and NASPOT7/2006/292, respectively. SPK004 (released in 2004 as Kakamega) was the female parent, 2006 was the initial year of clonal selection, and 1136 was the genotype number. This clone was officially released as ‘NASPOT 12 O’ [NASPOT = Namulonge sweetpotato,

12 = serial number according to the UPVRC and the National Agricultural Research Organization (NARO) nomenclature, O = orange fleshed]. Similarly, ‘NASPOT 13 O’ is a progeny of the released cultivar, with ‘NASPOT 7’ first selected in 2006 as genotype number 292, and given the UPVRC/NARO serial number 12, and officially released as ‘NASPOT 13 O’. Kakamega (SPK004) was the female parent of ‘NASPOT 7’; therefore, ‘NASPOT 7’, ‘NASPOT 12 O’, and ‘NASPOT 13 O’ are genetically related.

The polycross block from which the two cultivars originated was established in 2005/2006 at Namulonge with 24 parents (Table 1). The 24 parents in the polycross block consisted of three released and three common Ugandan landrace cultivars, two bred released Ugandan cultivars, three Ugandan breeding lines, and introductions from Kenya (2), Peru (8), and Rwanda (3). ‘NASPOT 12 O’ and ‘NASPOT 13 O’ are progenies of ‘Kakamega’ as the female parent, but because seed was open pollinated, their male pedigrees are unknown. The parents were included in the polycross nursery for improvement or as sources of one or a combination of genes for combining desirable traits such as orange-fleshed roots (provitamin A), high dry matter (≥30%), resistance to SPVD and *alternaria bataticola* stem blight, and early maturity (3 to 4 months).

## Description and Performance

The key standard morphological descriptors [International Potato Center (CIP), Asian Vegetable Research and Development Center, and International Board for Plant Genetic Resources, 1991] of the two released cultivars are listed in Table 2 and major differences are shown in Fig. 1. Both cultivars have semierect vines and vigorous growth. However, ‘NASPOT 12 O’ has dense foliage that suppresses aggressive weeds, and a balanced harvest index that makes it good for dual purpose use as animal feed and food (Claessens, 2009). The flower color in both cultivars is the same, pale purple limb with purple throat. Flowering and seed capsule set are sparse in both cultivars. Both cultivars have high storage root DMC (≥30%) and a dry texture with a sweet taste when cooked (Table 3). Storage root skin color is purple red in ‘NASPOT 12 O’ and cream in ‘NASPOT 13 O’. The storage root flesh color in both cultivars is deep orange but the intensity varies with age of the roots, location, and agro-climatic factors such as soil type, and wet or dry season.

Ssemakula et al. (2013) presented the data for official release of the two cultivars in Uganda. Details of the release information include descriptions of pedigree, cultivar, test sites, materials and methods, planting materials, on-station and on-farm trials, planting and harvesting dates, pest and disease evaluation procedures, farmer selection, acceptability evaluation, experimental designs, stability analysis, determination of dry matter

Received for publication 1 Oct. 2015. Accepted for publication 31 Dec. 2015.

This work was supported, in part, by a grant from USAID through HarvestPlus and from the McKnight Foundation.

<sup>1</sup>Corresponding author. E-mail: r.mwanga@cgiar.org.

This is an open access article distributed under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1. Origin and main attributes of 24 sweetpotato parents in the polycross nursery at Namulonge in 2005/2006 that gave rise to lines from which 'NASPOT 12 O' and 'NASPOT 13 O' were selected.<sup>z</sup>

| Parent                         | Origin of parent | Yr released/status | Desirable/undesirable trait                          |
|--------------------------------|------------------|--------------------|--|
| New Kawogo (CIP 441745)        | Uganda/landrace  | 1995               | HDM, high resistance to SPVD                         |
| Kyabafuruki                    | Uganda/landrace  | Germplasm          | High root yield, susceptible to SPVD                 |
| Otada                          | Uganda/landrace  | Germplasm          | HDM, HA  |
| 91/282-1-no. 1                 | Rwanda           | Breeding line      | High resistance to SPVD                              |
| 316/3                          | Uganda           | Breeding line      | OF, HDM, high resistance to SPVD                     |
| 91/282-1-no. 5                 | Rwanda           | Breeding line      | High resistance to SPVD                              |
| Beauregard (CIP 440132)        | CIP/Peru         | Germplasm          | OF, susceptible to SPVD                              |
| 316/2                          | Uganda           | Breeding line      | OF, High resistance to SPVD                          |
| Tainung 65 (CIP 440215)        | CIP/Peru         | Germplasm          | OF, low dry matter                                   |
| Tanzania (CIP 440166)          | Uganda/landrace  | 1995               | HDM, HA  |
| Excel (CIP 440016)             | CIP/Peru         | Germplasm          | OF, susceptible to SPVD                              |
| Karoti Dar es Salaam           | CIP/Nairobi      | Germplasm          | OF, susceptible to SPVD                              |
| Zapallo (CIP 420027)           | CIP/Peru         | Germplasm          | OF, LDM  |
| Tainung 64 (CIP 440159)        | CIP/Peru         | Germplasm          | OF, LDM, susceptible to SPVD                         |
| NASPOT5/58                     | Uganda           | Breeding line      | OF, HDM, high resistance to SPVD                     |
| Huarmeyano (CIP 420020)        | CIP/Peru         | Germplasm          | High resistance to SPFMV                             |
| NASPOT 1 (CIP 191133.1)        | Uganda/bred      | 1999               | HDM, HA, susceptible to alternaria bataticola blight |
| SPK004 (Kakamega) (CIP 441768) | Kenya            | 2004               | OF, HDM  |
| CC89.14.74 × OP (CIP 199004.3) | CIP/Peru         | Breeding line      | OF, susceptible to SPVD                              |
| Siliki                         | Uganda/landrace  | Germplasm          | HDM, HA, susceptible to SPVD                         |
| Ejumula (CIP 443750)           | Uganda/landrace  | 2004               | OF, HA, susceptible to SPVD                          |
| NASPOT 7 (CIP 100200.1)        | Uganda/bred      | 2007               | OF, HDM, high resistance to SPVD                     |
| 91/16-5                        | Rwanda           | Breeding line      | OF, susceptible to SPVD                              |
| Jewel (CIP 401562)             | CIP/Peru         | Germplasm          | OF, susceptible to SPVD                              |

<sup>z</sup>CIP = International Potato Center; HA = high acceptability; HDM = high dry matter; LDM = low dry matter; SPVD = sweetpotato virus disease; OF = orange fleshed; SPFMV = *Sweet potato feathery mottle virus*.

Table 2. Morphological descriptors of 'NASPOT 12 O' and 'NASPOT 13 O' released in Uganda in Nov. 2013.<sup>z</sup>

| Descriptor             | Cultivar                       |                             |                                  |
|------------------------|--------------------------------|-----------------------------|----------------------------------|
|                        | NASPOT 12 O                    | NASPOT 13 O                 | Dimbuka-Bukulula (local control) |
| Plant type             | Semierect                      | Semierect                   | Spreading                        |
|                        |                                | <i>Vine pigmentation</i>    |                                  |
| Predominant color      | Green                          | Green with few purple spots | Green                            |
| Secondary color        | Purple nodes                   | Absent                      | Absent                           |
|                        |                                | <i>Mature leaf shape</i>    |                                  |
| General leaf outline   | Triangular                     | Lobed                       | Triangular                       |
| Lobe type              | Very slight (teeth)            | Deep                        | No lateral lobes                 |
| Leaf lobe number       | Five                           | Three                       | One                              |
| Shape of central lobe  | Triangular                     | Lanceolate                  | Triangular                       |
|                        |                                | <i>Foliage color</i>        |                                  |
| Mature leaf            | Green                          | Green                       | Green                            |
| Immature leaf          | Mostly purple                  | Green                       | Green                            |
| Petiole pigmentation   | Green with purple at both ends | Green                       | Green                            |
|                        |                                | <i>Storage root</i>         |                                  |
| Storage root shape     | Round elliptic                 | Elliptic                    | Long irregular or curved         |
| Surface defects        | Longitudinal grooves           | Longitudinal grooves        | Shallow longitudinal grooves     |
| Predominant skin color | Purple red                     | Cream                       | Cream                            |
| Flesh color            | Dark orange                    | Dark orange                 | Cream                            |
|                        |                                | <i>Flowering</i>            |                                  |
| Flowering habit        | Sparse                         | Sparse                      | Sparse                           |
| Stigma exertion        | Inserted                       | Exerted                     | Inserted                         |
|                        |                                | <i>Capsule</i>              |                                  |
| Seed capsule set       | Sparse                         | Sparse                      | Sparse                           |

<sup>z</sup>Selected descriptors according to International Potato Center, Asian Vegetable Research and Development Center, and International Board for Plant Genetic Resources Descriptors (1991).

and beta-carotene, corresponding results, and cultivar maintenance. The following description is a summary of the cultivar release results. The released cultivars were tested for six seasons (Uganda has two rainy seasons per year when sweetpotatoes can be grown) on station and on farm during 2011 to 2013 in replicated, standard multilocation yield trials in 1) the warm, subhumid short grasslands where sweetpotato weevils and drought are important and frequent; 2) the warm, moist, tall grasslands where SPVD pressure is high; and 3) the cool, moist, southwestern highlands where alternaria bataticola blight and

low soil fertility problems are widespread. A total of eight multilocation on-station trials were conducted for four seasons, followed by 100 on-farm trial evaluation for two seasons under rain-fed conditions. The cultivars were routinely evaluated for resistance to SPVD, alternaria bataticola blight, and sweetpotato weevils, *Cylas puncticollis* (Boheman) and *Cylas brunneus* (Fabricius) (Table 3). Classification of the relative resistance to disease and weevil damage was based on field evaluation under natural disease pressure and weevil populations in each agroecology. Storage root

DMC, root yield, taste, and desirable agronomic attributes (such as earliness, root size, and shape) were also evaluated (Table 3). Data were subjected to analysis of variance using SAS statistical package (SAS V9.1; SAS Institute, Cary, NC). Treatment means were separated where appropriate, using Fisher's least significant difference test.

Results of the performance of the cultivars on station and on farm including palatability are presented in Tables 4 and 5. Both 'NASPOT 12 O' and 'NASPOT 13 O' have higher storage root and biomass yield, higher



Fig. 1. Morphological characters of 'NASPOT 12 O' (A) shoot and flower, (B) shoot tip, leaf and root, and 'NASPOT 13 O' (C) shoot and flower, and (D) shoot tip, leaf, and storage root.

harvest index, and higher SPVD field resistance than the control cultivar, Dimbuka-Bukulula (Table 4). For most traits, the two cultivars performed as well as or better than the local checks across districts (Table 5). The two cultivars have higher beta-carotene in storage roots than Dimbuka-Bukulula (Table 2). These two cultivars have moderate beta-carotene content compared with cultivars such as Resisto with high beta-carotene values (17–25 mg/100 g, on a fresh weight basis) (Grüneberg et al., 2015; Tumwegamire et al., 2014). Both cultivars have moderate field resistance to SPVD and alternaria blight, but are susceptible to weevils, though 'NASPOT 12 O' tends to be less susceptible because of the thick canopy that creates a humid microclimate that is not conducive to weevil colonization. Both cultivars are highly susceptible to the pests in no-choice tests under laboratory conditions. However, both cultivars are potentially valuable as sources of beta-carotene in a high DMC background, and they are already used as parents in the crossing block at Namulonge in Uganda. These cultivars are expected to

Table 3. Main agronomic disease and insect pest reaction and quality traits of two orange-fleshed sweetpotato cultivars released in Uganda in Nov. 2013.

| Attribute   | Cultivar         |                    |                                  |
|---|------------------|--------------------|----------------------------------|
|   | NASPOT 12 O      | NASPOT 13 O        | Dimbuka-Bukulula (local control) |
| Dry matter % (range)  | 31.7 (27.5–35.3) | 32.0 (26.7–34.1)   | 32.7 (29.7–35.1)                 |
| Cooked texture  | Somewhat dry     | Somewhat dry       | Somewhat dry                     |
| Sweetness   | Moderate         | Moderate           | Moderate                         |
| Field reaction to weevils <sup>z</sup>  | S                | S                  | S                                |
| Field reaction to SPVD <sup>yz</sup>  | MR               | MR                 | S                                |
| Field reaction to alternaria stem blight <sup>z</sup>   | MR               | MR                 | MR                               |
| Maturity (days)   | 105 (90–120)     | 120 (115–140)      | 128 (120–150)                    |
| Mean and (range <sup>x</sup> ) of storage root yields in various yield trials (t·ha <sup>-1</sup> ) | 26.5 (10.0–60.2) | 17.2 (4.5–56.8)    | 17.3 (2.9–41.9)                  |
| Mean storage root yield (% of local control)  | 145              | 94                 | 100                              |
| Beta-carotene content (mg/100 g on a fresh weight basis) <sup>w</sup>                               | 7.23 (5.71–8.81) | 11.03 (8.90–12.98) | 0.02 (0.013–0.024)               |

<sup>z</sup>Susceptible (S) = considerable damage or numbers present to severe damage or very high numbers present, respectively; moderately resistant (MR) = moderate damage or moderate numbers present (resistant = little or no apparent damage or few or no insects present).

<sup>y</sup>SPVD = sweetpotato virus disease.

<sup>x</sup>The wide variation in yield is attributed to variation in environmental factors such as erratic rain during some seasons, and differences in farm management and soil types in the different agroecologies.

<sup>w</sup>Sources: Ssemakula et al. (2013) and Tumwegamire et al. (2014).

Table 4. Performance of 'NASPOT 12 O' and 'NASPOT 13 O' in advanced yield trial at four on-station sites, Namulonge, Kachwekano, Ngetta, and Serere, for two seasons in 2011–12.<sup>z</sup>

| Clone             | Yield (t·ha <sup>-1</sup> ) <sup>y</sup> |      |         | Harvest index | Disease severity <sup>x</sup> |            | Dry matter content (%) |
|-------------------|--|------|---------|---------------|-------------------------------|------------|------------------------|
|                   | Total root                               | Vine | Biomass |               | SPVD                          | Alternaria |                        |
| NASPOT 1          | 33.2                                     | 33.8 | 67.0    | 0.50          | 4.0                           | 3.3        | 33.5                   |
| NASPOT7/2006/1185 | 18.9                                     | 38.8 | 57.7    | 0.33          | 2.8                           | 2.0        | 33.3                   |
| NASPOT 12 O       | 46.1                                     | 31.3 | 77.4    | 0.60          | 3.1                           | 2.6        | 30.3                   |
| Jewel(OP)/2005/6  | 17.3                                     | 41.0 | 58.3    | 0.30          | 3.0                           | 2.6        | 29.9                   |
| NASPOT 8          | 39.8                                     | 26.5 | 66.3    | 0.60          | 2.8                           | 2.3        | 34.4                   |
| New Kawogo        | 7.9                                      | 45.3 | 53.2    | 0.15          | 2.4                           | 2.7        | 31.2                   |
| NASPOT 13 O       | 27.8                                     | 41.7 | 69.5    | 0.40          | 3.1                           | 2.3        | 32.7                   |
| Dimbuka-Bukulula  | 18.3                                     | 39.6 | 57.9    | 0.32          | 3.8                           | 2.5        | 34.4                   |
| Mean              | 26.2                                     | 37.3 | 63.4    | 0.40          | 3.1                           | 2.5        | 32.5                   |
| LSD(0.05)         | 6.6                                      | 7.9  | 11.9    | 0.06          | 0.5                           | 0.5        | 3.6                    |
| CV (%)            | 35.1                                     | 25.9 | 24.3    | 18.90         | 21.3                          | 25.0       | 6.5                    |

<sup>z</sup>Sites: Namulonge is in the warm, moist, tall grasslands (high sweetpotato virus disease pressure agroecology); Kachwekano is in the cool, moist, southwestern highlands (with high alternaria blight pressure); and Ngetta and Serere are in the warm, subhumid short grasslands (with high weevil populations during dry periods).

<sup>y</sup>Mean of four replications. Plantings in each season was in a randomized complete block design (RCBD); 80 plants on five ridges (1 m by 0.3 m) per plot; only the 48 middle plants were harvested for yield determination.

<sup>x</sup>Sweetpotato virus disease (SPVD) symptoms/alternaria blight severity: scored 4 to 6 weeks after planting, and 1 month before harvest, on a scale from 1 to 9, where 1 = no virus/alternaria symptoms, 2 = unclear virus/alternaria symptoms, 3 = clear virus/alternaria symptoms at one plant per plot, 4 = clear virus/alternaria symptoms at two to three plants per plot, 5 = clear virus/alternaria symptoms at 5% to 10% of plants, 6 = clear virus/alternaria symptoms at 10–25% of plants, 7 = clear virus/alternaria symptoms at 25% to 50% of plants, 8 = clear virus/alternaria symptoms at nearly all plants per plot, 9 = clear virus/alternaria symptoms and clearly reduced growth in all plants.

LSD = least significant difference.

Table 5. Performance of 'NASPOT 12 O', 'NASPOT 13 O', and local checks (LC) during two seasons in on-farm sweetpotato trials in various districts of Uganda during 2012.

| District <sup>z</sup> /yr | Cultivar            | Yield <sup>y</sup> (t·ha <sup>-1</sup> ) |      |         | Disease severity <sup>x</sup> |            | Taste test rank <sup>w</sup> |
|---------------------------|---------------------|--|------|---------|-------------------------------|------------|------------------------------|
|                           |                     | Marketable                               | Root | Biomass | SPVD                          | Alternaria |                              |
| Isingiro 2012A            |                     |  |      |         |                               |            | (n = 21; m = 10, f = 11)     |
|                           | NASPOT 10 O         | 8.9                                      | 10.9 | 23.5    | 2.7                           | 1.2        | 2                            |
|                           | NASPOT 7/2006/1185  | 6.5                                      | 8.2  | 39.3    | 3.3                           | 1.2        | 6                            |
|                           | NASPOT 12 O         | 16.3                                     | 18.7 | 36.8    | 2.9                           | 1.7        | 1                            |
|                           | NASPOT 13 O         | 9.9                                      | 11.5 | 27.3    | 2.7                           | 1.6        | 3                            |
|                           | Kyebandira (LC)     | 9.2                                      | 11.3 | 28.7    | 3.4                           | 1.8        | 4                            |
|                           | Mean                | 10.2                                     | 12.7 | 31.1    | 3.0                           | 1.5        | NA                           |
|                           | LSD <sub>0.05</sub> | 3.8                                      | 4.0  | 9.1     | NS                            | 0.4        | NA                           |
| cv (%)                    | 38.9                | 34.5                                     | 30.8 | 28.3    | 28.2                          | NA         |                              |
| Buyende 2012A             |                     |  |      |         |                               |            | (n = 42; m = 16, f = 26)     |
|                           | NASPOT 10 O         | 11.5                                     | 12.6 | 28.2    | 2.3                           | 1.0        | 2                            |
|                           | NASPOT 7/2006/1185  | 4.1                                      | 5.6  | 38.9    | 3.0                           | 1.0        | 6                            |
|                           | NASPOT 12 O         | 17.5                                     | 19.5 | 40.2    | 2.6                           | 1.0        | 3                            |
|                           | NASPOT 13 O         | 8.1                                      | 9.2  | 28.2    | 2.1                           | 1.1        | 4                            |
|                           | Muwulu aduduma (LC) | 8.1                                      | 9.5  | 29.9    | 3.0                           | 1.0        | 1                            |
|                           | Mean                | 9.9                                      | 11.3 | 33.1    | 2.6                           | 1.0        | NA                           |
|                           | LSD <sub>0.05</sub> | 3.9                                      | 3.9  | 8.3     | 0.7                           | NS         | NA                           |
| cv (%)                    | 42.7                | 38.3                                     | 25.9 | 26.7    | 15.4                          | NA         |                              |
| Rakai 2012A               |                     |  |      |         |                               |            | (n = 49; m = 35, w = 14)     |
|                           | NASPOT 10 O         | 6.0                                      | 8.4  | 20.5    | 3.1                           | 1.0        | 2                            |
|                           | NASPOT 7/2006/1185  | 5.9                                      | 8.4  | 30.7    | 3.3                           | 1.0        | 6                            |
|                           | NASPOT 12 O         | 12.3                                     | 14.6 | 30.6    | 3.6                           | 1.0        | 3                            |
|                           | NASPOT 13 O         | 8.4                                      | 10.8 | 29.1    | 3.0                           | 1.0        | 1                            |
|                           | Mwolanfuzi (LC)     | 7.1                                      | 9.2  | 22.6    | 3.6                           | 1.0        | 4                            |
|                           | Mean                | 7.9                                      | 10.3 | 26.7    | 3.3                           | 1.0        | NA                           |
|                           | LSD <sub>0.05</sub> | 1.9                                      | 3.5  | NS      | NS                            | NS         | NA                           |
| cv (%)                    | 21.7                | 30.9                                     | 30.3 | 21.1    | NA                            | NA         |                              |
| Oyam 2012A                |                     |  |      |         |                               |            | (n = 89; m = 56, f = 33)     |
|                           | NASPOT 10 O         | 12.3                                     | 13.2 | 17.5    | 2.0                           | 1.0        | 2                            |
|                           | NASPOT 7/2006/1185  | 5.6                                      | 6.7  | 15.1    | 2.1                           | 1.0        | 5                            |
|                           | NASPOT 12 O         | 12.3                                     | 14.3 | 19.7    | 2.5                           | 1.5        | 1                            |
|                           | NASPOT 13 O         | 9.3                                      | 10.3 | 18.8    | 2.4                           | 1.4        | 3                            |
|                           | Liralira (LC)       | 8.9                                      | 10.5 | 32.2    | 3.0                           | 1.0        | 4                            |
|                           | Mean                | 9.7                                      | 11.0 | 20.6    | 2.4                           | 1.2        | NA                           |
|                           | LSD <sub>0.05</sub> | NS                                       | 4.4  | 8.3     | NS                            | 0.4        | NA                           |
| cv (%)                    | 41.4                | 35.9                                     | 36.3 | 26.7    | 27.6                          | NA         |                              |
| Kabale 2012A              |                     |  |      |         |                               |            | (n = 20; m = 5, f = 15)      |
|                           | NASPOT 10 O         | 6.0                                      | 8.1  | 22.1    | 1.6                           | 1.0        | 3                            |
|                           | NASPOT 7/2006/1185  | 1.4                                      | 2.7  | 32.7    | 2.1                           | 1.0        | 5                            |
|                           | NASPOT 12 O         | 10.6                                     | 13.8 | 34.5    | 2.0                           | 1.1        | 4                            |
|                           | NASPOT 13 O         | 8.8                                      | 11.4 | 30.0    | 2.1                           | 1.0        | 2                            |
|                           | Murungi (LC)        | 9.3                                      | 11.3 | 28.7    | 3.4                           | 1.7        | 1                            |
|                           | Mean                | 7.3                                      | 9.5  | 29.6    | 2.2                           | 1.2        | NA                           |
|                           | LSD <sub>0.05</sub> | 2.0                                      | 2.3  | NS      | 0.6                           | 0.3        | NA                           |
| cv (%)                    | 29.5                | 25.7                                     | 29.6 | 28.9    | 29.5                          | NA         |                              |
| Isingiro 2012B            |                     |  |      |         |                               |            | (n = 55; m = 14, f = 41)     |
|                           | NASPOT 10 O (LC)    | 6.7                                      | 8.2  | 24.9    | 2.1                           | 1.6        | 1                            |
|                           | NASPOT 7/2006/1185  | 6.9                                      | 8.8  | 32.9    | 2.0                           | 1.4        | 4                            |
|                           | NASPOT 12 O         | 11.6                                     | 15.3 | 37.8    | 2.3                           | 2.0        | 3                            |
|                           | NASPOT 13 O         | 9.1                                      | 11.7 | 31.6    | 2.4                           | 1.6        | 2                            |
|                           | Mean                | 8.6                                      | 11.0 | 31.8    | 2.2                           | 1.6        | NA                           |
|                           | LSD <sub>0.05</sub> | 3.2                                      | 3.9  | 6.9     | NS                            | NS         | NA                           |
|                           | cv (%)              | 33.4                                     | 32.0 | 19.6    | 31.6                          | 32.8       | NA                           |
| Buyende 2012B             |                     |  |      |         |                               |            | (n = 86; m = 33, w = 53)     |
|                           | NASPOT 10 O (LC)    | 9.7                                      | 10.8 | 22.1    | 1.7                           | 1.2        | 2                            |
|                           | NASPOT 7/2006/1185  | 2.5                                      | 3.6  | 28.5    | 1.9                           | 1.1        | 3                            |
|                           | NASPOT 12 O         | 11.7                                     | 13.5 | 32.0    | 2.1                           | 1.4        | 4                            |
|                           | NASPOT 13 O         | 4.0                                      | 4.9  | 25.0    | 2.6                           | 1.3        | 1                            |
|                           | Mean                | 6.9                                      | 8.2  | 26.9    | 2.1                           | 1.3        | NA                           |
|                           | LSD <sub>0.05</sub> | 2.4                                      | 2.6  | 5.1     | 0.5                           | NS         | NA                           |
|                           | cv (%)              | 37.3                                     | 34.1 | 21.0    | 24.1                          | 30.8       | NA                           |
| Rakai 2012B               |                     |  |      |         |                               |            | (n = 34; m = 14, w = 20)     |
|                           | NASPOT 10 O (LC)    | 6.5                                      | 8.6  | 17.9    | 2.7                           | 1.3        | 3                            |
|                           | NASPOT 7/2006/1185  | 3.8                                      | 5.6  | 25.6    | 2.6                           | 1.3        | 2                            |
|                           | NASPOT 12 O         | 8.3                                      | 10.9 | 27.4    | 2.7                           | 1.4        | 1                            |
|                           | NASPOT 13 O         | 4.3                                      | 5.7  | 21.6    | 2.7                           | 1.4        | 4                            |
|                           | Mean                | 5.7                                      | 7.7  | 23.1    | 2.8                           | 1.4        | NA                           |
|                           | LSD <sub>0.05</sub> | 1.6                                      | 1.6  | 4.9     | NS                            | NS         | NA                           |
|                           | cv (%)              | 29.6                                     | 25.7 | 42.1    | 28.9                          | 29.5       | NA                           |
| Oyam 2012B                |                     |  |      |         |                               |            | (n = 23; m = 5, f = 18)      |
|                           | NASPOT 10 O (LC)    | 12.3                                     | 13.2 | 17.5    | 2.0                           | 1.0        | 1                            |
|                           | NASPOT 7/2006/1185  | 5.6                                      | 6.7  | 15.1    | 2.1                           | 1.0        | 3                            |

(Continued on next page)

Table 5. (Continued) Performance of 'NASPOT 12 O', 'NASPOT 13 O', and local checks (LC) during two seasons in on-farm sweetpotato trials in various districts of Uganda during 2012.

| District <sup>z</sup> /yr | Cultivar            | Yield <sup>y</sup> (t·ha <sup>-1</sup> ) |      |         | Disease severity <sup>x</sup> |            | Taste test rank <sup>w</sup> |
|---------------------------|---------------------|--|------|---------|-------------------------------|------------|------------------------------|
|                           |                     | Marketable                               | Root | Biomass | SPVD                          | Alternaria |                              |
| Kabale 2012B              | NASPOT 12 O         | 12.3                                     | 14.3 | 19.7    | 2.6                           | 1.6        | 4                            |
|                           | NASPOT 13 O         | 9.3                                      | 10.3 | 18.8    | 2.4                           | 1.4        | 2                            |
|                           | Mean                | 9.9                                      | 11.1 | 17.8    | 2.3                           | 1.3        | NA                           |
|                           | LSD <sub>0.05</sub> | 3.2                                      | 3.1  | NS      | NS                            | 0.4        | NA                           |
|                           | cv (%)              | 28.9                                     | 24.7 | 28.6    | 26.5                          | 27.6       | NA                           |
|                           |                     |  |      |         |                               |            | (n = 22; m = 8, f = 14)      |
| Kabale 2012B              | NASPOT 10 O (LC)    | 9.3                                      | 11.6 | 25.5    | 2.9                           | 1.6        | 2                            |
|                           | NASPOT 7/2006/1185  | 4.5                                      | 6.5  | 23.6    | 2.3                           | 1.7        | 4                            |
|                           | NASPOT 12 O         | 10.4                                     | 14.6 | 29.8    | 2.9                           | 2.7        | 3                            |
|                           | NASPOT 13 O         | 8.2                                      | 10.9 | 24.4    | 2.3                           | 1.9        | 1                            |
|                           | Mean                | 8.1                                      | 10.9 | 25.8    | 2.6                           | 1.9        | NA                           |
|                           | LSD <sub>0.05</sub> | 3.0                                      | 3.6  | NS      | 0.7                           | 0.7        | NA                           |
|                           | cv (%)              | 29.6                                     | 25.7 | 42.1    | 28.9                          | 29.5       | NA                           |

<sup>z</sup>Rakai district represents the warm, moist tall grasslands; Isingiro, cool dry south western highlands; Buyende, the warm subhumid short grasslands; Oyam, the low plains and flat topped hills of northern Uganda; and Kabale, the cool moist southwest highlands; 2012A and 2012B, first and second rainy season in 2012.

<sup>y</sup>Yields based on 10–12 farms per district, gross plot was 30 m<sup>2</sup> (30 mounds), middle or net plot harvested was 18 m<sup>2</sup> (18 mounds of 54 plants); each farm in a district was treated as a replicate. NS = not significant.

<sup>x</sup>SPVD, Sweetpotato virus disease and alternaria bataticola blight severity in field evaluation, rating scale, 1 = no symptoms; 9 = very severe symptoms.

<sup>w</sup>Taste test rank was based on the aggregate pairwise comparison of the panel (farmers); n = number of farmers in the tasting panel, m = male, f = female; 1 = most preferred; 6 = least preferred.

NA = not applicable.

perform well in agroecologies with low-to-moderate SPVD pressure and with well-distributed rainfall for the first three months during growth.

#### Dissemination of Cultivars

'NASPOT 12 O' and 'NASPOT 13 O' are currently grown by local farmers where on-station and on-farm trials were conducted and in districts where HarvestPlus and collaborating partners promote dissemination of OFSP cultivars to alleviate vitamin A deficiency. A total of 100 households hosted the on-farm trials in six districts. Clonal propagation of the disseminated cultivars assures maintenance of genetic purity of sweetpotato cultivars at farm level. Some farmers and vine multipliers renew propagation material to avoid degeneration due to virus infection and indirectly maintain clonal purity. The districts of OFSP production currently include, Isingiro, Buyende, Rakai, Oyam, Bushenyi, Kabale, Kamwenge, Wakiso, Mukono, Mpigi, Soroti, Gulu, Lira, Kisoro, Mbarara, Masaka, Kibaale, Kole, and Kamuli. In the absence of promotions, the spread of the cultivars is mainly through farmer-to-farmer exchange or sale of planting materials.

#### Availability

The cultivars are maintained as pathogen-tested plants in the screenhouse at the Quarantine Station, Muguga, Kenya, and are maintained in the field by National Crops Resources Research Institute in Uganda. Requests for these cultivars should be addressed to: Seed Unit, CIP, P.O. Box 25171, Nairobi, Kenya. Requests for planting materials within Uganda should be directed to: Root

Crops Program, NaCRRI, P.O. Box 7084, Kampala.

#### Literature Cited

- Claessens, L., J.J. Stoolvogel, and J.M. Antle. 2009. Ex ante assessment of dual-purpose sweet potato in the crop-livestock systems of western Kenya: A minimum-data approach. *Agr. Syst.* 99:13–22.
- FAOSTAT. 2010. Roots, tubers, sweetpotato production. Food and Agriculture Organization of the United Nations, Rome, Italy. Sept. 2015. <<http://faostat3.fao.org/download/Q/QC/E>>.
- Grüneberg, W.J., D. Ma, R.O.M. Mwanga, E.E. Carey, K. Huamani, F. Diaz, R. Eyzaguirre, E. Guaf, M. Jusuf, A. Karuniawan, K. Tjintokohadi, Y.S. Song, S.R. Anil, M. Hossain, E. Rahaman, S.I. Attaluri, K. Somé, S.O. Afuape, K. Adofo, E. Lukonge, L. Karanja, J. Ndirigwe, G. Ssemakula, S. Agili, J.M. Randrianaivoarivony, M. Chiona, F. Chipungu, S.M. Laurie, J. Ricardo, M. Andrade, F. Rausch Fernandes, A.S. Mello, M.A. Khan, D.R. Labonte, and G.C. Yencho. 2015. Advances in sweetpotato breeding from 1993 to 2012. In: J. Low, M. Nyongesa, S. Quinn, and M. Parker (eds.). *Potato and sweetpotato in Africa: Transforming the value chains for food and nutrition security*. CAB International, Wallingford, UK.
- Hotz, C., C. Loechl, A. Lubowa, J.K. Tumwine, G. Ndezi, A.N. Masawi, R. Baingana, A. Carriquiry, A. Brauw, J.V. Meenakshi, and D.O. Gilligan. 2012. Introduction of  $\beta$ -carotene-rich orange sweet potato in rural Uganda resulted in increased vitamin A intakes among children and women and improved vitamin A status among children. *J. Nutr.* 142:1871–1880.
- International Potato Center, Asian Vegetable Research and Development Center, and the International Board for Plant Genetic Resources. 1991. Descriptors for sweetpotato. In: Z. Huaman (ed.). *International Board for Plant Genetic Resources*, Rome, Italy.
- Low, J.W., M. Arimond, N. Osman, B. Cuinguara, F. Zano, and D. Tschirley. 2007. A food-based approach introducing orange-fleshed sweet potatoes increased vitamin A intake and serum retinol concentrations in young children in rural Mozambique. *J. Nutr.* 137:1320–1327.
- Low, J., J. Lynam, B. Lemaga, C. Crissman, I. Barker, G. Thiele, S. Namanda, C. Wheatley, and A. Maria. 2009. Sweetpotato in sub-Saharan Africa. In: G. Loebenstein and G. Thottappilly (eds.). *The sweetpotato*. Springer, Berlin, Germany.
- Mwanga, R.O.M., C. Niringiye, A. Alajo, B. Kigozi, J. Namakula, I. Mpenbe, S. Tumwegamire, R.W. Gibson, and G.C. Yencho. 2011. 'NASPOT 11', a sweetpotato cultivar bred by a participatory plant-breeding approach in Uganda. *Hort-Science* 46:317–321.
- Ruel, M.T. 2001. Can food-based strategies help reduce vitamin A and iron deficiencies, a review of recent evidence. International Food Policy Research Institute Washington, DC.
- Ruel, M.T., H. Alderman, and the Maternal and Child Nutrition Study Group. 2013. Nutrition-sensitive interventions and programmes: How can they help to accelerate progress in improving maternal and child nutrition? *Lancet* 382:536–551.
- Ssemakula, G., C. Niringiye, B. Yada, M. Otema, G. Kyalo, J. Namakula, A. Alajo, B. Kigozi, R. Makumbi, G.C. Yencho, and R.O.M. Mwanga. 2013. Submission to the variety release Committee for the release of sweetpotato varieties. National Agricultural Research Organization (NARO)/National Crops Resources Research Institute (NaCRRI), Kampala, Uganda.
- Tumwegamire, S., R.O.M. Mwanga, M.I. Andrade, J.W. Low, G.N. Ssemakula, S.M. Laurie, F.P. Chipungu, J. Ndirigwe, S. Agili, L. Karanja, M. Chiona, J.C. Njoku, K. Mtunda, J. Ricardo, K. Adofo, E. Carey, and W.J. Grüneberg. 2014. Orange-fleshed sweetpotato for Africa. Catalogue 2014. 2nd ed. International Potato Center (CIP), Lima, Peru.