

Publ. Mat. **45** (2001), 467–475

## SMASH (CO)PRODUCTS AND SKEW PAIRINGS

J. N. ALONSO ALVAREZ, J. M. FERNÁNDEZ VILABOA AND  
R. GONZÁLEZ RODRÍGUEZ*Abstract*

---

Let  $\tau$  be an invertible skew pairing on  $(B, H)$ , where  $B$  and  $H$  are Hopf algebras in a symmetric monoidal category  $\mathcal{C}$  with (co)equalizers. Assume that  $H$  is quasitriangular. Then we obtain a new algebra structure such that  $B$  is a Hopf algebra in the braided category  ${}^H_H\mathcal{YD}$  and there exists a Hopf algebra isomorphism  $w: B \infty H \rightarrow B \bowtie_\tau H$  in  $\mathcal{C}$ , where  $B \infty H$  is a Hopf algebra with (co)algebra structure the smash (co)product and  $B \bowtie_\tau H$  is the Hopf algebra defined by Doi and Takeuchi.

---

### 1. Introduction

The smash product algebra and the smash coproduct coalgebra are well known in the theory of Hopf algebras. If  $B$  and  $H$  are Hopf algebras Radford [10] found necessary and sufficient conditions for the smash product algebra structure and the smash coproduct coalgebra structure on  $B \otimes H$  to afford  $B \otimes H$  a Hopf algebra structure. Also, in [10], Radford describes completely the structure of Hopf algebras with a projection. Assume  $B$  and  $H$  Hopf algebras and  $j: B \rightarrow H$  and  $f: H \rightarrow B$  Hopf algebra morphisms such that  $j \circ f = \text{id}_H$ . Then,  $B$  decomposes as a tensor product  $D \infty H$  where  $D$  is a Hopf algebra in the braided monoidal category of Yetter-Drinfeld modules and  $\infty$  denotes the smash (co)product. Afterwards, Majid in [8] and Bespalov in [2] obtain a braided interpretation of Radford's theorem. In [1] this result is proved in the context of braided categories, using the notions of  $H$ -Cleft comodule (module) algebras (coalgebras). On the other hand, Doi and Takeuchi in [6] studied the double crossproducts  $B \bowtie_\tau H$ . These are determined by an skew pairing  $\tau: B \otimes H \rightarrow K$  where  $B$  and  $H$  bialgebras. When  $B$  and  $H$  are Hopf algebras and  $\tau$  is a convolution invertible skew pairing,  $(B, \varphi_B =$

---

2000 *Mathematics Subject Classification.* 16W30, 17B37, 18D10, 81R50.

*Key words.* Quasitriangular Hopf algebra, monoidal category, skew pairing, smash (co)product.

$(\tau \otimes B \otimes \tau^{-1}) \circ (B \otimes H \otimes \delta_B \otimes H) \circ \delta_{B \otimes H} \circ c_{H, B}$ ) is a left  $H$ -module coalgebra and  $(H, \psi_H = ((\tau \circ c_{H, B}) \otimes H \otimes (\tau^{-1} \circ c_{H, B})) \circ (H \otimes B \otimes \delta_H \otimes B) \circ \delta_{H \otimes B})$  is a right  $B$ -module coalgebra. The construction of  $B \bowtie_{\tau} H$  is an example of Majid's double crossproduct  $B \bowtie H$  [9] because the Majid double crossproduct is defined as coalgebra  $B \bowtie H = B \otimes H$  and the product is equal to the one defined in [6] for  $B \bowtie_{\tau} H$ . In [6], it is shown that, if  $H$  is quasitriangular, there exists a Hopf algebra projection  $g: B \bowtie_{\tau} H \rightarrow H$  and therefore, using the Radford's theorem, it is possible to find  $D \in {}^H_H \mathcal{YD}$  such that  $D \infty H$  is isomorphic with  $B \bowtie_{\tau} H$ . In this paper we prove that the Hopf algebra  $B \bowtie_{\tau} H$ , defined by Doi and Takeuchi, is an  $H$ -Cleft comodule algebra and an  $H$ -Cleft module coalgebra and then using Theorem 3.2 of [1] we show that the object  $D$  such that  $D \infty H \approx B \bowtie_{\tau} H$  is  $B$  with a modified Hopf algebra structure. As a consequence, we obtain that  $(B, \varphi_B)$  is an  $H$ -module algebra too. Analogously, we have a similar result for the dual case studied by Caenepeel, Dăscălescu, Militaru and Panaite in [3]. The proof can be derived easily from the one developed in Theorem 4.1.

### 2. Preliminaries

We assume the reader is familiar with the machinery of monoidal categories. Details may be found in [7]. By  $(\mathcal{C}, \otimes, c, K)$  we denote a strict symmetric monoidal category with (co)equalizers where  $c$  is the natural isomorphism of symmetry and  $K$  is the base object. For every object  $A$  in  $\mathcal{C}$ ,  $\text{id}_A$  denotes the identity morphism.

An algebra in  $\mathcal{C}$  is a triple  $A = (A, \eta_A, \mu_A)$  where  $A$  is an object in  $\mathcal{C}$  and  $\eta_A: K \rightarrow A$ ,  $\mu_A: A \otimes A \rightarrow A$  are morphisms in  $\mathcal{C}$  such that  $\mu_A \circ (A \otimes \eta_A) = \text{id}_A = \mu_A \circ (\eta_A \otimes A)$ ,  $\mu_A \circ (A \otimes \mu_A) = \mu_A \circ (\mu_A \otimes A)$ . Given two algebras  $A = (A, \eta_A, \mu_A)$  and  $B = (B, \eta_B, \mu_B)$ ,  $f: A \rightarrow B$  is an algebra morphism if  $\mu_B \circ (f \otimes f) = f \circ \mu_A$ ,  $f \circ \eta_A = \eta_B$ . Also, if  $A, B$  are algebras in  $\mathcal{C}$ , the product algebra is  $AB = (A \otimes B, \eta_{A \otimes B} = \eta_A \otimes \eta_B, \mu_{A \otimes B} = (\mu_A \otimes \mu_B) \circ (A \otimes c_{B, A} \otimes B))$ .

A coalgebra in  $\mathcal{C}$  is a triple  $D = (D, \varepsilon_D, \delta_D)$  where  $D$  is an object in  $\mathcal{C}$  and  $\varepsilon_D: D \rightarrow K$ ,  $\delta_D: D \rightarrow D \otimes D$  are morphisms in  $\mathcal{C}$  such that  $(\varepsilon_D \otimes D) \circ \delta_D = \text{id}_D = (D \otimes \varepsilon_D) \circ \delta_D$ ,  $(\delta_D \otimes D) \circ \delta_D = (D \otimes \delta_D) \circ \delta_D$ . If  $D = (D, \varepsilon_D, \delta_D)$  and  $E = (E, \varepsilon_E, \delta_E)$  are coalgebras,  $f: D \rightarrow E$  is a coalgebra morphism if  $(f \otimes f) \circ \delta_D = \delta_E \circ f$ ,  $\varepsilon_E \circ f = \varepsilon_D$ . When  $D, E$  are coalgebras in  $\mathcal{C}$ , the product coalgebra is  $DE = (D \otimes E, \varepsilon_{D \otimes E} = \varepsilon_D \otimes \varepsilon_E, \delta_{D \otimes E} = (D \otimes c_{D, E} \otimes E) \circ (\delta_D \otimes \delta_E))$ .

Let  $D$  be a coalgebra and let  $A$  be an algebra. By  $\text{Reg}(D, A)$  we denote the set of invertible morphisms  $f: D \rightarrow A$  in  $\mathcal{C}$  respect to the

convolution operation  $f * g = \mu_A \circ (f \otimes g) \circ \delta_D$ .  $\text{Reg}(D, A)$  is a monoid where the unit element is  $\varepsilon_D \otimes \eta_A$ .

Let  $A$  be an algebra.  $(M, \psi_M)$  is a right  $A$ -module if  $M$  is an object in  $(\mathcal{C}, c)$  and  $\psi_M: M \otimes A \rightarrow M$  is a morphism in  $\mathcal{C}$  satisfying  $\psi_M \circ (M \otimes \eta_A) = \text{id}_M$ ,  $\psi_M \circ (\psi_M \otimes A) = \psi_M \circ (M \otimes \mu_A)$ . Given two right  $A$ -modules  $(M, \psi_M)$  and  $(N, \psi_N)$ ,  $f: M \rightarrow N$  is a morphism of right  $A$ -modules if  $\psi_N \circ (f \otimes A) = f \circ \psi_M$ . We denote the category of right  $A$ -modules by  $\mathcal{C}_A$ . In an analogous way we define the left  $A$ -modules and we denote this category by  ${}^A\mathcal{C}$ .

Let  $D$  be a coalgebra.  $(M, \rho_M)$  is a right  $D$ -comodule if  $M$  is an object in  $(\mathcal{C}, c)$  and  $\rho_M: M \rightarrow M \otimes D$  is a morphism in  $\mathcal{C}$  satisfying  $(M \otimes \varepsilon_D) \circ \rho_M = \text{id}_M$ ,  $(\rho_M \otimes D) \circ \rho_M = (M \otimes \delta_D) \circ \rho_M$ . Given two right  $D$ -comodules  $(M, \rho_M)$  and  $(N, \rho_N)$ ,  $f: M \rightarrow N$  is a morphism of right  $D$ -comodules if  $\rho_N \circ f = (f \otimes D) \circ \rho_M$ . We denote the category of right  $D$ -comodules by  $\mathcal{C}^D$ . Analogously,  ${}^D\mathcal{C}$  denotes the category of left  $D$ -comodules.

A bialgebra in  $\mathcal{C}$  is an object  $H$  with algebra and coalgebra structures, and such that  $\varepsilon_H$  and  $\delta_H$  are algebra morphisms (equivalently  $\eta_H$  and  $\mu_H$  are coalgebra morphisms). We say that  $H$  is a Hopf algebra if there exists a morphism  $\lambda_H: H \rightarrow H$  which is convolution inverse to the identical map. We call  $\lambda_H$  the antipode of  $H$ .

If  $H$  and  $B$  are bialgebras  $f: H \rightarrow B$  is a morphism of bialgebras if  $f$  is a morphism of algebras and coalgebras. Moreover, in this case if  $H$  and  $B$  are Hopf algebras, is not difficult to see that  $\lambda_B \circ f = f \circ \lambda_H$  and then we will say that  $f$  is a morphism of Hopf algebras.

Let  $H$  be a Hopf algebra in  $\mathcal{C}$ . Let  $M$  be an algebra (a coalgebra) such that  $(M, \psi_M)$  is in  $\mathcal{C}_H$  (resp.  $(M, \varphi_M)$  is in  ${}^H\mathcal{C}$ ). We say that  $M$  is a right (resp. left)  $H$ -module (co)algebra if  $\eta_M$  and  $\mu_M$  ( $\varepsilon_M$  and  $\delta_M$ ) are morphisms of right (resp. left)  $H$ -modules. If  $(M, \rho_M)$  is in  $\mathcal{C}^H$  (resp.  $(M, r_M)$  is in  ${}^H\mathcal{C}$ ). We say that  $M$  is a right (resp. left)  $H$ -comodule (co)algebra if  $\eta_M$  and  $\mu_M$  ( $\varepsilon_M$  and  $\delta_M$ ) are morphisms of right (resp. left)  $H$ -comodules.

We say that a right  $H$ -module coalgebra  $(M, \psi_M)$  is  $H$ -Cleft if there exists a cointegral  $g$ , i.e., an  $H$ -module morphism in  $\text{Reg}(M, H)$ . Replacing  $g$  by  $((\varepsilon_H \circ g^{-1}) \otimes g) \circ \delta_M$  we may assume that  $\varepsilon_H \circ g = \varepsilon_M$ . In this case, we will say that  $g$  is a total cointegral. Analogously, a right  $H$ -comodule algebra  $(M, \rho_M)$  is  $H$ -Cleft if there exists an integral  $f$ , i.e., an  $H$ -comodule morphism in  $\text{Reg}(H, M)$ . Replacing  $f$  by  $\mu_M \circ ((f^{-1} \circ \eta_H) \otimes f)$  we may assume that  $f \circ \eta_H = \eta_M$ . In this case, we will say that  $f$  is a total integral.

Let  $H$  be a bialgebra in  $\mathcal{C}$ ,  $(A; \varphi_A)$  be a left  $H$ -module algebra. We define

$$\begin{aligned}\eta_{A\sharp H} &= \eta_A \otimes \eta_H, \\ \mu_{A\sharp H} &= (\mu_A \otimes \mu_H) \circ (A \otimes \varphi_A \otimes H \otimes H) \circ (A \otimes H \otimes c_{H,A} \otimes H) \\ &\quad \circ (A \otimes \delta_H \otimes A \otimes H).\end{aligned}$$

It is well known that  $A\sharp H = (A \otimes H, \eta_{A\sharp H}, \mu_{A\sharp H})$  is an algebra in  $\mathcal{C}$ , called the smash product.

On the other hand, if  $(A, r_A)$  is a left  $H$ -comodule coalgebra, we have that  $A \bowtie H = (A \otimes H, \varepsilon_{A \bowtie H}, \delta_{A \bowtie H})$ , where  $\varepsilon_{A \bowtie H} = \varepsilon_A \otimes \varepsilon_H$  and  $\delta_{A \bowtie H} = (A \otimes \mu_H \otimes A \otimes H) \circ (A \otimes H \otimes c_{A,H} \otimes H) \circ (A \otimes r_A \otimes H \otimes H) \circ (\delta_A \otimes \delta_H)$ , is a coalgebra, called the smash coproduct.

### 3. Two-cocycles and skew pairings

Let  $H$  be a bialgebra in  $\mathcal{C}$ . A morphism  $\sigma \in \text{Reg}(H \otimes H, K)$  is a two cocycle if

$$\sigma \circ (H \otimes \mu_{\sigma H}) = \sigma \circ (\mu_{\sigma H} \otimes H)$$

where  $\mu_{\sigma H} = (\sigma \otimes \mu_H) \circ \delta_{H \otimes H}$ .

It is well known (see Theorem 1.6.a of [5]) that if  $\sigma$  is a two cocycle we have the next equalities:

- a)  $(\sigma \circ (\eta_H \otimes \eta_H)) * (\sigma^{-1} \circ (\eta_H \otimes \eta_H)) = \text{id}_K$ ,
- b)  $\sigma \circ (H \otimes \eta_H) = \varepsilon_H = \sigma \circ (\eta_H \otimes H)$ .

Moreover, if  $H$  is a Hopf algebra  $\sigma \circ (H \otimes \lambda_H) \in \text{Reg}(H \otimes H, K)$  has inverse  $\sigma^{-1} \circ (\lambda_H \otimes H)$ .

**Proposition 3.1.** *Let  $H$  be a bialgebra in  $\mathcal{C}$  and let  $\sigma$  be a two cocycle in  $\text{Reg}(H \otimes H, K)$ . Then:*

- a) *The triple  $(H, \eta_H, \mu_{\sigma H})$  is an algebra in  $\mathcal{C}$ .*
- b) *The triple  $(H, \eta_H, \mu_{H_{\sigma^{-1}}})$  is an algebra in  $\mathcal{C}$ , where  $\mu_{H_{\sigma^{-1}}} = (\mu_H \otimes \sigma^{-1}) \circ \delta_{H \otimes H}$ .*

*Proof:* We show b). The proof a) is analogous and we leave the details for the reader. Trivially,  $\mu_{H_{\sigma^{-1}}} \circ (\eta_H \otimes H) = \text{id}_H = \mu_{H_{\sigma^{-1}}} \circ (H \otimes \eta_H)$ . Finally,

$$\begin{aligned}\mu_{H_{\sigma^{-1}}} \circ (\mu_{H_{\sigma^{-1}}} \otimes H) &= \mu_H \circ (H \otimes \mu_H) \circ (H \otimes H \otimes H \otimes (\sigma^{-1} \circ (\mu_{H_{\sigma^{-1}}} \otimes H))) \circ \delta_{H \otimes H \otimes H} \\ &= \mu_H \circ (H \otimes \mu_H) \circ (H \otimes H \otimes H \otimes (\sigma^{-1} \circ (H \otimes \mu_{H_{\sigma^{-1}}})) \circ \delta_{H \otimes H \otimes H} \\ &= \mu_{H_{\sigma^{-1}}} \circ (H \otimes \mu_{H_{\sigma^{-1}}}).\end{aligned}\quad \square$$

**Proposition 3.2.** *Let  $H$  be a Hopf algebra in  $\mathcal{C}$  and let  $\sigma \in \text{Reg}(H \otimes H, K)$  be a two cocycle. Then*

$$H^\sigma = (H, \eta_H, \mu_{(\sigma_H)_{\sigma^{-1}}}, \varepsilon_H, \delta_H, \lambda_{H^\sigma})$$

*is a Hopf algebra in  $\mathcal{C}$  where  $\lambda_{H^\sigma} = (f \otimes \lambda_H \otimes f^{-1}) \circ (H \otimes \delta_H) \circ \delta_H$  and  $f = \sigma \circ (H \otimes \lambda_H) \circ \delta_H$  is a morphism in  $\text{Reg}(H, K)$  with inverse  $f^{-1} = \sigma^{-1} \circ (\lambda_H \otimes H) \circ \delta_H$ .*

*Proof:* See 1.6(b) of [5]. □

Let  $B$  and  $H$  be bialgebras in  $\mathcal{C}$ . A morphism  $\tau: B \otimes H \rightarrow K$  is called a skew pairing on  $(B, H)$  if:

- a)  $\tau \circ (\mu_B \otimes H) = (\tau \otimes \tau) \circ (B \otimes c_{B,H} \otimes H) \circ (B \otimes B \otimes \delta_H)$ ,
- b)  $\tau \circ (B \otimes \mu_H) = (\tau \otimes \tau) \circ (B \otimes c_{B,H} \otimes H) \circ (\delta_B \otimes c_{H,H})$ .

As a direct consequence of a) and b), if  $\tau$  is a convolution invertible skew pairing on  $(B, H)$ , then  $\tau \circ (\eta_B \otimes H) = \varepsilon_H$  and  $\tau \circ (B \otimes \eta_H) = \varepsilon_B$ .

Let  $\tau$  be a convolution invertible skew pairing on  $(B, H)$ . Then the morphism  $\sigma_\tau = \varepsilon_B \otimes (\tau \circ c_{H,B}) \otimes \varepsilon_H$  is a two cocycle in  $\text{Reg}(B \otimes H \otimes B \otimes H, K)$ , with inverse  $\sigma_\tau^{-1} = \varepsilon_B \otimes (\tau^{-1} \circ c_{H,B}) \otimes \varepsilon_H$ , called the two cocycle associated with  $\tau$  (1.5 of [6]).

Let  $H, B$  be Hopf algebras in  $\mathcal{C}$  and let  $\tau$  be a convolution invertible skew pairing on  $(B, H)$ . Then  $(B, \varphi_B)$  is a left  $H$ -module coalgebra and  $(H, \psi_H)$  is a right  $B$ -module coalgebra, where

$$\begin{aligned} \varphi_B &= (\tau \otimes B \otimes \tau^{-1}) \circ (B \otimes H \otimes \delta_B \otimes H) \circ \delta_{B \otimes H} \circ c_{H,B}, \\ \psi_H &= ((\tau \circ c_{H,B}) \otimes H \otimes (\tau^{-1} \circ c_{H,B})) \circ (H \otimes B \otimes \delta_H \otimes B) \circ \delta_{H \otimes B}. \end{aligned}$$

The object  $B \bowtie_\tau H = (B \otimes H, \eta_{B \otimes H}, \mu_{B \bowtie_\tau H}, \varepsilon_{B \otimes H}, \delta_{B \otimes H}, \lambda_{B \bowtie_\tau H})$  is a Hopf algebra in  $\mathcal{C}$  where

$$\begin{aligned} \mu_{B \bowtie_\tau H} &= (\mu_B \otimes \mu_H) \circ (B \otimes \varphi_B \otimes \psi_H \otimes H) \circ (B \otimes \delta_{H \otimes B} \otimes H), \\ \lambda_{B \bowtie_\tau H} &= (\varphi_B \otimes \psi_H) \circ \delta_{H \otimes B} \circ (\lambda_H \otimes \lambda_B) \circ c_{B,H}. \end{aligned}$$

Moreover, if  $\sigma_\tau$  is the two cocycle associated with  $\tau$  then the Hopf algebras  $(B \otimes H)^{\sigma_\tau}$  and  $B \bowtie_\tau H$  are the same [6] and [4].

#### 4. The Hopf algebras $B \bowtie_\tau H$ and $B \infty H$

A quasitriangular Hopf algebra in  $\mathcal{C}$  is a pair  $(H, \mathcal{R})$  where  $H$  is a Hopf algebra in  $\mathcal{C}$  and  $\mathcal{R}$  is a morphism in  $\text{Reg}(K, H \otimes H)$  such that:

- a)  $(\delta_H \otimes H) \circ \mathcal{R} = (H \otimes H \otimes \mu_H) \circ (H \otimes c_{H,H} \otimes H) \circ (\mathcal{R} \otimes \mathcal{R})$ .
- b)  $(H \otimes \delta_H) \circ \mathcal{R} = (\mu_H \otimes c_{H,H}) \circ (H \otimes c_{H,H} \otimes H) \circ (\mathcal{R} \otimes \mathcal{R})$ .

c)  $\mu_{H \otimes H} \circ ((c_{H,H} \circ \delta_H) \otimes \mathcal{R}) = \mu_{H \otimes H} \circ (\mathcal{R} \otimes \delta_H)$ .

If  $(H, \mathcal{R})$  is a quasitriangular Hopf algebra in  $\mathcal{C}$ , it is not difficult to prove that the morphism  $\mathcal{R}$  obeys:

- 1)  $(\varepsilon_H \otimes H) \circ \mathcal{R} = \eta_H = (H \otimes \varepsilon_H) \circ \mathcal{R}$
- 2)  $(\mu_H \otimes \mu_H \otimes \mu_H) \circ (H \otimes c_{H,H} \otimes c_{H,H} \otimes H) \circ (\mathcal{R} \otimes \mathcal{R} \otimes \mathcal{R})$   
 $= (\mu_H \otimes H \otimes (\mu_H \circ c_{H,H})) \circ (H \otimes H \otimes c_{H,H} \otimes H)$   
 $\circ (H \otimes c_{H,H} \otimes (\mu_H \circ c_{H,H}) \otimes H) \circ (\mathcal{R} \otimes \mathcal{R} \otimes \mathcal{R})$
- 3)  $\mathcal{R}^{-1} = (\lambda_H \otimes H) \circ \mathcal{R}$
- 4)  $(\lambda_H \otimes \lambda_H) \circ \mathcal{R} = \mathcal{R}$
- 5)  $(H \otimes \lambda_H) \circ \mathcal{R}^{-1} = \mathcal{R}$
- 6)  $(\mu_{H \otimes H} \otimes H) \circ (H \otimes H \otimes \delta_H \otimes H) \circ (\mathcal{R} \otimes \mathcal{R})$   
 $= (H \otimes \mu_{H \otimes H}) \circ (H \otimes \mathcal{R} \otimes \delta_H) \circ \mathcal{R}$ .

Let  $H$  be a Hopf algebra in  $(\mathcal{C}, c)$ . Let  $(M, \varphi_M)$  be in  ${}^H\mathcal{C}$  and  $(M, r_M)$  be in  ${}^H\mathcal{C}$ . We say that  $(M, \varphi_M, r_M)$  is in  ${}^H_H\mathcal{YD}$  if  $\varphi_M$  and  $r_M$  satisfies the compatibility condition:

$$(\mu_H \otimes M) \circ (H \otimes c_{M,H}) \circ ((r_M \circ \varphi_M) \otimes H) \circ (H \otimes c_{H,M}) \circ (\delta_H \otimes M) = (\mu_H \otimes \varphi_M) \circ (H \otimes c_{H,H} \otimes M) \circ (\delta_H \otimes r_M).$$

If  $B$  is a Hopf algebra in  ${}^H_H\mathcal{YD}$  with antipode  $\lambda_B$  then

$$B \bowtie H = (B \otimes H, \eta_{B \sharp H}, \mu_{B \sharp H}, \varepsilon_{B \bowtie H}, \delta_{B \bowtie H})$$

is a Hopf algebra in  $\mathcal{C}$  with antipode

$$\lambda_{B \bowtie H} = (\varphi_B \otimes H) \circ (H \otimes c_{H,B}) \circ (\delta_H \otimes B) \circ (\lambda_H \otimes \lambda_B) \circ (\mu_H \otimes B) \circ (H \otimes c_{B,H}) \circ (r_B \otimes H)$$

(see [2] for more details).

**Theorem 4.1.** *Let  $B$  and  $H$  be Hopf algebras in  $\mathcal{C}$ . Assume that  $H$  is quasitriangular. Let  $\tau \in \text{Reg}(B \otimes H, K)$  be a skew pairing on  $(B, H)$ . Then  $B$  is a Hopf algebra in the category  ${}^H_H\mathcal{YD}$  and there exists a Hopf algebra isomorphism  $w: B \bowtie H \rightarrow B \bowtie_\tau H$ .*

*Proof:* By 2.5 of [6],  $g: B \bowtie_\tau H \rightarrow H$ , defined by

$$g = (\tau \otimes (\mu_H \circ c_{H,H})) \circ (B \otimes c_{H,H} \otimes H) \circ (B \otimes H \otimes \mathcal{R})$$

and  $f = \eta_B \otimes H: H \rightarrow B \bowtie_\tau H$  are Hopf algebra morphisms such that  $g \circ f = \text{id}_H$ . Then  $(B \bowtie_\tau H, \rho_{B \bowtie_\tau H})$  where

$$\rho_{B \bowtie_\tau H} = (B \otimes H \otimes \tau \circ (\mu_H \circ c_{H,H})) \circ (B \otimes H \otimes B \otimes c_{H,H} \otimes H) \circ (\delta_{B \otimes H} \otimes \mathcal{R})$$

is a right  $H$ -Cleft comodule algebra with total integral  $f$  and  $(B \bowtie_{\tau} H, \psi_{B \bowtie_{\tau} H} = B \otimes \mu_H)$  is a right  $H$ -Cleft module coalgebra with total coin-tegral  $g$ . Thus, applying 3.2 of [1], the object  $(B \bowtie_{\tau} H)_0$  defined by the equalizer diagram

$$(B \bowtie_{\tau} H)_0 \xrightarrow{i_{B \bowtie_{\tau} H}} B \bowtie_{\tau} H \xrightarrow[\begin{smallmatrix} \rho_{B \bowtie_{\tau} H} \\ B \bowtie_{\tau} H \otimes \eta_H \end{smallmatrix}]{\rho_{B \bowtie_{\tau} H}} B \bowtie_{\tau} H \otimes H$$

is a Hopf algebra in  ${}^H_H\mathcal{YD}$  such that  $(B \bowtie_{\tau} H)_0 \infty H$  and  $B \bowtie_{\tau} H$  are isomorphic Hopf algebras. Moreover,  $(B \bowtie_{\tau} H)_0$  defined by the coequalizer of  $\psi_{B \bowtie_{\tau} H}$  and  $B \otimes H \otimes \varepsilon_H$  is such that  $(B \bowtie_{\tau} H)_0 = (B \bowtie_{\tau} H)_0$  and therefore  $(B \bowtie_{\tau} H)_0 = (B \bowtie_{\tau} H)_0 = B$  because the coequalizer morphism of  $\psi_{B \bowtie_{\tau} H}$  and  $B \otimes H \otimes \varepsilon_H$  is  $p = B \otimes \varepsilon_H$ . As a consequence,  $(B, \eta_B, m_B, \varepsilon_B, \delta_B, s_B)$  where  $m_B = \mu_B \circ (B \otimes \varphi_B) \circ (i_{B \bowtie_{\tau} H} \otimes B)$  and  $s_B = (\tau \otimes \varphi_B) \circ (B \otimes \mathcal{R} \otimes \lambda_B) \circ \delta_B$  is a Hopf algebra in  ${}^H_H\mathcal{YD}$ .

The isomorphism  $w$  between  $B \infty H$  and  $B \bowtie_{\tau} H$  is  $w = \mu_{B \bowtie_{\tau} H} \circ (i_{B \bowtie_{\tau} H} \otimes f)$ . Note that  $w = (B \otimes \tau \otimes \mu_H) \circ (\delta_B \otimes H \otimes \lambda_H \otimes H) \circ (B \otimes \mathcal{R} \otimes H)$  because

$$\begin{aligned} i_{B \bowtie_{\tau} H} &= i_{B \bowtie_{\tau} H} \circ p \circ (B \otimes \eta_H) \\ &= \mu_{B \bowtie_{\tau} H} \circ (B \bowtie_{\tau} H \otimes f^{-1}) \circ \rho_{B \bowtie_{\tau} H} \circ (B \otimes \eta_H) \\ &= \mu_{B \bowtie_{\tau} H} \circ (B \otimes H \otimes \eta_B \otimes \lambda_H) \circ \rho_{B \bowtie_{\tau} H} \circ (B \otimes \eta_H) \\ &= (B \otimes \tau \otimes \lambda_H) \circ (\delta_B \otimes \mathcal{R}). \end{aligned}$$

Finally, we compute  $\varphi_{(B \bowtie_{\tau} H)_0}$  and  $r_{(B \bowtie_{\tau} H)_0}$ ,

$$\begin{aligned} &i_{B \bowtie_{\tau} H} \circ \varphi_{(B \bowtie_{\tau} H)_0} \\ &= \mu_{B \bowtie_{\tau} H} \circ (B \otimes H \otimes (\mu_{B \bowtie_{\tau} H} \circ c_{B \otimes H, B \otimes H})) \\ &\quad \circ (f \otimes f^{-1} \otimes i_{B \bowtie_{\tau} H}) \circ (\delta_H \otimes B) \\ &= (B \otimes \mu_H) \circ (\varphi_B \otimes \psi_H \otimes \tau \otimes \mu_H) \circ (\delta_{H \otimes B} \otimes B \otimes H \otimes \lambda_H \otimes H) \\ &\quad \circ (H \otimes \delta_B \otimes \mathcal{R} \otimes H) \circ (H \otimes c_{H, B}) \otimes (H \otimes \lambda_H \otimes B) \circ (\delta_H \otimes B) \\ &= (B \otimes \mu_H) \circ (\varphi_B \otimes \psi_H \otimes \tau \otimes H) \circ (\delta_{H \otimes B} \otimes B \otimes H \otimes (\lambda_H \circ \mu_H \circ c_{H, H})) \\ &\quad \circ (H \otimes \delta_B \otimes \mathcal{R} \otimes H) \circ (H \otimes c_{H, B}) \circ (\delta_H \otimes B) \\ &= i_{B \bowtie_{\tau} H} \circ \varphi_B. \end{aligned}$$

Thus,  $\varphi_{(B \bowtie_{\tau} H)_0} = p \circ i_{B \bowtie_{\tau} H} \circ \varphi_{(B \bowtie_{\tau} H)_0} = p \circ i_{B \bowtie_{\tau} H} \circ \varphi_B = \varphi_B$  and, as a consequence, the left  $H$ -module coalgebra  $(B, \varphi_B)$  is a left  $H$ -module algebra too.

On the other hand,

$$\begin{aligned}
 & r_{(B \bowtie_{\tau} H)_0} \\
 &= (g \otimes p) \circ \delta_{B \otimes H} \circ i_{B \bowtie_{\tau} H} \\
 &= (\tau \otimes c_{B,H}) \circ (B \otimes H \otimes B \otimes \tau \otimes \mu_H) \circ (B \otimes H \otimes \delta_B \otimes c_{H,H} \otimes \lambda_H) \\
 &\circ (B \otimes c_{B,H} \otimes H \otimes H \otimes H) \circ (\delta_B \otimes \mathcal{R} \otimes \mathcal{R}). \quad \square
 \end{aligned}$$

**Acknowledgements.** The authors has been supported by the Xunta de Galicia, Project: PGIDT00PXI20706PR, and by the Ministerio de Ciencia y Tecnología, Project: BFM2000-0513-C02-02.

### References

- [1] J. N. ALONSO ALVAREZ AND J. M. FERNÁNDEZ VILABOA, Cleft extensions in braided categories, *Comm. Algebra* **28(7)** (2000), 3185–3196.
- [2] Y. BESPALOV, Crossed modules and quantum groups in braided categories, *Appl. Categ. Structures* **5(2)** (1997), 155–204.
- [3] S. CAENEPEEL, S. DĂSCĂLESCU, G. MILITARU AND F. PANAITTE, Coalgebra deformations of bialgebras by Harrison cocycles, copairings of Hopf algebras and double crosscoproducts, *Bull. Belg. Math. Soc. Simon Stevin* **4(5)** (1997), 647–671.
- [4] HUI-XIANG CHEN, Braiding structures of double crossproducts, *Tsukuba J. Math* **22(3)** (1998), 589–609.
- [5] Y. DOI, Braided bialgebras and quadratic bialgebras, *Comm. Algebra* **21(5)** (1993), 1731–1749.
- [6] Y. DOI AND M. TAKEUCHI, Multiplication alteration by two cocycles —the quantum version. *Comm. Algebra* **22(14)** (1994), 5715–5732.
- [7] S. MAC LANE, “*Categories for the working mathematicien*”, Graduate Texts in Mathematics **5**, Springer-Verlag, New York-Berlin, 1971.
- [8] S. MAJID, Braided matrix structure of the Sklyanin algebra and of the quantum Lorentz group, *Comm. Math. Phys.* **156(3)** (1993), 607–638.
- [9] S. MAJID, “*Foundations of quantum group theory*”, Cambridge University Press, Cambridge, 1995.
- [10] D. RADFORD, The structure of Hopf algebras with a projection, *J. Algebra* **92(2)** (1985), 322–347.



J. N. Alonso Alvarez:  
Departamento de Matemáticas  
Universidad de Vigo  
Campus Universitario Lagoas-Marcosende  
E-36280, Vigo  
Spain  
*E-mail address:* `jnalonso@uvigo.es`

J. M. Fernández Vilaboa:  
Departamento de Alxebra  
Universidad de Santiago de Compostela  
E-15771, Santiago de Compostela  
Spain  
*E-mail address:* `vilaboa@zmat.usc.es`

R. González Rodríguez:  
Departamento de Matemática Aplicada  
Universidad de Vigo  
Campus Universitario Lagoas-Marcosende  
E-36280, Vigo  
Spain  
*E-mail address:* `rgon@dma.uvigo.es`

Primera versió rebuda el 3 de novembre de 2000,  
darrera versió rebuda el 3 de juliol de 2001.