




Digestion in the cattle rumen

The digestive tract

Cattle belong to the group of animals referred to as ruminants. These animals have a “complex” stomach comprising four different compartments, which enable them to utilize various roughages efficiently and to obtain nutrients from them.



The four compartments are rumen, reticulum, omasum and abomasum. The abomasum is the true stomach and is comparable to the “simple” stomach of the non-ruminants. The other three are the “fore” stomachs.

The **reticulum** is a pouch-like structure, and the tissues are arranged in a network resembling a honeycomb. A small fold of tissue lies between the reticulum and the **rumen**, but the two are not actually separate compartments. Collectively they are called the **rumino-reticulum**.

The omasum is globe-shaped structure and contains leaves of tissue (like pages in a book). The omasum absorbs water and other substances from digestive contents.

The **abomasum** is the only compartment (also called the true stomach) with a glandular lining. Hydrochloric acid and digestive enzymes, needed for the breakdown of feeds, are secreted into the abomasum. The abomasum is comparable to the stomach of the non-ruminant.


The small intestine is about 40 metres long and has three parts, the duodenum, jejunum and ileum. The small intestine receives the secretions of the pancreas and the gall bladder, which aid digestion. Most of the digestive process is completed here, and many nutrients are absorbed through

the villi (small finger-like projections) into the blood and lymphatic systems.

The caecum is the large area located at the junction of the small and large intestine, where some previously undigested fibre may be broken down.

The large intestine is the last segment of the tract through which undigested feedstuffs pass. Absorption of water is the primary digestive activity occurring in the large intestine.

The digestion process



Within the rumen are billions of micro-organisms, both bacteria and protozoa. These micro-organisms initiate the process of digestion by:

- converting the carbohydrates (e.g. sugars, starches, cellulose etc.) to volatile fatty acids (VFA);
- breaking down the proteins into amino acids and even further into ammonia, carbon dioxide and VFA;
- forming new amino acids (including the “essential” amino acids) and more proteins by multiplying themselves.


The bodies of the micro-organisms contain proteins; more proteins are formed when they multiply; the proteins are made of amino acids - both essential and non-essential)

The micro-organisms also produce (synthesize) vitamins of the "B" group, which are absorbed and utilized by ruminants)

The most important features of the ruminant digestive process are:

- The ease with which roughages are converted into VFAs, which are then absorbed and utilized by the animals as a source of energy (and production of fat);
- The formation of essential amino acids (or proteins containing them, which are broken down into the respective amino acids in the abomasum) from non-protein nitrogen sources e.g. urea and proteins which do not contain any essential amino acids. The amino acids are subsequently absorbed and utilized to form proteins or as a source of energy.

To prevent reduction in intake and digestion of cellulose, rumen pH should not fall below 6 for long periods of time. The problem is less severe at low level of concentrate feeding than at high levels.




Rumen microbes synthesize vitamins of the B complex, vitamin C and K

Energy metabolism

Ruminants derive about 70% of their energy from the VFAs produced in the rumen. VFAs comprise of more than 95% of the acids produced in the rumen. In general, 65% is acetic, 20% propionic and 15% butyric.

Fibrous carbohydrates → Increased acetate → increased milk fat and decreased milk yield.

Non-fibrous carbohydrates → more VFAs → Increased propionate → Increased milk yield (increased glucose synthesis) and decreased milk fat.




Protein metabolism

Proteins are largely broken down in the rumen.

Non-protein nitrogen (eg urea in Urea-Molasses Blocks) is degraded into ammonia, which is used by bacteria to produce bacterial protein. This process requires easily available carbohydrates (eg Molasses in Urea-Molasses Blocks).

Bacterial protein is digested in the intestine.



An initial version of this factsheet was developed through the Agricultural Innovation Program for Pakistan. It was made possible by the support of the American people through the United States Agency for International Development (USAID). The contents are the sole responsibility of ILRI and do not necessarily reflect the views of USAID or the United States Government.

It was prepared by M.N.M Ibrahim (m.ibrahim@cgiar.org).



RESEARCH PROGRAM ON Livestock and Fish



INTERNATIONAL LIVESTOCK RESEARCH INSTITUTE

ilri.org

better lives through livestock

ILRI is a member of the CGIAR Consortium

Box 30709, Nairobi 00100, Kenya
Phone: +254 20 422 3000
Fax: +254 20 422 3001
Email: ILRI-Kenya@cgiar.org

Box 5689, Addis Ababa, Ethiopia
Phone: +251 11 617 2000
Fax: +251 11 617 2001
Email: ILRI-Ethiopia@cgiar.org

