



## Incidence and distribution of cassava mosaic begomoviruses in Côte d'Ivoire

Marie N. Y. Toualy<sup>1,2\*</sup>, Segun A. Akinbade<sup>2,3</sup>, Séka Koutoua<sup>1</sup>, H. Atta Diallo<sup>1</sup>, P. Lava Kumar<sup>2</sup>

<sup>1</sup>Université Nangui Abrogoua, Unité de Formation et de Recherche des Sciences de la Nature (UFR-SN), Laboratoire de Biologie et Amélioration des Productions Végétales, 02 BP 801 Abidjan 02, Côte d'Ivoire

<sup>2</sup>International Institute of Tropical Agriculture (IITA), PMB 5320, Ibadan, Nigeria

<sup>3</sup>Current address: Washington State Department of Agriculture, Prosser, United States

Article published on June 29, 2014

**Key words:** Cassava, cassava mosaic begomovirus, Côte d'Ivoire, Multiplex-PCR.

### Abstract

Cassava mosaic disease (CMD) caused by the whitefly-transmitted begomoviruses (family *Geminiviridae*) is a major threat to production of cassava (*Manihot esculenta* Crantz) in Côte d'Ivoire. A survey was conducted in the major production zones in Côte d'Ivoire to assess the incidence, severity, and distribution of cassava viral diseases. At each survey site, up to ten plants were assessed for symptom severity; incidence and samples were taken for virus testing. Techniques based on polymerase chain reaction (PCR) were used for the detection of cassava mosaic begomoviruses (CMBs) in the sampled leaves. Incidence of CMD varied from 0 to 100% and symptom severity from 1 to 5. Incidence differed significantly between the various agro-ecological zones ( $P < 0.001$ ), but severity was the same in those zones. Out of the 335 samples tested, *African cassava mosaic virus* (ACMV) was detected in 43.3%, *East African cassava mosaic Cameroon virus* (EACMCV) in 5.7%, and both ACMV and EACMCV in 31.3%; 19.7% of the samples analyzed were negative to all the viruses tested. None of the samples was tested positive to the *East African cassava mosaic virus-Uganda* (EACMV-Ug). These results suggest high incidence of CMD in the cassava production zones in Côte d'Ivoire and underscores a need for implementation of control measures including phytosanitary measures with utilization of CMD-free materials for planting and adoption of resistant varieties.

\* Corresponding Author: Marie N. Y. Toualy ✉ [nmalyt@gmail.com](mailto:nmalyt@gmail.com)

## Introduction

Cassava (*Manihot esculenta* Crantz, family *Euphorbiaceae*) is the third largest source of carbohydrates in the world and an important food staple crop in sub-Saharan Africa (Fargette *et al.*, 1994; Legg and Fauquet, 2004). The starchy tuberous roots are a source of food and income for more than 800 million people in Africa, Asia, and Latin America. Africa contributes more than 56% to the world's production (262.6 million tons) (FAO, 2014). Cassava is moving towards an industrialized system in which plant material is used for a variety of products including starch, flour, and animal feed (Thresh, 2006). Côte d'Ivoire is ranked no. 10 in area (360,000 ha) and no. 14 in production (2.4 million tons) among 40 cassava-producing countries in Africa (FAO, 2014). Most families consume cassava in various processed forms, such as attiéké (cassava couscous), foutou (pounded cassava mixed with pounded plantain), placali (paste), and gari (toasted granules). Human consumption of cassava leaves is popular only in the western part of the country. The demand for cassava and cassava-based foods is increasing in the country. However, productivity at 6.7 t/ha is very low compared with the average yield of 9.8 t/ha in Africa. This growing demand is mainly being met by an expansion in the cropping area which has increased by about 25% from 267,616 ha in 2002 to 360,000 ha in 2012 (FAO, 2014). Pests and diseases, especially cassava mosaic disease (CMD) caused by whitefly-transmitted begomoviruses (family *Geminiviridae*), are among the major factors for low yields. CMD is known to seriously decrease yields (Alabi *et al.*, 2011), and the effects are further exacerbated by the widespread cultivation of susceptible landraces such as Yacé and Bonoua (N'Zué *et al.*, 2005).

Nine different begomovirus species, commonly referred as cassava mosaic begomoviruses (CMBVs), have been identified in the CMD etiology in different regions of Africa (Alabi *et al.*, 2011; Harimalala *et al.*, 2012; Tiendrébéogo *et al.*, 2012). Of the various CMBVs, *African cassava mosaic virus* (ACMV), *East African cassava mosaic virus* (EACMV), and *East African cassava mosaic Cameroon virus* (EACMCV)

are known to be widely prevalent in sub-Saharan Africa (Patil and Fauquet, 2009). Several strains of CMBVs have also been identified; most notable of these is EACMV-Uganda (EACMV-Ug) which was responsible for the devastating pandemic in East Africa in the 1990s (Legg *et al.*, 2006). All these viruses are vectored by whitefly, *Bemisia tabaci* Gennadius (Hemiptera: *Aleyrodidae*), and also spread through the cuttings used routinely for vegetative propagation (Legg *et al.*, 2011).

CMD in Côte d'Ivoire was first reported by Hedin in 1931. This disease was known to be endemic in the coastal areas and also in the northern parts (Walter, 1980). Past studies have identified the occurrence of ACMV (Walter, 1980) and EACMCV (Pita *et al.*, 1999). This study was conducted to provide comprehensive information on the distribution and incidence of CMBVs and the severity of CMD in Côte d'Ivoire so that the complexity of disease situation could be understood and to contribute to the development of appropriate control measures.

## Materials and methods

### Survey

The survey was conducted in 2009 in 72 localities (farms) covering all the major cassava-growing areas. At each survey site, geo-reference points were taken using a GPS reader and details were recorded of location, varieties grown, and the incidence and severity of disease. Leaf samples for virus testing were taken from a minimum of five plants per field, wrapped in aluminium foil and stored in a cool box, and then transported to the laboratory for virus testing.

Disease incidence per field was calculated using the formula below:

$$\text{Incidence per field (\%)} = \frac{\text{Number of infected (symptomatic) plants} \times 100}{20}$$

The severity of CMD on symptomatic plants was assessed by rating plants on a 1 to 5 scale, as described by Hahn *et al.* (1980), where 1 = unaffected shoots (no symptoms); 2 = mild chlorosis, mild

distortions at bases of most leaves while the remaining parts of the leaves and leaflets appear green and healthy (symptoms on about 25% of the leaves); 3 = pronounced mosaic pattern on most leaves, narrowing and distortion of the lower one-third of the leaflets (symptoms on about 50% of the leaves); 4 = severe mosaic distortion of two-thirds of most leaves and a general reduction of leaf size and stunting of shoots (symptoms on about 75% of the leaves); and 5 = very severe mosaic symptoms on all leaves, distortion, twisting, and severe reduction of most leaves, accompanied by the severe stunting of plants (symptoms on about 100% of the leaves).

#### *DNA extraction*

Total DNA was isolated from the leaf samples according to the protocol described by Dellaporta *et al.* (1983). About 50 to 100 mg of the leaf sample was ground in 500  $\mu$ L of extraction buffer [100 mM Tris (pH 8.0) 8.5 mM EDTA and 10 mM  $\beta$ -mercaptoethanol]. Each extract was transferred into a 1.5 mL sterile microfuge tube and 33  $\mu$ L of 20% SDS (Sodium dodecyl sulfate) was added in each tube. The mixture was vortexed briefly and incubated at 65°C in a water bath for 10 min. The tubes were allowed to cool to room temperature, and 160  $\mu$ L of 5M potassium acetate was added to the mixture. The tubes were vortexed thoroughly and centrifuged at 10,000 g for 10 min. The supernatant was collected into a separate sterile microfuge tube and 200  $\mu$ L of cold iso-propanol was added to the tube and incubated at 4°C for 20 min. The solution was centrifuged at 10,000 g for 10 min to precipitate DNA. The supernatant was carefully removed and the DNA pellet was washed with 500  $\mu$ L of 70% ethanol and air dried at room temperature. The DNA pellet was dissolved in 50  $\mu$ L of TE buffer and stored at -20°C until further use.

#### *Polymerase Chain Reaction (PCR)-based detection of viruses*

PCR assays with oligonucleotide primers specific to ACMV, EACMV-like viruses, EACMCV, and EACMV-Ug were used to detect cassava mosaic begomoviruses in the leaf samples collected from the field (Table 1).

First, a multiplex-PCR assay developed by Alabi *et al.* (2008a) was used to test all the samples for the detection of ACMV and EACMV-like viruses using the primer CMBrep/F+ACMVrep/R+EACMVrep/R. All the samples that tested positive to EACMV were further analyzed for EACMCV using VNF031+VNF032 (Fondong *et al.*, 2000) and EACMV-Ug using specific primers UV-AL1/F1 and ACMV-CP/R3 (Zhou *et al.*, 1997).

PCR reaction composition was as follows: 2.5  $\mu$ L of PCR reaction buffer (5x), 0.25  $\mu$ L of 10 mM dNTPs (Promega, USA), 0.75  $\mu$ L of 25 mM MgCl<sub>2</sub>, 0.25  $\mu$ L of 20 pM of each primer, 1 U *Taq* DNA Polymerase (Promega, USA), 2  $\mu$ L of 1:50 (v/v) diluted DNA and sterile distilled water to a final volume of 12.5  $\mu$ L. PCR assays were performed in a GeneAmp PCR System 9700 (Applied Biosystems, CA, USA) using one cycle of 94°C for 1 min; 52°C for 2 min, and 72°C for 3 min, followed by 36 cycles, in which each cycle consisted of 94°C at 1 min, 52°C for 2 min and 72°C for 1.33 min with a final extension at 72°C for 5 min. The PCR amplified products were resolved by agarose gel electrophoresis, stained with ethidium bromide, and visualized under UV light using a gel documentation system (Biorad Universal Hood, Biorad Laboratories, Milan, Italy).

#### *Statistical analysis*

Statistical analysis of the incidence and symptoms severity by zone were conducted with the analysis of variance (ANOVA) with one criteria of classification. In the case of difference between the means, they were compared using the LSD test at 5%. The software used was Statistica 7.1.

## **Results**

### *Symptomatology*

A total of 335 leaf samples were collected in the 72 farms surveyed. Different types of symptom phenotypes occurred in the different locations in all the surveyed fields. Severe or mild mosaic symptoms were observed on 14.9% of the symptomatic plants; coiling, shoe-string, leaf distortion, stunting, and leaf

reduction were also observed on 78.9% of the plants (Fig. 1)



**Fig. 1.** Different types of virus symptoms on cassava leaves. (A) Asymptomatic leaves, (B) mild mosaic, (C) severe mosaic, (D) severe mosaic, distortion, leaf curl, (E) severe mosaic, severe distortion, severe reduction of leaf area, stunting.

#### Detection of viruses

In multiplex-PCR, CMBRep/ F+ACMVrep/ R+EACMVrep/ R primers amplified expected DNA fragments of 400 bp corresponding to ACMV in

74.6% of the leaf samples and 650 bp corresponding to EACMV in 37.01%. Mixed infection of ACMV and EACMV was detected in 31.34% of the samples. All the 124 samples that tested positive to EACMV in multiplex-PCR were re-tested with specific primers for EACMCV and EACMV-Ug. EACMCV was detected in 121 samples (97.58%), indicating that EACMCV is the prevailing EACMV-type of virus in the country. None of the samples was tested positive to EACMV-Ug. Altogether, 80.3% of the leaf samples were tested positive to viruses (ACMV, EACMCV or both). Out of the 335 leaf samples 43.3% were found to be infected by ACMV alone; 5.7% by EACMCV alone, and 31.3% with both ACMV and EACMCV; 19.7% of samples analyzed were negative to all the viruses tested. ACMV was detected in 71 and EACMCV in 49 of the 72 locations surveyed (Fig. 2). Only 6 of the 24 asymptomatic samples were tested positive to ACMV or EACMCV.

**Table 1.** Primers used for the detection of cassava mosaic begomoviruses.


Target virus	Primer	Primer sequence (5'→3')	Reference
ACMV+EACMV	CMB Rep-F ACMV Rep-R EACMV Rep-R	CRTCAATGACGTTGTACCA CAGCGGMAGTAAGTCMGA GGTTTGCAGAGAACTACATC	Alabi <i>et al.</i> , 2008a
EACMCV	VNF031 VNF032	GGATACAGATAGGGTCCAC GACGAGGACAAGAATTCCAAT	Fondong <i>et al.</i> , 2000
EACMV-Ug	UV-AL1/F1 ACMV-CP/R3	TGTCTTCTGGGACTTGTGTG TGCCTCCTGATGATTATATGTC	Zhou <i>et al.</i> , 1997

**Table 2.** Incidence and distribution of begomoviruses infecting cassava in Côte d'Ivoire.


Zones	Viruses (%)				
	ACMV	EACMV	ACMV+ EACMV	EACMCV	EACMV-Ug
Central	28.8	8.47	23.7	32.2	0
Central-West	33.3	9.1	27.3	30.3	0
East	22.2	5.6	69.4	75	0
North-East	69.2	0	19.2	19.2	0
North	47.4	7.7	17.9	24.3	0
South	52.4	2.9	36.9	39.8	0

NB: EACMCV percentage was calculated after detection by multiplex-PCR for ACMV and EACMV; all samples which were positive for EACMV were used for EACMCV and EACMV-Ug detection (124 tested samples after multiplex-PCR were positives for EACMV)





**Fig. 3.** Frequency (%) of fields by CMD incidence.



**Fig. 4.** Frequency (%) of leaf samples by CMD severity.

### Discussion

Findings of this study confirm that CMD is an endemic problem in all agro-ecological zones of Côte d'Ivoire. The incidence of disease varied from one zone to another but severity was similar in all zones. The incidence was high in the southern part of the country and relatively low in the North. The high incidence rates observed in various fields suggests that stem cuttings are the likely origin of the virus. Traditionally, farmers reuse as planting materials stems from their own farms which are often infected by viruses. This explains why CMD is widely disseminated and may be prevalent in areas where disease spread by vectors is limited. In addition, the two widely grown cultivars, Yacé and Bonoua, were found to be highly susceptible to the viruses that cause CMD. Most of the planting materials in the fields are already infected thus creating a dearth of CMD-free material. This leads to the perpetuation of viruses through infected stems (N'zué *et al.*, 2005). Previous studies have reported a similar situation with regard to CMD in several countries in sub-Saharan Africa and suggest that symptoms depend on

the virus species, strains, and mixed infections (Fauquet and Fargette, 1990; Harrison *et al.*, 1997; Otim-Nape *et al.*, 1997; Fondong *et al.*, 2000; Pita *et al.*, 2001a, b; Ogbe *et al.*, 2003; Alabi *et al.*, 2008b).

Analysis of infected cassava leaf samples confirmed the presence of ACMV and EACMCV but not EACMV-Ug. ACMV was the most prevalent begomovirus infecting cassava in all the zones of Côte d'Ivoire. Similar observations were reported by Harrison *et al.* (1997) in Uganda and Karakacha (2000) in Kenya. ACMV and EACMV occur in infected plants in Africa either alone or as mixed infections of different combinations (Fondong *et al.*, 2000; Berry and Rey, 2001; Ogbe *et al.*, 2003; Were *et al.*, 2004; Bull *et al.*, 2006). The proportion of single infections by EACMCV was lower (5.67%) than co-infections with ACMV (31%). Sources of inoculum are naturally infected plants when used as planting materials in successive years and also other herbaceous hosts of begomoviruses (Alabi *et al.*, 2008b). Together this may explain the severity observed on cassava in the survey. The majority of the new fields were planted with cuttings of plants harvested in previous fields, and probably infected. Also, the activities of insect vectors have an effect on CMD incidence and the transmission of begomoviruses (Patil and Fauquet, 2009). Most infections of EACMCV in Côte d'Ivoire were observed in the South.

Ogbe in his studies on begomoviruses on cassava in Nigeria (Ogbe *et al.*, 2001) observed that EACMCV was found in the humid forest, derived/coastal and southern Guinea savannas. EACMV-Ug was not detected in this study. However, occurrence of this strain in Burkina Faso underscores the need for vigilance against its spread in the country (Tiendrébéogo *et al.*, 2009).

Forty-eight (14.32%) samples from some symptomatic leaves were tested negative. No further investigations were made to determine the reasons for this negative reaction in PCR. Some samples from asymptomatic leaves were tested positive for viruses. This indicates that the absence of virus infection

cannot be assumed from the absence of visual symptoms on leaves.

### Conclusion

The study confirms the occurrence of two cassava mosaic begomoviruses, ACMV and EACMV, causing CMD infection in Côte d'Ivoire. ACMV and mixed infections of ACMV and EACMV were the most frequently occurring viruses in the plants infected by CMD. Characterization of EACMV using species specific primers indicated that EACMV species prevalent in the country is EACMCV. The high levels of disease incidence and severity found in the surveyed fields warrant the wider introduction of CMD resistant varieties.

### Acknowledgements

We gratefully acknowledge financial support from International Institute of Tropical Agriculture (IITA) Ibadan, Nigeria and West Africa Agricultural Productivity Program-Côte d'Ivoire (WAAPP-CI).

### References

- Alabi OJ, Kumar PL, Naidu RA.** 2011. Cassava mosaic disease: a curse to food security in sub-Saharan Africa. *APSnetFeatures* (ISSN: 2153-0297). doi:10.1094/APSnetFeature-2011-0701. <http://www.apsnet.org/publications/apsnetfeatures/Pages/cassava.aspx>
- Alabi OJ, Kumar PL, Naidu RA.** 2008a. Multiplex PCR method for the detection of *African cassava mosaic virus* and *East African cassava mosaic Cameroon virus* in cassava. *Journal of Virological Methods* **154**, 111-120.
- Alabi OJ, Ogbe FO, Bandyopadhyay R, Kumar PL, Dixon AGO, Hughes Jd'A, Naidu RA.** 2008b. Alternative hosts of *African cassava mosaic virus* and *East African cassava mosaic Cameroon virus* in Nigeria. *Archives of Virology* **153**, 1743-1747.
- Berry S, Rey MEC.** 2001. Molecular evidence for diverse populations of cassava infecting begomoviruses in southern Africa. *Archives of Virology* **146**, 1795-1802.
- Bull SE, Briddon RW, Sserubombwe S, Ngugi K, Markham PG, Stanley J.** 2006. Genetic diversity and phylogeography of cassava mosaic viruses in Kenya. *Journal of General Virology* **87**, 3053-3065.
- Dellaporta SL, Wood J, Hicks JB.** 1983. A plant DNA miniprep: version II. *Plant Molecular Biology Reporter* **1**, 19-21.
- F.A.O.** 2014. Crop Production data 2013. FAOSTAT. Food and Agriculture Organization of the United Nations, Rome, Italy. <http://www.fao.org>.
- Fargette D, Jeger M, Fauquet C, Fishpool LD.** 1994. Analysis of Temporal Disease Progress of *African cassava mosaic virus*. *Phytopathology* **54** (1), 91-98.
- Fauquet C, Fargette D.** 1990. *African cassava mosaic virus*: etiology, epidemiology, and control. *Plant Disease* **74**, 404-411.
- Fondong VN, Pita JS, Rey MEC, de Kochko A, Beachy RN, Fauquet CM.** 2000. Evidence of synergism between *African cassava mosaic virus* and a new double-recombinant geminivirus infecting cassava in Cameroon. *Journal of General Virology* **81**, 287-297.
- Hahn SK, Terry ER, Leuschner K.** 1980. Breeding cassava for resistance to cassava mosaic disease. *Euphytica* **29**, 673-683.
- Harimalala M, De Bruyn A, Hoareau M, Andrianjaka A, Ranomenjanahary S, Lefeuvre P, Lett JM.** 2012. Molecular characterization of a new alpha satellite associated with a cassava mosaic geminivirus in Madagascar. *Archives of Virology* **158**, 1829-1832.
- Harrison BD, Zhou X, Otim-Nape GW, Lui Y, Robinson DJ.** 1997. Role of a novel type of double infection in the geminivirus induced epidemic of severe cassava mosaic in Uganda. *Annals of Applied Biology* **131**, 437-448.

- Hedin L.** 1931. Culture du manioc en Côte d'Ivoire: Observations complémentaires sur la mosaïque. *Revue de Botanique Appliquée à l'Agriculture Coon* **11**, 558.
- Karakacha HW.** 2001. Serological and molecular characterization of begomoviruses infecting cassava (*Manihot esculenta* Crantz) in Africa. PhD **dissertation**, University of Hanover, Germany, 187 pp.
- Legg J, Fauquet C.** 2004. Cassava mosaic geminiviruses in Africa. *Plant Molecular Biology* **56**, 585-599.
- Legg JP, Owor B, Sseruwagi P, Ndunguru J.** 2006. Cassava mosaic virus disease in East and Central Africa: epidemiology and management of a regional pandemic. *Advances in Virus Research* **67**, 355-418.
- Legg JP, Jeremiah SC, Obiero HM, Maruthi MN, Ndyetabula I, Okao-Okuja G, Bouwmeester H, Bigirimana S, Tata-Hangy W, Gashaka G, Mkamilo G, Alicai T, Kumar PL.** 2011. Comparing the regional epidemiology of cassava mosaic and cassava brown streak virus pandemics in Africa. *Virus Research* **159**, 161-170.
- N'zué B, Zohouri, GP, Yapi-Gnaoré V.** 2005. Bien cultiver le manioc en Côte d'Ivoire. Centre National de Recherche Agronomique (CNRA), Abidjan, 4 pp.
- Ogbe FO, Thottappilly G, Dixon AGO, Mignouna HD, Quin FM.** 2001. Evidence of double infection and random occurrence of cassava begomoviruses in sub-Saharan Africa. In: *Roots Crops in the 21<sup>st</sup> Century*. Akoroda MO, Ngeve JM (eds.), *Proc. Seventh Triennial Symp. Intern/Soc. Tropical Roots Crop African Branch (ISTRAC-AB)*, Cotonou, Benin Republic, pp. 524-529.
- Ogbe FO, Thottappilly G, Dixon AGO, Mignouna HD.** 2003. Variants of *East African cassava mosaic virus* and its distribution in double infections with *African cassava mosaic virus* in Nigeria. *Plant Disease* **87**, 229-232.
- Otim-Nape GW, Bua A, Baguma Y, Thresh JM.** 1997. Epidemic of severe cassava mosaic disease in Uganda and efforts to control it. *African Journal of Root and Tuber Crops* **2**, 42-43.
- Patil B, Fauquet C.** 2009. Cassava mosaic geminiviruses: actual knowledge and perspectives. *Molecular Plant Pathology* **10**, 685-701.
- Pita JS, Fondong VN, Sangaré A, Otim-Nape GW, Ogwal S, Fauquet CM.** 1999. Biodiversity of Cassava Mosaic Disease in East and West Africa: Special Cases in Uganda and Ivory Coast. In: *Proc 4th Intl Sci Meeting of the Cassava Biotechnology Network*, Salvador, Bahia, Brazil. 3-7 November 1999.
- Pita JS, Fondong VN, Sangaré A, Kokora RNN, Fauquet CM.** 2001a. Genomic and biological diversity of the African cassava geminiviruses. *Euphytica* **120**, 115-125.
- Pita JS, Fondong VN, Sangaré A, Otim-Nape GW, Ogwal S, Fauquet CM.** 2001b. Recombination, pseudorecombination and synergism of geminiviruses are determinant keys to the epidemic of severe cassava mosaic disease in Uganda. *Journal of General Virology* **82**, 655-665.
- Thresh J.** 2006. Control of tropical plant virus diseases. *Virus Research* **67**, 245-295.
- Tiendrébéogo F, Lefeuvre P, Hoareau M, Traoré EVS, Barro N, Reynaud B, Traoré AS, Konaté G, Traoré O, Lett JM.** 2009. Occurrence of *East African cassava mosaic virus-Uganda* (EACMV-UG) in Burkina Faso. *Plant Pathology* **58**, 783.
- Tiendrébéogo F, Pierre Lefeuvre P, Hoareau M, Harimalala MA, De Bruyn A, Villemot J, Traoré VSE, Konaté G, Traoré AS, Nicolas Barro N, Reynaud B, Traoré O, Lett JM.** 2012. Evolution of *African cassava mosaic virus* by



recombination between bipartite and monopartite begomoviruses. *Virology Journal* **9**, 67.

**Walter B.** 1980. Isolation and purification of a virus transmitted from mosaic-diseased cassava in the Ivory Coast. *Plant Disease* **64**, 1040-1042.

**Were HK, Winter S, Maiss E.** 2004. Occurrence and distribution of cassava begomoviruses in Kenya. *Annals of Applied Biology* **145**, 175-184.

**Zhou X, Liu Y, Calvert L, Munoz C, Otim-Nape GW, Robinson DJ, Bryan D, Harrison BD.** 1997. Evidence that DNA-A of a geminivirus associated with severe cassava mosaic disease in Uganda has arisen by interspecific recombination. *Journal of General Virology* **78**, 2101-2111.