



# *Agrobacterium*-mediated genetic transformation of yam (*Dioscorea rotundata*): an important tool for functional study of genes and crop improvement

Evans Nyaboga, Jandra N. Tripathi, Rajesh Manoharan and Leena Tripathi\*

International Institute of Tropical Agriculture, Nairobi, Kenya

## Edited by:

Soren K. Rasmussen, University of Copenhagen, Denmark

## Reviewed by:

Surinder Singh, Commonwealth Scientific and Industrial Research Organisation, Australia  
Ahmad Parveez Ghulam Kadir, Malaysian Palm Oil Board, Malaysia

## \*Correspondence:

Leena Tripathi, International Institute of Tropical Agriculture, P. O. Box 30709-00100, Nairobi, Kenya  
e-mail: l.tripathi@cgiar.org

Although genetic transformation of clonally propagated crops has been widely studied as a tool for crop improvement and as a vital part of the development of functional genomics resources, there has been no report of any existing *Agrobacterium*-mediated transformation of yam (*Dioscorea* spp.) with evidence of stable integration of T-DNA. Yam is an important crop in the tropics and subtropics providing food security and income to over 300 million people. However, yam production remains constrained by increasing levels of field and storage pests and diseases. A major constraint to the development of biotechnological approaches for yam improvement has been the lack of an efficient and robust transformation and regeneration system. In this study, we developed an *Agrobacterium*-mediated transformation of *Dioscorea rotundata* using axillary buds as explants. Two cultivars of *D. rotundata* were transformed using *Agrobacterium tumefaciens* harboring the binary vectors containing selectable marker and reporter genes. After selection with appropriate concentrations of antibiotic, shoots were developed on shoot induction and elongation medium. The elongated antibiotic-resistant shoots were subsequently rooted on medium supplemented with selection agent. Successful transformation was confirmed by polymerase chain reaction, Southern blot analysis, and reporter genes assay. Expression of *gusA* gene in transgenic plants was also verified by reverse transcription polymerase chain reaction analysis. Transformation efficiency varied from 9.4 to 18.2% depending on the cultivars, selectable marker genes, and the *Agrobacterium* strain used for transformation. It took 3–4 months from *Agro*-infection to regeneration of complete transgenic plant. Here we report an efficient, fast and reproducible protocol for *Agrobacterium*-mediated transformation of *D. rotundata* using axillary buds as explants, which provides a useful platform for future genetic engineering studies in this economically important crop.

**Keywords:** *Dioscorea rotundata*, *Agrobacterium*-mediated transformation, axillary buds, selection marker gene, reporter gene

## INTRODUCTION

Yam (*Dioscorea* spp.) is an economically important food crop in many tropical countries especially in West Africa, South Asia, and the Caribbean. It is the second most important root and tuber crop in the world after cassava in terms of production (Jova et al., 2005; Adegbite et al., 2006). Yam tubers are nutritionally rich and a major source of dietary fiber, carbohydrates, vitamin C, and essential minerals (Charles et al., 2005; Polycarp et al., 2012). In addition, they are also known for their secondary metabolites (steroidal saponins, diterpenoids, and alkaloids) which have been exploited for pharmaceutical products (Mignouna et al., 2008). There are 600 *Dioscorea* species, however, only 10 of about 90 edible species are regularly cultivated for food. *Dioscorea rotundata* and *D. cayenensis* (both known as Guinea yam) are the most popular and economically important yams in West and Central Africa, where they are indigenous (Mignouna et al., 2003; Adegbite et al., 2006; Quain et al., 2011), while *D. alata* (referred to as water or greater yam) is the most widely distributed species globally. The consumer demand for yam is very high in sub-Saharan Africa,

but the yam production is declining in this region due to factors including diseases and pests, high costs of planting material, and decreasing soil fertility.

Diseases caused by viruses, fungi and bacteria and nematode pests either singly or in combination are responsible for yield losses (Nwankiti and Arene, 1978; Onwueme, 1978; Ng, 1992; Hughes et al., 1997). Nematodes are of particular concern because, apart from causing significant reduction in tuber yield and quality, they facilitate fungal and bacterial attacks. A major economic pest of yam is *Scutellonema bradys*, known as the yam nematode and causal agent of dry rot. This nematode occurs mostly in West Africa, where yam is its principal host, but is also recorded on yams from parts of South and Central America and Asia (Bridge et al., 2005). The nematode affects all the main cultivated yam species and cultivars in West Africa, mainly on mature tubers and during storage (Kwoseh, 2000; Coyne et al., 2006). Plant parasitic nematodes damage is also a critical factor in tuber quality reduction and yield loss in yam both in the field and storage (Adegbite et al., 2006). Yam nematodes reproduce and build up large populations

in stored tubers causing severe damage and facilitating fungal and bacterial attacks that cause anthracnose disease, dry rot, soft rot, and wet rot.

The most important field pathogen of yam is the foliar anthracnose-causing fungus *Colletotrichum gloeosporioides*, which is a major threat to yam cultivation, in all yam-producing areas (Abang et al., 2002). The disease causes leaf necrosis and shoots dieback of yams, thus reducing the photosynthetic efficiency of the plant, which results in yield losses of over 90% in susceptible genotypes (Egesi et al., 2007). Yam viral diseases also constitute a major pathological problem in yam production in all growing regions of the world. The use of infected vegetative propagules and uncontrolled introductions of infected germplasm by farmers through porous land borders have resulted in the presence of yam viruses in all yam-growing areas of West Africa (Goudou-Urbino et al., 1996; Hughes et al., 1997). Viruses reported to infect yams in West Africa include yam mosaic virus (YMV), yam mild mosaic virus (YMMV), *D. dumetorum* virus, *D. alata* bacilliform virus (DaBV), cucumber mosaic virus (CMV), *D. mottle* virus (DMoV), and *D. sansibarensis* virus (DsBV; Seal and Muller, 2007). Yam viruses are of substantial economic importance not only because of yield losses they cause, but also due to the high cost of preventive measures (Degras, 1993).

Efforts have been made in the form of conventional breeding toward the development of pest and disease resistant and high yielding varieties. Transfer of desirable genes from the secondary gene pool of wild relatives to the cultivated primary gene pool remains difficult in many crops, including yams (Spillane and Gepts, 2001). Genetic improvement of yam through breeding programs face challenges due to constraints such as the long breeding cycle, dioecious, poor flowering nature, polyploidy, vegetative propagation, and heterozygous genetic background (Mignouna et al., 2008). Genetic engineering has emerged as a valuable alternative and complementary approach to improve crops including yam. Because of the difficulties surrounding conventional breeding of yam, the use of transgenic approaches to improve this crop is particularly compelling. However, the capacity to achieve successful genetic transformation depends largely on efficient plant regeneration systems. Regeneration systems of *D. rotundata* and *D. alata* have been established (Adeniyi et al., 2008; Tripathi et al., unpublished). Recently, direct shoot organogenesis was also reported on petiole explants of *D. rotundata*, *D. cayenensis*, and *D. alata* (Anike et al., 2012). These regeneration systems have not been evaluated for the amenability to transformation.

*Agrobacterium*-mediated transformation is the gene delivery system, which is most preferred by plant biotechnologists because of its easy accessibility, tendency to transfer low copies of DNA fragments carrying the genes of interest at higher efficiencies with lower cost and the transfer of very large DNA fragments with minimal rearrangement (Shibata and Liu, 2000; Gelvin, 2003). Therefore, plant transformation through *Agrobacterium*-mediated DNA transfer has become a favored approach for many crop species (Barampuzam and Zhang, 2011). To date, there are only few reports of transient transformation of *D. alata* by particle bombardment using a reporter gene. Tör et al. (1993) successfully transformed cell suspension of *D. alata* by particle bombardment

and found that the foreign gene (*gusA*) could be stably expressed in the transgenic cells; however, transgenic plants were not produced from transformed cells. Tör et al. (1998) further demonstrated that foreign genes could be delivered to protoplasts of *D. alata* using a polyethylene glycol-mediated uptake method. However, regeneration of transgenic plants was not reported. Quain et al. (2011) also reported transient transformation of *D. rotundata* using *Agrobacterium*; however, it cannot be applied for crop improvement since no transgenic plant was regenerated. As efficient transformation system for yam is currently not available, therefore, the main objective of this study was to establish an efficient *Agrobacterium*-mediated transformation system for *D. rotundata*.

## MATERIALS AND METHODS

### YAM CULTIVARS AND EXPLANT PREPARATION


Yam cultivars of Tropical *D. rotundata* (TDr) 2579 and 2436 were obtained as plantlets from *in vitro* germplasm collection at International Institute of Tropical Agriculture (IITA)-Ibadan, Nigeria. All the cultivars were maintained *in vitro* and multiplied as shoot cultures on yam basic medium (YBM) containing Murashige and Skoog medium (MS) salts and vitamins, 0.05 mg/l 6-Benzylaminopurine (BAP), 0.02 mg/l Naphthaleneacetic acid (NAA), 25 mg/l Ascorbic acid, 30 g/l sucrose, 2.4 g/l gelrite. The pH of the medium was adjusted to 5.8 prior to autoclaving. The cultures were incubated in growth room at 28°C with 16/8 h photoperiod. The nodal explants (3–5 mm) containing axillary buds were excised from young *in vitro* shoots.

### SENSITIVITY OF AXILLARY BUD EXPLANTS TO ANTIBIOTICS

Prior to transformation experiments, the sensitivity tests to selective agents (hygromycin and kanamycin) were carried out in order to find an effective inhibitory concentration, which arrests the formation of shoot buds and shoots from nodal explants. The sensitivity to antibiotics was determined by culturing nodal explants having axillary buds on shoot bud induction medium (SBM; MS salts and vitamins, 1 mg/l BAP, 0.318 mg/l Copper sulfate, 20 g/l sucrose, 2.4 g/l gelrite) supplemented with different concentrations of hygromycin (0–15 mg/l) or kanamycin (0–250 mg/l). The cultures were transferred to a fresh medium containing the same level of antibiotic every 2 weeks and then scored for the frequency of regeneration after 8 weeks. The minimal inhibitory concentration of antibiotics was used in all the transformation experiments.

### Agrobacterium STRAINS AND BINARY VECTORS USED FOR TRANSFORMATION

*Agrobacterium tumefaciens* strains LBA4404 and EHA105 were used in this study. The binary vectors pCAMBIA1301, pCAMBIA2301 (CAMBIA Company, Australia) and pCAMBIA2300-gfp were used for transformation (Figure 1). The pCAMBIA1301 contained hygromycin phosphotransferase (*hpt*) gene as selection marker, while pCAMBIA2301 and pCAMBIA2300-gfp contained neomycin phosphotransferase II (*nptII*) as selectable



marker. Plasmids pCAMBIA1301 and pCAMBIA2301 contained the intron-containing *gusA* reporter gene, while plasmid pCAMBIA2300-gfp contained *gfp* as reporter gene. The binary vectors were transformed into *A. tumefaciens* strains LBA4404 and EHA105 by electroporation. Single colonies from Luria Bertani (LB) agar (10 g/l Tryptone, 5 g/l Yeast extract, 10 g/l NaCl, 15 g/l Agar, pH 7.5) plates containing kanamycin (50 mg/l), rifampicin (50 mg/l) and streptomycin (100 mg/l) were used to initiate 2 ml LB medium starter cultures. After 48 h shaking at 150 rpm at 28°C, this suspension was used to inoculate a 20 ml LB medium containing the same antibiotics, and grown overnight on a shaking platform at 150 rpm to reach an OD<sub>600</sub> of 1.0. Bacterial culture was centrifuged at 3500 rpm for 15 min and pellet was re-suspended in liquid SBM medium supplemented with 200 μM acetosyringone (Sigma Chemical Co.) and

grown further for 2–3 h at 25°C with shaking at 100 rpm. The optical density (OD<sub>600</sub>) of culture was checked and adjusted to 0.5. The bacterium suspension was used for transformation experiments.

#### INOCULATION OF EXPLANTS WITH *A. tumefaciens* AND CO-CULTIVATION

The explants of *D. rotundata* were immersed in bacterial suspension and vacuum infiltrated for 5 min followed by gentle shaking at 45 rpm for 30 min at room temperature. After inoculation, explants were blotted on sterile paper towels and co-cultivated for 3 days under dark condition at 28°C, in petri dishes containing SBM medium supplemented with 100 μM acetosyringone. Fifty explants were used in each experiment and transformation efficiency was compared with two *Agrobacterium* strains (LBA4404

and EHA105) and plasmids with different selection marker genes (*hpt* and *nptII*). Experiments were repeated twice.

### SELECTION AND REGENERATION OF TRANSGENIC PLANTS

Following co-cultivation, the explants were rinsed three to four times with liquid SBM medium supplemented with 500 mg/l carbenicillin and blotted dry on sterile filter paper and placed onto SBM supplemented with 250 mg/l carbenicillin for 1 week of recovery at 28°C 16/8 h photoperiod. After 1 week of incubation, the explants were transferred to fresh SBM medium supplemented with 250 mg/l carbenicillin and 7.5 mg/l hygromycin or 100 mg/l kanamycin depending on the plasmid used and incubated for 14 days at 28°C under 16/8 h photoperiod. This step was repeated twice with gradually increasing the antibiotic selection to 10 and 15 mg/l for hygromycin selection or 125 and 150 mg/l for kanamycin selection. The elongated shoots were separated and transferred to YBM containing 250 mg/l carbenicillin and 15 mg/l for hygromycin or 150 mg/l for kanamycin for 1 month for rooting. The well rooted plantlets were transferred to peat pellets, covered with transparent polythene bag and placed in a glasshouse at 28°C. After 3–4 weeks, each peat pellet is fragmented and plantlets transferred into pots containing sterile soil and covered with plastic bags. When plants have reached 30–50 cm in height the plastic bags were opened to allow further growth. The putative transgenic plants regenerated on selective medium were subjected to  $\beta$ -glucuronidase (GUS) histochemical assay or fluorescent microscopy and molecular analysis.

### HISTOCHEMICAL GUS ASSAY

Transient and stable histochemical GUS assay was carried out in different tissues as described by Jefferson et al. (1987) with modifications. Tissues were immersed in a buffer containing 2 mM X-Gluc, 50 mM phosphate, 50 mM potassium ferrocyanide and 5% Triton X-100 at pH 7.0 and vacuum infiltrated for 10 min, and then incubated overnight at 37°C for 24 h. Tissues containing chlorophyll were repeatedly soaked in 95% ethanol until chlorophyll was removed. Transient expression of *gusA* gene was examined in *Agro*-infected explants after 3 days of co-cultivation, while stable expression of the reporter gene was analyzed in leaves, shoots and roots isolated from putative transgenic plants regenerated on selective medium.

### VISUALIZATION OF GFP FLUORESCENCE

Transient and stable GFP expression was analyzed using a Nikon SMZ1500 stereomicroscope with GFP-Plus fluorescence module. The images were recorded in TIFF format using a digital camera. All plants putatively transformed with pCAMBIA2300-gfp were tested for GFP expression.

### GENOMIC DNA ISOLATION AND PCR ANALYSIS OF TRANSGENIC LINES

Plant genomic DNA for polymerase chain reaction (PCR) was extracted from regenerated putative transgenic young leaves using a DNeasy kit (Qiagen, GmbH, Germany). Specific primers used for *gusA* were: forward 5'-TTTAACTATGCCGGGATCCATCGC-3' and reverse 5'-CCAGTCGAGCATCTCTCAGCGTA-3'. Specific primers for *hpt* were: forward 5'-CCACTATCGGCGAGTACTTCTACACAGC-3' and 5'-GCCTGAACCTACCCGCGACGTCTGTGTC-3'.

PCR was conducted in a total volume of 20  $\mu$ l, containing 100 ng template DNA, 2  $\mu$ l 10  $\times$  buffer, 0.5  $\mu$ l of 10 mM dNTP, 0.5  $\mu$ l of 10  $\mu$ M primers, 1 unit of Taq DNA polymerase (Qiagen, GmbH, Germany). The PCR conditions were: initial denaturation at 94°C for 10 min, 35 cycles of denaturation at 94°C for 15 s, annealing at 62°C for 40 s for the *gusA* gene, 58°C for 40 s for the *hpt* gene, and extension at 72°C for 50 s, followed by final extension at 72°C for 7 min and holding at 4°C. The amplified PCR products were separated by electrophoresis on 0.8% (w/v) agarose gel stained with GelRed™ (Biotium) and visualized under a UV transilluminator and photographs were taken by the gel documentation system.

### RNA EXTRACTION AND RT-PCR ANALYSIS

Total RNA was extracted from 100 mg young leaf tissue of 10 transgenic lines and non-transgenic control plants using the RNeasy plant mini kit (Qiagen, GmbH, Hilden, Germany) and treated with DNase (RNeasy Plant Mini kit, Qiagen). The quantity and quality ( $A_{260/230}$  and  $A_{260/280}$ ) of total RNA were determined using the Nanodrop 2000. RNA was checked with PCR for absence of genomic DNA. Complementary DNA (cDNA) was synthesized using 1  $\mu$ g of total RNA and reverse transcriptase of the Maxima H Minus First Strand cDNA synthesis kit with oligoDT primers (Thermo scientific). For reverse transcriptase polymerase chain reaction (RT-PCR), 2  $\mu$ l of each cDNA synthesized was used. PCR cycling conditions included initial denaturation of 94°C for 10 min, followed by 35 cycles of 94°C for 15 s, 62°C for 40 s and 72°C for 50 s and final extension for 7 min. RT-PCR was performed with primers specific to the *gusA* gene as described above and housekeeping gene actin primers forward 5'-ACCGAAGCCCCTCTTAACCC-3' and reverse 5'-GTATGGCTGACACCATCACC-3'. The amplified RT-PCR products were separated and visualized as described in the PCR section above.

### DOT BLOT AND SOUTHERN BLOT ANALYSIS

The integration of the transgene into the genome of yam was analyzed using dot blot and Southern hybridization. Genomic DNA for dot blot analysis was extracted from twelve PCR positive transgenic lines using a DNeasy kit (Qiagen, GmbH, Germany). About 200 ng of genomic DNA in triplicate for each transgenic line was denatured at 98°C for 10 min, immediately chilled on ice for 5 min and immobilized onto a positively charged nylon membrane (Roche Applied Sciences, Mannheim, Germany) using a BIORAD Bio-Dot Microfiltration apparatus following the manufacturer's protocols and recommendations. The DNA samples were fixed on the membrane by cross-linking in a STRATA-LINK™ UV cross-linker. A *gusA*-specific probe was labeled with DIG-dUTP using PCR DIG Probe Synthesis Kit (Roche Applied Sciences, Mannheim, Germany). Hybridization, stringency washes and detection was carried out using a DIG Luminescent Detection Kit for Nucleic Acids (Roche Diagnostics, UK) according to the manufacturer's instructions.

For Southern blot analysis, genomic DNA was isolated from *in vitro* grown plants using cetyltrimethylammonium bromide (CTAB) method developed by Sharma et al. (2008) with modifications. The genomic DNA (20  $\mu$ g) of transgenic lines and non-transgenic control plant was digested with *Hind*III (New



England Biolabs, USA) for overnight at 37°C. The plasmid DNA digested with *Hind*III was used as positive control. Restricted DNA was separated on a 0.8% (w/v) agarose gel at 40 V for 6 h and transferred to a positively charged nylon membrane (Roche Applied Sciences, Mannheim, Germany) by capillary transfer method and fixed by cross-linking in a STRATA-LINK™ UV cross-linker. Hybridization and detection was performed as described above.

### STATISTICAL ANALYSIS

Data were subjected to significance by analysis of variance (ANOVA) and mean separation by Duncan's multiple range tests (DMRTs;  $p < 0.05$ ) using SPSS 11.09 software for Windows.

## RESULTS AND DISCUSSION

### SELECTION OF SELECTABLE MARKER SUITABLE FOR YAM TRANSFORMATION

An effective selection strategy is very important for developing an efficient genetic transformation procedure. This can be achieved by the use of a selective agent which prevents non-transformed tissues from regenerating, while permitting the development of transformed cells into shoots without any lethality of the explant tissues (Song et al., 2012). The choice of selection agent depends on the plant nature and each plant species responds differently to the selection agent. The bacterial *nptII* and *hpt* are the most frequently used selectable marker gene used for generating transgenic plants. These enzymes detoxify aminoglycoside antibiotics by phosphorylation, thereby permitting cell growth and development of transformed plant cells into shoots in the presence of antibiotics. Therefore, optimization of the dose of selection pressure using hygromycin or kanamycin is important, as a sub-optimal dose results in high frequency of escapes (Datta et al., 1990). On the other hand unnecessary high antibiotic doses not only kill untransformed tissues, but also inhibit growth of transformed cells, leading to delay in the regeneration process (Wilkinson and Dons, 1993). Therefore, optimization of the aminoglycoside concentration was based on the minimal antibiotic concentration sufficient to prevent regeneration of untransformed tissues. The effective antibiotic concentration is another important factor for selection and regeneration of transgenic plant cells. Antibiotics decay in the plant tissue media due to various factors such as light, pH, temperature (Padilla and Burgos, 2010) as well as the antibiotic degradation in the vicinity of transgenic cells able to inactivate the antibiotic (Rosellini et al., 2007). Regular subculturing on fresh selection media as described in our study increases the effective inhibitory action of the antibiotic used.


It has been reported that monocotyledonous plants are sensitive to hygromycin, but not to kanamycin (Hauptmann et al., 1988; Eady and Lister, 1998; Chin et al., 2007). Tör et al. (1993) also reported that the suspension cells of the *D. alata* showed a high tolerance to kanamycin and no growth inhibition even at a concentration of 500 µg/ml. However, our results indicated that shoot induction of *D. rotundata* is sensitive to both hygromycin and kanamycin (Table 1, Figure 2). Shoot bud induction and plant regeneration from axillary buds of nodal explants were completely inhibited on a medium containing 10 mg/l of hygromycin or 150 mg/l kanamycin. No study has been performed so far on the hygromycin-based selection for the transformation

**Table 1 | Effects of different concentrations of hygromycin and kanamycin on shoot bud induction and plant regeneration of *D. rotundata* using nodal explants.**

Antibiotic	Concentration (mg/l)	Explant development/regeneration response
Hygromycin	0	Green shoots regenerated
	5	Green shoots regenerated
	7.5	Axillary bud induced but no shoot production
	10	No shoots regenerated
	15	No shoots regenerated and explants were bleached
Kanamycin	0	Green shoots regenerated
	50	Green shoots regenerated
	75	Green shoots regenerated with pale patches
	100	Shoots regenerated and turned completely white
	125	Albino shoots regenerated, arrested development
	150	No shoots regenerated
	200	No shoots regenerated
250	No shoots regenerated	

of yam. The effective inhibitory concentration of hygromycin and kanamycin determined through this study will assist with the design of selection conditions for both *hpt* and *nptII* gene-based plasmids in the future for effective transformation of yam and will also be useful to engineer yam with multiple T-DNA insertion.

Efficient selection of transformed tissues was accomplished by increasing the selection pressure in a step wise manner from 7.5 to 15 mg/l and 100 to 150 mg/l for hygromycin and kanamycin, respectively. This process allows transformed explants to express effectively the antibiotic-resistance gene and initiate cell division, thus improving regeneration of explants to produce plants (Bull et al., 2009). Final selection on higher concentration of antibiotics also eliminates generation of false positive or escape plants. The use of low antibiotic concentrations in regeneration medium at early stages promotes transformed cell recovery, while a subsequent gradual increase of antibiotic concentration effectively eliminates non-transformed cells (Burgos and Albuquerque, 2003). Such pattern of selection was previously reported to be effective for cassava (Zhang and Puonti-Kaerlas, 2000), castor (Sujatha and Sailaja, 2005), dendrobium (Suwanaketchanatit et al., 2007), *Lotus corniculatus* (Nikolicć et al., 2007), grapevine (Fan et al., 2008), rapeseed (Liu et al., 2011), and spinach (Milojević et al., 2012) and also for jute, under kanamycin selection (Sarker et al., 2008).




### TRANSFORMATION, SELECTION, AND REGENERATION OF TRANSGENIC PLANTS

After co-cultivation, the explants were subjected to a resting period of 5–7 days in carbenicillin supplemented medium lacking selection agent to improve the regeneration of *Agro*-infected explants. It is reported that direct transfer to selective medium after co-cultivation could result in tissue necrosis of the explants (Khanna et al., 2007). *Agro*-infected nodal explants began to form axillary buds after 7 days on selective medium (Figure 3). The

induced buds started producing shoots 4–6 weeks after *Agro*-infection on selective regeneration medium supplemented with gradual increase of antibiotics every 2 weeks (Figure 3). In 8–10 weeks some of the shoots elongated and turned green and other shoots turn white or chimeric. In this study, a clear difference was observed during kanamycin selection process of transformed (green) and non-transformed (bleached) developing shoot buds. In order to eliminate possible chimeric plants, the shoots produced were sub-cultured several times with the same level of selection





pressure (15 mg/l or 150 mg/l kanamycin). After three subcultures, the chimeric shoots completely bleached and died while the transgenic shoots continued to survive and grow normally. All the shoots generated on selective medium produced roots when transferred onto YBM containing 15 mg/l hygromycin or 150 mg/l of kanamycin. In this rooting assay, only transformed shoots survived to rooting, whereas the escaping shoots did not produce roots. The putative transgenic plants generated were validated by PCR and proved to be transgenic by GUS assay and GFP fluorescence. A generalized scheme for stable genetic transformation protocol is shown in **Figure 4**.

The effect of cultivars, *Agrobacterium* strains and selection marker genes was examined on transformation efficiency using axillary buds as explants. Significant variation in transformation efficiency was observed among different cultivars based on *Agrobacterium* strains and antibiotic selection marker used (**Table 2**). We observed transformation efficiency of 9.4–18.2% depending on different transformation factors including the yam cultivars, *Agrobacterium* strains and antibiotic selection marker. Differences are known to exist between transformation efficiencies of different genotypes, expression vectors, selection marker

SBM; (C) proliferation of shoots within 8 weeks of culture on SBM; (D) rooting of elongated transformed shoot; (E) acclimatized transgenic plant maintained in glasshouse; (F) non-transgenic plant in soil in the glasshouse.

genes, and the strain of *Agrobacterium* as well as the tissue culture conditions (Cheng et al., 2004). Among these factors, the cultivar of the explants is considered as a crucial one that can hardly be overcome or complemented through optimizing other external factors, for example by manipulating highly virulent strains (Hansen et al., 1994) or by optimizing plant culture conditions (Zuo et al., 2002).

In the initial experiment we compared the effect of two *Agrobacterium* strains (EHA105 and LBA4404) harboring plasmid pCAMBIA1301 on the transformation efficiency of two cultivars of *D. rotundata*. No significant difference was observed in transformation efficiency due to various *Agrobacterium* strains used in the transformation (**Table 2**). However, there was a significant difference ( $p < 0.05$ ) in transformation efficiency among the two cultivars transformed. The transformation efficiency was higher (16–18%) for cultivar TDr 2436 in comparison to cultivar 2579 (12–14%). As there is no significant effect of *Agrobacterium* strain used on transformation efficiencies, only one strain EHA105 was used for further studies. Our results suggest that different cultivars of the same species may differ remarkably in their susceptibility to *Agrobacterium* infection. The biochemical basis of




FIGURE 4 | Schematic diagram showing various steps of stable genetic transformation of *D. rotundata* using nodal explants.



**Table 2 | Effect of *Agrobacterium* strains, cultivars, selectable marker, and reporter genes on transformation efficiency of *D. rotundata*.**

Plasmid construct	Cultivar	<i>Agrobacterium</i> strain	No. of regenerated plants on selective media	Transformation efficiency (%)
pCAMBIA1301	TDr2436	EHA105	9.4 ± 0.50 <sup>a</sup>	18.2 ± 0.37 <sup>a</sup>
		LBA4404	8.4 ± 0.24 <sup>ab</sup>	16.2 ± 0.37 <sup>ab</sup>
	TDr2579	EHA105	6.2 ± 0.37 <sup>cde</sup>	12.8 ± 0.37 <sup>cd</sup>
		LBA4404	7.6 ± 0.50 <sup>bc</sup>	14.4 ± 0.24 <sup>c</sup>
pCAMBIA2301	TDr2436	EHA105	7.4 ± 0.40 <sup>bc</sup>	14.8 ± 0.37 <sup>c</sup>
	TDr2579		4.8 ± 0.37 <sup>e</sup>	10.2 ± 0.37 <sup>e</sup>
pCAMBIA2300-gfp	TDr2436	EHA105	7.2 ± 0.80 <sup>bcd</sup>	14 ± 0.31 <sup>c</sup>
	TDr2579		5.8 ± 0.37 <sup>de</sup>	9.4 ± 0.24 <sup>e</sup>

The transformation experiments were repeated twice and 50 nodal explants were used for each transformation. Each value represents mean ± S.E. of two independent transformation experiments. Mean values followed by same letters within a column are not significantly different according to Duncan's multiple range test (DMRT) at 5% level.

$$\text{Transformation efficiency (\%)} = \frac{\text{Number of transgenic plants regenerated on selective medium}}{\text{Total number of explants transformed}} \times 100\%$$

these variations involves more complex mechanisms, as has been extensively reviewed by many authors (McCullen and Binns, 2006; Citovsky et al., 2007; Gelvin, 2010), that the transfer of DNA from *A. tumefaciens* to plant genome is a complex process involving a number of discrete, essential steps. The difference in the susceptibility of cultivars to *Agrobacterium* could be due to the presence of inhibitory metabolites to *Agrobacterium* sensory machinery (Liu and Nester, 2006; Maresh et al., 2006). Plant host defense response stimulated by *Agrobacterium* infection may be another factor influencing the susceptibility of plant cells to *Agrobacterium* (Ditt et al., 2005; Zipfel et al., 2006; Anand et al., 2008).

Selectable marker genes are required for establishment of efficient transformation in plants. In most cases, selection is based on antibiotic (kanamycin or hygromycin) or herbicide (phosphinothricin) resistance (Miki and McHugh, 2004). Selectable marker genes allow the plant cells that carry them to regenerate in media containing selective agents, while non-transformed cells die. The choice of a selectable marker gene depends on its efficiency, applicability to a wide range of plants, availability for researchers and its market acceptance (Kraus, 2010). In this study, we compared two selection marker genes (*hpt* and *nptII*) for transformation efficiency of two cultivars of *D. rotundata*. There was significant difference ( $p < 0.05$ ) in transformation efficiency using *hpt* and *nptII* as selectable marker genes (Table 2). Although there was a significant difference between *hpt* and *nptII* selectable markers, our study demonstrates that hygromycin as well as kanamycin selection are efficient and can be used for the recovery of transgenic yam tissues and plants. No escape plants were obtained with any of the selection agents used. The *hpt*-hygromycin system has been reported to be more efficient than the *nptII*-kanamycin and the phosphinothricin acetyl transferase (PAT)-phosphinothricin systems (Song et al., 2012). However, the availability of multiple resistance gene/antibiotic selection systems that allow for efficient selection of transgenic yam is essential to generate multiple improved traits from independent T-DNA cassettes. From a regulatory perspective, *nptII* is particularly interesting since it is present in a large proportion of commercialized genetically modified crops

(Miki and McHugh, 2004) and several independent studies have demonstrated its safe use in transgenic crops (Fuchs et al., 1993; Ramessar et al., 2007).

Other selection strategies that are free of antibiotic and/or herbicide-resistance genes can also be used to select transgenic plants. One of these strategies is the use of visual markers such as GUS (Jefferson et al., 1987) and green fluorescent protein (GFP; Prasher et al., 1992). In this study, we also compared two reporter genes (*gusA* and *gfp*) for the transformation efficiency of two cultivars of *D. rotundata*. There was no significant difference in transformation efficiency using *gusA* and *gfp* reporter genes for both cultivars (Table 2). Our results demonstrate that both *gusA* and *gfp* can be used as reporter genes for developing transgenic yam. The use of the *gusA* gene as a marker for transformation is effective and is widely applied in many species, including monocots such as rice (Wakasa et al., 2012) and orchard grass (Lee et al., 2006), and dicots such as common bean (Mukeshimana et al., 2013) and alfalfa (Duque et al., 2007). The presence of an intron in the *gusA* gene guards against false positives that may result from expression of the gene in *A. tumefaciens* (<http://www.cambia.org>). However, the GUS assay involves destruction of the tested tissues of transgenic explants; therefore, it should be performed at a later stage of the transformation study and presents a bottleneck for verification strategies in large-scale plant transformation protocols. GFP fluorescence visualization is also a useful tool for selecting transgenic yam plants. It has been widely used in the transformation of many plant species such as pepper (Jung et al., 2011), *Petunia hybrida* (Mußmann et al., 2011), and alfalfa (Duque et al., 2007). The fluorescent marker GFP is a highly versatile reporter gene, because the *gfp* gene expression can be monitored any time in living cells under a fluorescence microscope in a non-destructive manner (Chalfie et al., 1994). The same tissues may then be used for regeneration of stable transformants, which is not possible with other marker genes requiring destructive or toxic enzyme assays. Visual marker genes like *gfp* can even be used for yam transformation without using antibiotics or herbicides as the selection agents.

Complete transgenic plantlets ready for transfer to the greenhouse were produced within 3–4 months after *Agro*-infection. The rooted plants grew normally after transplanting to soil in the glasshouse (Figure 3) confirming that selection marker (*hpt* or *nptII*) or/and the reporter genes (*gusA* or *gfp*) does not have any apparent adverse effect on the normal development and morphology of the transgenic yam plants.

#### GUS EXPRESSION AND VISUALIZATION OF GFP FLUORESCENCE IN TRANSGENIC PLANTS

The use of both *gfp* and *gusA* genes as visual markers provides a useful method for confirming putative transformed plants (Padilla et al., 2006; Duque et al., 2007). The putative transgenic lines were verified by GUS histochemical assay or GFP fluorescence at different levels of plant development. The GUS staining was observed in nodal explants and emerging axillary buds indicating successful *Agro*-infection and transient expression of *gusA* gene (Figure 5). Transgenic plants displayed intense blue coloration in the leaf, stem, root, in contrast with non-transformed plant tissues, indicating the stable integration of the *gusA* gene into the genome and its expression (Figure 5).


In this study, we examined explants for GFP fluorescence at different stages, including axillary bud induction and *in vitro* plantlet (Figure 5). GFP expression was observed in both the axillary buds and leaves of transgenic plantlets (Figure 5). An advantage of visualizing GFP expression in our system was to enable us to select

transformation events at an early stage thus avoiding the transfer of non-transgenic shoot that survived the kanamycin selection, saving both time and labor.

#### MOLECULAR ANALYSIS OF TRANSGENIC PLANTS


To confirm the presence of foreign genes into the genome of transgenic plants, antibiotic-resistant plants were analyzed by PCR and RT-PCR (Figure 6). PCR analysis was performed with genomic DNA of putative transgenic and control non-transgenic plants in order to confirm the presence of transgene. The amplified product of about 500 base pairs corresponding to the internal fragment of *gusA* gene was observed from genomic DNA of all the transgenic plants tested using *gusA* gene-specific primers confirming the presence *gusA* transgene in transgenic plants. An amplified fragment of 958 base pairs was also observed from all tested transgenic plants using *hpt* specific primers confirming the presence of *hpt* gene. The amplified products were observed in all the plants tested, confirming the presence of both transgenes *gusA* and *hpt*, without any escape plant. No amplified product was observed in case of non-transgenic control plants.

The transgenic lines were analyzed using RT-PCR in order to verify the expression of *gusA* gene. The *gusA* transcript amplification of the expected fragment size (~500 base pair) was observed from samples of all the transgenic lines tested (Figure 6). Specific *Actin* transcript amplification was detected from all plants as an




**FIGURE 5 | Expression of reporter genes in tissues of putative transgenic plants of *D. rotundata*.** (A) Transient expression of *gusA* gene in emerging axillary buds 1 week after *Agro*-infection; (B) control non-transgenic plantlets; (C) stable expression of the *gusA* gene in transgenic plantlets; (D) transient

expression of *gfp* gene in emerging axillary buds 3 days after *Agro*-infection; (E) gene expression in transgenic buds produced ~1–2 weeks after *Agro*-infection; (F) leaf of transgenic plantlets viewed under UV light using GFP filter.



internal control for cDNA synthesis. A gDNA control was included in the assay with actin primers and showed the larger unspliced fragments, indicating that DNA contamination was below PCR detection levels in RNA samples. The results indicated that target genes were successfully incorporated into plant genome and were expressed in transgenic plants.

Polymerase chain reaction positive transgenic lines were further analyzed by dot blot and Southern blot hybridization using *gusA* probe to confirm integration of the transgene into the genome of yam. Genomic DNA of all the 12 transgenic lines tested by dot blot analysis were confirmed to contain the *gusA* gene (**Figure 7A**). Three transgenic lines were further tested with Southern blot





hybridization. Their unique hybridization patterns indicated that each transgenic line resulted from an independent transformation event (Figure 7B). No hybridization signal was detected in the non-transgenic control plant.

### FUTURE PROSPECTS OF GENETIC ENGINEERING FOR YAM IMPROVEMENT

Pests and diseases are among the most important of the many factors that have deleterious effects on yam tuber yield and quality and over time these constraints have become more severe (Amusa et al., 2003; Baimey et al., 2006; Aidoo et al., 2011). The progress made in this study in establishing an efficient genetic transformation system of *D. rotundata* could open up many avenues to produce disease resistant yams, through pathogen-derived resistance strategies, that would not be possible using conventional breeding approaches. Host plant resistance to anthracnose has been proposed as a viable alternative to the use of chemical fungicides in controlling the disease. However, studies have shown that there are no genotypes tolerant or resistant to the disease (Abang et al., 2002). Therefore, the most attractive strategy for anthracnose control in yam is probably the production of disease resistant plants through the transgenic approach. These approaches could include the expression of genes encoding plant, fungal or bacterial hydrolytic enzymes (Lorito et al., 1998), genes encoding elicitors of defense response (Keller et al., 1999) and antimicrobial peptides (AMPs; Broekaert et al., 1997). AMPs have a broad-spectrum antimicrobial activity against fungi as well as bacteria and most are non-toxic to plant and mammalian cells.

Use of resistant varieties can be an effective strategy in controlling yam nematodes, but there are no varieties known to be tolerant to nematodes. The use of transgenic plants will be an alternative approach to improve the nematode resistance of yam. Several transgenes confer plant resistance to both tropical and temperate plant parasitic nematodes (Atkinson et al., 2003). Cystatins inhibit nematode digestive cysteine proteinase activity, suppressing the growth and multiplication of these pests (Urwin et al., 1997) and is one of the transgenes that has been successfully used to control plant nematodes. It has been found that the cystatins confers the improved resistance to a range of nematodes in different crops like potato, sweet potato, rice, tomato, and plantain (Atkinson et al., 1996; Vain et al., 1998; Urwin et al., 2001; Chan et al., 2010; Gao et al., 2011; Roderick et al., 2012) and have proven efficacy under field conditions (Urwin et al., 2001, 2003). Such an approach could also be used to enhance resistance of yam against nematodes in the near future.

### CONCLUSION

We have established a highly efficient and simple *Agrobacterium*-mediated transformation protocol for *D. rotundata* using axillary buds as explants. Stable transgenic plantlets which showed presence, integration, and expression of transgenes were successfully regenerated within 3–4 months from axillary bud explants. To the best of our knowledge, this is the first report of *Agrobacterium*-mediated transformation of yam with experimental evidence of stable integration of T-DNA in *D. rotundata* genotypes. This protocol opens up an avenue for future genetic improvement of *D.*

*rotundata* with candidate genes of proven agronomic importance to attain sustainable production.

### ACKNOWLEDGMENTS

The authors would like to thank the Bill & Melinda Gates Foundation and IITA for financial support to this study and the Biosciences Eastern and Central Africa (BeCA) – International Livestock Research Institute, Kenya for providing the laboratory facilities.

### REFERENCES

- Abang, M. M., Winter, S., Green, K. R., Hoffmann, P., Mignouna, H. D., and Wolf, G. A. (2002). Molecular identification of *Colletotrichum gloeosporioides* causing anthracnose of yam in Nigeria. *Plant Pathol.* 51, 63–71. doi: 10.1046/j.0032-0862.2001.00655.x
- Adegbite, A. A., Saka, J. O., Agbaje, G. O., Owolade, O. F., Olaifa, G. O., Lawal, A., et al. (2006). Survey of plant parasitic nematodes associated with yams in Edo, Ekiti and Oyo States of Nigeria. *Afr. J. Agric. Res.* 1, 125–130. doi: 10.2478/v10045-008-0051-4
- Adeniyi, O. J., Adetimirin, V. O., Ingelbrecht, I., and Asiedu, R. (2008). Shoot and plantlet regeneration from meristems of *Dioscorea rotundata* Poir and *Dioscorea alata* L. *Afr. J. Biotechnol.* 7, 1003–1008.
- Aidoo, R., Nimoh, F., Bakang, J. E. A., Ohene-Yankyer, K., Fialor, S. C., and Abaidoo, R. C. (2011). Economics of small-scale seed yam production in Ghana: implications for commercialization. *J. Sustain. Dev. Afr.* 13, 65–78.
- Amusa, N., Adegbite, A., Mohammed, S., and Baiyewu, R. (2003). Yam diseases and its management in Nigeria. *Afr. J. Biotechnol.* 2, 497–502.
- Anand, A., Uppalapati, S. R., Ryu, C. M., Allen, S. N., Kang, L., Tang, Y. H., et al. (2008). Salicylic acid and systemic acquired resistance play a role in attenuating crown gall disease caused by *Agrobacterium tumefaciens*. *Plant Physiol.* 146, 703–715. doi: 10.1104/pp.107.111302
- Anike, F. N., Konan, K., Olivier, K., and Dodo, H. (2012). Efficient shoot organogenesis in petioles of yam (*Dioscorea* spp.). *Plant Cell Tissue Organ Cult.* 111, 303–313. doi: 10.1007/s11240-012-0195-9
- Atkinson, H. J., Urwin, P. E., Clarke, M. C., and McPherson, M. J. (1996). Image analysis of the growth of *Globodera pallida* and *Meloidogyne incognita* on transgenic tomato roots expressing cystatins. *J. Nematol.* 28, 209–215.
- Atkinson, H. J., Urwin, P. E., and McPherson, M. J. (2003). Engineering plants for nematode resistance. *Annu. Rev. Phytopathol.* 41, 615–639. doi: 10.1146/annurev.phyto.41.052002.095737
- Baimey, H., Coyne, D., and Labuschagne, N. (2006). Effect of fertilizer application on yam nematode (*Scutellonema bradys*) multiplication and consequent damage to yam (*Dioscorea* spp.) under field and storage conditions in Benin. *Int. J. Pest Manag.* 52, 63–70. doi: 10.1080/09670870600552380
- Barampuram, S., and Zhang, Z. (2011). “Recent advances in plant transformation,” in *Plant Chromosome Engineering: Methods and Protocols, Methods in Molecular Biology*, Vol. 701, ed. J. A. Birchler (Berlin: Springer), 1–35. doi: 10.1007/978-1-61737-957-4\_1
- Bridge, J., Coyne, D., and Kwoseh, C. K. (2005). “Nematode parasites of tropical root and tuber crops,” in *Plant Parasitic Nematodes in Subtropical and Tropical Agriculture, Revised 2nd Edn*, eds M. Luc, R. Sikora, and J. Bridge (Wallingford: CAB International), 221–258.
- Broekaert, W. F., Cammue, B. P. A., De Bolle, M. F. C., Thevissen, K., De Samblanx, G. W., and Osborn, R. W. (1997). Antimicrobial peptides from plants. *Crit. Rev. Plant Sci.* 16, 297–323. doi: 10.1080/07352689709701952
- Bull, S. E., Owiti, J. A., Niklaus, M., Beeching, J. R., Gruijssem, W., and Vanderschuren, H. (2009). *Agrobacterium*-mediated transformation of friable embryonic calli and regeneration of transgenic cassava. *Nat. Protoc.* 4, 1845–1854. doi: 10.1038/nprot.2009.208
- Burgos, L., and Alburquerque, N. (2003). Ethylene inhibitors and low kanamycin concentrations improve adventitious regeneration from apricot leaves. *Plant Cell Rep.* 21, 1167–1174. doi: 10.1007/s00299-003-0625-6
- Chalfie, M., Tu, Y., Euskirchen, G., Ward, W. W., and Prasher, D. C. (1994). Green fluorescent protein as a marker for gene expression. *Science* 263, 802–805. doi: 10.1126/science.8303295
- Chan, Y. L., Yang, A. H., Chen, J. T., Yeh, K. W., and Chan, M. T. (2010). Heterologous expression of taro cystatin protects transgenic tomato against *Meloidogyne*

- incognita* infection by means of interfering sex determination and suppressing gall formation. *Plant Cell Rep.* 29, 231–238. doi: 10.1007/s00299-009-0815-y
- Charles, A. L., Sriroth, K., and Huang, T. C. (2005). Proximate composition, mineral contents, hydrogen cyanide and phytic acid of 5 cassava genotypes. *Food Chem.* 92, 615–620. doi: 10.1016/j.foodchem.2004.08.024
- Cheng, M., Lowe, B. A., Spencer, T. M., Ye, X. D., and Armstrong, C. L. (2004). Factors influencing *Agrobacterium*-mediated transformation of monocotyledonous species. *In Vitro Cell. Dev. Biol. Plant* 40, 31–45. doi: 10.1079/IVP2003501
- Chin, D. P., Mishiba, K. I., and Mii, M. (2007). *Agrobacterium*-mediated transformation of protocorm-like bodies in *Cymbidium*. *Plant Cell Rep.* 26, 735–743. doi: 10.1007/s00299-006-0284-5
- Citovsky, V., Kozlovsky, S. V., Lacroix, B., Zaltsman, A., Dafny-Yelin, M., Vyas, S., et al. (2007). Biological systems of the host cell involved in *Agrobacterium* infection. *Cell. Microbiol.* 9, 9–20. doi: 10.1111/j.1462-5822.2006.00830.x
- Coyne, D. L., Williamson, V. M., and Banna, A. B. (2006). Analysis of the pathogenic, morphological and genetic variability of *Scutellonema bradys* on yam in West Africa. *Afr. Plant Prot.* 12:1.
- Datta, S. K., Peterhans, A., Datta, K., and Potrykus, I. (1990). Genetically engineered fertile Indica rice recovered from protoplasts. *Nat. Biotechnol.* 8, 736–740. doi: 10.1038/nbt0890-736
- Degras, L. (1993). *The Yam: A Tropical Root Crop*. London: Macmillan.
- Ditt, R. F., Nester, E., and Comai, L. (2005). The plant cell defense and *Agrobacterium tumefaciens*. *FEMS Microbiol. Lett.* 247, 207–213. doi: 10.1016/j.femsle.2005.05.010
- Duque, A. S., Araujo, S. S., Cordeiro, M. A., Santos, D. M., and Fevereiro, M. P. (2007). Use of fused *gfp* and *gus* reporters for the recovery of transformed *Medicago truncatula* somatic embryos without selective pressure. *Plant Cell Tissue Organ Cult.* 90, 325–330. doi: 10.1007/s11240-007-9268-6
- Eady, C. C., and Lister, C. E. (1998). A comparison of four selective agents for use with *Allium cepa* L. immature embryos and immature embryo derived cultures. *Plant Cell Rep.* 18, 117–121. doi: 10.1007/s002990050542
- Egesi, C. N., Onyeka, T. J., and Asiedu, R. (2007). Severity of anthracnose and virus diseases of water yam (*Dioscorea alata* L.) in Nigeria I: effects of yam genotype and date of planting. *Crop Prot.* 26, 1259–1265. doi: 10.1016/j.cropro.2006.10.025
- Fan, C., Pu, N., Wang, X., Wang, Y., Fang, L., Xu, W., et al. (2008). *Agrobacterium*-mediated genetic transformation of grapevine (*Vitis vinifera* L.) with a novel stilbene synthase gene from Chinese wild *Vitis pseudoreticulata*. *Plant Cell Tissue Organ Cult.* 92, 197–206. doi: 10.1007/s11240-007-9324-2
- Fuchs, R. L., Ream, J. E., Hammond, B. G., Naylor, M. W., Leimgruber, R. M., and Berberich, S. A. (1993). Safety assessment of the neomycin phosphotransferase II (NPTII) protein. *Biotechnol.* 11, 1543–1547. doi: 10.1038/nbt1293-1543
- Gao, S., Yu, B., Zhai, H., He, S.-Z., and Liu, Q.-C. (2011). Enhanced stem nematode resistance of transgenic sweetpotato plants expressing *oryzacystatin-I* gene. *Agric. Sci. China* 10, 519–525. doi: 10.1016/S1671-2927(11)60032-1
- Gelvin, S. B. (2003). *Agrobacterium*-mediated plant transformation: the biology behind the “Gene-Jockeying” tool. *Microbiol. Mol. Biol. Rev.* 67, 16–37. doi: 10.1128/MMBR.67.1.16-37.2003
- Gelvin, S. B. (2010). Plant proteins involved in *Agrobacterium*-mediated genetic transformation. *Annu. Rev. Phytopathol.* 48, 45–68. doi: 10.1146/annurev-phyto-080508-081852
- Goudou-Urbino, C., Givord, L., Quiot, J. B., Boeglin, M., Konate, G., and Dubern, J. (1996). Differentiation of yam mosaic virus isolates using symptomatology, Western blot assay and monoclonal antibody. *J. Phytopathol.* 144, 235–240. doi: 10.1111/j.1439-0434.1996.tb01522.x
- Hansen, G., Das, A., and Chilton, M. D. (1994). Constitutive expression of the virulence genes improves the efficiency of plant transformation by *Agrobacterium*. *Proc. Natl. Acad. Sci. U.S.A.* 91, 7603–7607. doi: 10.1073/pnas.91.16.7603
- Hauptmann, R. M., Vasil, V., Ozias-Akins, P., Tabaeizadeh, Z., Rogers, S. G., Fraley, R. T., et al. (1988). Evaluation of selectable markers for obtaining stable transformants in the Gramineae. *Plant Physiol.* 86, 602–606. doi: 10.1104/pp.86.2.602
- Hughes, J. D. A., Dongo, L. N., and Atiri, G. I. (1997). Viruses infecting cultivated yams (*Dioscorea alata* and *D. rotundata*) in Nigeria. *Phytopathology* 87:S45.
- Jefferson, R. A., Kavanagh, T. A., and Bevan, M. W. (1987). GUS fusions: beta-glucuronidase as a sensitive and versatile gene fusion marker in higher plants. *EMBO J.* 6, 3901–3907.
- Jova, M. C., Kosky, R. G., Perez, M. B., Pino, A. S., Vega, V. M., Torres, J. L., et al. (2005). Production of yam microtubers using a temporary immersion system. *Plant Cell Tissue Organ Cult.* 83, 103–107. doi: 10.1007/s11240-005-4853-z
- Jung, M., Shin, S. H., Park, J. M., Lee, S. N., Lee, M. Y., Ryu, K. H., et al. (2011). Detection of transgene in early developmental stage by GFP monitoring enhances the efficiency of genetic transformation of pepper. *Plant Biotechnol. Rep.* 5, 157–167. doi: 10.1007/s11816-011-0168-1
- Keller, H., Pamboukdjian, N., Ponchet, M., Poupet, A., Delon, R., Verrier, J. L., et al. (1999). Pathogen-induced elicitor production in transgenic tobacco generates a hypersensitive response and nonspecific disease resistance. *Plant Cell* 11, 223–235. doi: 10.1105/tpc.11.2.223
- Khanna, H. K., Paul, J. Y., Harding, R. M., Dickman, M. B., and Dale, J. L. (2007). Inhibition of *Agrobacterium*-induced cell death by antiapoptotic gene expression leads to very high transformation efficiency of banana. *Mol. Plant Microbe Interact.* 20, 1048–1054. doi: 10.1094/MPMI-20-9-1048
- Kraus, J. (2010). “Concepts of marker genes for plants,” in *Genetic Modification of Plants, Biotechnology in Agriculture and Forestry*, Vol. 64, eds F. Kempken and C. Jung (Berlin: Springer), 39–60.
- Kwoseh, C. K. (2000). *Identification of Resistance to Major Nematode Pests of Yams (Dioscorea spp.) in West Africa*. Ph.D. thesis, University of Reading, Reading.
- Lee, S. H., Lee, D. G., Woo, H. S., Lee, K. W., Kim, D. H., Kwak, S. S., et al. (2006). Production of transgenic orchard grass via *Agrobacterium*-mediated transformation of seed-derived callus tissues. *Plant Sci.* 171, 408–414. doi: 10.1016/j.plantsci.2006.05.006
- Liu, H., Guo, X., Naem, M. S., Liu, D., Xu, L., Zhang, W., et al. (2011). Transgenic *Brassica napus* L. lines carrying a two gene construct demonstrate enhanced resistance against *Plutella xylostella* and *Sclerotinia sclerotiorum*. *Plant Cell Tissue Organ Cult.* 106, 143–151. doi: 10.1007/s11240-010-9902-6
- Liu, P., and Nester, E. W. (2006). Indoleacetic acid, a product of transferred DNA, inhibits vir gene expression and growth of *Agrobacterium tumefaciens* C58. *Proc. Natl. Acad. Sci. U.S.A.* 103, 4658–4662. doi: 10.1073/pnas.0600366103
- Lorito, M., Woo, S. L., Garcia, I., Colucci, G., Harman, G. E., Pintor-Toro, J. A., et al. (1998). Genes from mycoparasitic fungi as a source for improving plant resistance to fungal pathogens. *Proc. Natl. Acad. Sci. U.S.A.* 95, 7860–7865. doi: 10.1073/pnas.95.14.7860
- Maresh, J., Zhang, J., and Lynn, D. G. (2006). The innate immunity of maize and the dynamic chemical strategies regulating two-component signal transduction in *Agrobacterium tumefaciens*. *ACS Chem. Biol.* 1, 165–175. doi: 10.1021/cb600051w
- McCullen, C. A., and Binns, A. N. (2006). *Agrobacterium tumefaciens* and plant cell interactions and activities required for interkingdom macromolecular transfer. *Annu. Rev. Cell Dev. Biol.* 22, 101–127. doi: 10.1146/annurev.cellbio.22.011105.102022
- Mignouna, H. D., Abang, M. M., and Asiedu, R. (2003). Harnessing modern biotechnology for tropical tuber crop improvement: Yam (*Dioscorea* spp.) molecular breeding. *Afr. J. Biotechnol.* 2, 478–485.
- Mignouna, H. D., Abang, M. M., and Asiedu, R. (2008). “Genomics of yams, a common source of food and medicine in the tropics,” in *Plant Genetics and Genomics: Crops and Models*, eds P. Moore and R. Ming (Berlin: Springer), 549–570.
- Miki, B., and McHugh, S. (2004). Selectable marker genes in transgenic plants: applications, alternatives and biosafety. *J. Biotechnol.* 107, 193–232. doi: 10.1016/j.jbiotec.2003.10.011
- Milojević, J., Tubić, L., Nolić, V., Mitić, N., Čalić-Drăgošavac, D., Vinterhalter, B., et al. (2012). Hygromycin promotes somatic embryogenesis in spinach. *Plant Cell Tissue Organ Cult.* 109, 573–579. doi: 10.1007/s11240-012-0117-x
- Mußmann, V., Serek, M., and Winkelmann, T. (2011). Selection of transgenic *Petunia* plants using the green fluorescent protein (GFP). *Plant Cell Tissue Organ Cult.* 107, 483–492. doi: 10.1007/s11240-011-9998-3
- Mukeshimana, G., Ma, Y., Walworth, A. E., Song, G. Q., and Kelly, J. D. (2013). Factors influencing regeneration and *Agrobacterium tumefaciens* mediated transformation of common bean *Phaseolus vulgaris* L. *Plant Biotechnol. Rep.* 7, 59–70. doi: 10.1007/s11816-012-0237-0
- Ng, S. Y. C. (1992). Micropropagation of white yam (*Dioscorea rotundata* Poir.). *Biotechnol. Agric. For.* 19, 135–159. doi: 10.1007/978-3-662-07770-2\_9
- Nikolić, R., Mitić, N., Ninković, S., and Nešković, M. (2007). Efficient genetic transformation of *Lotus corniculatus* L. using a direct shoot regeneration protocol, stepwise hygromycin B selection, and a super-binary *Agrobacterium tumefaciens* vector. *Arch. Biol. Sci.* 59, 311–317. doi: 10.2298/ABS0704311N

- Nwankiti, A. O., and Arene, O. B. (1978). Disease of yam in Nigeria. *PANS* 24, 468–496.
- Onwueme, I. C. (1978). *The Tropical Tuber Crops: Yams, Cassava, Sweet Potato and Cocoyams*. Chichester: Wiley.
- Padilla, I. M., and Burgos, L. (2010). Aminoglycoside antibiotics: structure, functions and effects on in vitro plant culture and genetic transformation protocols. *Plant Cell Rep.* 29, 1203–1213. doi: 10.1007/s00299-010-0900-2
- Padilla, I. M. G., Golis, A., Gentile, A., Damiano, C., and Scorza, R. (2006). Evaluation of transformation in peach *Prunus persica* explants using green fluorescent protein (GFP) and beta-glucuronidase (GUS) reporter genes. *Plant Cell Tissue Organ Cult.* 84, 309–314. doi: 10.1007/s11240-005-9039-1
- Polycarp, D., Afoakwa, E. O., Budu, A. S., and Otoo, E. (2012). Characterization of chemical composition and anti-nutrition factors in seven species within the Ghanaian yam (*Dioscorea*) germplasm. *Int. Food Res. J.* 19, 985–992.
- Prasher, D., Eckenrode, V., Ward, W., Prendergast, F., and Cormier, M. (1992). Primary structure of the *Aequorea victoria* green fluorescent protein. *Gene* 111, 229–233. doi: 10.1016/0378-1119(92)90691-H
- Quain, M. D., Egnin, M., Bey, B., Thompson, R., and Bonsi, C. (2011). Transgenic potential of *Dioscorea rotundata*, using *Agrobacterium*-mediated genetic transformation. *Asp. Appl. Biol.* 110, 71–79.
- Ramessar, K., Peremarti, A., Gomez-Galera, S., Naqvi, S., Moralejo, M., Munoz, P., et al. (2007). Biosafety and risk assessment framework for selectable marker genes in transgenic crop plants: a case of the science not supporting the politics. *Transgenic Res.* 16, 261–280. doi: 10.1007/s11248-007-9083-1
- Roderick, H., Tripathi, L., Babirye, A., Wang, D., Tripathi, J. N., Urwin, P. E., et al. (2012). Generation of transgenic plantain (*Musa* spp.) with resistance to plant pathogenic nematodes. *Mol. Plant Pathol.* 13, 842–851. doi: 10.1111/j.1364-3703.2012.00792.x
- Rosellini, D., Capomaccio, S., Ferradini, N., Sardaro, M. L. S., Nicolia, A., and Veronesi, F. (2007). Non-antibiotic, efficient selection for alfalfa genetic engineering. *Plant Cell Rep.* 26, 1035–1044. doi: 10.1007/s00299-007-0321-z
- Sarker, R. H., Al-Amin, G. M., Hassan, F., and Hoque, M. I. (2008). *Agrobacterium*-mediated genetic transformation of two varieties of jute (*Corchorus capsularis* L.). *Plant Tissue Cult. Biotechnol.* 18, 7–16.
- Seal, S., and Muller, E. (2007). Molecular analysis of a full-length sequence of a new yam badnavirus from *Dioscorea sansibarensis*. *Arch. Virol.* 152, 819–825. doi: 10.1007/s00705-006-0888-7
- Sharma, K., Mishra, A. K., and Misra, R. S. (2008). A simple and efficient method for extraction of genomic DNA from tropical tuber crops. *Afr. J. Biotechnol.* 7, 1018–1022.
- Shibata, D., and Liu, Y. G. (2000). *Agrobacterium*-mediated plant transformation with large DNA fragments. *Trends Plant Sci.* 5, 354–357. doi: 10.1016/S1360-1385(00)01689-7
- Song, G. Q., Walworth, A., and Hancock, J. F. (2012). Factors influencing *Agrobacterium*-mediated transformation of switchgrass cultivars. *Plant Cell Tissue Organ Cult.* 108, 445–453. doi: 10.1007/s11240-011-0056-y
- Spillane, C., and Gepts, P. (2001). “Evolutionary and genetic perspectives on the dynamics of crop gene pools,” in *Broadening the Genetic Base of Crop Production*, eds D. Cooper, C. Spillane, and T. Hodgkin (Wallingford: CABI), 25–70. doi: 10.1079/9780851994116.0025
- Sujatha, M., and Sailaja, M. (2005). Stable genetic transformation of castor (*Ricinus communis* L.) via *Agrobacterium tumefaciens*-mediated gene transfer using embryo axes from mature seeds. *Plant Cell Rep.* 23, 803–810. doi: 10.1007/s00299-004-0898-4
- Suwanaketchanati, C., Piluek, J., Peyachoknagu, S., and Huehne, P. S. (2007). High efficiency of stable genetic transformation in *Dendrobium* via microprojectile bombardment. *Biol. Plant.* 51, 720–727. doi: 10.1007/s10535-007-0148-z
- Tör, M., Ainsworth, C. C., and Mantell, S. H. (1993). Stable transformation of the food yam (*Dioscorea alata* L.) by particle bombardment. *Plant Cell Rep.* 12, 468–473. doi: 10.1007/BF00234714
- Tör, M., Twyford, C. T., Funes, I., Boccon-Gibod, J., Ainsworth, C. C., and Mantell, S. H. (1998). Isolation and culture of protoplasts from immature leaves and embryogenic cell suspensions of *Dioscorea* yams: tools for transient gene expression studies. *Plant Cell Tissue Organ Cult.* 53, 113–125. doi: 10.1023/A:1006028406641
- Urwin, P. E., Green, J., and Atkinson, H. J. (2003). Expression of a plant cystatin confers partial resistance to *Globodera*, full resistance is achieved by pyramiding a cystatin with natural resistance. *Mol. Breed.* 12, 263–269. doi: 10.1023/A:1026352620308
- Urwin, P. E., Lilley, C. J., McPherson, M. J., and Atkinson, H. J. (1997). Resistance to both cyst and root-knot nematodes conferred by transgenic *Arabidopsis* expressing a modified plant cystatin. *Plant J.* 12, 455–461. doi: 10.1046/j.1365-313X.1997.12020455.x
- Urwin, P. E., Troth, K. M., Zubko, E. I., and Atkinson, H. J. (2001). Effective transgenic resistance to *Globodera pallida* in potato field trials. *Mol. Breed.* 8, 95–101. doi: 10.1023/A:1011942003994
- Vain, P., Worland, B., Clarke, M. C., Richard, G., Beavis, M., Liu, H., et al. (1998). Expression of an engineered cysteine proteinase inhibitor (oryzacystatin-I<sub>D86</sub>) for nematode resistance in transgenic rice plants. *Theor. Appl. Genet.* 96, 266–271. doi: 10.1007/s001220050735
- Wakasa, Y., Ozawa, K., and Takaiwa, F. (2012). *Agrobacterium*-mediated co-transformation of rice using two selectable marker genes derived from rice genome components. *Plant Cell Rep.* 31, 2075–2084. doi: 10.1007/s00299-012-1318-9
- Wilmink, A., and Dons, J. J. M. (1993). Selective agents and marker genes for use in transformation of monocotyledonous plant. *Plant Mol. Biol. Rep.* 11, 165–185. doi: 10.1007/BF02670474
- Zhang, P., and Puonti-Kaerlas, J. (2000). PIG-mediated cassava transformation using positive and negative selection. *Plant Cell Rep.* 19, 1041–1048. doi: 10.1007/s002990000245
- Zipfel, C., Kunze, G., Chinchilla, D., Caniard, A., Jones, J. D. G., Boller, T., et al. (2006). Perception of the bacterial PAMP EF-Tu by the receptor EFR restricts *Agrobacterium*-mediated transformation. *Cell* 125, 749–760. doi: 10.1016/j.cell.2006.03.037
- Zuo, J. R., Niu, Q. W., Ikeda, Y., and Chua, N. H. (2002). Marker-free transformation: increasing transformation frequency by the use of regeneration-promoting genes. *Curr. Opin. Biotechnol.* 13, 173–180. doi: 10.1016/S0958-1669(02)00301-4

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 09 July 2014; accepted: 26 August 2014; published online: 15 September 2014.  
Citation: Nyaboga E, Tripathi JN, Manoharan R and Tripathi L (2014) *Agrobacterium*-mediated genetic transformation of yam (*Dioscorea rotundata*): an important tool for functional study of genes and crop improvement. *Front. Plant Sci.* 5:463. doi: 10.3389/fpls.2014.00463

This article was submitted to *Plant Biotechnology*, a section of the journal *Frontiers in Plant Science*.

Copyright © 2014 Nyaboga, Tripathi, Manoharan and Tripathi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.