

# Geographic priorities for research and development on dryland cereals and legumes

**ABSTRACT** - Dryland cereal and legume crops have often received less attention than maize, wheat and rice in terms of research and development priorities. But these crops are important globally because they serve populations living in poverty and particular socioeconomic and environmental niches. Compared to other crops, less is known about the global distribution of dryland cereal and legume crops and the conditions where they are grown. This research reports on an international effort to compile geographic information on cereal and legume crops and the conditions under which they are cultivated. The study suggested that dryland cereal and legume crops should be given priority in 18 farming systems worldwide, representing 160 million ha. The priority regions include the drier areas of South Asia, West and East Africa, Middle East and North Africa, Central America and other parts of Asia. These regions are prone to drought and heat stress, among other biotic and abiotic constraints. They represent 60% of the global poor and malnourished and make up half of the global population.

**INTRODUCTION** - The CGIAR (hereafter referred to as DCL) requested an analysis of the principal commodities of their proposed program and the farming systems in which they are found. The 12 priority crops of the Dryland Cereals and Legumes Agri-Food System research program are chickpea, common bean, cowpea, faba bean, groundnut, lentil, pigeon pea, soybean, barley, pearl millet, small millet and sorghum (DCL, 2015). The research builds on a global classification of farming systems, on maps of the spatial distribution of all 12 DCL crop commodities, on socioeconomic data on population, poverty, malnutrition, on market access, and on soil and climatic data.

- | The analysis identifies   | The study examines  | The analysis and resulting database provides   |
|---|---|--|
| <ul style="list-style-type: none"> <li>Where these crops occur in the context of constraints and opportunities for their development</li> <li>How can DCL technologies be geographically targeted for reducing poverty and malnutrition?</li> </ul> | <ul style="list-style-type: none"> <li>The spatial extents of key constraints to DCL crop production, using the most recent spatial data available</li> </ul> | <ul style="list-style-type: none"> <li>The first global farming systems information resource for specifically evaluating priorities for DCL crop improvement and management</li> </ul> |

**RESULTS**

- DCL crops should be given priority in 18 farming systems worldwide where they cover 160 million ha.
- These dryland system areas are home to the majority of the world's poor and food insecure.

FARMING SYSTEMS	REGION	2010 Population (000)	2005 Population (000)	2000 Rural Population (000)	2000 Urban Population (000)	Stunting Children (000)	Poverty headcount (US\$1.90) (000)	Poverty headcount (US\$1.25) (000)	DCL Crop Area (1000 ha)
Cereal-root crop mixed	SSA	116,472	84,100	69,199	14,961	6,300	39	52,965	73,618
Maize mixed	SSA	126,279	96,684	79,837	23,847	6,314	411	51,910	68,998
Agro-pastoral milw/sorghum	SSA	10,864	8,484	7,080	1,404	3,103	57	33,889	40,389
Pastoral	SSA	51,662	38,105	29,677	10,027	3,288	26.5	13,369	20,971
Rice-wheat	SA	613,964	491,399	365,499	125,901	28,292	515	237,306	440,256
Rainfed mixed	SA	400,921	356,767	249,337	107,430	24,541	62.6	157,816	286,661
Dry wheat	SA	47,017	45,000	33,544	12,056	3,610	65.5	18,074	22,620
Highland mixed	MENA	22,919	27,102	31,038	36,987	1,072	22.4	3,446	11,284
Rainfed mixed	MENA	47,798	38,815	18,852	24,963	499	16.3	1,666	6,415
Dryland mixed	MENA	56,966	47,204	18,093	29,132	750	18.7	1,128	4,390
Pastoral	MENA	38,441	33,845	16,798	17,047	1,068	21.9	988	4,444
Maize-beans (Mesoamerica)	LAC	88,137	76,106	29,688	47,420	2,038	35.9	4,084	9,278
Large scale cereal-vegetable	EECA	63,105	65,093	28,474	37,119	319	8.7	1,501	1,178
Small scale cereal-vegetable	EECA	19,852	19,898	8,763	11,135	382	19.6	658	2,175
Extensive cereal-livestock	EECA	90,121	95,405	36,044	67,361	70	3.7	1,639	2,848
Lowland rice	EAP	851,280	785,701	496,073	289,627	13,360	31.8	117,021	284,030
Upland intensive mixed	EAP	501,857	502,323	358,539	143,783	15,427	33.6	84,484	193,653
Temperate mixed	EAP	285,074	260,574	138,989	121,585	2,594	21.6	36,416	82,207
TOTAL		3,543,606	3,159,275	2,023,332	1,136,441	114,917		815,472	1,546,593

FARMING SYSTEMS	REGION	DCL Crop Area (ha)	Potential Drought Impact Index	Temperature Change 2050
Cereal-root crop mixed	SSA	21,327,541	2,971,040	2.48
Maize mixed	SSA	7,606,508	1,592,730	2.47
Agro-pastoral milw/sorghum	SSA	18,691,342	7,644,810	2.77
Pastoral	SSA	10,808,337	7,409,830	2.73
Rice-wheat	SA	11,262,839	4,431,820	2.83
Rainfed mixed	SA	20,769,078	7,858,182	2.48
Dry rainfed	SA	8,695,308	2,868,150	2.36
Highland mixed	MENA	2,961,344	98,050	3.01
Rainfed mixed	MENA	1,588,829	123,471	2.64
Dryland mixed	MENA	3,340,374	104,013	2.39
Pastoral	MENA	1,000,516	10,959	2.93
Maize-beans (Mesoamerica)	LAC	1,749,799	391,401	2.36
Large scale cereal-vegetable	EECA	6,947,991	86,032	2.82
Small scale cereal-vegetable	EECA	2,550,258	1,849	2.82
Extensive cereal-livestock	EECA	9,160,822	17,198	3.31
Lowland rice	EAP	8,778,265	982,407	2.25
Upland intensive mixed	EAP	7,868,661	1,065,610	2.42
Temperate mixed	EAP	6,530,133	1,088,910	2.91




Figure 4. Combination of Crops. Many of the DCL crops are found in environment prone to heat and drought stress – two constraint key to crop improvement efforts.




Figure 2. Once the database was organized we could graph relationships between variables, such as this comparison of rainfall and drought




Figure 3. All the data from the project was compiled and developed within an online Atlas (<http://www.eatlasdcl.cgiar.org/>) available for download.

**MATERIALS AND METHODS** - The present study uses a spatial overlay, biophysical and socioeconomic information are organized according to the 63 Dixon farming systems (Dixon et al., 2001) but with a focus on the 12 principal commodities and farming systems of DCL. A key advantage of this research was that instead of analyzing crop information by country (250 in total), subnational estimates of crop distribution are generated based on pixel level data (Hyman et al., 2008). Spatial overlay was used to organize the data into spatial units according to farming system and combinations of farming systems and country. The result of the overlay procedure is a set of database files (dBase format) organized by farming system region and combination of farming system region and country. The process facilitated an analysis of DCL crops in 18 farming systems where these crops are concentrated.




Figure 1. Crop combinations with large areas

**DISCUSSION**

- South Asia and Sub-Saharan Africa are the most important regions for crop improvement and adapted crop management practices
- Adverse biotic and abiotic constraints and socioeconomic conditions set the context for research and development in these priority systems
- Future geographic research is needed to update maps to latest conditions, improve spatial resolution and carry out genotype-by-environment analysis.

**REFERENCES** – See our website at <http://www.eatlasdcl.cgiar.org/>, or our published paper: Hyman, G., Barona, E., Biradar, C., Guevara, E., Dixon, J., Beebe, S., Castano, S.E., Alabi, T., Gumma, M.K., Sivasankar, S. and Rivera, O., 2016. Priority regions for research on dryland cereals and legumes. *F1000 Research*, 5(885), pp.01-18. <https://f1000research.com/articles/5-885/v2>.



<sup>1</sup>International Center for Tropical Agriculture (CIAT), Cali, Colombia  
<sup>2</sup>Independent Researcher, Miami, USA  
<sup>3</sup>International Center for Agricultural Research in the Dry Areas (ICARDA), Beirut, Lebanon  
<sup>4</sup>Australian Centre for International Agricultural Research, Canberra, Australia  
<sup>5</sup>International Institute of Tropical Agriculture (IITA), Ibadan, Nigeria  
<sup>6</sup>International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), Patancheru, India