

Technical University of Denmark



Thermophoretic transport of water nanodroplets coned in carbon nanotubes: the role of friction

Oyarzua, Elton; Walther, Jens Honore; Zambrano, Harvey

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):

Oyarzua, E., Walther, J. H., & Zambrano, H. (2017). Thermophoretic transport of water nanodroplets coned in carbon nanotubes: the role of friction. Abstract from 70th Annual Meeting of the American Physical Society Division of Fluid Dynamics (DFD17), Denver, United States.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Abstract Submitted
for the DFD17 Meeting of
The American Physical Society

Thermophoretic transport of water nanodroplets confined in carbon nanotubes: the role of friction.¹ ELTON OYARZUA, Universidad de Concepcion, JENS H. WALTHER, Technical University of Denmark, HARVEY A. ZAMBRANO, Universidad de Concepcion — The development of efficient nanofluidic devices requires driving mechanisms that provide controlled transport of fluids through nanoconduits. Temperature gradients have been proposed as a mechanism to drive particles, fullerenes and nanodroplets inside carbon nanotubes (CNTs). In this work, molecular dynamics (MD) simulations are conducted to study thermophoresis of water nanodroplets inside CNTs. To gain insight into the interplay between the thermophoretic force acting on the droplet and the retarding liquid-solid friction, sets of constrained and unconstrained MD simulations are conducted. The results indicate that the thermophoretic motion of a nanodroplet displays two kinetic regimes: an initial regime characterized by a decreasing acceleration and afterwards a terminal regime with constant velocity. During the initial regime, the magnitude of the friction force increases linearly with the droplet velocity whereas the thermophoretic force has a constant magnitude defined by the magnitude of the thermal gradient and the droplet size. Subsequently, in the terminal regime, the droplet moves at constant velocity due to a dynamic balance between the thermophoretic force and the retarding friction force.

¹We acknowledge partial support from CONICYT (Chile) under scholarship No. 21140427.

Elton Oyarzua
Universidad de Concepcion

Date submitted: 06 Jul 2017

Electronic form version 1.4