

Technical University of Denmark



Fungi & Health: can polysaccharides from the fungus inonotus obliquus (CHAGA) inhibit tumor growth?

Wold, C. W. ; Corthay, A. ; Kjeldsen, Christian; Duus, Jens Øllgaard; Christensen, B. E.; Inngjerdingen, K. T.

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Wold, C. W., Corthay, A., Kjeldsen, C., Duus, J. Ø., Christensen, B. E., & Inngjerdingen, K. T. (2017). Fungi & Health: can polysaccharides from the fungus inonotus obliquus (CHAGA) inhibit tumor growth?. Abstract from 19th European Carbohydrate Symposium, Barcelona, Spain.

DTU Library

Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



FUNGI & HEALTH: CAN POLYSACCHARIDES FROM THE FUNGUS *INONOTUS OBLIQUUS* (CHAGA) INHIBIT TUMOR GROWTH?

Wold C W^a, Corthay A^b, Kjeldsen C^c, Duus J Ø^c, Christensen B E^d, Inngjerdningen K T^a

^a Department of Pharmaceutical Chemistry, School of Pharmacy, University of Oslo, Norway

^b Tumor Immunology Group, Rikshospitalet, Oslo University Hospital, Norway

^c Department of Chemistry, Organic Chemistry, Technical University of Denmark, Denmark

^d Department of Biotechnology and Food Science, Norwegian University of Science and Technology, Norway
e-mail: c.w.wold@farmasi.uio.no

Inonotus obliquus (Chaga) – a white rot fungus found on birch trees in the northern hemisphere – has been used in traditional medicine in Europe and Asia for centuries [1]. Native peoples have made use of Chaga by brewing it as a tea to treat gastro-intestinal problems, to heal wounds and even to treat cancer. The last few decades, studies have found Chaga to contain biologically active substances such as polysaccharides, triterpenoids, polyphenols and melanin [2]. *In vivo* effects such as tumor growth inhibition have been observed in mice receiving various Chaga extracts [2,3]. The main hypothesis behind the tumor inhibiting effect is two-fold: i) fungal polysaccharides may inhibit tumor growth indirectly by activating certain immune cells such as macrophages and ii) triterpenoids and other steroids from Chaga may give a direct cytotoxic effect against cancer cells [3,4]. While triterpenoids from Chaga have been extensively characterized, detailed analysis of the polysaccharides is lacking. The present work has aimed to isolate and characterize the polysaccharides in Chaga, by e.g. column chromatography (ion-exchange/gel filtration), GC-MS, SEC-MALLS and extensive NMR analysis. The water-soluble polysaccharides were found to be complex hetero-polysaccharides, with a structure dominated by (1→3/1→6)-β-glucan and (1→6)-α-galactan, with β-xylose, α-mannose and α-galacturonic acid present in significant amounts. 3-O-methyl α-galactose was reported in Chaga for the first time. The polysaccharide fractions obtained were screened in *in vitro* bioassays for their potential as immunomodulators. Several fractions showed promising results by activating murine bone-marrow derived macrophages to inhibit the growth of Lewis lung carcinoma cells *in vitro*. The results suggest further studies to be conducted on immune cell activation and *in vivo* tumor growth inhibition.

References:

- [1] Babitskaya V et al. Melanin Complex from Medicinal Mushroom *Inonotus obliquus* (Pers.:Fr.) Pilát (Chaga), *Int J Med Mushrooms* **2002**, *4*: 139-145.
- [2] Zheng W. et al., Chemical diversity of biologically active metabolites in the sclerotia of *Inonotus obliquus* and submerged culture strategies for up-regulating their production, *Appl Microbiol Biotechnol* **2010**, *87*: 1237-1254.
- [3] Chen Y. et al., Purification, characterization and biological activity of a novel polysaccharide from *Inonotus obliquus*, *Int J Biol Macromol* **2015**, *79*: 587-594.
- [4] Zhao F. et al., Triterpenoids from *Inonotus obliquus* and their antitumor activities, *Fitoterapia* **2015**, *101*: 34-40.