

gram(s) used to refine structure: *SHELXTL*. Molecular graphics: *SHELXTL*. Software used to prepare material for publication: *SHELXTL*.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and complete geometry have been deposited with the IUCr (Reference: HA1178). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, S1–S19.
- Howard, J. A. K., Hoy, V. J., O'Hagan, D. & Smith, G. M. (1996). *Tetrahedron*, **52**, 12613–12622.
- Murray-Rust, P., Stallings, W. C., Monti, C. T., Preston, R. K. & Glusker, J. P. (1983). *J. Am. Chem. Soc.* **105**, 3206–3214.
- Sheldrick, G. M. (1995). *SHELXTLS.04/VMS. An Integrated System for Solving, Refining and Displaying Crystal Structures from Diffraction Data*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Shimoni, L. & Glusker, J. P. (1994). *Struct. Chem.* **5**, 383–397.
- Siemens (1996a). *SMART. Data Collection Software*. Version 4.050. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Siemens (1996b). *SAINT. Data Reduction Software*. Version 4.050. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.

Acta Cryst. (1997). **C53**, 126–128

6-Oxo-6-phenyl-6-phospha-3,9-dithiabicyclo[9.4.0]pentadeca-1(11),12,14-triene

RAIKKO KIVEKÄS,^a JOSÉ ANTONIO MUÑOZ,^b LUIS ESCRICHE,^b JAUME CASABÓ^b AND REIJO SILLANPÄÄ^c

^aDepartment of Chemistry, University of Helsinki, PO Box 55, FIN-00014, Finland, ^bDepartament de Química, Universitat Autònoma de Barcelona, 08193 Bellaterra, Barcelona, Spain, and ^cDepartment of Chemistry, University of Turku, FIN-20014 Turku, Finland. E-mail: sillanpa@sara.uu.fi

(Received 8 July 1996; accepted 30 September 1996)

Abstract


The title molecule, 6-phenyl-6-phospha-3,9-dithiabicyclo[9.4.0]pentadeca-1(11),12,14-triene 6-oxide, C₁₈H₂₁OPS₂, has an exdentate conformation, with the S and P atoms oriented away from the centre of the macrocyclic cavity.

Comment

Tricoordinated P atom-containing macrocycles are of potential use as ligands in catalysis and ion-selective

transport (Caminade & Majoral, 1994). It has been reported that the introduction of S donor atoms into these systems could modulate their complexing abilities towards transition metal ions (Kyba, Clubb, Larson, Schueler & Davis, 1985). These mixed macrocycles containing phosphorus and sulfur can be easily transformed into their phosphane or sulfoxide equivalents which are useful in molecular recognition and the complexation of anions (Izatt, Pawlak, Brandshaw & Bruening, 1991).

We have reported recently that S₂P(=O)-containing macrocycles show good selectivity properties as neutral carriers in ion-selective electrodes (Casabó *et al.*, 1996). In this context and in order to clarify the relationships between the molecular conformations and selectivity properties, we report here the synthesis of the S₂P-containing macrocycle 6-phenyl-6-phospha-3,9-dithiabicyclo[9.4.0]pentadeca-1(11),12,14-triene, (1), and its phosphine oxide derivative, (2), together with the crystal structure determination of the latter.



The bond lengths and angles in compound (2) (Fig. 1) are normal. A comparison of the torsion angles of (2) with those of the 6-S derivative reveals only minor differences (at most 10°) and thus, the conformations of the two compounds are essentially similar (Lockhart *et al.*, 1992). Also, the intramolecular distances between the heavy atoms are almost the same. The longest S...S distance is 5.795 (1) Å in (2) and 5.772 (1) Å in the 6-S




Fig. 1. The molecular structure of the title compound showing the atomic numbering scheme. Displacement ellipsoids are drawn at the 30% probability level.

compound. The S···P distances in (2) are 4.342 (1) and 4.474 (1) Å, while the comparable short S···S distances in the 6-S derivative are 4.398 (1) and 4.421 (1) Å, respectively. These results indicate that the influence of the S and P=O groups on the conformations of these macrocycles are almost the same.

The title compound and the corresponding 6-O substituted compound display considerable differences in their ring conformations (Kivekäs *et al.*, 1994), for example, in the latter compound, the S···S distance is only 4.501 (1) Å compared with the long S···S distance of 5.795 (1) Å in (2).

The crystal packing shows two C—H···O hydrogen bonds, which are from H9A and H11A to the O atom. The C9···Oⁱ and C11···Oⁱ [symmetry code: (i) $x - 1, y, z$] distances are 3.329 (4) and 3.293 (5) Å, respectively. The C—H···O angles are almost linear, C9—H9A···Oⁱ being 173 (3) and C11—H11A···Oⁱ being 175 (3)°.

Experimental

Dehydrated and deoxygenated THF was used in the synthesis. Dilithium 3-phenyl-3-phosphapentane-1,5-dithiolate was obtained as described by Muñoz, Escriche, Casabó, Kivekäs & Sillanpää (1996). The macrocycle (1) was synthesized, under high dilution conditions, by mixing a solution of dilithium 3-phenyl-3-phosphapentane-1,5-dithiolate (4.40 g, 18.1 mmol) in THF (100 ml) with a solution of α, α' -dichloro-*o*-xylene (3.17 g, 18.1 mmol) in THF (100 ml) over THF (500 ml) at a rate of 20 ml h⁻¹. After addition was complete, the reaction mixture was stirred at room temperature for 48 h and then evaporated to dryness. The resulting oily material was treated with diethyl ether (50 ml), affording a white solid (LiCl) which was filtered off. The ether solution was then evaporated to 25 ml and cyclohexane (20 ml) was added. A white solid of (1) appeared which was filtered off and vacuum dried. Yield: 1.40 g (23%). Analysis calculated for C₁₈H₂₁PS₂: C 65.03, H 6.37, S 19.29%; found: C 64.7, H 6.1, S 18.9%. ¹H NMR (400 MHz, CDCl₃): δ 1.61 (*t*, 4H, ³J_{HH} = 7.5 Hz, SCH₂), 2.66 (*m*, 4H, CCH₂P), 3.85 (*d*, 2H, ³J_{HH} = 14.4 Hz, SCH₂Ph), 4.00 (*d*, 2H, ³J_{HH} = 14.4 Hz, SCH₂Ph), 7.25–7.50 (*m*, 9H, Ph). ¹³C{¹H} NMR (100 MHz, CDCl₃): δ 27.47, 27.72, 32.26 (PhCH₂SCH₂CH₂P), 127.49, 128.21, 128.39, 130.55, 131.37, 136.93, 138.33 (Ph). ³¹P{¹H} NMR (101 MHz, CDCl₃): δ -21.49. The macrocycle (1) was dissolved in dichloromethane and this solution was left in an open vessel for 48 h. After this time, colourless prismatic crystals of 6-oxo-6-phenyl-6-phospha-3,9-dithiabicyclo[9.4.0]-pentadeca-1(11),12,14-triene, (2), appeared. This material was suitable for X-ray diffraction analysis. Analysis calculated for C₁₈H₂₁OPS₂: C 61.98, H 6.02, S 18.36%; found: C 61.5, H 6.0, S 18.3%. ¹H NMR (400 MHz, CDCl₃): δ 2.01 (*m*, 4H), 2.64 (*m*, 4H), 3.92 (*d*, ²J_{HH} = 15 Hz, 2H), 4.10 (*d*, ²J_{HH} = 15 Hz), 7.3 (*m*, 9H, Ph). ¹³C{¹H} NMR (100 MHz, CDCl₃): δ 27.26 (*d*, ¹J_{PC} = 5.98 Hz) 27.87, (*d*, ²J_{PC} = 7.84 Hz), 32.34 (*s*), 127–138 (Ph). ³¹P{¹H} NMR (101 MHz, CDCl₃): δ 41.

Crystal data

C₁₈H₂₁OPS₂
M_r = 348.44

Mo K α radiation
 λ = 0.71069 Å

Monoclinic
P2₁/c
a = 5.644 (2) Å
b = 13.624 (2) Å
c = 22.617 (1) Å
 β = 95.31 (1)°
V = 1731.6 (7) Å³
Z = 4
D_x = 1.337 Mg m⁻³
D_m not measured

Data collection

Rigaku AFC-5S diffractometer
 ω -2 θ scans
Absorption correction:
empirical *via* ψ scans
(North, Phillips & Mathews, 1968)
T_{min} = 0.880, T_{max} = 0.916
3622 measured reflections
3544 independent reflections

Refinement

Refinement on F²
R(F) = 0.0425
wR(F²) = 0.1620
S = 1.041
3034 reflections
262 parameters
w = 1/[\sigma²(F_o²) + 1.1268P]
where P = (F_o² + 2F_c²)/3
(Δ/σ)_{max} = -0.002

Cell parameters from 25 reflections
 θ = 13.2–17.6°
 μ = 0.399 mm⁻¹
T = 293 (2) K
Prism
0.38 × 0.30 × 0.22 mm
Colourless

1960 observed reflections
[I > 2 σ (I)]
R_{int} = 0.0370
 θ _{max} = 25°
h = 0 → 7
k = 0 → 16
l = -27 → 27
3 standard reflections monitored every 150 reflections
intensity decay: none

$\Delta\rho_{max}$ = 0.264 e Å⁻³
 $\Delta\rho_{min}$ = -0.296 e Å⁻³
Extinction correction: none
Atomic scattering factors
from *International Tables for Crystallography* (1992, Vol. C, Tables 4.2.6.8 and 6.1.1.4)

Table 1. Fractional atomic coordinates and equivalent isotropic displacement parameters (Å²)

$$U_{eq} = (1/3)\sum_i \sum_j U_{ij} a_i^* a_j^* \mathbf{a}_i \cdot \mathbf{a}_j$$

	x	y	z	U _{eq}
S1	0.1671 (2)	0.86625 (7)	0.50311 (4)	0.0533 (3)
S2	-0.0123 (2)	0.73363 (8)	0.25963 (4)	0.0559 (3)
P	0.49263 (15)	0.62949 (7)	0.41079 (4)	0.0389 (2)
O	0.7520 (4)	0.6439 (2)	0.42218 (11)	0.0598 (7)
C1	0.1382 (6)	0.9316 (2)	0.3840 (2)	0.0433 (9)
C2	0.0863 (6)	0.9092 (3)	0.3236 (2)	0.0467 (9)
C3	0.2187 (8)	0.9546 (3)	0.2825 (2)	0.0610 (11)
C4	0.3988 (9)	1.0192 (3)	0.2992 (2)	0.0687 (13)
C5	0.4504 (8)	1.0417 (3)	0.3580 (2)	0.0617 (12)
C6	0.3206 (7)	0.9984 (3)	0.3997 (2)	0.0512 (10)
C7	0.0022 (7)	0.8858 (3)	0.4316 (2)	0.0503 (10)
C8	0.4005 (6)	0.7870 (3)	0.4814 (2)	0.0464 (9)
C9	0.3231 (6)	0.6814 (3)	0.4672 (2)	0.0386 (8)
C10	0.3762 (6)	0.6878 (3)	0.3421 (2)	0.0412 (8)
C11	0.1250 (7)	0.6576 (3)	0.3188 (2)	0.0480 (9)
C12	-0.1058 (7)	0.8387 (3)	0.3004 (2)	0.0636 (12)
C13	0.4085 (6)	0.5025 (2)	0.40216 (15)	0.0397 (8)
C14	0.5488 (7)	0.4426 (3)	0.3703 (2)	0.0530 (10)
C15	0.4893 (9)	0.3455 (3)	0.3594 (2)	0.0633 (12)
C16	0.2912 (8)	0.3067 (3)	0.3804 (2)	0.0576 (11)
C17	0.1547 (7)	0.3638 (3)	0.4134 (2)	0.0602 (11)
C18	0.2097 (7)	0.4614 (3)	0.4236 (2)	0.0566 (11)

Table 2. Selected geometric parameters (Å, °)

S1—C8	1.806 (4)	P—O	1.476 (3)
S1—C7	1.809 (4)	P—C13	1.800 (3)
S2—C12	1.808 (4)	P—C9	1.808 (3)
S2—C11	1.808 (4)	P—C10	1.813 (4)

C8—S1—C7	100.0 (2)	C1—C7—S1	116.3 (3)
C12—S2—C11	101.6 (2)	C9—C8—S1	114.7 (3)
O—P—C13	113.20 (15)	C8—C9—P	110.9 (3)
O—P—C9	114.0 (2)	C11—C10—P	115.5 (3)
C13—P—C9	107.5 (2)	C10—C11—S2	115.1 (3)
O—P—C10	111.5 (2)	C2—C12—S2	116.6 (3)
C13—P—C10	104.9 (2)	C18—C13—P	124.5 (3)
C9—P—C10	105.0 (2)	C14—C13—P	117.7 (3)
C7—C1—C2—C12	-0.1 (5)	C9—P—C10—C11	-70.0 (3)
C2—C1—C7—S1	148.3 (3)	P—C10—C11—S2	167.7 (2)
C8—S1—C7—C1	-59.0 (3)	C12—S2—C11—C10	-79.0 (3)
C7—S1—C8—C9	-74.7 (3)	C1—C2—C12—S2	-120.3 (4)
S1—C8—C9—P	148.5 (2)	C11—S2—C12—C2	75.0 (4)
C10—P—C9—C8	-73.9 (3)		

H atoms were located from $\Delta\rho$ maps and refined with U_{eq} fixed to $1.2U_{eq}$ of the host atom.

Data collection: *MSCIAFC Diffractometer Control Software* (Molecular Structure Corporation, 1995). Cell refinement: *MSCIAFC Diffractometer Control Software*. Data reduction: *TEXSAN* (Molecular Structure Corporation, 1989). Program(s) used to solve structure: *SHELXS86* (Sheldrick, 1985). Program(s) used to refine structure: *SHELXL93* (Sheldrick, 1993). Molecular graphics: *ORTEPII* (Johnson, 1976). Software used to prepare material for publication: *SHELXL93*.

This work is partly financed by Comision Interministerial de Ciencia y Tecnologia from the Spanish Government through project MAT94-0668. RK thanks Ministerio de Educacion y Ciencia of the Spanish Government for grant SAB95-0249 and the Oskar Öflund's Foundation for financial support.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and complete geometry have been deposited with the IUCr (Reference: AB1413). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Caminade, A. M. & Majoral, J. P. (1994). *Chem. Rev.* **94**, 1183–1213.
- Casabó, J., Escriche, L., Pérez-Jiménez, C., Muñoz, J. A., Teixidor, F., Baucells, J. & Errachid, A. (1996). *Anal. Chim. Acta*, **320**, 63–68.
- Izatt, R. M., Pawlak, K., Brandshaw, J. S. & Bruening, R. L. (1991). *Chem. Rev.* **91**, 1721–2085.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Kivekäs, R., Sillanpää, R., Escriche, L., Almajano, M. P., Teixidor, F. & Casabó, J. (1994). *Acta Cryst.* **C50**, 2047–2049.
- Kyba, E. P., Clubb, C. N., Larson, S. B., Schueler, V. J. & Davis, R. E. (1985). *J. Am. Chem. Soc.* **107**, 2141–2148.
- Lockhart, J. C., Mousley, D. P., Stuart Hill, M. N., Tomkinson, N. P., Teixidor, F., Almajano, M. P., Escriche, L., Casabó, J. F., Sillanpää, R. & Kivekäs, R. (1992). *J. Chem. Soc. Dalton Trans.* pp. 2889–2897.
- Molecular Structure Corporation (1989). *TEXSAN. Single Crystal Structure Analysis Software*. Version 5.0. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
- Molecular Structure Corporation (1995). *MSCIAFC Diffractometer Control Software*. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
- Muñoz, J. A., Escriche, L., Casabó, J., Kivekäs, R. & Sillanpää, R. (1996). *Inorg. Chim. Acta*. In the press.
- North, A. C. T., Phillips, D. C. & Mathews, F. C. (1968). *Acta Cryst.* **A24**, 351–359.

Sheldrick, G. M. (1985). *SHELXS86. Program for the Solution of Crystal Structures*. University of Göttingen, Germany.

Sheldrick, G. M. (1993). *SHELXL93. Program for the Refinement of Crystal Structures*. University of Göttingen, Germany.

Acta Cryst. (1997). **C53**, 128–130

Diguanidinium Hexafluorosilicate

ALICJA WAŚKOWSKA

*Institute of Low Temperature and Structure Research,
Polish Academy of Sciences, Okolna str. 2, 50 950 Wrocław,
Poland. E-mail: waskowsk@highscreen.int.pan.wroc.pl*

(Received 23 April 1996; accepted 20 September 1996)

Abstract

The crystal structure of diguanidinium hexafluorosilicate, [C(NH₂)₃]₂SiF₆, consists of regular octahedral SiF₆²⁻ anions linked through a system of hydrogen bonds to the guanidinium cations, which have a planar conformation. The striking feature of the structure is the high degree of hydrogen bonding. All H atoms of the guanidinium units are involved in hydrogen bonds of the type N—H...F of length 2.901 (2), 2.908 (2) and 2.891 (1) Å. Unlike other materials belonging to the family of compounds of general formula R_x(NH_{4-x})₂MX₃, the crystals of the present compound do not display ferroelectric/ferroelastic properties.

Comment

This paper is a continuation of earlier reports on structural studies of the complex compounds of metal halogenides with various N-atom containing organic ligands (e.g. Morimoto & Lingafelter, 1970; Adams & Small, 1974; Waśkowska, Czaplá & Janczak, 1993). The general formula is R_x(NH_{4-x})₂MX₃ and (R_xNH_{4-x})₂MX₄, where R is an aliphatic or aromatic hydrocarbon, M is a metal atom and X is Cl, Br, I or F. Most of these compounds show phase transitions of ferroelectric or ferroelastic character which are related to the dynamics of the constituent organic or inorganic subunits (Doudin & Chapuis, 1988; Miyazaki, Irokawa, Komukae, Osaka & Makita, 1991; Katusiak & Szafranski, 1994). The title compound could be a candidate for behaviour of this type, but our specific heat measurements by differential scanning calorimetry (DSC), performed in the temperature range 106–410 K, showed no anomalies which could be related to a phase transition. The X-ray diffraction study of the title crystal structure, (I), has been undertaken in order to try to provide a structural explanation for the distinct physical properties of this