



# Phase retrieval with random Gaussian sensing vectors by alternating projections

Irène Waldspurger

## ► To cite this version:

Irène Waldspurger. Phase retrieval with random Gaussian sensing vectors by alternating projections. IEEE Transactions on Information Theory, Institute of Electrical and Electronics Engineers, In press. hal-01645081

HAL Id: hal-01645081

<https://hal.archives-ouvertes.fr/hal-01645081>

Submitted on 22 Nov 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Phase retrieval with random Gaussian sensing vectors by alternating projections

Irène Waldspurger

**Abstract**—We consider a phase retrieval problem, where we want to reconstruct a  $n$ -dimensional vector from its phaseless scalar products with  $m$  sensing vectors, independently sampled from complex normal distributions. We show that, with a suitable initialization procedure, the classical algorithm of alternating projections (Gerchberg-Saxton) succeeds with high probability when  $m \geq Cn$ , for some  $C > 0$ . We conjecture that this result is still true when no special initialization procedure is used, and present numerical experiments that support this conjecture.

## I. INTRODUCTION

A *phase retrieval* problem consists in recovering an unknown vector  $x_0 \in \mathbb{C}^n$  from  $m$  phaseless linear measurements, of the form

$$b_k = |\langle a_k, x_0 \rangle|, \quad k = 1, \dots, m.$$

Such problems naturally appear in various applications, notably in optical imaging [1]. A lot of efforts have thus been made to design efficient algorithms that could numerically solve these problems.

The oldest reconstruction algorithms [2], [3] were iterative: they started from a random initial guess of  $x_0$ , and tried to iteratively refine it by various heuristics. These methods sometimes succeed, but can also fail to converge towards the correct solution: they may get stuck in *stagnation points*, whose existence is due to the non-convexity of the problem. When they converge and when they do not is not clearly understood.

To overcome convergence problems, convexification methods have been introduced [4], [5]. Their principle is to lift the non-convex problem to a matricial space, where it can be approximated by a convex problem. These methods provably reconstruct the unknown vector  $x_0$  with high probability if the sensing vectors  $a_k$  are “random enough” [6]–[8]. Numerical experiments show that they also perform well on more structured, non-random phase retrieval problems [9], [10].

Unfortunately, this good precision comes at a high computational cost: optimizing an  $n \times n$  matrix is much slower than directly reconstructing a  $n$ -dimensional vector. More recently, a new family of algorithms has thus been developed, which enjoy similar theoretical guarantees as convexified methods, but have a much smaller computational complexity. The algorithms of this family rely on the following two-step scheme:

- (1) an initialization step, that returns a point close to the solution;

- (2) a gradient descent (with possible additional refinements) over a well-chosen non-convex cost function.

The intuitive reason why this scheme works is that the cost function, although globally non-convex, enjoys some good geometrical property in a neighborhood of the solution (like convexity or a weak form of it [11]). When the initial point belongs to this neighborhood, the gradient descent then converges to the correct solution.

A preliminary form of this scheme appeared in [12], with an alternating minimization in step (2) instead of a gradient descent. [13] then proved, for a specific cost function, that this two-step scheme was able to exactly reconstruct the unknown  $x_0$  with high probability, in the regime  $m = O(n \log n)$ , if the sensing vectors were independent and Gaussian. In [14], [15], the same result was shown in the regime  $m = O(n)$  for a slightly different cost function, with additional truncation steps. In [16], [17], it was extended to a different, non-smooth, cost function.

These new methods enjoy much stronger theoretical guarantees than “traditional” algorithms. However, it is not clear whether they really perform better in applications, or whether they actually behave similarly, and are simply easier to theoretically study. Traditional algorithms are well-known, simple and very easy to implement; understanding how they compare to more modern methods is of much value for applications.

In this article, we take a first step towards this goal, by considering the very classical alternating projections algorithm, introduced by Gerchberg and Saxton in [2], arguably the most simple and widely used method for phase retrieval. We show that, in the setting where sensing vectors are independent and Gaussian, it performs as well as gradient descent over a suitable cost function: it converges linearly to the true solution with high probability, provided that it is correctly initialized.

**Theorem** (See Corollary III.8). *There exist absolute constants  $C_1, C_2, M > 0, \delta \in ]0; 1[$  such that, if  $m \geq Mn$ , if the sensing vectors are independently chosen according to complex normal distributions, and if the alternating projections are carefully initialized, for example with the method described in [14], then the sequence of iterates  $(z_t)_{t \in \mathbb{N}}$  they produce satisfies*

$$\forall t \in \mathbb{N}^*, \quad \inf_{\phi \in \mathbb{R}} \|e^{i\phi} x_0 - z_t\| \leq \delta^t \|x_0\|,$$

with probability at least

$$1 - C_1 \exp(-C_2 m).$$

Several authors have already tried to establish properties of this kind, but, compared to ours, their results were significantly suboptimal in various respects. Using transversality arguments,

[18], [19] have shown the local convergence of alternating projections for relatively general families of sensing vectors. Unfortunately, transversality arguments give no control on the convergence radius of the algorithm, which can be extremely small. Lower bounding the radius requires using the statistical properties of the sensing vectors. This was first attempted in [12]. The result obtained by these authors was very similar to ours, but required resampling the sensing vectors at each step of the algorithm, an operation that is almost never done in practice. For a non resampled version, a preliminary result was given in [20], but this result did not capture the correct scaling of the convergence radius as a function of  $m$  and  $n$ . As a consequence, it only established global convergence to the correct solution for a suboptimal number of measurements ( $m = O(n \log^2 n)$ ), and with a complex initialization procedure.

To have theory and practice exactly coincide, the role of the initialization procedure should also be examined: in applications, alternating projections are often used with a random initialization, and not with a carefully chosen one. Our numerical experiments indicate that removing the careful initialization does not significantly alter the convergence of the algorithm (still in the setting of Gaussian independent sensing vectors). This fact is related to the observations of [21], who proved that, at least in the regime  $m \geq O(n \log^3 n)$  and for a specific cost function, the initialization part of the two-step scheme is not necessary in order for the algorithm to converge. In the context of alternating projections, we were however not able to prove a similar result<sup>1</sup>, and leave it as a conjecture.

**Conjecture** (See Conjecture IV.1). *Let any  $\epsilon > 0$  be fixed. When  $m \geq Cn$ , for  $C > 0$  large enough, alternating projections, starting from a random initialization (chosen according to a rotationally invariant distribution), converge to the true solution with probability at least  $1 - \epsilon$ .*

The article is organized as follows. Section II precisely defines phase retrieval problems and the alternating projections algorithm. Section III states and proves the main result: the global convergence of alternating projections, with proper initialization, for  $m = O(n)$  independent Gaussian measurements. Section IV contains numerical experiments, and presents our conjecture about the non-necessity of the initialization procedure.

This article only considers the most simple setting, where sensing vectors are independent and Gaussian, and measurements are not noisy. We made this choice in order to keep the technical content simple, but we hope that our results extend to more realistic settings, and future work should examine this issue.

### A. Notations

For any  $z \in \mathbb{C}$ ,  $|z|$  is the modulus of  $z$ . We extend this notation to vectors: if  $z \in \mathbb{C}^k$  for some  $k \in \mathbb{N}^*$ , then  $|z|$  is the

<sup>1</sup>The proof technique used by [21] collapses when gradient descent is replaced with alternating projections, as detailed in a companion study of this article, available on arXiv [22].

vector of  $(\mathbb{R}^+)^k$  such that

$$|z|_i = |z_i|, \quad \forall i = 1, \dots, k.$$

For any  $z \in \mathbb{C}$ , we set  $E_{\text{phase}}(z)$  to be the following subset of  $\mathbb{C}$ :

$$E_{\text{phase}}(z) = \begin{cases} \left\{ \frac{z}{|z|} \right\} & \text{if } z \in \mathbb{C} - \{0\}; \\ \{e^{i\phi}, \phi \in \mathbb{R}\} & \text{if } z = 0. \end{cases}$$

We extend this definition to vectors  $z \in \mathbb{C}^k$ :

$$E_{\text{phase}}(z) = \prod_{i=1}^k E_{\text{phase}}(z_i).$$

For any  $z \in \mathbb{C}$ , we define  $\text{phase}(z)$  by

$$\text{phase}(z) = \begin{cases} \frac{z}{|z|} & \text{if } z \in \mathbb{C} - \{0\}; \\ 1 & \text{if } z = 0, \end{cases}$$

and extend this definition to vectors  $z \in \mathbb{C}^k$ , as for the modulus.

We denote by  $\odot$  the pointwise product of vectors: for all  $a, b \in \mathbb{C}^k$ ,  $(a \odot b)$  is the vector of  $\mathbb{C}^k$  such that

$$(a \odot b)_i = a_i b_i, \quad \forall i = 1, \dots, k.$$

We define the operator norm of any matrix  $A \in \mathbb{C}^{n_1 \times n_2}$  by

$$\|A\| = \sup_{v \in \mathbb{C}^{n_2}, \|v\|=1} \|Av\|.$$

We denote by  $A^\dagger$  its Moore-Penrose pseudo-inverse. We note that  $AA^\dagger$  is the orthogonal projection onto  $\text{Range}(A)$ .

## II. PROBLEM SETUP

### A. Phase retrieval problem

Let  $n, m$  be positive integers. The goal of a phase retrieval problem is to reconstruct an unknown vector  $x_0 \in \mathbb{C}^n$  from  $m$  measurements with a specific form.

We assume  $a_1, \dots, a_m \in \mathbb{C}^n$  are given; they are called the *sensing vectors*. We define a matrix  $A \in \mathbb{C}^{m \times n}$  by

$$A = \begin{pmatrix} a_1^* \\ \vdots \\ a_m^* \end{pmatrix}.$$

This matrix is called the *measurement matrix*. The associated *phase retrieval problem* is:

$$\text{reconstruct } x_0 \text{ from } b \stackrel{\text{def}}{=} |Ax_0|. \quad (1)$$

As the modulus is invariant to multiplication by unitary complex numbers, we can never hope to reconstruct  $x_0$  better than *up to multiplication by a global phase*. So, instead of exactly reconstructing  $x_0$ , we want to reconstruct  $x_1$  such that

$$x_1 = e^{i\phi} x_0, \quad \text{for some } \phi \in \mathbb{R}.$$

In all this article, we assume the sensing vectors to be independent realizations of centered Gaussian variables with identity covariance:

$$(a_i)_j \sim \mathcal{N}\left(0, \frac{1}{2}\right) + \mathcal{N}\left(0, \frac{1}{2}\right) i, \quad \forall 1 \leq i \leq m, 1 \leq j \leq n. \quad (2)$$

The measurement matrix is in particular independent from  $x_0$ .

[23] and [24] have proved that when  $m \geq 4n - 4$ , Problem (1) always has a unique solution, up to a global phase, for all measurement matrices  $A$  belonging to some open dense subset of  $\mathbb{C}^{m \times n}$ . So when  $A$  is chosen according to a probability density that is absolutely continuous with respect to Lebesgue measure, Problem (1) has a unique solution with probability 1 (because the complement of an open dense subset has zero Lebesgue measure). In particular, with our measurement model (2), the reconstruction is guaranteed to be unique, with probability 1, when  $m \geq 4n - 4$ .

### B. Alternating projections

The alternating projections method has been introduced for phase retrieval problems by [2]. It focuses on the reconstruction of  $Ax_0$ ; if  $A$  is injective, this then allows to recover  $x_0$ .

To reconstruct  $Ax_0$ , it is enough to find  $z \in \mathbb{C}^m$  in the intersection of the following two sets.

- (1)  $z \in \{z' \in \mathbb{C}^m, |z'| = b\}$ ;
- (2)  $z \in \text{Range}(A)$ .

Indeed, when the solution to Problem (1) is unique,  $Ax_0$  is the only element of  $\mathbb{C}^m$  that simultaneously satisfies these two conditions (up to a global phase).

A natural heuristic to find such a  $z$  is to pick any initial guess  $z_0$ , then to alternatively project it on the two constraint sets. In this context, we call *projection* on a closed set  $E \subset \mathbb{C}^m$  a function  $P : \mathbb{C}^m \rightarrow E$  such that, for any  $x \in \mathbb{C}^m$ ,

$$\|x - P(x)\| = \inf_{e \in E} \|x - e\|.$$

The two sets defining constraints (1) and (2) admit projections with simple analytical expressions, which leads to the following formulas:

$$y'_k = b \odot \text{phase}(y_k); \quad (\text{Projection onto set (1)}) \quad (3a)$$

$$y_{k+1} = (AA^\dagger)y'_k. \quad (\text{Projection onto set (2)}) \quad (3b)$$

If, for each  $k$ , we define  $z_k$  as the vector such that  $y_k = Az_k$ <sup>2</sup>, an equivalent form of these equations is:

$$z_{k+1} = A^\dagger(b \odot \text{phase}(Az_k)).$$

The hope is that the sequence  $(y_k)_{k \in \mathbb{N}}$  converges towards  $Ax_0$ . Unfortunately, it can get stuck in *stagnation points*. The following proposition (proven in Appendix A) characterizes these stagnation points.

**Proposition II.1.** *For any  $y_0$ , the sequence  $(y_k)_{k \in \mathbb{N}}$  is bounded. Any accumulation point  $y_\infty$  of  $(y_k)_{k \in \mathbb{N}}$  satisfies the following property:*

$$\exists u \in E_{\text{phase}(y_\infty)}, \quad (AA^\dagger)(b \odot u) = y_\infty.$$

In particular, if  $y_\infty$  has no zero entry,

$$(AA^\dagger)(b \odot \text{phase}(y_\infty)) = y_\infty.$$

Despite the relative simplicity of this characteristic property, it is extremely difficult to exactly compute the stagnation points, determine their attraction basin or avoid them when the algorithm happens to run into them.

<sup>2</sup>which exists and is unique, because  $y_k$  belongs to  $\text{Range}(A)$  and  $A$  is injective with probability 1

## III. ALTERNATING PROJECTIONS WITH GOOD INITIALIZATION

In this section, we prove our main result: for  $m = O(n)$  Gaussian independent sensing vectors, the method of alternating projections converges to the correct solution with high probability, if it is carefully initialized.

### A. Local convergence of alternating projections

We begin with a key result, that we will need to establish our statement. This result is a local contraction property of the alternating projections operator  $x \rightarrow A^\dagger(b \odot \text{phase}(Ax))$ .

**Theorem III.1.** *There exist  $\epsilon, C_1, C_2, M > 0$ , and  $\delta \in ]0; 1[$  such that, if  $m \geq Mn$ , then, with probability at least*

$$1 - C_1 \exp(-C_2 m),$$

*the realizations of  $A$  and  $b = |Ax_0|$  are such that the following property holds: for any  $x \in \mathbb{C}^n$  such that*

$$\inf_{\phi \in \mathbb{R}} \|e^{i\phi} x_0 - x\| \leq \epsilon \|x_0\|,$$

*we have*

$$\inf_{\phi \in \mathbb{R}} \|e^{i\phi} x_0 - A^\dagger(b \odot \text{phase}(Ax))\| \leq \delta \inf_{\phi \in \mathbb{R}} \|e^{i\phi} x_0 - x\|. \quad (4)$$

*Proof.* For any  $x \in \mathbb{C}^n$ , we can write  $Ax$  as

$$Ax = \lambda_x(Ax_0) + \mu_x v^x, \quad (5)$$

where  $\lambda_x \in \mathbb{C}, \mu_x \in \mathbb{R}^+$ , and  $v^x \in \text{Range}(A)$  is a unitary vector orthogonal to  $Ax_0$ .

**Outline of the proof:** As we will see, the norm of  $\|A^\dagger\|$  can be upper bounded by a number arbitrarily close to 1, so to prove Inequality (4), it is enough to focus on the following quantity:

$$\begin{aligned} & \inf_{\phi \in \mathbb{R}} \|e^{i\phi} Ax_0 - b \odot \text{phase}(Ax)\| \\ &= \inf_{\phi \in \mathbb{R}} \|e^{i\phi} |Ax_0| \odot \text{phase}(Ax_0) - |Ax_0| \odot \text{phase}(Ax)\|. \end{aligned}$$

As  $Ax$  is close to  $Ax_0$  (up to a global phase), we can use a kind of mean value inequality, formalized by Lemma III.2. This will yield the upper bound

$$\begin{aligned} & 2 \left\| |Ax_0| \odot \mathbb{1}_{6|\mu_x/\lambda_x||v^x| \geq |Ax_0|} \right\| \\ & + \frac{6}{5} \left\| \text{Im} \left( \left( \frac{\mu_x}{\lambda_x} v^x \right) \odot \overline{\text{phase}(Ax_0)} \right) \right\|. \quad (6) \end{aligned}$$

The imaginary part comes from the derivative of the phase. The term involving an indicator function can be thought of as a second order term.

For the first term of Equation (6), we will see that, when  $\epsilon$  is small enough, the norm of  $6|\mu_x/\lambda_x||v^x|$  is very small compared to the norm of  $Ax_0$ , so the vector  $\mathbb{1}_{6|\mu_x/\lambda_x||v^x| \geq |Ax_0|}$  is very sparse. Using this sparsity, we will be able to upper bound the first term in Equation (6) by

$$12\eta \frac{|\mu_x|}{|\lambda_x|}, \quad (7)$$

where  $\eta > 0$  can be arbitrarily close to zero if  $\epsilon$  is small enough.

For the second term, if we imagined that  $v^x$  was a random vector, chosen uniformly in the sphere independently from  $Ax_0$  (which is of course not true), elementary computations would show that

$$\left\| \operatorname{Im} \left( \left( \frac{\mu_x}{\lambda_x} v^x \right) \odot \overline{\operatorname{phase}(Ax_0)} \right) \right\| \approx \frac{1}{\sqrt{2}} \frac{|\mu_x|}{|\lambda_x|}.$$

Making this intuition rigorous, we will show that the second term in (6) can be upper bounded by

$$\frac{24}{25} \frac{|\mu_x|}{|\lambda_x|}. \quad (8)$$

Combining Equations (7) and (8), we will then upper bound (6) by

$$\delta \frac{|\mu_x|}{|\lambda_x|}$$

for some  $\delta < 1$ .

From the definition of  $\lambda_x$  and  $\mu_x$ , we will see that  $|\lambda_x| \approx 1$  and  $|\mu_x| \lesssim \inf_{\phi \in \mathbb{R}} \|e^{i\phi} Ax_0 - Ax\|$ , which leads us to

$$\begin{aligned} & \inf_{\phi \in \mathbb{R}} \|e^{i\phi} Ax_0 - b \odot \operatorname{phase}(Ax)\| \\ & \lesssim \delta \inf_{\phi \in \mathbb{R}} \|e^{i\phi} Ax_0 - Ax\|. \end{aligned}$$

Applying  $A^\dagger$  on both sides concludes.

Let us now give the details.

**Proof of Equation (6):** The following lemma is proven in Paragraph B-A.

**Lemma III.2.** For any  $z_0, z \in \mathbb{C}$ ,

$$|\operatorname{phase}(z_0 + z) - \operatorname{phase}(z_0)| \leq 2 \cdot \mathbb{1}_{|z| \geq |z_0|/6} + \frac{6}{5} \left| \operatorname{Im} \left( \frac{z}{z_0} \right) \right|.$$

From the lemma, for any  $i = 1, \dots, m$ ,

$$\begin{aligned} & |\operatorname{phase}(\lambda_x)(Ax_0)_i - (b \odot \operatorname{phase}(Ax))_i| \\ & = |\operatorname{phase}(\lambda_x)(Ax_0)_i - |Ax_0|_i \operatorname{phase}(\lambda_x(Ax_0)_i + \mu_x(v^x)_i)| \\ & = |Ax_0|_i \left| \operatorname{phase}(Ax_0)_i - \operatorname{phase} \left( (Ax_0)_i + \frac{\mu_x}{\lambda_x} (v^x)_i \right) \right| \\ & \leq 2 \cdot |Ax_0|_i \mathbb{1}_{|\mu_x/\lambda_x| |v^x| \geq |Ax_0|_i/6} + \frac{6}{5} |Ax_0|_i \left| \operatorname{Im} \left( \frac{\frac{\mu_x}{\lambda_x} v^x}{(Ax_0)_i} \right) \right| \\ & = 2 \cdot |Ax_0|_i \mathbb{1}_{|\mu_x/\lambda_x| |v^x| \geq |Ax_0|_i/6} + \frac{6}{5} \left| \operatorname{Im} \left( \frac{\frac{\mu_x}{\lambda_x} v^x}{\operatorname{phase}((Ax_0)_i)} \right) \right|. \end{aligned}$$

As a consequence,

$$\begin{aligned} & \left\| \operatorname{phase}(\lambda_x)(Ax_0) - b \odot \operatorname{phase}(Ax) \right\| \\ & \leq \left\| 2 \cdot |Ax_0| \odot \mathbb{1}_{|\mu_x/\lambda_x| |v^x| \geq |Ax_0|/6} \right. \\ & \quad \left. + \frac{6}{5} \left| \operatorname{Im} \left( \left( \frac{\mu_x}{\lambda_x} v^x \right) \odot \overline{\operatorname{phase}(Ax_0)} \right) \right| \right\| \\ & \leq 2 \left\| |Ax_0| \odot \mathbb{1}_{6|\mu_x/\lambda_x| |v^x| \geq |Ax_0|} \right\| \\ & \quad + \frac{6}{5} \left\| \operatorname{Im} \left( \left( \frac{\mu_x}{\lambda_x} v^x \right) \odot \overline{\operatorname{phase}(Ax_0)} \right) \right\|. \quad (9) \end{aligned}$$

**Upper bound for  $\frac{|\mu_x|}{|\lambda_x|}$ :** We use a classical result, that allows us to control the norms of  $A$  and  $A^\dagger$ .

**Proposition III.3** ([25], Thm II.13). If  $A$  is chosen according to Equation (2), then, for any  $t$ , with probability at least

$$1 - 2 \exp(-mt^2),$$

we have, for any  $x \in \mathbb{C}^n$ ,

$$\sqrt{m} \left( 1 - \sqrt{\frac{n}{m}} - t \right) \|x\| \leq \|Ax\|;$$

$$\|Ax\| \leq \sqrt{m} \left( 1 + \sqrt{\frac{n}{m}} + t \right) \|x\|.$$

Let us set  $\epsilon^x = \inf_{\phi \in \mathbb{R}} \frac{\|e^{i\phi} x_0 - x\|}{\|x_0\|} \leq \epsilon$ . We have

$$\begin{aligned} \epsilon^x \| \|A\| \|x_0\| & \geq \inf_{\phi \in \mathbb{R}} \|e^{i\phi} Ax_0 - Ax\| \\ & = \sqrt{(1 - |\lambda_x|)^2 \|Ax_0\|^2 + |\mu_x|^2}. \end{aligned}$$

For any two numbers  $\alpha$  and  $\beta$ ,  $\sqrt{\alpha^2 + \beta^2} \geq \max(\alpha, \beta)$ . Hence, the previous inequality implies

$$\begin{aligned} |\lambda_x| & \geq 1 - \frac{\epsilon^x \| \|A\| \|x_0\|}{\|Ax_0\|} \\ \text{and } |\mu_x| & \leq \epsilon^x \| \|A\| \|x_0\|. \end{aligned}$$

From there, we get, upper bounding  $\| \|A\| \|x_0\|$  and lower bounding  $\|Ax_0\|$  according to Proposition III.3,

$$\begin{aligned} \frac{|\mu_x|}{|\lambda_x|} & \leq \frac{\epsilon^x \| \|A\| \|x_0\|}{1 - \frac{\epsilon^x \| \|A\| \|x_0\|}{\|Ax_0\|}} \\ & \leq \frac{\epsilon^x \sqrt{m} \left( 1 + \sqrt{n/m} + t \right) \|x_0\|}{1 - \frac{\epsilon^x \sqrt{m} \left( 1 + \sqrt{n/m} + t \right)}{\sqrt{m} \left( 1 - \sqrt{n/m} - t \right)}} \\ & \leq \frac{\epsilon^x \sqrt{m} \|x_0\|}{\left( 1 - \sqrt{n/m} - t \right) - \epsilon^x \left( 1 + \sqrt{n/m} + t \right)}. \end{aligned}$$

So when  $m \geq 20000n$  and  $\epsilon \leq 10^{-3}$ , setting  $t = 10^{-3}$ , we have

$$\frac{|\mu_x|}{|\lambda_x|} \leq 1.01 \sqrt{m} \epsilon^x \|x_0\|, \quad (10)$$

with probability at least  $1 - 2 \exp(-C_2 m)$ , for  $C_2 = t^2 = 10^{-6}$ .

**Upper bounds for the two terms of Equation (9) (corresponds to Equations (7) and (8) in the outline):** Each term is handled in a separate lemma. The first lemma is proved in Paragraph B-B, the second one in Paragraph B-C.

**Lemma III.4.** For any  $\eta > 0$ , there exists  $C_1, C_2, M, \gamma > 0$  such that the inequality

$$\| |Ax_0| \odot \mathbb{1}_{|v| \geq |Ax_0|} \| \leq \eta \|v\|$$

holds for any  $v \in \operatorname{Range}(A)$  such that  $\|v\| < \gamma \|Ax_0\|$ , with probability at least

$$1 - C_1 \exp(-C_2 m),$$

when  $m \geq Mn$ .

**Lemma III.5.** When  $m \geq 20000n$ , the property

$$\| \operatorname{Im} (v \odot \overline{\operatorname{phase}(Ax_0)}) \| \leq \frac{4}{5} \|v\| \quad (11)$$

holds for any  $v \in \text{Range}(A) \cap \{Ax_0\}^\perp$ , with probability at least

$$1 - 4 \exp(-5 \cdot 10^{-5} m).$$

**Upper bound for Equation (9):** Let us set  $\eta = \frac{1}{600}$ , which ensures

$$12\eta + \frac{24}{25} \leq 0.98.$$

We define  $\gamma > 0$  as in Lemma III.4. Using Lemmas III.4 (applied to  $v = 6(\mu_x/\lambda_x)v^x$ ) and III.5, when  $m \geq Mn$  for  $M$  large enough, we can upper bound Equation (9) by

$$\begin{aligned} & \|\text{phase}(\lambda_x)(Ax_0) - b \odot \text{phase}(Ax)\| \\ & \leq \left(12\eta + \frac{24}{25}\right) \left| \frac{\mu_x}{\lambda_x} \right| \\ & \leq 0.98 \times 1.01 \sqrt{m} \epsilon^x \|x_0\| \\ & \leq 0.99 \sqrt{m} \epsilon^x \|x_0\|. \end{aligned}$$

This holds uniformly over all  $x$  such that  $\inf_{\phi \in \mathbb{R}} \|e^{i\phi} x_0 - x\| \leq \epsilon \|x_0\|$ , with probability at least

$$1 - C'_1 \exp(-C'_2 m). \quad (12)$$

(From Equation (10) and Proposition III.3, still assuming  $m \geq 20000n$ ,  $\epsilon \leq 10^{-3}$  and  $t = 10^{-3}$ , the condition  $\|v\| < \gamma \|Ax_0\|$  in Lemma III.4 is satisfied when the event described in Proposition III.3 is realized.)

**Conclusion:** From the last inequality, we have

$$\begin{aligned} & \inf_{\phi \in \mathbb{R}} \|e^{i\phi} x_0 - A^\dagger(b \odot \text{phase}(Ax))\| \\ & \leq \|\text{phase}(\lambda_x)x_0 - A^\dagger(b \odot \text{phase}(Ax))\| \\ & \leq \|A^\dagger\| \|\text{phase}(\lambda_x)(Ax_0) - b \odot \text{phase}(Ax)\| \\ & \leq 0.99 \|A^\dagger\| \sqrt{m} \epsilon^x \|x_0\| \\ & = 0.99 \|A^\dagger\| \sqrt{m} \inf_{\phi \in \mathbb{R}} \|e^{i\phi} x_0 - x\|. \end{aligned}$$

When the event of Proposition III.3 is realized, still for  $t = 10^{-3}$ ,

$$\begin{aligned} \| \|A^\dagger\| \| & \leq \frac{1}{\sqrt{m}(0.999 - \sqrt{n/m})} \\ & \leq \frac{0.999}{0.99\sqrt{m}}. \end{aligned}$$

So, with the probability of Equation (12), the following property holds uniformly over all  $x$  satisfying  $\inf_{\phi \in \mathbb{R}} \|e^{i\phi} x_0 - x\| \leq \epsilon \|x_0\|$ :

$$\begin{aligned} & \inf_{\phi \in \mathbb{R}} \|e^{i\phi} x_0 - A^\dagger(b \odot \text{phase}(Ax))\| \\ & \leq 0.999 \inf_{\phi \in \mathbb{R}} \|e^{i\phi} x_0 - x\|. \end{aligned}$$

□

**Remark III.6.** From the proof (and Remark B.3 in appendix, evaluated at  $\eta = \frac{1}{600}$ ), we can deduce explicit values for the constants  $\epsilon, C_1, C_2, M$  in Theorem III.1:

$$\begin{aligned} \delta &= 0.999; \\ \epsilon &= 10^{-3}; \\ M &= 3.10^9; \\ C_1 &= 12; \\ C_2 &= 10^{-21}. \end{aligned}$$

## B. Global convergence

In the last paragraph, we have seen that the alternating projections operator is contractive, with high probability, in an  $\epsilon \|x_0\|$ -neighborhood of the solution  $x_0$ . This implies that, if the starting point of alternating projections is at distance at most  $\epsilon \|x_0\|$  from  $x_0$ , alternating projections converge to  $x_0$ . So if we have a way to find such an initial point, we obtain a globally convergent algorithm.

Several initialization methods have been proposed that achieve the precision we need with an optimal number of measurements, that is  $m = O(n)$ . Let us mention the truncated spectral initialization by [14] (improving upon the slightly suboptimal spectral initializations introduced by [12] and [13]), the null initialization by [19] and the method described by [26]. All these methods consist in computing the largest or smallest eigenvector of

$$\sum_{i=1}^m \alpha_i a_i a_i^*,$$

where the  $\alpha_1, \dots, \alpha_m$  are carefully chosen coefficients, that depend only on  $b$ .

The method of [14], for example, has the following guarantees.

**Theorem III.7** (Proposition 3 of [14]). *Let  $\epsilon > 0$  be fixed.*

*There exist  $C_1, C_2, M > 0$  such that, when  $m \geq Mn$ , with probability at least*

$$1 - C_1 \exp(-C_2 m),$$

*the realizations of  $A$  and  $b = |Ax_0|$  are such that the main eigenvector of*

$$\frac{1}{m} \sum_{i=1}^m \left( b_i^2 \mathbb{1}_{b_i^2 \leq \frac{\epsilon}{m} \sum_{j=1}^m b_j^2} \right) a_i a_i^*, \quad (13)$$

*that we denote by  $z$ , obeys*

$$\inf_{\phi \in \mathbb{R}, \lambda \in \mathbb{R}_+^*} \|e^{i\phi} x_0 - \lambda z\| \leq \epsilon \|x_0\|.$$

Combining this initialization procedure with alternating projections, we get Algorithm 1. As shown by the following corollary, it converges towards the correct solution, at a linear rate, with high probability, for  $m = O(n)$ .

**Input** :  $A \in \mathbb{C}^{m \times n}, b = |Ax_0| \in \mathbb{R}^m, T \in \mathbb{N}^*$ .

**Initialization:** set  $z_0$  to be the main eigenvector of the matrix in Equation (13).

**for**  $t = 1$  **to**  $T$  **do**

    | Set  $z_t \leftarrow A^\dagger(b \odot \text{phase}(Az_{t-1}))$ .

**end**

**Output:**  $z_T$ .

**Algorithm 1:** Alternating projections with truncated spectral initialization

**Corollary III.8.** *There exist  $C_1, C_2, M > 0, \delta \in ]0; 1[$  such that, with probability at least*

$$1 - C_1 \exp(-C_2 m),$$

the realizations of  $A$  and  $b = |Ax_0|$  are such that, for any  $T \in \mathbb{N}^*$ , Algorithm 1 satisfies

$$\forall t \leq T, \quad \inf_{\phi \in \mathbb{R}} \|e^{i\phi} x_0 - z_t\| \leq \delta^t \|x_0\|, \quad (14)$$

provided that  $m \geq Mn$ .

*Proof.* Let us fix  $\epsilon, \delta \in ]0; 1[$  as in Theorem III.1. Let us assume that the properties described in Theorems III.1 and III.7 hold; it happens on an event of probability at least

$$1 - C_1 \exp(-C_2 m),$$

provided that  $m \geq Mn$ , for some constants  $C_1, C_2, M > 0$ .

Let us prove that, on this event, Equation (14) also holds.

We proceed by recursion. From Theorem III.7, there exist  $\phi \in \mathbb{R}, \lambda \in \mathbb{R}_+^*$  such that

$$\|e^{i\phi} x_0 - \lambda z_0\| \leq \epsilon \|x_0\|.$$

So, from Theorem III.1, applied to  $x = \lambda z_0$ ,

$$\begin{aligned} \inf_{\phi \in \mathbb{R}} \|e^{i\phi} x_0 - z_1\| &= \inf_{\phi \in \mathbb{R}} \|e^{i\phi} x_0 - A^\dagger(b \odot \text{phase}(z_0))\| \\ &= \inf_{\phi \in \mathbb{R}} \|e^{i\phi} x_0 - A^\dagger(b \odot \text{phase}(\lambda z_0))\| \\ &\leq \delta \inf_{\phi \in \mathbb{R}} \|e^{i\phi} x_0 - \lambda z_0\| \\ &\leq \epsilon \delta \|x_0\|. \end{aligned}$$

This proves Equation (14) for  $t = 1$ . Let us stress that it is critical here that the property of Theorem III.1 holds *uniformly* over all  $x$  that are  $\epsilon \|x_0\|$ -close to  $x_0$ : if this property was only true for vectors  $x$  independent from  $A$  and  $b$ , we could not apply it to  $\lambda z_0$ , since  $\lambda z_0$  depends on  $A$  and  $b$  through the initialization procedure.

The same reasoning can be reapplied to also prove the equation for  $t = 2, 3, \dots$   $\square$

**Remark III.9.** From our computations based on the proof in [14], explicit values for the constants in Theorem III.7 can be chosen as follows:

$$\begin{aligned} C_1 &= 6; \\ C_2 &= 5.10^{-12} \epsilon^4; \\ M &= 4.10^{12} \epsilon^{-4}. \end{aligned}$$

Combined with Remark III.6, this guarantees that Corollary III.8 holds for

$$\begin{aligned} \delta &= 0.999; \\ C_1 &= 18; \\ C_2 &= 5.10^{-24}; \\ M &= 4.10^{24}. \end{aligned}$$

*These theoretical values are absurdly high. They do not match our numerical experiments (Section IV), which suggest that Algorithm 1 succeeds with probability close to 1 as soon as  $m \geq 4n$ .*

*For Theorem III.7, much tighter constants could probably be deduced from the recent work [27]. However, for Corollary III.8, it does not seem possible to make the constants close to tight without significantly improving the proof technique.*

*One particular source of non-tightness in the current technique is that several inequalities that are required hold “uniformly over some large set” (the property in Theorem III.1, for example, holds “for all  $x \in \mathbb{C}^n$  that are  $\epsilon$ -close to  $x_0$ ”). This requirement is a priori too strong: it is actually sufficient that the inequalities hold for “all the points in the large set that are visited by the algorithm”. However, since the points visited by the algorithm do not have a simple expression, it is unclear how to prove only the weaker version.*

### C. Complexity

Let  $\eta > 0$  be the relative precision that we want to achieve:

$$\inf_{\phi \in \mathbb{R}} \|e^{i\phi} x_0 - z_T\| \leq \eta \|x_0\|.$$

Let us compute the number of operations that Algorithm 1 requires to reach this precision.

The main eigenvector of the matrix defined in Equation (13) can be computed - up to precision  $\eta$  - in approximately  $O(\log(1/\eta) + \log(n))$  power iterations. Each power iteration is essentially a matrix-vector multiplication, and thus requires  $O(mn)$  operations.<sup>3</sup> As a consequence, the complexity of the initialization is

$$O(mn (\log(1/\eta) + \log(n))).$$

Then, at each step of the  $\text{f}\circ\text{r}$  loop, the most costly operation is the multiplication by  $A^\dagger$ . When performed with the conjugate gradient method, it requires  $O(mn \log(1/\eta))$  operations. To reach a precision equal to  $\eta$ , we need to perform  $O(\log(1/\eta))$  iterations of the loop. So the total complexity of Algorithm 1 is

$$O(mn (\log^2(1/\eta) + \log(n))).$$

As a comparison, Truncated Wirtinger flow, which is currently the most efficient known method for phase retrieval from Gaussian measurements, has an identical complexity, up to a  $\log(1/\eta)$  factor (see Figure 1).

Let us note that, when  $A$  has some special structure (it encodes Fourier masks [7], for example), the complexity of the algorithm can be further reduced, and exactly matches the one of Truncated Wirtinger flow.

## IV. NUMERICAL EXPERIMENTS

In this section, we numerically validate Corollary III.8. We formulate a conjecture about the convergence of alternating projections with random initialization, in the regime  $m = O(n)$ .

The code used to generate Figures 2, 3 and 4 is available at [http://www-math.mit.edu/~waldspur/code/alternating\\_projections\\_code.zip](http://www-math.mit.edu/~waldspur/code/alternating_projections_code.zip).

<sup>3</sup>These matrix-vector multiplications can be computed without forming the whole matrix (which would require  $O(mn^2)$  operations), because this matrix factorizes as

$$\frac{1}{m} A^* \text{Diag}(b^2 \odot I) A,$$

where  $I \in \mathbb{R}^m$  is such that  $\forall i \leq m, I_i = \mathbb{1}_{b_i^2 \leq \frac{\epsilon}{m} \sum_{j=1}^m b_j^2}$ .

	Alternating projections	Truncated Wirtinger flow
No special structure for $A$	$O(mn(\log^2(1/\eta) + \log(n)))$	$O(mn(\log(1/\eta) + \log(n)))$
Fourier masks	$O(m \log(n)(\log(1/\eta) + \log(n)))$	$O(m \log(n)(\log(1/\eta) + \log(n)))$

Figure 1: Complexity of alternating projections with initialization, and truncated Wirtinger flow.




Figure 2: Probability of success for Algorithm 1, as a function of  $n$  and  $m$ . Black points indicate a probability equal to 0, and white points a probability equal to 1. The line serves as a reference: it is the line  $m = 4n$ .

#### A. Alternating projections with careful initialization

Corollary III.8 states that alternating projections succeed with high probability, when they start from a good initial point, in the regime where the number of measurements is linear in the problem dimension ( $m = O(n)$ ).

We use the initialization method described in [14], as presented in Algorithm 1. We run the algorithm many times, for various choices of  $n$  and  $m$ . This allows us to compute an empirical probability of success, for each value of  $(n, m)$ .

The results are presented in Figure 2. They confirm that, when  $m = Cn$ , for a sufficiently large constant  $C > 0$ , the success probability can be arbitrarily close to 1.

#### B. Alternating projections without careful initialization

In a second experiment, we measure the probability that alternating projections succeed, when started from a random initial point (sampled from the unit sphere with uniform probability).

As previously, we numerically compute this probability for various pairs  $(m, n)$ . The results, presented in Figure 3 seem to indicate that alternating projections behave similarly with and without a careful initialization procedure: they converge towards the correct solution as soon as  $m \geq Cn$ , for a suitable constant  $C > 0$ . Only the value of  $C$  changes. Figure 4 displays similar computations, under a different form: it shows how many measurements  $m$  are needed so that alternating projections, with careful or with random initialization, succeeds

with probability  $1/2$ . This number apparently grows linearly with  $n$  whatever the initialization method.

As a consequence, we have the following conjecture.

**Conjecture IV.1.** *Let any  $\epsilon > 0$  be fixed. When  $m \geq Cn$ , for  $C > 0$  large enough, alternating projections with a random rotationally invariant initialization succeed with probability at least  $1 - \epsilon$ .*

This conjecture is in the same spirit as several recent works, which show that non-convex iterative algorithms, starting from random initial points, succeed with high probability for a number of problems. These works cover matrix sensing [28], matrix completion [29], robust PCA [30], and synchronization [31]. The closest one to our conjecture is [21], that also considers phase retrieval with Gaussian sensing vectors: in the almost optimal regime  $m = O(n \log^3 n)$ , it shows that gradient descent over a (specific) non-convex function succeeds with high probability, even when randomly initialized.

Unfortunately, the proof method of these works does not seem to adapt our case. Indeed, it consists in showing that the iterative routine has no attractive stagnation point. But in our case, complementary experiments (not shown here), suggest that, in the regime  $m = O(n)$ , there are attractive stagnation points. These points seem to have small attraction basins, so they are not necessarily an obstacle for the success of the algorithm with high probability, but rigorously establishing this property seems difficult.

## APPENDIX A

### PROOF OF PROPOSITION II.1

**Proposition (Proposition II.1).** *For any  $y_0$ , the sequence  $(y_k)_{k \in \mathbb{N}}$  is bounded. Any accumulation point  $y_\infty$  of  $(y_k)_{k \in \mathbb{N}}$  satisfies the following property:*

$$\exists u \in E_{\text{phase}}(y_\infty), \quad (AA^\dagger)(b \odot u) = y_\infty.$$

*In particular, if  $y_\infty$  has no zero entry,*

$$(AA^\dagger)(b \odot \text{phase}(y_\infty)) = y_\infty.$$

*Proof of Proposition II.1.* The boundedness of  $(y_k)_{k \in \mathbb{N}}$  is a consequence of the fact that  $\|y'_k\| = \|b\|$  for all  $k$ , so  $\|y_{k+1}\| \leq \|AA^\dagger\| \|b\|$ .

Let us show the second part of the statement. Let  $y_\infty$  be an accumulation point of  $(y_k)_{k \in \mathbb{N}}$ , and let  $\phi : \mathbb{N} \rightarrow \mathbb{N}$  be an extraction such that

$$y_{\phi(n)} \rightarrow y_\infty \quad \text{when } n \rightarrow +\infty.$$

By compactity, as  $(y'_{\phi(n)})_{n \in \mathbb{N}}$  and  $(y_{\phi(n)+1})_{n \in \mathbb{N}}$  are bounded sequences, we can assume, even if we have to consider replace  $\phi$  by a subextraction, that they also converge. We denote by  $y'_\infty$  and  $y_\infty^{+1}$  their limits:

$$y'_{\phi(n)} \rightarrow y'_\infty \quad \text{and} \quad y_{\phi(n)+1} \rightarrow y_\infty^{+1} \quad \text{when } n \rightarrow +\infty.$$






Figure 3: Probability of success for alternating projections with a random Gaussian initialization, as a function of  $n$  and  $m$ . Black points indicate a probability equal to 0, and white points a probability equal to 1. The line serves as a reference: it is the line  $m = 4n$ .




Figure 4: As a function of  $n$ , the smallest value of  $m$  for which alternating projections succeeds with probability 50%. The solid line is when the algorithm is initialized with the method in [14]; the dashed line is with random Gaussian initialization.

Let us define

$$E_b = \{y' \in \mathbb{C}^m, |y'| = b\}.$$

We observe that, for any  $k$ ,

$$d(y'_{k-1}, \text{Range}(A)) \geq d(y_k, E_b) \geq d(y'_k, \text{Range}(A)).$$

Indeed, because the operators  $y \rightarrow b \odot \text{phase}(y)$  and  $y \rightarrow (AA^\dagger)y$  are projections,

$$\begin{aligned} d(y'_{k-1}, \text{Range}(A)) &= d(y'_{k-1}, y_k) \geq d(y_k, E_b); \\ d(y_k, E_b) &= d(y_k, y'_k) \geq d(y'_k, \text{Range}(A)). \end{aligned}$$

So the sequences  $(d(y_k, E_b))_{k \in \mathbb{N}}$  and  $(d(y'_k, \text{Range}(A)))_{k \in \mathbb{N}}$  converge to the same non-negative limit, that we denote by  $\delta$ . In particular,

$$d(y_\infty, E_b) = \delta = d(y'_\infty, \text{Range}(A)).$$

If we pass to the limit the equalities  $d(y_{\phi(n)}, E_b) = \|y_{\phi(n)} - y'_{\phi(n)}\|$  and  $d(y'_{\phi(n)}, \text{Range}(A)) = \|y'_{\phi(n)} - y_{\phi(n)+1}\|$ , we get

$$\|y_\infty - y'_\infty\| = \|y'_\infty - y_\infty^{+1}\| = \delta = d(y'_\infty, \text{Range}(A)).$$

As  $\text{Range}(A)$  is convex, the projection of  $y'_\infty$  onto it is uniquely defined. This implies

$$y_\infty = y_\infty^{+1},$$

and, because  $\forall n, y_{\phi(n)+1} = (AA^\dagger)y'_{\phi(n)}$ ,

$$y_\infty = y_\infty^{+1} = (AA^\dagger)y'_\infty.$$

To conclude, we now have to show that  $y'_\infty = b \odot u$  for some  $u \in E_{\text{phase}(y_\infty)}$ . We use the fact that, for all  $n$ ,  $y'_{\phi(n)} = b \odot \text{phase}(y_{\phi(n)})$ .

For any  $i \in \{1, \dots, m\}$ , if  $(y_\infty)_i \neq 0$ , phase is continuous around  $(y_\infty)_i$ , so  $(y'_\infty)_i = b_i \text{phase}((y_\infty)_i)$ . We then set  $u_i = \text{phase}((y_\infty)_i)$ , and we have  $(y'_\infty)_i = b_i u_i$ .

If  $(y_\infty)_i = 0$ , we set  $u_i = \text{phase}((y'_\infty)_i) \in E_{\text{phase}(0)} = E_{\text{phase}((y_\infty)_i)}$ . We then have  $y'_\infty = |y'_\infty| u_i = b_i u_i$ .

With this definition of  $u$ , we have, as claimed,  $y'_\infty = b \odot u$  and  $u \in E_{\text{phase}(y_\infty)}$ .  $\square$

## APPENDIX B

### TECHNICAL LEMMAS FOR SECTION III

#### A. Proof of Lemma III.2

**Lemma (Lemma III.2).** For any  $z_0, z \in \mathbb{C}$ ,

$$|\text{phase}(z_0 + z) - \text{phase}(z_0)| \leq 2 \cdot \mathbb{1}_{|z| \geq |z_0|/6} + \frac{6}{5} \left| \text{Im} \left( \frac{z}{z_0} \right) \right|.$$

*Proof.* The inequality holds if  $z_0 = 0$ , so we can assume  $z_0 \neq 0$ . We remark that, in this case,

$$|\text{phase}(z_0 + z) - \text{phase}(z_0)| = |\text{phase}(1 + z/z_0) - 1|.$$

It is thus enough to prove the lemma for  $z_0 = 1$ , so we make this assumption.

When  $|z| \geq 1/6$ , the inequality is valid. Let us now assume that  $|z| < 1/6$ . Let  $\theta \in ]-\pi/2; \pi/2[$  be such that

$$e^{i\theta} = \text{phase}(1 + z).$$

Then

$$\begin{aligned} |\text{phase}(1 + z) - 1| &= |e^{i\theta} - 1| \\ &= 2|\sin(\theta/2)| \\ &\leq |\tan \theta| \\ &= \frac{|\text{Im}(1 + z)|}{|\text{Re}(1 + z)|} \\ &\leq \frac{|\text{Im}(z)|}{1 - |z|} \\ &\leq \frac{6}{5} |\text{Im}(z)|. \end{aligned}$$

So the inequality is also valid.  $\square$

### B. Proof of Lemma III.4

**Lemma** (Lemma III.4). *For any  $\eta > 0$ , there exists  $C_1, C_2, M, \gamma > 0$  such that the inequality*

$$\| |Ax_0| \odot \mathbb{1}_{|v| \geq |Ax_0|} \| \leq \eta \|v\|$$

*holds for any  $v \in \text{Range}(A)$  such that  $\|v\| < \gamma \|Ax_0\|$ , with probability at least*

$$1 - C_1 \exp(-C_2 m),$$

when  $m \geq Mn$ .

*Proof.* For any  $S \subset \{1, \dots, m\}$ , we denote by  $\mathbb{1}_S$  the vector of  $\mathbb{C}^m$  such that

$$\begin{aligned} (\mathbb{1}_S)_j &= 1 \text{ if } j \in S \\ &= 0 \text{ if } j \notin S. \end{aligned}$$

We use the following two lemmas, proven in Paragraphs B-B1 and B-B2.

**Lemma B.1.** *Let  $\beta \in ]0; 1/2[$  be fixed. With probability at least*

$$1 - 2 \exp(-\beta^3 m/e),$$

*the following property holds: for any  $S \subset \{1, \dots, m\}$  such that  $\text{Card}(S) \geq \beta m$ ,*

$$\| |Ax_0| \odot \mathbb{1}_S \| \geq \beta^{3/2} e^{-1/2} \|Ax_0\|. \quad (15)$$

**Lemma B.2.** *Let  $\beta \in ]0; \frac{1}{100}[$  be fixed. For any  $M \geq \frac{3}{\beta}$ , if  $m \geq Mn$ , then, with probability at least*

$$1 - 4 \exp\left(-3\beta \log\left(\frac{1}{\beta}\right) m\right),$$

*the following property holds: for any  $S \subset \{1, \dots, m\}$  such that  $\text{Card}(S) < \beta m$  and for any  $y \in \text{Range}(A)$ ,*

$$\|y \odot \mathbb{1}_S\| \leq 10\sqrt{\beta \log(1/\beta)} \|y\|. \quad (16)$$

Let  $\beta > 0$  be such that  $10\sqrt{\beta \log(1/\beta)} \leq \eta$  (we can for instance choose  $\beta = \min\left(0.01, \frac{\eta^2}{300 \log(1/\eta)}\right)$ ). Let  $M$  be larger than  $\frac{3}{\beta}$ , as in Lemma B.2. We set

$$\gamma = \beta^{3/2} e^{-1/2}.$$

We assume that Equations (15) and (16) hold; from the lemmas, this occurs with probability at least

$$1 - 6 \exp(-\beta^3 m/e),$$

provided that  $m \geq Mn$ . (We have used the fact that, for  $\beta \leq 0.01$ ,  $\beta^3/e \leq 3\beta \log(1/\beta)$ .)

On this event, for any  $v \in \text{Range}(A)$  such that  $\|v\| < \gamma \|Ax_0\|$ , if we set  $S_v = \{i \text{ s.t. } |v_i| \geq |Ax_0|_i\}$ , we have that

$$\text{Card } S_v < \beta m.$$

Indeed, if it was not the case, we would have, by Equation (15),

$$\begin{aligned} \|v\| &\geq \|v \odot \mathbb{1}_{S_v}\| \\ &\geq \| |Ax_0| \odot \mathbb{1}_{S_v} \| \\ &\geq \beta^{3/2} e^{-1/2} \|Ax_0\| \\ &= \gamma \|Ax_0\|, \end{aligned}$$

which is in contradiction with the way we have chosen  $v$ .

So we can apply Equation (16), and we get

$$\begin{aligned} \| |Ax_0| \odot \mathbb{1}_{|v| \geq |Ax_0|} \| &\leq \|v \odot \mathbb{1}_S\| \\ &\leq 10\sqrt{\beta \log(1/\beta)} \|v\| \\ &\leq \eta \|v\|. \end{aligned}$$

□

**Remark B.3.** *We can choose the values of  $C_1, C_2, M, \gamma$  as follows:*

$$\begin{aligned} C_1 &= 6; \\ C_2 &= \frac{10^{-6}}{e} \min\left(1, \frac{\eta^6}{27 \log^3(1/\eta)}\right); \\ M &= \frac{300}{\min\left(1, \frac{\eta^2}{3 \log(1/\eta)}\right)}; \\ \gamma &= 10^{-3} e^{-1/2} \min\left(1, \frac{\eta^3}{3^{3/2} \log^{3/2}(1/\eta)}\right). \end{aligned}$$

#### 1) Proof of Lemma B.1:

*Proof of Lemma B.1.* We first assume  $S$  fixed, with cardinality  $\text{Card } S \geq \beta m$ . We use the following lemma.

**Lemma B.4** ([32], Lemma 2.2). *Let  $k_1 < k_2$  be natural numbers. Let  $X \in \mathbb{C}^{k_2}$  be a random vector whose coordinates are independent, Gaussian, of variance 1. Let  $Y$  be the projection of  $X$  onto its  $k_1$  first coordinates. Then, for any  $t > 0$ ,*

$$\begin{aligned} \text{Proba}\left(\frac{\|Y\|}{\|X\|} \leq \sqrt{\frac{tk_1}{k_2}}\right) &\leq \exp(k_1(1-t+\log t)) \quad \text{if } t < 1; \\ \text{Proba}\left(\frac{\|Y\|}{\|X\|} \geq \sqrt{\frac{tk_1}{k_2}}\right) &\leq \exp(k_1(1-t+\log t)) \quad \text{if } t > 1. \end{aligned}$$

From this lemma, for any  $t \in ]0; 1[$ , because  $Ax_0$  has independent Gaussian coordinates,

$$P\left(\frac{\| |Ax_0| \odot \mathbb{1}_S \|}{\|Ax_0\|} \leq \sqrt{t\beta}\right) \leq \exp(-\beta m(t-1-\ln t)).$$

In particular, for  $t = \frac{\beta^2}{e}$ ,

$$\begin{aligned} P\left(\frac{\| |Ax_0| \odot \mathbb{1}_S \|}{\|Ax_0\|} \leq \beta^{3/2} e^{-1/2}\right) \\ \leq \exp\left(-\beta m \left(\frac{\beta^2}{e} - 2 \ln \beta\right)\right). \end{aligned} \quad (17)$$

As soon as  $m \geq \frac{1}{2\beta}$ , the number of subsets  $S$  of  $\{1, \dots, m\}$  with cardinality  $\lceil \beta m \rceil$  satisfies

$$\begin{aligned} \binom{m}{\lceil \beta m \rceil} &\leq \left(\frac{em}{\lceil \beta m \rceil}\right)^{\lceil \beta m \rceil} \\ &\leq \exp\left(2m\beta \log \frac{1}{\beta}\right). \end{aligned} \quad (18)$$

(The first inequality is a classical result regarding binomial coefficients.)

We combine Equations (17) and (18): Property (15) is satisfied for any  $S$  of cardinality  $\lceil \beta m \rceil$  with probability at least

$$1 - \exp\left(-\frac{\beta^3}{e}m\right) \geq 1 - 2\exp\left(-\frac{\beta^3}{e}m\right),$$

provided that  $m \geq \frac{1}{2\beta}$ .

When  $m < \frac{1}{2\beta}$ , since  $\beta < 1/2$ ,

$$\begin{aligned} 1 - 2\exp(-\beta^3 m/e) &< 1 - 2\exp(-\beta^2/(2e)) \\ &< 1 - 2\exp(-1/(8e)) \\ &< 0, \end{aligned}$$

so the property also holds with probability larger than  $1 - 2\exp(-\beta^3 m/e)$ .

If it is satisfied for any  $S$  of cardinality  $\lceil \beta m \rceil$ , then it is satisfied for any  $S$  of cardinality larger than  $\beta m$ , which implies the result.  $\square$

## 2) Proof of Lemma B.2:

*Proof of Lemma B.2.* We first assume  $S$  to be fixed, of cardinality exactly  $\lceil \beta m \rceil$ .

Any vector  $y \in \text{Range}(A)$  is of the form  $y = Av$ , for some  $v \in \mathbb{C}^n$ . Inequality (16) can then be rewritten as:

$$\|A_S v\| = \|\text{Diag}(\mathbf{1}_S)Av\| \leq 10\sqrt{\beta \log(1/\beta)}\|Av\|, \quad (19)$$

where  $A_S$ , by definition, is the submatrix obtained from  $A$  by extracting the rows whose indexes are in  $S$ .

We apply Proposition III.3 to  $A$  and  $A_S$ , respectively for  $t = \frac{1}{2}$  and  $t = 3\sqrt{\log(1/\beta)}$ . It guarantees that the following properties hold:

$$\begin{aligned} \inf_{v \in \mathbb{C}^n} \frac{\|Av\|}{\|v\|} &\geq \sqrt{m} \left( \frac{1}{2} - \sqrt{\frac{n}{m}} \right); \\ \sup_{v \in \mathbb{C}^n} \frac{\|A_S v\|}{\|v\|} &\leq \sqrt{\text{Card } S} \left( 1 + \sqrt{\frac{n}{\text{Card } S}} + 3\sqrt{\log(1/\beta)} \right), \end{aligned}$$

with respective probabilities at least

$$\begin{aligned} 1 - 2\exp\left(-\frac{m}{4}\right); \\ \text{and } 1 - 2\exp(-9(\text{Card } S)\log(1/\beta)) \\ \geq 1 - 2\exp(-9\beta \log(1/\beta)m). \end{aligned}$$

Assuming  $m \geq Mn$  for some  $M > 0$ , we deduce from these inequalities that, for all  $v \in \mathbb{C}^n$ ,

$$\begin{aligned} \|A_S v\| &\leq \sqrt{\frac{\text{Card } S}{m}} \left( \frac{1 + \sqrt{\frac{n}{\text{Card } S}} + 3\sqrt{\log(1/\beta)}}{\frac{1}{2} - \sqrt{\frac{n}{m}}} \right) \|Av\| \\ &\leq \sqrt{\beta + \frac{1}{m}} \left( \frac{1 + \sqrt{\frac{1}{\beta M}} + 3\sqrt{\log(1/\beta)}}{\frac{1}{2} - \sqrt{\frac{1}{M}}} \right) \|Av\|, \end{aligned} \quad (20)$$

with probability at least

$$1 - 2\exp(-9\beta \log(1/\beta)m) - 2\exp\left(-\frac{m}{4}\right).$$

If we choose  $M \geq \frac{3}{\beta}$ , remember that  $\beta \leq 0.01$  and  $m \geq Mn \geq M$ , we can deduce from Equation (20) that

$$\begin{aligned} \|A_S v\| &\leq \sqrt{\beta} \sqrt{1 + \frac{1}{3}} \left( \frac{1 + \sqrt{\frac{1}{3}} + 3\sqrt{\log(1/\beta)}}{\frac{1}{2} - \sqrt{\frac{1}{300}}} \right) \|Av\| \\ &\leq 10\sqrt{\beta \log(1/\beta)}\|Av\|, \end{aligned}$$

which is Equation (19).

As in the proof of Lemma B.1, there are at most

$$\exp\left(2m\beta \log \frac{1}{\beta}\right)$$

subsets of  $\{1, \dots, m\}$  with cardinality  $\lceil \beta m \rceil$  (note that  $m \geq M \geq 3/\beta \geq 1/(2\beta)$ ). As a consequence, Equation (19) holds for any  $v \in \mathbb{C}^n$  and  $S$  of cardinality  $\lceil \beta m \rceil$  with probability at least

$$1 - 2\exp(-7\beta \log(1/\beta)m) - 2\exp\left(-\left(\frac{1}{4} - 2\beta \log \frac{1}{\beta}\right)m\right).$$

When  $\beta \leq \frac{1}{100}$ , we have

$$\frac{1}{4} - 2\beta \log \frac{1}{\beta} > 3\beta \log \frac{1}{\beta},$$

so the resulting probability is larger than

$$1 - 4\exp\left(-3\beta \log \left(\frac{1}{\beta}\right)m\right).$$

This ends the proof. Indeed, if Equation (19) holds for any set of cardinality  $\lceil \beta m \rceil$ , it also holds for any set of cardinality  $\text{Card } S < \beta m$ , because  $\|A_{S'} v\| \leq \|A_S v\|$  whenever  $S' \subset S$ . This implies Equation (16).  $\square$

## C. Proof of Lemma III.5

**Lemma (Lemma III.5).** *When  $m \geq 20000n$ , the property*

$$\|\text{Im}(v \odot \overline{\text{phase}(Ax_0)})\| \leq \frac{4}{5}\|v\| \quad (21)$$

holds for any  $v \in \text{Range}(A) \cap \{Ax_0\}^\perp$ , with probability at least

$$1 - 4\exp(-5.10^{-5}m).$$

*Proof.* If we multiply  $x_0$  by a positive real number, we can assume  $\|x_0\| = 1$ . Moreover, as the law of  $A$  is invariant under right multiplication by a unitary matrix, we can assume that

$$x_0 = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}.$$

Then, if we write  $A_1$  the first column of  $A$ , and  $A_{2:n}$  the submatrix of  $A$  obtained by removing this first column,

$$\begin{aligned} &\text{Range}(A) \cap \{Ax_0\}^\perp \\ &= \left\{ w - \frac{\langle w, A_1 \rangle}{\|A_1\|^2} A_1, w \in \text{Range}(A_{2:n}) \right\}. \end{aligned} \quad (22)$$

We first observe that

$$\sup_{w \in \text{Range}(A_{2:n}) - \{0\}} \frac{|\langle w, A_1 \rangle|}{\|w\|}$$

is the norm of the orthogonal projection of  $A_1$  onto  $\text{Range}(A_{2:n})$ . The  $(n-1)$ -dimensional subspace  $\text{Range}(A_{2:n})$  has a rotationally invariant distribution in  $\mathbb{C}^m$ , and is independent of  $A_1$ . Thus, from Lemma B.4 coming from [32], for any  $t > 1$ ,

$$\sup_{w \in \text{Range}(A_{2:n}) - \{0\}} \frac{|\langle w, A_1 \rangle|}{\|w\| \|A_1\|} < \sqrt{\frac{t(n-1)}{m}},$$

with probability at least

$$1 - \exp(-(n-1)(t-1-\log t)).$$

We take  $t = \frac{m}{n-1}(0.04)^2$  (which is larger than 1 when  $m \geq Mn$  with  $M > 625$ ), and it implies that

$$\sup_{w \in \text{Range}(A_{2:n}) - \{0\}} \frac{|\langle w, A_1 \rangle|}{\|w\| \|A_1\|} < 0.04 \quad (23)$$

with probability at least

$$\begin{aligned} & 1 - \exp(-(n-1)(t-1-\log t)) \\ & \geq 1 - \exp(-(n-1)t/4) \\ & = 1 - \exp(-4.10^{-4}m). \end{aligned}$$

provided that  $m \geq 2500n$ . (We have used the fact that, when  $M \geq 2500$ , then  $t \geq 4$ , and when  $t \geq 4$ , then  $t-1-\log t \geq t/4$ .)

Second, as  $A_{2:n}$  is a random matrix of size  $m \times (n-1)$ , whose entries are independent and distributed according to the law  $\mathcal{N}(0, 1/2) + \mathcal{N}(0, 1/2)i$ , we deduce from Proposition III.3 applied with  $t = 0.01$  that, with probability at least

$$1 - 2 \exp(-10^{-4}m),$$

we have, for any  $x \in \mathbb{C}^{n-1}$ ,

$$\begin{aligned} \|A_{2:n}x\| & \geq \sqrt{m} \left( 1 - \sqrt{\frac{(n-1)}{m}} - 0.01 \right) \|x\| \\ & \geq 0.98\sqrt{m}\|x\|, \end{aligned} \quad (24)$$

provided that  $m \geq 10000n$ .

We now set

$$C = \text{Diag}(\overline{\text{phase}(A_1)})A_{2:n}.$$

The matrix  $(\text{Im } C \quad \text{Re } C)$  has size  $m \times (2(n-1))$ ; its entries are independent and distributed according to the law  $\mathcal{N}(0, 1/2)$ . So by [25, Thm II.13] (applied with  $t = 0.01$ ), with probability at least

$$1 - \exp(-5.10^{-5}m),$$

we have, for any  $x \in \mathbb{R}^{2(n-1)}$ ,

$$\begin{aligned} \|(\text{Im } C \quad \text{Re } C)x\| & \leq \sqrt{\frac{m}{2}} \left( 1 + \sqrt{\frac{2(n-1)}{m}} + 0.01 \right) \|x\| \\ & \leq 1.02\sqrt{\frac{m}{2}}\|x\|, \end{aligned} \quad (25)$$

provided that  $m \geq 20000n$ .

When Equations (24) and (25) are simultaneously valid, any  $w = A_{2:n}w'$  belonging to  $\text{Range}(A_{2:n})$  satisfies:

$$\begin{aligned} \left\| \text{Im}(w \odot \overline{\text{phase}(Ax_0)}) \right\| & = \|\text{Im}(Cw')\| \\ & = \left\| (\text{Im } C \quad \text{Re } C) \begin{pmatrix} \text{Re } w' \\ \text{Im } w' \end{pmatrix} \right\| \\ & \leq 1.02\sqrt{\frac{m}{2}} \left\| \begin{pmatrix} \text{Re } w' \\ \text{Im } w' \end{pmatrix} \right\| \\ & = 1.02\sqrt{\frac{m}{2}}\|w'\| \\ & \leq \frac{1.02}{0.98\sqrt{2}}\|A_{2:n}w'\| \\ & = \frac{1.02}{0.98\sqrt{2}}\|w\| \\ & \leq 0.75\|w\|. \end{aligned} \quad (26)$$

We now conclude. Equations (23), (24) and (25) hold simultaneously with probability at least

$$1 - 4 \exp(-5.10^{-5}m),$$

provided that  $m \geq 20000n$ . Let us show that, on this event, Equation (21) also holds. Any  $v \in \text{Range}(A) \cap \{Ax_0\}^\perp$ , from Equality (22), can be written as

$$v = w - \frac{\langle w, A_1 \rangle}{\|A_1\|^2} A_1,$$

for some  $w \in \text{Range}(A_{2:n})$ . Using Equation (23), then Equation (26), we get:

$$\begin{aligned} \left\| \text{Im}(v \odot \overline{\text{phase}(Ax_0)}) \right\| & \leq \left\| \text{Im}(w \odot \overline{\text{phase}(Ax_0)}) \right\| \\ & \quad + \left\| \frac{\langle w, A_1 \rangle}{\|A_1\|^2} A_1 \right\| \\ & \leq \left\| \text{Im}(w \odot \overline{\text{phase}(Ax_0)}) \right\| \\ & \quad + 0.04\|w\| \\ & \leq 0.79\|w\|. \end{aligned}$$

But then, by Equation (23) again,

$$\|v\|^2 = \|w\|^2 - \frac{\langle w, A_1 \rangle^2}{\|A_1\|^2} \geq (1 - (0.04)^2)\|w\|^2.$$

So

$$\begin{aligned} \left\| \text{Im}(v \odot \overline{\text{phase}(Ax_0)}) \right\| & \leq 0.79\|w\| \\ & \leq \frac{0.79}{\sqrt{1 - (0.04)^2}}\|v\| \\ & \leq \frac{4}{5}\|v\|. \end{aligned}$$

□

## REFERENCES

- [1] Y. Schechtman, Y. C. Eldar, O. Cohen, H. N. Chapman, J. Miao, and M. Segev, "Phase retrieval with application to optical imaging: a contemporary overview," *IEEE Signal processing magazine*, vol. 32, no. 3, pp. 87–109, 2015.
- [2] R. Gerchberg and W. Saxton, "A practical algorithm for the determination of phase from image and diffraction plane pictures," *Optik*, vol. 35, pp. 237–246, 1972.

- [3] J. R. Fienup, "Phase retrieval algorithms: a comparison," *Applied Optics*, vol. 21, no. 15, pp. 2758–2769, 1982.
- [4] A. Chai, M. Moscoso, and G. Papanicolaou, "Array imaging using intensity-only measurements," *Inverse Problems*, vol. 27, no. 1, 2011.
- [5] E. J. Candès, T. Strohmer, and V. Voroninski, "Phaselift: exact and stable signal recovery from magnitude measurements via convex programming," *Communications in Pure and Applied Mathematics*, vol. 66, no. 8, pp. 1241–1274, 2013.
- [6] E. J. Candès and X. Li, "Solving quadratic equations via phaselift when there are about as many equations as unknowns," *Foundations of Computational Mathematics*, vol. 14, no. 5, pp. 1017–1026, 2014.
- [7] E. J. Candès, X. Li, and M. Soltanolkotabi, "Phase retrieval from coded diffraction patterns," *Applied and Computational Harmonic Analysis*, vol. 39, no. 2, pp. 277–299, 2015.
- [8] D. Gross, F. Krahermer, and R. Kueng, "A partial derandomization of PhaseLift using spherical designs," *Journal of Fourier Analysis and Applications*, vol. 21, no. 2, pp. 229–266, 2015.
- [9] D. L. Sun and J. O. Smith, "Estimating a signal from a magnitude spectrogram via convex optimization," in *Audio Engineering Society 133rd Convention*, 2012.
- [10] I. Waldspurger, A. d'Aspremont, and S. Mallat, "Phase recovery, maxcut and complex semidefinite programming," *Mathematical Programming*, vol. 149, no. 1–2, pp. 47–81, 2015.
- [11] C. D. White, S. Sanghavi, and R. Ward, "The local convexity of solving systems of quadratic equations," *preprint*, 2015, <http://arxiv.org/abs/1506.07868>.
- [12] P. Netrapalli, P. Jain, and S. Sanghavi, "Phase retrieval using alternating minimization," in *Advances in Neural Information Processing Systems 26*, 2013, pp. 1796–2804.
- [13] E. J. Candès, X. Li, and M. Soltanolkotabi, "Phase retrieval via wirtinger flow: theory and algorithms," *IEEE Transactions of Information Theory*, vol. 61, no. 4, pp. 1985–2007, 2015.
- [14] Y. Chen and E. J. Candès, "Solving random quadratic systems of equations is nearly as easy as solving linear systems," *To appear in Communications on Pure and Applied Mathematics*, 2015.
- [15] R. Kolté and A. Özgür, "Phase retrieval via incremental truncated Wirtinger flow," *preprint*, 2016, [arxiv.org/abs/1606.03196](http://arxiv.org/abs/1606.03196).
- [16] H. Zhang and Y. Liang, "Reshaped Wirtinger flow for solving quadratic systems of equations," in *Advances in Neural Information Processing Systems 29*, 2016.
- [17] G. Wang, G. B. Giannakis, and Y. C. Eldar, "Solving random systems of quadratic equations via truncated generalized gradient flow," *preprint*, 2016, <https://arxiv.org/abs/1605.08285v1>.
- [18] D. Noll and A. Rondepierre, "On local convergence of the method of alternating projections," *Foundations of Computational Mathematics*, vol. 16, no. 2, pp. 425–455, 2016.
- [19] P. Chen, A. Fannjiang, and G.-R. Liu, "Phase retrieval with one or two diffraction patterns by alternating projection with null initialization," *preprint*, 2016, <http://arxiv.org/abs/1510.07379>.
- [20] M. Soltanolkotabi, "Algorithms and theory for clustering and nonconvex quadratic programming," Ph.D. dissertation, Stanford University, 2014.
- [21] J. Sun, Q. Qu, and J. Wright, "A geometric analysis of phase retrieval," *preprint*, 2016, <http://arxiv.org/abs/1602.06664>.
- [22] I. Waldspurger, "Phase retrieval with random gaussian sensing vectors by alternating projections," *preprint*, 2016, <https://arxiv.org/abs/1609.03088>.
- [23] R. Balan, P. Casazza, and D. Edidin, "On signal reconstruction without noisy phase," *Applied and Computational Harmonic Analysis*, vol. 20, pp. 345–356, 2006.
- [24] A. Conca, D. Edidin, M. Hering, and C. Vinzant, "Algebraic characterization of injectivity in phase retrieval," *Applied and Computational Harmonic Analysis*, vol. 32, no. 2, pp. 346–356, 2015.
- [25] K. R. Davidson and S. J. Szarek, "Local operator theory, random matrices and Banach spaces," in *Handbook of the geometry of Banach spaces, volume 1*, W. B. Johnson and J. Lindenstrauss, Eds. Elsevier, 2001, pp. 317–366.
- [26] B. Gao and Z. Xu, "Gauss-Newton method for phase retrieval," *preprint*, 2016, <http://arxiv.org/abs/1606.08135>.
- [27] Y. M. Lu and G. Li, "Phase transitions of spectral initialization for high-dimensional nonconvex estimation," *preprint*, 2017, <https://arxiv.org/abs/1702.06435>.
- [28] S. Bhojanapalli, B. Neyshabur, and N. Srebro, "Global optimality of local search for low rank matrix recovery," in *Advances in Neural Information Processing Systems 29*, 2016.
- [29] R. Ge, J. D. Lee, and T. Ma, "Matrix completion has no spurious local minimum," in *Advances in Neural Information Processing Systems 29*. Curran Associates, Inc., 2016.
- [30] R. Ge, C. Jin, and Y. Zheng, "No spurious local minima in nonconvex low rank problems: a unified geometric analysis," in *to appear in the Proceedings of the 34th International Conference on Machine Learning*, 2017.
- [31] N. Boumal, "Nonconvex phase synchronization," *SIAM Journal on Optimization*, vol. 26, no. 4, pp. 2355–2377, 2016.
- [32] S. Dasgupta and A. Gupta, "An elementary proof of a theorem of Johnson and Lindenstrauss," *Random Structures and Algorithms*, vol. 22, no. 1, pp. 60–65, 2003.