

- 1 *Discussiones Mathematicae*
2 *Probability and Statistics* xx (xxxx) 1–14

3 ESTIMATING THE EXTREMAL INDEX THROUGH THE
4 TAIL DEPENDENCE CONCEPT

5 MARTA FERREIRA

6 *Center of Mathematics*
7 *University of Minho, Portugal*

8 e-mail: msferreira@math.uminho.pt

9 **Abstract**

10 The extremal index θ is an important parameter in extreme value anal-
11 ysis when extending results from independent and identically distributed
12 sequences to stationary ones. A connection between the extremal index and
13 the tail dependence coefficient allows the introduction of new estimators.
14 The proposed ones are easy to compute and we analyze their performance
15 through a simulation study. Comparisons with other existing methods are
16 also presented. Case studies within environment are considered in the end.

17 **Keywords:** Extreme value theory, extremal index, tail dependence coeffi-
18 cient.

19 **2010 Mathematics Subject Classification:** 62G32.

20 1. INTRODUCTION

The central result in classical Extreme Value Theory states that, for an i.i.d. se-
quence, $\{X_n\}_{n \geq 1}$, having common distribution function (d.f.) F , if there are
constants $a_n > 0$ and $b_n \in \mathbb{R}$ such that,

$$P(\max(X_1, \dots, X_n) \leq a_n x + b_n) \xrightarrow[n \rightarrow \infty]{} G(x), \quad (1)$$

for some non degenerate function G , then it must be the Generalized Extreme
Value function (*GEV*),

$$G(x) = \exp(-(1 + \gamma x)^{-1/\gamma}), \quad 1 + \gamma x > 0, \quad \gamma \in \mathbb{R},$$

21 ($G(x) = \exp(-e^{-x})$ for $\gamma = 0$) and we say that F belongs to the max-domain of
22 attraction of G , in short, $F \in \mathcal{D}(G)$. The parameter γ , known as the tail index,

23 is a shape parameter determining the tail behavior of F : if $\gamma > 0$ we are in the
 24 domain of attraction Fréchet corresponding to a heavy tail, $\gamma < 0$ indicates the
 25 Weibull domain of attraction of light tails and $\gamma = 0$ means a Gumbel domain of
 26 attraction and an exponential tail.

27 In a multivariate context, it is possible to extend the convergence given in
 28 (1), but the class of models in the limit is much wider than model GEV. For
 29 simplicity, we consider the bivariate case, but everything can be rewritten for the
 30 more general d -variate case, $d \geq 2$. More precisely, let $\{(X_1^{(n)}, X_2^{(n)})\}_{n \geq 1}$ be a
 31 sequence of i.i.d. copies of the random pair (X_1, X_2) , with common d.f. \mathbf{F} , and
 32 let $M_j^{(n)} = \max_{1 \leq i \leq n} X_j^{(i)}$, $j = 1, 2$, be the maximum of each marginal. If there
 33 exist sequences of real constants $a_j^{(n)} > 0$ and $b_j^{(n)}$, for $j = 1, 2$ and $n \geq 1$, and a
 34 d.f. \mathbf{G} with non-degenerate margins, such that,

$$\begin{aligned} & P(M_1^{(n)} \leq a_1^{(n)}x_1 + b_1^{(n)}, M_2^{(n)} \leq a_2^{(n)}x_2 + b_2^{(n)}) \\ &= \mathbf{F}^n(a_1^{(n)}x_1 + b_1^{(n)}, a_2^{(n)}x_2 + b_2^{(n)}) \\ &\xrightarrow[n \rightarrow \infty]{} \mathbf{G}(x_1, x_2), \end{aligned}$$

35 for every continuity points of \mathbf{G} , then this latter is said to be a bivariate extreme
 36 value distribution (BEV) and is defined by expression

$$\mathbf{G}(x_1, x_2) = \exp[-l\{-\log G_1(x_1), -\log G_2(x_2)\}], \quad (2)$$

37 for some bivariate function l , where G_j , $j = 1, 2$, is the marginal d.f. of \mathbf{G} . In this
 38 case, we have that \mathbf{F} belongs to the max-domain of attraction of \mathbf{G} , in short $\mathbf{F} \in$
 39 $\mathcal{D}(\mathbf{G})$. The function l in (2), usually called *stable tail dependence function* is con-
 40 vex and homogeneous of order 1, and we have $\max(x_1, x_2) \leq l(x_1, x_2) \leq x_1 + x_2$,
 41 for all $(x_1, x_2) \in [0, \infty)^2$, where the upper limit corresponds to independence and
 42 the lower one means complete dependence (see, e.g. Beirlant *et al.* [2], Section
 43 8.2.2).

44
 45 The result in (1) may also be extended to study the maximum of a wide
 46 class of dependent processes, a more realistic assumption for several data. Here
 47 we concentrate on stationary sequences where the dependence is restricted by
 48 distributional mixing conditions.

The condition $D(u_n)$ of Leadbetter ([14], 1983), providing a short range depen-
 dence for which at long lags the extremes are independent, is sufficient to
 extend the result in (1) to stationary sequences. More precisely, for a stationary
 sequence $\{X_n\}_{n \geq 1}$ satisfying $D(u_n)$ with $u_n = a_n x + b_n$, we have that

$$P(\max(X_1, \dots, X_n) \leq u_n) \xrightarrow[n \rightarrow \infty]{} G^\theta(x), \quad (3)$$

49 where $0 \leq \theta \leq 1$ is the extremal index. The extremal index is the primary measure
 50 of extremal dependence in such processes, with $\theta = 1$ indicating independence at
 51 asymptotically high levels.

There are different interpretations of the extremal index. This concept, origi-
 nated in papers by Loynes ([15], 1965), O'Brien ([17], 1974) and developed in
 detail by Leadbetter ([14], 1983), reflects the effect of clustering of extreme ob-
 servations on the limiting distribution of the maximum. O'Brien (1987) proved
 that the presence of clustering affects the limiting distribution of block maxima:

$$P(\max(X_2, \dots, X_{r_n}) \leq u_n | X_1 > u_n) \xrightarrow{n \rightarrow \infty} \theta, \quad (4)$$

with r_n such that $r_n \rightarrow \infty$ and $r_n = o(n)$. Under a mixing condition slightly
 restrictive than $D(u_n)$, Hsing *et al.* ([13], 1988) showed that the limiting mean
 number of exceedances of u_n in an interval of length r_n is the inverse of the
 extremal index:

$$E\left[\sum_{i=1}^{r_n} \mathbb{1}_{\{X_i > u_n\}} \mid \sum_{i=1}^{r_n} \mathbb{1}_{\{X_i > u_n\}} \geq 1\right] \rightarrow \theta^{-1}, \quad (5)$$

with $\mathbb{1}(\cdot)$ the indicator function. By stationarity this property is satisfied for any
 block of r_n consecutive elements defined in the sequence. By rewriting (3) as

$$P(\max(X_1, \dots, X_n) \leq u_n) \xrightarrow{n \rightarrow \infty} e^{-\theta\tau(x)}, 0 < \tau(x) < \infty,$$

Ferro and Segers ([9], 2003) found that the process of inter-exceedance times nor-
 malized by exceedances of u_n follows a mixture of a point mass and an exponential
 distribution $Exp(\theta^{-1})$, i.e.,

$$P(\overline{F}(u_n)T(u_n) > t) \xrightarrow{n \rightarrow \infty} \theta e^{-\theta t}, t > 0, \quad (6)$$

52 with $T(u_n) = \min\{n \geq 1 : X_{n+1} > u_n | X_1 > u_n\}$, also under a slightly stricter
 53 mixing condition than $D(u_n)$.

54

Inference about θ has been extensively studied, with the most popular esti-
 mators being the runs method obtained from equation (4), the blocks method
 derived from (5) and the intervals method developed from (6). More precisely,
 the runs estimator is given by

$$\widehat{\theta}^{(R)} = (N)^{-1} \sum_{i=1}^{n-1} \mathbb{1}_{\{X_i > u\}} \mathbb{1}_{\{X_{i+1} \leq u\}} \cdots \mathbb{1}_{\{X_{i+r} \leq u\}},$$

where N is the total number of exceedances of a high threshold u . The blocks
 estimator for a sample divided into b blocks of length r (so $n \approx br$), can be stated
 as

$$\widehat{\theta}^{(B)} = \frac{\log(1 - C_n(u)/b)}{r \log(1 - N/n)}$$

where $C_n(u)$ is the number of blocks in which at least one exceedance of u occurs. After some considerations, the result in (6) yields the intervals estimator

$$\widehat{\theta}^{(I)} = \begin{cases} 1 \wedge \frac{2(\sum_{i=1}^{N-1} T_i)^2}{(N-1) \sum_{i=1}^{N-1} T_i^2} & , \text{ if } \max\{T_i : 1 \leq i \leq N-1\} \leq 2 \\ 1 \wedge \frac{2(\sum_{i=1}^{N-1} (T_i-1))^2}{(N-1) \sum_{i=1}^{N-1} (T_i-1)(T_i-2)} & , \text{ if } \max\{T_i : 1 \leq i \leq N-1\} > 2, \end{cases}$$

55 with T_i denoting the i th inter-exceedance time, $i = 1, \dots, N-1$. For a survey,
56 see for instance, Ancona-Navarrete and Tawn ([1], 2000) and Beirlant *et al.* ([2],
57 2004).

58

Imposing some convenient local dependence condition may eliminate the need for a cluster identification scheme as in the case of the blocks or the runs estimators. An example of such condition is the local dependence condition $D^{(2)}(u_n)$ of Chernick *et al.* (1991), which holds whenever

$$nP(X_j > u_n, X_{j+1} \leq u_n, M_{j+2, r_n} > u_n) \rightarrow 0, n \rightarrow \infty,$$

59 with $M_{i,j} = \max\{X_i, \dots, X_j\}$, for $i \leq j$ ($M_{i,j} = -\infty$ if $i > j$), the block sizes
60 sequence $\{r_n\}$ is such that $n/r_n \rightarrow \infty$ and condition $D(u_n)$ is simultaneously
61 satisfied. Condition $D^{(2)}(u_n)$ restricts the occurrence of an observation again ex-
62 ceeding the high threshold u_n after dropping below it within a cluster.

63

Under $D^{(2)}(u_n)$, and considering a log-likelihood based on the limiting d.f. obtained in (6), Süveges ([22], 2007) presents the maximum likelihood estimator

$$\widehat{\theta}^{(ML)} = \frac{\sum_{i=1}^{N-1} qS_i + N - 1 + N_C - \left[\left(\sum_{i=1}^{N-1} qS_i N - 1 + N_C \right)^2 - 8N_C \sum_{i=1}^{N-1} qS_i \right]^{1/2}}{2 \sum_{i=1}^{N-1} qS_i},$$

64 where q is the estimate of $\overline{F}(u)$, $S_i = T_i - 1$ and $N_C = \sum_{i=1}^{N-1} \mathbb{1}_{\{S_i \neq 0\}}$.

65

Considering a lightly stronger condition $D''(u_n)$ that restricts the occurrence of two or more upcrossings by imposing that $n \sum_{j=2}^{r_n-1} P(X_1 > u_n, X_j \leq u_n < X_{j+1}) \rightarrow 0$, as $n \rightarrow \infty$, Nandagopalan ([16], 1990) derives the estimator

$$\widehat{\theta}^{(N)} = \frac{\sum_{j=1}^{n-1} \mathbb{1}_{\{X_j \leq u < X_{j+1}\}}}{\sum_{j=1}^n \mathbb{1}_{\{X_j > u\}}},$$

66 for a suitable high threshold u . This is a special case of the runs estimator when
67 $r = 1$.

68

69 A recent result in Ferreira and Ferreira ([7], 2012a), allow us to state $\theta = 1 - \lambda$
 70 under condition $D^{(2)}(u_n)$, where λ is the tail dependence coefficient introduced
 71 by Sibuya ([21], 1960). Here we shall analyze the estimation of θ based on some λ
 72 estimation methodologies of the literature. This will be done through a simulation
 73 study. The performance of our approach will be also assessed by comparing with
 74 the simulation results obtained for the above exposed existing estimators of the
 75 extremal index. At the end, we illustrate with applications to real environmental
 76 data.

77 2. TAIL DEPENDENCE

78 The *tail-dependence coefficient* (TDC), usually denoted λ and first introduced in
 79 Sibuya ([21], 1960), measures the probability of occurring extreme values for one
 80 random variable (r.v.) given that another assumes an extreme value too, i.e.,

$$\lambda = \lim_{t \rightarrow \infty} P(F_1(X_1) > 1 - 1/t | F_2(X_2) > 1 - 1/t), \quad (7)$$

81 where F_1 and F_2 are the distribution functions (d.f.'s) of r.v.'s X_1 and X_2 , re-
 82 spectively. It characterizes the dependence in the tail of a random pair (X_1, X_2) ,
 83 in the sense that, $\lambda > 0$ corresponds to tail dependence whereas $\lambda = 0$ means tail
 84 independence.

85
 86 The relation $\theta = 1 - \lambda$ stated in Proposition 4 of Ferreira and Ferreira ([7],
 87 2012a) under the local dependence condition $D^{(2)}$, lead to new estimators for
 88 θ through the TDC. A wide study concerning TDC estimation is presented in
 89 Frahm *et al.* (2005). Parametric estimators are more accurate but may have
 90 disastrous performances under wrong model assumptions. Here we will focus on
 91 nonparametric approach.

92
 93 Schmidt and Stadtmüller ([19], 2006) considered the estimator based on (7)
 94 by plugging-in the respective empirical counterparts,

$$\widehat{\lambda}^{(SS)} \equiv \widehat{\lambda}^{(SS)}(k_n) = \frac{1}{k_n} \sum_{i=1}^n \mathbf{1}_{\{\widehat{F}_1(X_1) > 1 - \frac{k_n}{n}, \widehat{F}_2(X_2) > 1 - \frac{k_n}{n}\}}, \quad (8)$$

where \widehat{F}_j is the empirical d.f. of F_j , $j = 1, 2$, and $\{k_n\}$ is an intermediate sequence,
 i.e., $k_n \rightarrow \infty$ and $k_n/n \rightarrow 0$, as $n \rightarrow \infty$. Concerning estimation accuracy, some
 modifications of this latter may be used, like replacing the denominator n by $n+1$,
 i.e., considering

$$\widehat{F}_j(u) = \frac{1}{n+1} \sum_{k=1}^n \mathbf{1}_{\{X_j^{(k)} \leq u\}}$$

95 (for a discussion on this topic see, for instance, Beirlant et al. 2004). The choice
 96 of the value k in the sequence $\{k_n\}$ that allows the better trade-off between bias
 97 and variance is of major difficulty, since small values of k come along with a large
 98 variance whenever an increasing k results in a strong bias. The true value is
 99 usually located at a stable region of the plot $(k, \widehat{\lambda}^{(SS)}(k))$, for $1 \leq k < n$.

100 In order to avoid the variance-bias problem, we will use an heuristic procedure
 101 presented in Frahm *et al.* ([10], 2005), consisting on a “plateau finding algorithm”
 102 applied to a smoothed version of $(k, \widehat{\lambda}^{(SS)}(k))$, $1 \leq k < n$.

103

104 Based on the approach considered in Capéraà *et al.* ([3], 1997), which assumes
 105 that the underlying distribution approximates a BEV model given in (2), Frahm
 106 *et al.* ([10], 2005) have proposed the following estimator:

$$\widehat{\lambda}^{(CFG)} = 2 - 2 \exp \left\{ \frac{1}{n} \sum_{i=1}^n \log \left(\frac{\sqrt{\log \widehat{F}_1(X_1) \log \widehat{F}_2(X_2)}}{\log(\widehat{F}_1(X_1) \vee \widehat{F}_2(X_2))^{-2}} \right) \right\}, \quad (9)$$

107 where $x \vee y = \max(x, y)$. Another estimator developed in Ferreira and Ferreira
 108 ([8], 2012b) under the same assumption but with a simpler form, is given by

$$\widehat{\lambda}^{(FF)} = 3 - (1 - \overline{\widehat{F}_1(X_1) \vee \widehat{F}_2(X_2)})^{-1},$$

109 where $\overline{\widehat{F}_1(X_1) \vee \widehat{F}_2(X_2)}$ is the sample mean of $\widehat{F}_1(X_1) \vee \widehat{F}_2(X_2)$, i.e.,

$$\overline{\widehat{F}_1(X_1) \vee \widehat{F}_2(X_2)} = \frac{1}{n} \sum_{i=1}^n [\widehat{F}_1(X_1^{(i)}) \vee \widehat{F}_2(X_2^{(i)})].$$

110 For a discussion about the asymptotic properties of these estimators see, respec-
 111 tively, Genest and Segers ([11], 2009) and Ferreira ([6], 2013).

112

113 From now on, we will use notation $\widehat{\theta}^{(SS)}$, $\widehat{\theta}^{(CFG)}$ and $\widehat{\theta}^{(FF)}$, whenever we refer
 114 to estimators $\widehat{\lambda}^{(SS)}$, $\widehat{\lambda}^{(CFG)}$ and $\widehat{\lambda}^{(FF)}$, that is,

$$\widehat{\theta}^{(SS)} = 1 - \widehat{\lambda}^{(SS)}, \quad \widehat{\theta}^{(CFG)} = 1 - \widehat{\lambda}^{(CFG)} \quad \text{and} \quad \widehat{\theta}^{(FF)} = 1 - \widehat{\lambda}^{(FF)}.$$

115

3. SIMULATION STUDY

116 We are going to analyze the performance of the estimators described above,
 117 through a simulation study based on the following models:

- 118 • Independent sequence which have $\theta = 1$ (with unit Fréchet margins).
- 119 • Markov Gaussian dependence process, $Z_j = \alpha Z_{j-1} + \epsilon_j$, where the ϵ_j are
 120 i.i.d. $N(0, 1 - \alpha^2)$ r.v.'s, for $j \geq 2$ and Z_1 is $N(0, 1)$ distributed. This process
 121 has $\theta = 1$ and shall be denoted AR .

- Bivariate extreme value Markov process with logistic dependence function, i.e.,

$$P(X_j \leq x, X_{j+1} \leq y) = \exp(-(x^{1/\alpha} + y^{1/\alpha})^\alpha).$$

122 As in Ancona-Navarrete and Tawn ([1], 2000), we consider the dependence
123 parameter $\alpha = 0.5$ which gives $\theta = 0.328$, and denote the process *BEV*.

- Autoregressive maximum process, $X_i = \alpha X_{i-1} \vee \epsilon_i$, where $0 < \alpha < 1$ and $\{\epsilon_i\}$ are i.i.d. r.v.'s with d.f. $F_\epsilon(x) = \exp(-(1 - \alpha)/x)$, $x > 0$. This process has $\theta = 1 - \alpha$. We consider $\alpha = 0.5$ and hence $\theta = 0.5$, and denote the process *MAR*.

- Moving maxima process, $X_i = \bigvee_{j=0, \dots, m} \alpha_j \epsilon_{i-j}$, with $\sum_{j=0}^m \alpha_j = 1$ and $\alpha_j \geq 0$, $\{\epsilon_i\}$ are i.i.d. unit F chet r.v.'s. This process has $\theta = \bigvee_{j=0, \dots, m} \alpha_j$. We consider $m = 3$, $\alpha_1 = \alpha_2 = 0.2$, $\alpha_0 = \alpha_3 = 0.3$ and so $\theta = 0.3$, and denote the process *MM*.

132 We consider samples of size $n = 10000$ and compare the estimators using
133 the absolute mean bias and the root mean square error (rmse) criteria, obtained
134 using 200 independent replications of the estimation procedures. The results
135 of the proposed estimators, $\hat{\theta}^{(FF)}$, $\hat{\theta}^{(CFG)}$ and $\hat{\theta}^{(SS)}$, are presented in Table 1.
136 For comparison, we also include the simulation results obtained from estimators
137 $\hat{\theta}^{(ML)}$ and $\hat{\theta}^{(N)}$ derived under similar local dependence conditions, i.e., $D^{(2)}$ and
138 D'' , respectively (see Table 2). The estimates derived from the runs, the blocks
139 and the intervals methods were also computed and can be found in Table 3. We
140 remark that the values considered for the number of blocks/runs were derived
141 through additional simulation studies conducted in Ancona-Navarrete and Tawn,
142 ([1], 2000).

143 Observe that the worst performance of the estimators coincides with the AR
144 process. In this case, estimator $\hat{\theta}^{(SS)}$ followed by $\hat{\theta}^{(ML)}$, $\hat{\theta}^{(N)}$, $\hat{\theta}^{(B)}$ and $\hat{\theta}^{(I)}$
145 for $u = q_{0.99}$ exceed the remaining. In particular, the bad performance of the
146 proposed estimators $\hat{\theta}^{(FF)}$ and $\hat{\theta}^{(CFG)}$ is due to the bad behavior of the respective
147 tail dependence coefficient estimators $\hat{\lambda}^{(FF)}$ in (8) and $\hat{\lambda}^{(CFG)}$ in (9) under tail
148 independent non-BEV models, i.e., models for which $\lambda = 0$ and whose dependence
149 structure for consecutive pairs can not be formulated as in (2), such as the case of
150 AR (see Ferreira, [6] 2013). Indeed, estimators $\hat{\theta}^{(FF)}$ and $\hat{\theta}^{(CFG)}$ are not robust.
151 They present the worst performances also within the BEV and MM processes,
152 missing the $D^{(2)}$ condition. Therefore, concerning robustness, the best of the
153 three here proposed estimators is $\hat{\theta}^{(SS)}$, which only demands the $D^{(2)}$ condition
154 and behaves better whenever this latter is violated (see the results for BEV and
155 AR in Table 1). All the estimators behave quite well in the MAR process, with
156 the best performances occurring for our proposals $\hat{\theta}^{(FF)}$ and $\hat{\theta}^{(CFG)}$, as well as,
157 for $\hat{\theta}^{(ML)}$ and $\hat{\theta}^{(N)}$ with $u = q_{0.99}$. We remark that this process satisfies condition

158 $D^{(2)}$ as well as the BEV dependence assumption (see, e.g., Ferreira and Ferreira
 159 [7] 2012a and Ancona-Navarrete and Tawn [1] 2000). Regarding the MM case,
 160 the best performance lies with the runs, blocks and intervals estimators, which is
 161 not surprising since it is easy to identify independent clusters in this process.

Table 1. Sample absolute mean bias and rmse (in brackets) of estimators $\hat{\theta}^{(FF)}$, $\hat{\theta}^{(CFG)}$
 and $\hat{\theta}^{(SS)}$.

	$\hat{\theta}^{(FF)}$	$\hat{\theta}^{(CFG)}$	$\hat{\theta}^{(SS)}$
Indep.	0.00 (0.010)	0.00 (0.010)	0.05 (0.050)
AR	0.40 (0.403)	0.36 (0.364)	0.12 (0.131)
BEV	0.09 (0.088)	0.09 (0.089)	0.06 (0.063)
MAR	0.00 (0.010)	0.00 (0.010)	0.03 (0.041)
MM	0.10 (0.100)	0.10 (0.101)	0.07 (0.073)

Table 2. Sample absolute mean bias and rmse (in brackets) of estimators $\hat{\theta}^{(ML)} \equiv \hat{\theta}_u^{(ML)}$
 and $\hat{\theta}^{(N)} \equiv \hat{\theta}_u^{(N)}$, by considering thresholds $u = q_{0.95}, q_{0.99}$, respectively, the empirical
 quantiles 0.95 and 0.99.

	$\hat{\theta}_{q_{0.95}}^{(ML)}$	$\hat{\theta}_{q_{0.99}}^{(ML)}$	$\hat{\theta}_{q_{0.95}}^{(N)}$	$\hat{\theta}_{q_{0.99}}^{(N)}$
Indep.	0.05 (0.045)	0.01 (0.000)	0.05 (0.055)	0.01 (0.000)
AR	0.24 (0.237)	0.13 (0.130)	0.24 (0.245)	0.13 (0.134)
BEV	0.08 (0.089)	0.10 (0.114)	0.08 (0.077)	0.09 (0.114)
MAR	0.01 (0.032)	0.00 (0.045)	0.02 (0.032)	0.00 (0.045)
MM	0.10 (0.095)	0.11 (0.118)	0.09 (0.089)	0.11 (0.114)

162 3.1. Case studies

163 3.1.1. Wooster temperatures

164 We consider the daily minimum temperatures (in degrees Fahrenheit) at
 165 Wooster (Ohio), from 1983 to 1988, more precisely, the period of November-
 166 February winter months in order to achieve some stationarity (see Figure 1).
 167 This series was analyzed in Coles ([5], 2001) and blocks estimates were computed

Table 3. Sample absolute mean bias and rmse (in brackets) of runs estimator $\widehat{\theta}^{(R)} \equiv \widehat{\theta}_u^{(R)}$, blocks estimator $\widehat{\theta}^{(B)} \equiv \widehat{\theta}_u^{(B)}$ and intervals estimator $\widehat{\theta}^{(I)} \equiv \widehat{\theta}_u^{(I)}$ by considering thresholds $u = q_{0.95}, q_{0.99}$, respectively, the empirical quantiles 0.95 and 0.99. In the blocks and runs estimators it was used the suggested number of runs/blocks in Ancona-Navarrete and Tawn ([1], 2000).

	$\widehat{\theta}_{q_{0.95}}^{(R)}$	$\widetilde{\theta}_{q_{0.99}}^{(R)}$	$\widehat{\theta}_{q_{0.95}}^{(B)}$	$\widehat{\theta}_{q_{0.99}}^{(B)}$	$\widetilde{\theta}_{q_{0.95}}^{(I)}$	$\widehat{\theta}_{q_{0.99}}^{(I)}$
Indep.	0.05 (0.055)	0.01 (0.000)	0.00 (0.008)	0.01 (0.014)	0.01 (0.000)	0.03 (0.055)
AR	0.37 (0.370)	0.19 (0.183)	0.24 (0.241)	0.13 (0.135)	0.22 (0.224)	0.13 (0.155)
BEV	0.03 (0.028)	0.04 (0.063)	0.07 (0.064)	0.03 (0.090)	0.04 (0.055)	0.03 (0.084)
MAR	0.02 (0.032)	0.00 (0.045)	0.03 (0.044)	0.02 (0.034)	0.03 (0.045)	0.03 (0.084)
MM	0.03 (0.027)	0.00 (0.031)	0.02 (0.030)	0.03 (0.041)	0.03 (0.045)	0.02 (0.055)

168 for the extremal index. In particular, it was considered the threshold $u = -10$
 169 with number of blocks $b = 20, 31$ leading to, respectively, $\widehat{\theta}^{(B)} = 0.27, 0.42$.

170 Since we have a sample of minimum values we assume that an approximation
 171 to a BEV model dependence structure between consecutive pairs is plausible. In
 172 order to check condition $D^{(2)}$, we use the empirical methodology of Süveges ([22],
 173 2007) by calculating the proportion of anti- $D^{(2)}$ events among the exceedances for
 174 a range of block sizes and thresholds:

$$p(u, r) = \frac{\sum_{j=1}^n \mathbb{1}_{\{X_j > u, X_{j+1} \leq u, M_{j+2, r} > u\}}}{\sum_{j=1}^n \mathbb{1}_{\{X_j > u\}}}.$$

175 Observe in Figure 2 that $p(u, r) \approx 0$ as u and r increase, which leads to an informal
 176 validation of $D^{(2)}$. Thus we assume the validity of estimators $\widehat{\theta}^{(ML)}$ and $\widehat{\theta}^{(N)}$, as
 177 well as the here presented $\widehat{\theta}^{(FF)}$, $\widehat{\theta}^{(CFG)}$ and $\widehat{\theta}^{(SS)}$.

178 In Figure 3 are plotted, for several thresholds, the obtained estimates from
 179 $\widehat{\theta}^{(B)}$ (for $b = 20, 31$), $\widehat{\theta}^{(R)}$ (for $r = 2, 4$) and $\widehat{\theta}^{(I)}$ (left), and from $\widehat{\theta}^{(ML)}$ and
 180 $\widehat{\theta}^{(N)}$ (right). Considering again $u = -10$, we have $\widehat{\theta}^{(R)} = 0.35, 0.23$, for $r =$
 181 $2, 4$, respectively, $\widehat{\theta}^{(I)} = 0.26$, $\widehat{\theta}^{(ML)} = 0.43$ and $\widehat{\theta}^{(N)} = 0.4$. By applying our
 182 estimators, we have $\widehat{\theta}^{(FF)} = 0.36$, $\widehat{\theta}^{(CFG)} = 0.38$ and $\widehat{\theta}^{(SS)} = 0.38$, more closer to
 183 the ones obtained for $\widehat{\theta}^{(ML)}$, $\widehat{\theta}^{(N)}$, $\widehat{\theta}^{(B)}$ with $b = 31$ and $\widehat{\theta}^{(R)}$ with $r = 2$.

184 3.1.2. Ozone pollution

185 We now consider $n = 120$ weekly maxima of hourly averages of ozone concen-
 186 trations measured in parts per million, in the San Francisco bay area, San Jose,
 187 available in the package Xtremes (Reiss and Thomas, [18] 2007). These data have




Figure 1. Negated Wooster daily minimum temperatures (in degrees Fahrenheit) on the left, and considering winters only on the right.




Figure 2. The observed proportion of anti $D^{(2)}(u_n)$ condition for winters negated Wooster daily minimum temperatures (in degrees Fahrenheit).

188 been analyzed in Gomes *et al.* ([12], 2008) and Sebastião *et al.* ([20], 2013). We as-
 189 sume stationarity as in the latter reference (see also Figure 4). Gomes *et al.* ([12],
 190 2008) argued the plausibility of condition $D^{(2)}$ to hold, based on the fact that
 191 these type of meteorological data is usually modeled by processes that satisfy this
 192 latter. See also Figure 5 and the conclusions in Sebastião *et al.* ([20], 2013) which
 193 corroborates this assumption. A sample of maxima makes us comfortable with
 194 the hypothesis of an underlying model approximately BEV for consecutive pairs
 195 of observations. The extremal index was evaluated in 0.7 in Gomes *et al.* ([12],
 196 2008). In what concerns estimators $\hat{\theta}^{(FF)}$, $\hat{\theta}^{(CFG)}$ and $\hat{\theta}^{(SS)}$, we have obtained,




Figure 3. The blocks, runs and intervals estimators (left) and the maximum likelihood and Nandagopalan estimators (right), against threshold, for winters negated Wooster daily minimum temperatures (in degrees Fahrenheit).

197 respectively, 0.74, 0.74 and 0.75. In analyzing Figure 6, the value 0.7 is a possible
 198 estimate, except in the case of the blocks estimator.




Figure 4. Weekly maxima of hourly averages of ozone concentrations (in parts per million), in the San Francisco bay area, San Jose.




Figure 5. The observed proportion of anti $D^{(2)}(u_n)$ condition for weekly maxima of hourly averages of ozone concentrations (in parts per million), in the San Francisco bay area, San Jose.




Figure 6. The blocks, runs and intervals estimators against threshold for weekly maxima of hourly averages of ozone concentrations (in parts per million), in the San Francisco bay area, San Jose.

4. CONCLUDING REMARKS

200 Here we have considered new estimators for the extremal index based on the
 201 tail dependence coefficient estimation, under the validity of condition $D^{(2)}(u_n)$ of
 202 Chernick *et al.* ([4], 1991). Estimators $\hat{\theta}^{(FF)}$ and $\hat{\theta}^{(CFG)}$ also require that the un-

203 derlying distribution of consecutive random pairs can be approximated by a BEV
 204 model dependence structure. These latter are not robust whenever one of the two
 205 assumptions is breached. On the other hand, estimator $\hat{\theta}^{(SS)}$ presents compara-
 206 ble biases and rmse's to estimators $\hat{\theta}^{(ML)}$ and $\hat{\theta}^{(N)}$ which were also derived under
 207 condition $D^{(2)}(u_n)$, in some cases, even outperforming these two latter. Estimator
 208 $\hat{\theta}^{(SS)}$ has also comparable performances to the ones of the runs and the blocks
 209 estimators in some models. Observe that it depends on only one parameter (the
 210 number k of observations to consider in the estimation), while the runs and blocks
 211 estimators depend on a high threshold u and the number of runs r or blocks b ,
 212 respectively. Since $D^{(2)}(u_n)$ is a crucial requisite in the new approach, it is impor-
 213 tant to develop a more reliable diagnostic statistical tool for this condition. This
 214 will be the aim of a future work.
 215

216 Acknowledgements

217

218 The author was financed by FEDER Funds through "Programa Operacional Fac-
 219 tores de Competitividade - COMPETE" and by Portuguese Funds through FCT -
 220 "Fundação para a Ciência e a Tecnologia", within the project PEst-OE/MAT/UI0013/2014.

221

REFERENCES

- 222 [1] M.A. Ancona-Navarrete and J.A. Tawn, *A comparison of methods for esti-*
 223 *inating the extremal index*, *Extremes* **3** (2000) 5–38.
- 224 [2] J. Beirlant, Y. Goegebeur, J. Segers, J. Teugels, *Statistics of Extremes: The-*
 225 *ory and Application* (John Wiley, 2004).
- 226 [3] P. Capéraà, A.L. Fougères and C. Genest, *A nonparametric estimation pro-*
 227 *cedure for bivariate extreme value copulas*, *Biometrika* **84** (1997) 567–577.
- 228 [4] M.R. Chernick, T. Hsing and W.P. McCormick, *Calculating the extremal*
 229 *index for a class of stationary sequences*, *Adv. Appl. Probab.* **23** (1991) 835–
 230 850.
- 231 [5] S.G. Coles, *An Introduction to Statistical Modelling of Extreme Values* (Lon-
 232 don, Springer, 2001).
- 233 [6] M. Ferreira, *Nonparametric estimation of the tail dependence coefficient*,
 234 *REVSTAT* **11(1)** (2013) 1–16.
- 235 [7] M. Ferreira and H. Ferreira, *On extremal dependence: some contributions*,
 236 *TEST* **21(3)** (2012a) 566–583.

- 237 [8] H. Ferreira and M. Ferreira, *On extremal dependence of block vectors*, Ky-
238 bernetika **48(5)** (2012b) 988–1006.
- 239 [9] C.A. Ferro and J. Segers, *Inference for clusters of extremes*, J. R. Stat. Soc.
240 Ser. B Stat. Methodol. **65** (2003) 545–556.
- 241 [10] G. Frahm, M. Junker and R. Schmidt, *Estimating the tail-dependence coeffi-*
242 *cient: properties and pitfalls*, Insurance Math. Econom. **37(1)** (2005) 80–100.
- 243 [11] C. Genest and J. Segers J., *Rank-based inference for bivariate extreme-value*
244 *copulas*, Ann. Statist. **37** (2009) 2990–3022.
- 245 [12] M.I. Gomes, A. Hall and C. Miranda, *Subsampling techniques and the jack-*
246 *knife methodology in the estimation of the extremal index*, J. Stat. Comput.
247 Simul. **52** (2008) 2022–2041.
- 248 [13] T. Hsing, J. Husler and M.R. Leadbetter, *On the exceedance point process*
249 *for a stationary sequence*, Probab. Theory Related Fields **78** (1988) 97–112.
- 250 [14] M.R. Leadbetter, *Extremes and local dependence in stationary sequences*, Z.
251 Wahrsch. Ver. Geb. **65** (1983) 291–306.
- 252 [15] R.M. Loynes, *Extreme Values in Uniformly Mixing Stationary Stochastic Pro-*
253 *cesses*, Annals of Mathematical Statistics **36** (1965) 993–999.
- 254 [16] S. Nandagopalan, *Multivariate extremes and estimation of the extremal index*
255 (Ph.D. Thesis, University of North Carolina at Chapel Hill, 1990).
- 256 [17] G.L. O’Brien, *The maximum term of uniformly mixing stationary sequences*,
257 Z. Wahrsch. Ver. Geb. **30** (1974) 57–63.
- 258 [18] R.D. Reiss, M. Thomas, *Statistical analysis of extreme values with appli-*
259 *cations to insurance, finance, hydrology and other fields* (Birkhäuser, Basel,
260 2007).
- 261 [19] R. Schmidt and U. Stadtmüller, *Nonparametric estimation of tail dependence*,
262 Scandinavian J. Statist. **33** (2006) 307–335.
- 263 [20] J.R. Sebastião, A.P. Martins, H. Ferreira and L. Pereira, *Estimating the*
264 *upcrossings index*, TEST **22(4)** (2013) 549–579.
- 265 [21] M. Sibuya, *Bivariate extreme statistics*, Ann. Inst. Stat. Math. **11** (1960)
266 195–210.
- 267 [22] M. Süveges, *Likelihood estimation of the extremal index*, Extremes **10** (2007)
268 41–55.