

ORBIT DECIDABILITY AND THE CONJUGACY PROBLEM FOR SOME EXTENSIONS OF GROUPS

O. BOGOPOLSKI, A. MARTINO AND E. VENTURA

ABSTRACT. Given a short exact sequence of groups with certain conditions, $1 \rightarrow F \rightarrow G \rightarrow H \rightarrow 1$, we prove that G has solvable conjugacy problem if and only if the corresponding action subgroup $A \leq \text{Aut}(F)$ is orbit decidable. From this, we deduce that the conjugacy problem is solvable, among others, for all groups of the form $\mathbb{Z}^2 \rtimes F_m$, $F_2 \rtimes F_m$, $F_n \rtimes \mathbb{Z}$, and $\mathbb{Z}^n \rtimes_A F_m$ with virtually solvable action group $A \leq \text{GL}_n(\mathbb{Z})$. Also, we give an easy way of constructing groups of the form $\mathbb{Z}^4 \rtimes F_n$ and $F_3 \rtimes F_n$ with unsolvable conjugacy problem. On the way, we solve the twisted conjugacy problem for virtually surface and virtually polycyclic groups, and give an example of a group with solvable conjugacy problem but unsolvable twisted conjugacy problem. As an application, an alternative solution to the conjugacy problem in $\text{Aut}(F_2)$ is given.

1. INTRODUCTION

Let G be a group, and $u, v \in G$. The symbol \sim will be used to denote standard conjugacy in G ($u \sim v$ if there exists $x \in G$ such that $v = x^{-1}ux$). In this paper, we shall work with a twisted version, which is another equivalence relation on G . Given an automorphism $\varphi \in \text{Aut}(G)$, we say that u and v are φ -twisted conjugated, denoted $u \sim_\varphi v$, if there exists $x \in G$ such that $v = (x\varphi)^{-1}ux$. Of course, \sim_{Id} coincides with \sim . Reidemeister was the first author considering the relation \sim_φ (see [35]), which has an important role in modern Nielsen fixed point theory. A few interesting references can be found in [17], where it is proven that the number of φ -twisted conjugacy classes in a non-elementary word-hyperbolic group is always infinite; [6], where an algorithm is given for recognizing φ -twisted conjugacy classes in free groups; and [18], where the notion of twisted conjugacy separability is analyzed.

Precisely, the recognition of twisted conjugacy classes is one of the main problems focused on in the present paper. The twisted conjugacy problem for a group G consists on finding an algorithm which, given an automorphism $\varphi \in \text{Aut}(G)$ and two elements $u, v \in G$, decides whether $v \sim_\varphi u$

or not. Of course, a positive solution to the twisted conjugacy problem automatically gives a solution to the (standard) conjugacy problem, which in turn provides a solution to the word problem. The existence of a group G with solvable word problem but unsolvable conjugacy problem is well known (see [30]). In this direction, one of the results here is the existence of a group with solvable conjugacy problem, but unsolvable twisted conjugacy problem (see Corollary 4.9 below).

Let us mention the real motivation and origin of the present work. Several months ago, the same three authors (together with O. Maslakova) wrote the paper [6], where they solved the twisted conjugacy problem for finitely generated free groups and, as a corollary, a solution to the conjugacy problem for free-by-cyclic groups was also obtained. A key ingredient in this second result was Brinkmann's theorem, saying that there is an algorithm such that, given an automorphism α of a finitely generated free group and two elements u and v , decides whether v is conjugate to some iterated image of u , $v \sim u\alpha^k$ (see [10]). At some point, Susan Hermiller brought to our attention an article due to Miller, [30], where he constructed a free-by-free group with unsolvable conjugacy problem. It turns out that the full proof in [6] extends perfectly well to the bigger family of free-by-free groups without any problem, except for a single step in the argumentation: at the point in [6] where we used Brinkmann's result, a much stronger problem arises, which we call *orbit decidability* (see below for definitions). This way, orbit decidability is really the unique obstruction when extending the arguments from free-by-cyclic to free-by-free groups.

Hence, we can deduce that a free-by-free group has solvable conjugacy problem if and only if the corresponding orbit decidability problem is solvable. This idea is formally expressed in Theorem 3.1, the main result of the present paper. In fact, Theorem 3.1 works not only for free-by-free groups, but for an even bigger family of groups. And, of course, when we restrict ourselves to the free-by-cyclic case, orbit decidability becomes exactly the algorithmic problem already solved by Brinkmann's theorem.

More precisely, Theorem 3.1 talks about a short exact sequence $1 \rightarrow F \rightarrow G \rightarrow H \rightarrow 1$ with some conditions on F and H . And states that the group G has solvable conjugacy problem if and only if the action subgroup $A_G \leq \text{Aut}(F)$ is *orbit decidable*. Here, A_G is the group of elements in G acting by right conjugation on F ; and being orbit decidable means that, given $u, v \in F$, we can algorithmically decide whether $v \sim u\alpha$ for some $\alpha \in A_G$. Of course, if we take F to be free and H to be infinite cyclic, we get free-by-cyclic groups for G ; and the action subgroup being orbit decidable is precisely Brinkmann's theorem (see Subsection 6.2). The conditions imposed to the groups F and H are the twisted conjugacy problem for F ,

and the conjugacy problem plus a technical condition about centralizers for H .

In light of Theorem 3.1, it becomes interesting, first, to collect groups F where the twisted conjugacy problem can be solved. And then, for every such group F , to study the property of orbit decidability for subgroups of $Aut(F)$: every orbit decidable (undecidable) subgroup of $Aut(F)$ will correspond to extensions of F having solvable (unsolvable) conjugacy problem. After proving Theorem 3.1, this is the main goal of the paper. We first show how to solve the twisted conjugacy problem for surface and polycyclic groups (in fact, the solution of these problems follows easily from the solution of the conjugacy problem for polycyclic and for fundamental groups of 3-manifolds, respectively). We also prove that the solvability of this problem goes up to finite index for these classes of groups. Then, we concentrate on the simplest examples of groups with solvable twisted conjugacy problem, namely finitely generated free (say F_n) and free abelian (\mathbb{Z}^n). On the positive side, we find several orbit decidable subgroups of $Aut(F_n)$ and $Aut(\mathbb{Z}^n) = GL_n(\mathbb{Z})$, with the corresponding free-by-free and free abelian-by-free groups with solvable conjugacy problem. Probably the most interesting result in this direction is Proposition 6.9 saying that virtually solvable subgroups of $GL_n(\mathbb{Z})$ are orbit decidable; or, say in a different way, orbit undecidable subgroups of $GL_n(\mathbb{Z})$ must contain F_2 . On the negative side, we establish a source of orbit undecidability, certainly related with a special role of F_2 inside the automorphism group. In particular, this will give orbit undecidable subgroups in $Aut(F_n)$ for $n \geq 3$, which will correspond to Miller's free-by-free groups with unsolvable conjugacy problem; and orbit undecidable subgroups in $GL_n(\mathbb{Z})$ for $n \geq 4$, which will correspond to the first known examples of \mathbb{Z}^4 -by-free groups with unsolvable conjugacy problem.

All over the paper, F_n denotes the free group of finite rank $n \geq 0$; and F , G and H stand for arbitrary groups. Having Theorem 3.1 in mind, we will use one or another of these letters to make clear which position in the short exact sequence the reader should think about. For example, typical groups to put in the first position of the sequence are free groups (F_n), and in the third position are hyperbolic groups (H). We will write morphisms as acting on the right, $x \mapsto x\varphi$. In particular, the inner automorphism given by right conjugation by $g \in G$ is denoted $\gamma_g: G \rightarrow G$, $x \mapsto x\gamma_g = g^{-1}xg$. As usual, $End(G)$ denotes the monoid of endomorphisms of G , $Aut(G)$ the group of automorphisms of G , $Inn(G)$ denotes the group of inner automorphisms, and $Out(G) = Aut(G)/Inn(G)$. Finally, we write $[u, v] = u^{-1}v^{-1}uv$, and $C_G(u) = \{g \in G \mid gu = ug\} \leq G$ for the centralizer of an element u .

As usual, given a group $F = \langle X \mid R \rangle$ and m automorphisms $\varphi_1, \dots, \varphi_m \in \text{Aut}(F)$, the free extension of F by $\varphi_1, \dots, \varphi_m$ is the group

$$F \rtimes_{\varphi_1, \dots, \varphi_m} F_m = \langle X, t_1, \dots, t_m \mid R, t_i^{-1} x t_i = x \varphi_i \quad (x \in X, i = 1, \dots, m) \rangle.$$

Such a group is called *F-by-[f.g. free]*. In particular, if $m = 1$ we call it *F-by-cyclic* and denote by $F \rtimes_{\varphi_1} \mathbb{Z}$. It is well known that a group G is *F-by-[f.g. free]* if and only if it has a normal subgroup isomorphic to F , with finitely generated free quotient G/F (i.e. if and only if it is the middle term in a short exact sequence of the form $1 \rightarrow F \rightarrow G \rightarrow F_n \rightarrow 1$). Above, when talking about “free-by-free” and “free abelian-by-free” we meant *[f.g. free]-by-[f.g. free]* and *[f.g. abelian]-by-[f.g. free]*, respectively. To avoid confusions, we shall keep these last names; they make an appropriate reference to finite generation, and they stress the fact that parenthesis are relevant (note, for example, that surface groups are both free-by-cyclic and finitely generated, but most of them are not *[f.g. free]-by-cyclic*).

The paper is structured as follows. In Section 2 we discuss some preliminaries concerning algorithmic issues, setting up the notation and names used later. Section 3 contains the first part of the work, developing the relation between the conjugacy problem and the concept of orbit decidability. In Section 4 the applicability of Theorem 3.1 is enlarged, by finding more groups with solvable twisted conjugacy problem (Subsection 4.1), and by proving that many hyperbolic groups have small centralizers in an algorithmic sense (Subsection 4.2). Then, Section 5 is dedicated to solve the conjugacy problem for $\text{Aut}(F_2)$; this problem was already known to be solvable (see [3] and [4]), but we present here an alternative solution to illustrate an application of the techniques developed in the paper. Sections 6 and 7 are dedicated, respectively, to several positive and negative results, namely orbit decidable subgroups corresponding to extensions with solvable conjugacy problem, and orbit undecidable subgroups corresponding to extensions with unsolvable conjugacy problem. Both, in the free abelian case, and in the free case. Finally, Section 8 is dedicated to summarize and comment on some questions and open problems.

2. ALGORITHMIC PRELIMINARIES

From the algorithmic point of view, the sentence “let G be a group” is not precise enough: the algorithmic behaviour of G may depend on how G is given to us. For the purposes of this paper, we assume that a group will always be given to us in an *algorithmic* way: elements must be represented by finite objects and multiplication and inversion must be doable in an algorithmic fashion; also, morphisms between groups, $\alpha: F \rightarrow G$, are to be

represented by a finite amount of information in such a way that one can algorithmically compute images of elements in F .

If G is finitely presented, a natural way (though not the unique one) consists in giving the group G by a finite presentation, $G = \langle x_1, \dots, x_n \mid r_1, \dots, r_m \rangle$; here, elements of G are represented by words in the x_i 's, multiplication and inversion are the standard ones in the free group (modulo relations), and morphisms are given by images of generators.

Let G and $\phi: G \rightarrow G$ be a group and an automorphism given in an algorithmic way. Also, let $F \leq G$ be a subgroup. The following are interesting algorithmic problems in group theory (the first two typically known as *Dehn problems*):

- the **word problem** for G , denoted $WP(G)$: given an element $w \in G$, decide whether it represents the trivial element of G . It is well known that there exists finitely presented groups with unsolvable word problem (see [31] or [7]).
- the **conjugacy problem** for G , denoted $CP(G)$: given two elements $u, v \in G$, decide whether they are conjugate to each other in G . Clearly, a solution for the $CP(G)$ immediately gives a solution for the $WP(G)$, and it is well known the existence of finitely presented groups with solvable word problem but unsolvable conjugacy problem (see, for example, [11] or [30]).
- the **ϕ -twisted conjugacy problem** for G , denoted $TCP_\phi(G)$: given two elements $u, v \in G$, decide whether they are ϕ -twisted conjugate to each other in G (i.e. whether $v = (g\phi)^{-1}ug$ for some $g \in G$). Note that $TCP_{Id}(G)$ is $CP(G)$.
- the **(uniform) twisted conjugacy problem** for G , denoted $TCP(G)$: given an automorphism $\phi \in Aut(G)$ and two elements $u, v \in G$, decide whether they are ϕ -twisted conjugate to each other in G . This is part of a more general problem posted by G. Makanin in Question 10.26(a) of [23]. Obviously, a solution for the $TCP(G)$ immediately gives a solution for the $TCP_\phi(G)$ for all $\phi \in Aut(G)$ (in particular, $CP(G)$ and $WP(G)$). In section 4.1, we give an example of a finitely presented group with solvable conjugacy problem but unsolvable twisted conjugacy problem.
- the **membership problem** for F in G , denoted $MP(F, G)$: given an element $w \in G$ decide whether it belongs to F or not. There are well-known pairs (F, G) with unsolvable $MP(F, G)$ (see [29] or [30]).

Conjugacy and twisted conjugacy problems have the “search” variants, respectively called the **conjugacy search problem**, $CSP(G)$, and the

twisted conjugacy search problem, $\text{TCSP}(G)$, for G . They consist on additionally finding a conjugator (or twisted-conjugator) in case it exists.

When groups are given by finite presentations and morphisms by images of generators, the “yes” parts of the listed problems are always solvable. Let $G = \langle x_1, \dots, x_n \mid r_1, \dots, r_m \rangle$ and $H = \langle y_1, \dots, y_{n'} \mid s_1, \dots, s_{m'} \rangle$ be two finitely presented groups, $\phi: G \rightarrow H$ be a morphism (given by $x_i \mapsto w_i(y_1, \dots, y_{n'})$, $i = 1, \dots, n$), and let $F \leq G$ be a subgroup given by a finite set of generators $\{f_1(x_1, \dots, x_n), \dots, f_t(x_1, \dots, x_n)\}$. We have:

- the **“yes” part of the word problem** for G , denoted $\text{WP}^+(G)$: given an element of G as a (reduced) word on the generators, $w(x_1, \dots, x_n) \in G$, which is known to be trivial in G , find an expression of w as a product of conjugates of the relations r_i . No matter if $\text{WP}(G)$ is solvable or not, $\text{WP}^+(G)$ is always solvable by brute force: enumerate the normal closure $R = \langle\langle r_1, \dots, r_m \rangle\rangle$ in the free group $\langle x_1, \dots, x_n \mid \rangle$, and check one by one whether its elements equal w as a word after reduction; since we know that $w =_G 1$, the process will eventually terminate. Note that, without the assumption, this is not an algorithm because if $w \neq_G 1$ then it works forever without giving any answer.
- the **“yes” part of the conjugacy problem** for G , denoted $\text{CP}^+(G)$: given two elements $u(x_1, \dots, x_n), v(x_1, \dots, x_n) \in G$ which are known to be conjugate to each other in G , find $w(x_1, \dots, x_n)$ such that $w^{-1}uw =_G v$. Again, no matter if $\text{CP}(G)$ or even $\text{WP}(G)$ are solvable or not, $\text{CP}^+(G)$ is always solvable by brute force: enumerate the elements in the free group $F = \langle x_1, \dots, x_n \mid \rangle$ and for each one $w(x_1, \dots, x_n)$ apply $\text{WP}^+(G)$ to $v^{-1}w^{-1}uw$. We know that if $v^{-1}w^{-1}uw \neq_G 1$ this process will never terminate. But we can start them for several w 's and, while running, keep opening similar processes for new w 's; eventually, one of them will stop telling us which w satisfies $w^{-1}uw =_G v$. As before, note that, without the assumption, this is not an algorithm because if u and v are not conjugate to each other in G then it works forever without giving any answer.
- the **“yes” part of the twisted conjugacy problem** for G , denoted $\text{TCP}^+(G)$: it is defined, and solved by brute force, in the exact same way as $\text{CP}^+(G)$.
- the **“yes” part of the membership problem** for F in G , denoted $\text{MP}^+(F, G)$: given an element $w(x_1, \dots, x_n) \in G$ known to belong to F , express w as a word on the f_i 's. In a similar way as above, even if $\text{MP}(F, G)$ is unsolvable, $\text{MP}^+(F, G)$ is always solvable by brute force.

Note that, using these brute force arguments, the conjugacy problem is solvable if and only if the conjugacy search problem is solvable. Similarly, the twisted conjugacy problem is solvable if and only if the twisted conjugacy search problem is solvable. However, the corresponding complexities may be rather different.

Finally, let us state few more problems, which will be considered in the present paper. We have:

- the **coset intersection problem** for G , denoted $\text{CIP}(G)$: given two finite sets of elements $\{a_1, \dots, a_r\}$ and $\{b_1, \dots, b_s\}$ in G , and two elements $x, y \in G$ decide whether the coset intersection $xA \cap yB$ is empty or not, where $A = \langle a_1, \dots, a_r \rangle \leq G$ and $B = \langle b_1, \dots, b_s \rangle \leq G$.

Let $F \trianglelefteq G$ be a normal subgroup of G . By normality, two elements $u, v \in G$ conjugate to each other in G , either both belong to F or both outside F . Accordingly, $\text{CP}(G)$ splits into two parts, one of them relevant for our purposes:

- the **conjugacy problem for G restricted to F** , denoted $\text{CP}_F(G)$: given two elements $u, v \in F$, decide whether they are conjugate to each other in G . Note that a solution to $\text{CP}(G)$ automatically gives a solution to $\text{CP}_F(G)$, and that this is not, in general, the same problem as $\text{CP}(F)$.

Now, let F be a group given in an algorithmic way, and A be a subgroup of $\text{Aut}(F)$. We have:

- the **orbit decidability problem** for A , denoted $\text{OD}(A)$: given two elements $u, v \in F$, decide whether there exists $\varphi \in A$ such that $u\varphi$ and v are conjugate to each other in F . As will be seen in Section 7, there are finitely generated subgroups $A \leq \text{Aut}(F)$ with unsolvable $\text{OD}(A)$.

Note that orbit decidability for $A \leq \text{Aut}(F)$ is equivalent to the existence of an algorithm which, for any two elements $u, v \in F$, decides whether there exists $\varphi \in A \cdot \text{Inn}(F)$ such that $u\varphi = v$. In particular, if two subgroups $A, B \leq \text{Aut}(F)$ satisfy $A \cdot \text{Inn}(F) = B \cdot \text{Inn}(F)$ then $\text{OD}(A)$ and $\text{OD}(B)$ are the same problem (in particular, A is orbit decidable if and only if B is orbit decidable). This means that orbit decidability is, in fact, a property of subgroups of $\text{Out}(F)$. However, we shall keep talking about $\text{Aut}(F)$ for notational convenience.

To finish the section, let us consider an arbitrary short exact sequence of groups,

$$1 \longrightarrow F \xrightarrow{\alpha} G \xrightarrow{\beta} H \longrightarrow 1.$$

Such a sequence is said to be *algorithmic* if: (i) the groups F , G and H and the morphisms α and β are given to us in an algorithmic way, i.e. we can effectively operate in F , G and H , and compute images under α and β , (ii) we can compute at least one pre-image in G of any element in H , and (iii) we can compute pre-images in F of elements in G mapping to the trivial element in H .

The typical example (though not the unique one) of an algorithmic short exact sequence is that given by finite presentations and images of generators. In fact, (i) is immediate, and we can use $\text{MP}^+(G\beta, H)$ to compute pre-images in G of elements in H , and use $\text{MP}^+(F\alpha, G)$ to compute pre-images in F of elements in G mapping to 1_H .

3. ORBIT DECIDABILITY AND THE CONJUGACY PROBLEM

Consider a short exact sequence of groups,

$$1 \longrightarrow F \xrightarrow{\alpha} G \xrightarrow{\beta} H \longrightarrow 1.$$

Since $F\alpha$ is normal in G , for every $g \in G$, the conjugation γ_g of G induces an automorphism of F , $x \mapsto g^{-1}xg$, which will be denoted $\varphi_g \in \text{Aut}(F)$ (note that, in general, φ_g does not belong to $\text{Inn}(F)$). It is clear that the set of all such automorphisms,

$$A_G = \{\varphi_g \mid g \in G\},$$

is a subgroup of $\text{Aut}(F)$ containing $\text{Inn}(F)$. We shall refer to it as the *action subgroup* of the given short exact sequence.

Assuming some hypothesis on the sequence and the groups involved on it, the following theorem shows that the solvability of the conjugacy problem for G is equivalent to the orbit decidability of the action subgroup $A_G \leq \text{Aut}(F)$.

Theorem 3.1. *Let*

$$1 \longrightarrow F \xrightarrow{\alpha} G \xrightarrow{\beta} H \longrightarrow 1.$$

be an algorithmic short exact sequence of groups such that

- (i) *F has solvable twisted conjugacy problem,*
- (ii) *H has solvable conjugacy problem, and*
- (iii) *for every $1 \neq h \in H$, the subgroup $\langle h \rangle$ has finite index in its centralizer $C_H(h)$, and there is an algorithm which computes a finite set of coset representatives, $z_{h,1}, \dots, z_{h,t_h} \in H$,*

$$C_H(h) = \langle h \rangle z_{h,1} \sqcup \dots \sqcup \langle h \rangle z_{h,t_h}.$$

Then, the following are equivalent:

- (a) *the conjugacy problem for G is solvable,*

- (b) *the conjugacy problem for G restricted to F is solvable,*
(c) *the action subgroup $A_G = \{\varphi_g \mid g \in G\} \leq \text{Aut}(F)$ is orbit decidable.*

Proof. As usual, we shall identify F with $F\alpha \leq G$. By definition, $x\varphi_g = g^{-1}xg$ for every $g \in G$ and $x \in F$. So, given two elements $x, x' \in F$, finding $g \in G$ such that $x' = g^{-1}xg$ is the same as finding $\varphi \in A_G$ such that $x' = x\varphi$. Since $A_G = A_G \cdot \text{Inn}(F)$, this is solving $\text{OD}(A_G)$. Hence, (b) and (c) are equivalent. It is also obvious that (a) implies (b). So, the relevant implication is (b) \Rightarrow (a).

Assume that (b) holds, let $g, g' \in G$ be two given elements in G and let us decide whether they are conjugate to each other in G .

Map them to H . Using (ii), we can decide whether $g\beta$ and $g'\beta$ are conjugate to each other in H . If they are not, then g and g' cannot either be conjugate to each other in G . Otherwise, (ii) gives us an element of H conjugating $g\beta$ into $g'\beta$. Compute a pre-image $u \in G$ of this element. It satisfies $(g^u)\beta = (g\beta)^{u\beta} = g'\beta$. Now, changing g to g^u , we may assume that $g\beta = g'\beta$. If this is the trivial element in H (which we can decide because of (ii)) then g and g' lie in the image of α , and applying (b) we are done. Hence, we are restricted to the case $g\beta = g'\beta \neq_H 1$.

Now, compute $f \in F$ such that $g' = gf$ (this is the α -pre-image of $g^{-1}g'$). Since $g\beta \neq_H 1$, we can use (iii) to compute elements $z_1, \dots, z_t \in H$ such that $C_H(g\beta) = \langle g\beta \rangle z_1 \sqcup \dots \sqcup \langle g\beta \rangle z_t$, and then compute a pre-image $y_i \in G$ for each z_i , $i = 1, \dots, t$. Note that, by construction, the β -images of g and y_i (respectively $g\beta$ and z_i) commute in H so, $y_i^{-1}gy_i = gp_i$ for some computable $p_i \in F$.

Since $g\beta = g'\beta$, every possible conjugator of g into g' must map to $C_H(g\beta)$ under β so, it must be of the form $g^r y_i x$ for some integer r , some $i \in \{1, \dots, t\}$, and some $x \in F$. Hence,

$$gf = g' = (x^{-1}y_i^{-1}g^{-r})g(g^r y_i x) = x^{-1}(y_i^{-1}gy_i)x = x^{-1}gp_i x.$$

Thus, deciding whether g and g' are conjugate to each other in G amounts to decide whether there exists $i \in \{1, \dots, t\}$ and $x \in F$ satisfying $gf = x^{-1}gp_i x$, which is equivalent to $f = (g^{-1}x^{-1}g)p_i x$ and so to $f = (x\varphi_g)^{-1}p_i x$. Since i takes finitely many values and the previous equation means precisely $f \sim_{\varphi_g} p_i$, we can algorithmically solve this problem by hypothesis (i). This completes the proof. \square

Remark 3.2. Note that in the proof of Theorem 3.1 we did not use the full power of hypothesis (i). In fact, we used a solution to $\text{TCP}_\phi(F)$ only for the automorphisms in the action subgroup, $\phi \in A_G$. For specific examples, this may be a weaker assumption than a full solution to $\text{TCP}(F)$.

Theorem 3.1 gives us a relatively big family of groups G for which the conjugacy problem reduces to its restriction to a certain normal subgroup.

Now, we point out some examples of groups F and H satisfying hypotheses (i)-(iii) of the Theorem, and hence providing a family of groups G for which the characterization is valid.

Suppose that F is a finitely generated abelian group. For any given $\phi \in \text{Aut}(F)$ and $u, v \in F$, we have $u \sim_\phi v$ if and only if $u \equiv v$ modulo $\text{Im}(\phi - \text{Id})$. Hence, $\text{TCP}(F)$ reduces to solving a system of linear equations (some over \mathbb{Z} and some modulo certain integers). So it is solvable. On the other hand, Theorem 1.5 in [6] states that $\text{TCP}(F)$ is also solvable when F is finitely generated free. So, finitely generated abelian groups, and finitely generated free groups satisfy (i).

With respect to conditions (ii) and (iii), it is well known that in a free group H , the centralizer of an element $1 \neq h \in H$ is cyclic and generated by the *root* of h (i.e., the unique non proper power $\hat{h} \in H$ such that h is a positive power of \hat{h}), which is computable. Clearly, then, finitely generated free groups satisfy hypotheses (ii) and (iii).

Focusing on these families of groups, Theorem 3.1 is talking about solvability of the conjugacy problem for [f.g. free]-by-[f.g. free] and [f.g. abelian]-by-[f.g. free] groups, and can be restated as follows.

Theorem 3.3. *Let $F = \langle x_1, \dots, x_n \mid R \rangle$ be a (finitely generated) free or abelian group, and let $\varphi_1, \dots, \varphi_m \in \text{Aut}(F)$. Then, the [f.g. free]-by-[f.g. free] or [f.g. abelian]-by-[f.g. free] group*

$$G = \langle x_1, \dots, x_n, t_1, \dots, t_m \mid R, t_j^{-1} x_i t_j = x_i \varphi_j, i = 1, \dots, n, j = 1, \dots, m \rangle$$

has solvable conjugacy problem if and only if $\langle \varphi_1, \dots, \varphi_m \rangle \leq \text{Aut}(F)$ is orbit decidable. \square

In the following section, applicability of Theorem 3.1 will be enlarged by finding more groups with solvable twisted conjugacy problem (see Subsection 4.1), and finding more groups which satisfy condition (iii) (see Subsection 4.2).

At this point, we want to remark that the study of the conjugacy problem in the families of [f.g. free]-by-[f.g. free] and [f.g. free abelian]-by-[f.g. free] groups was the authors' original motivation for developing the present research. One can interpret Theorem 3.3 by saying that orbit decidability is the unique possible obstruction to the solvability of the conjugacy problem within these families of groups. So, by finding examples of orbit decidable subgroups in $\text{Aut}(F_n)$ or $\text{Aut}(\mathbb{Z}^n) = \text{GL}_n(\mathbb{Z})$ one is in fact solving the conjugacy problem in some [f.g. free]-by-[f.g. free] or [f.g. free abelian]-by-[f.g. free] groups; we develop this in Section 6.

But in the literature there are known examples of [f.g. free]-by-[f.g. free] groups with unsolvable conjugacy problem: Miller's example of such a group

(see Miller [30]) automatically gives us an example of a finitely generated subgroup of the automorphism group of a free group, which is orbit undecidable. We discuss this in Section 7, where we also find orbit undecidable subgroups in $GL_n(\mathbb{Z})$ for $n \geq 4$; these last examples correspond to the first known [f.g. free abelian]-by-[f.g. free] groups with unsolvable conjugacy problem.

To conclude the section, we point out the following consequence of Theorem 3.1.

Corollary 3.4. *Consider two algorithmic short exact sequences*

$$\begin{aligned} 1 &\longrightarrow F \longrightarrow G_1 \longrightarrow H_1 \longrightarrow 1 \\ 1 &\longrightarrow F \longrightarrow G_2 \longrightarrow H_2 \longrightarrow 1 \end{aligned}$$

both satisfying the hypotheses of Theorem 3.1, and sharing the same first term. If the two action subgroups coincide, $A_{G_1} = A_{G_2} \leq \text{Aut}(F)$, then G_1 has solvable conjugacy problem if and only if G_2 has solvable conjugacy problem. \square

Note that, in the situation of Corollary 3.4, A_{G_1} and A_{G_2} can be equal, even with G_1 and G_2 being far from isomorphic (for example, choose two very different sets of generators for $A \leq \text{Aut}(F)$, and consider two extensions of F by a free group with free generators acting on F as the chosen automorphisms). A nice example of this fact is the case of doubles of groups: the *double of G over F* $\leq G$ is the amalgamated product of two copies of G with the corresponding F 's identified in the natural way.

Corollary 3.5. *Let $1 \rightarrow F \rightarrow G \rightarrow H \rightarrow 1$ be a short exact sequence satisfying the hypotheses of Theorem 3.1. Then G has solvable conjugacy problem if and only if the double of G over F has solvable conjugacy problem.*

Proof. Let $1 \rightarrow F' \rightarrow G' \rightarrow H' \rightarrow 1$ be another copy of the same short exact sequence, and construct the double of G over F , say $G *_{F=F'} G'$. We have the natural short exact sequence for it,

$$1 \longrightarrow F = F' \longrightarrow G *_{F=F'} G' \longrightarrow H * H' \longrightarrow 1,$$

whose action subgroup clearly coincides with $A_G \leq \text{Aut}(F)$. So, Corollary 3.4 gives the result. \square

4. ENLARGING THE APPLICABILITY OF THEOREM 3.1

In this section we shall enlarge the applicability of Theorem 3.1, by solving the twisted conjugacy problem for more groups F other than free and free abelian (Subsection 4.1), and by finding more groups H with conditions (ii) and (iii), other than free (Subsection 4.2).

4.1. More groups with solvable twisted conjugacy problem. Let us begin by proving a partial converse to Theorem 3.1, in the case where $H = \mathbb{Z}$. In this case, conditions (ii) and (iii) are obviously satisfied, and the relevant part of the statement says that, for $\varphi \in \text{Aut}(F)$, solvability of $\text{TCP}_{\varphi^r}(F)$ for every $r \in \mathbb{Z}$ and of $\text{OD}(\langle \varphi \rangle)$ does imply solvability of $\text{CP}(F \rtimes_{\varphi} \mathbb{Z})$, see Remark 3.2. A weaker version of the convers is true as well.

Proposition 4.1. *Let F be a finitely generated group and let $\varphi \in \text{Aut}(F)$, both given in an algorithmic way (and so $F \rtimes_{\varphi} \mathbb{Z}$). If $\text{CP}(F \rtimes_{\varphi} \mathbb{Z})$ is solvable then $\text{TCP}_{\varphi}(F)$ is solvable, and $\langle \varphi \rangle \leq \text{Aut}(F)$ is orbit decidable.*

Proof. Let us assume that $F \rtimes_{\varphi} \mathbb{Z}$ has solvable conjugacy problem. Then, $\langle \varphi \rangle \leq \text{Aut}(F)$ is orbit decidable by Theorem 3.1 (a) \Rightarrow (c) (which uses non of the hypothesis there) applied to the short exact sequence $1 \rightarrow F \rightarrow F \rtimes_{\varphi} \mathbb{Z} \rightarrow \mathbb{Z} \rightarrow 1$.

On the other hand, let $u, v \in F$. For $x \in F$, the equality $(x\varphi)^{-1}ux = v$ holds in F if and only if $x^{-1}(tu)x = tv$ holds in $F \rtimes_{\varphi} \mathbb{Z}$. So, $u \sim_{\varphi} v$ if and only if tu and tv are conjugated to each other in $F \rtimes_{\varphi} \mathbb{Z}$ by some element of F . By hypothesis, we know how to decide whether tu and tv are conjugated to each other in $F \rtimes_{\varphi} \mathbb{Z}$ by an arbitrary element, say $t^k x$. And,

$$(t^k x)^{-1}(tu)(t^k x) = x^{-1}t^{-k}tut^k x = x^{-1}t(u\varphi^k)x.$$

Hence, tu and tv are conjugated in $F \rtimes_{\varphi} \mathbb{Z}$ if and only if, for some integer k , $t(u\varphi^k)$ and tv are conjugated to each other in $F \rtimes_{\varphi} \mathbb{Z}$ by some element of F . That is, if and only if, $u\varphi^k \sim_{\varphi} v$ for some integer k . But equation $u = (u\varphi)^{-1}(u\varphi)u$ tells us that $u \sim_{\varphi} u\varphi$ (see Lemma 1.7 in [6]). So, solving the conjugacy problem in $F \rtimes_{\varphi} \mathbb{Z}$ for tu and tv , is the same as deciding whether $u \sim_{\varphi} v$ in F . Thus, F has solvable φ -twisted conjugacy problem. \square

Combining Theorem 3.1 and this weak convers, we have the following remarkable consequence.

Corollary 4.2. *Let F be a finitely generated group given in an algorithmic way. All cyclic extensions $F \rtimes_{\varphi} \mathbb{Z}$ have solvable conjugacy problem if and only if both F has solvable twisted conjugacy problem and all cyclic subgroups of $\text{Aut}(F)$ are orbit decidable.* \square

This corollary is useful in order to find more groups with solvable twisted conjugacy problem. For example, using a recent result by J.P. Preaux, we can easily prove that surface groups have solvable twisted conjugacy problem, and that Brinkmann's result (see [10] or Theorem 6.17 in section 6.2 below) is also valid for surface groups.

Theorem 4.3. *Let F be the fundamental group of a closed surface. Then, $\text{TCP}(F)$ is solvable and all cyclic subgroups of $\text{Aut}(F)$ are orbit decidable.*

Proof. By Theorem 1.6 in [34], $F \rtimes_{\varphi} \mathbb{Z}$ has solvable conjugacy problem for every automorphism $\varphi \in \text{Aut}(F)$. Now, Corollary 4.2 gives the result. \square

The same argument also provides an alternative proof of the recent result that polycyclic groups have solvable twisted conjugacy problem (see [18]). We remind that a group F is called *polycyclic* when there exists a finite sequence of subgroups $F = K_0 \triangleright K_1 \triangleright \cdots \triangleright K_{n-1} \triangleright K_n = 1$, each normal in the previous one, and with every quotient K_i/K_{i+1} being cyclic (finite or infinite). If all these quotients are infinite cyclic we say that F is *poly- \mathbb{Z}* .

Theorem 4.4. *Let F be a polycyclic group. Then, F has solvable twisted conjugacy problem, and all cyclic subgroups of $\text{Aut}(F)$ are orbit decidable.*

Proof. Let φ be an arbitrary automorphism of F . Since F is normal in $F \rtimes_{\varphi} \mathbb{Z}$ with cyclic quotient, the group $F \rtimes_{\varphi} \mathbb{Z}$ is again polycyclic. So, it has solvable conjugacy problem (see [36]). Again, Corollary 4.2 gives the result. \square

Let us study now how the twisted conjugacy problem behaves under finite index extensions. To do this, we shall need the classical Todd-Coxeter algorithm, and a technical lemma about computing finite index characteristic subgroups.

Theorem 4.5 (Todd-Coxeter, Chapter 8 in [22]). *Let $L = \langle X \mid R \rangle$ be a finite presentation and $K \leq L$ a finite index subgroup given by a finite set of generators. There is an algorithm to compute a set of left coset representatives $\{1 = g_0, g_1, \dots, g_q\}$ of K in L (so, $L = K \sqcup g_1 K \sqcup \cdots \sqcup g_q K$), plus the information on how to multiply them by generators of L on the left, say $x_i g_j \in g_{d(i,j)} K$. Moreover, one can algorithmically write any given $g \in L$ in the form $g = g_p k$ for some (unique) $p = 0, \dots, q$, and some $k \in K$ expressed as a word on the generators of K . In particular, $MP(K, L)$ is solvable.*

Lemma 4.6. *Let F be a group given by a finite presentation $\langle X \mid R \rangle$, and suppose we are given a set of words $\{w_1, \dots, w_r\}$ on X such that $K = \langle w_1, \dots, w_r \rangle \leq F$ is a finite index subgroup. Then, the finite index characteristic subgroup $K_0 = \bigcap_{[F:K'] = s} K' \leq F$, where $s = [F : K]$, has a computable set of generators.*

Proof. Think F as a quotient of the free group on X , $\pi : F(X) \twoheadrightarrow F$ (write $N = \ker \pi$). Using the elementary fact that, for every finite index subgroup $L \leq F(X)$, $[F(X) : L] \geq [F : L\pi]$ with equality if and only if $N \leq L$, we

have

$$K_0 = \bigcap_{\substack{K' \leq F \\ [F:K'] = s}} K' = \bigcap_{\substack{N \leq K'' \leq F(X) \\ [F(X):K''] = s}} (K''\pi) = \left(\bigcap_{\substack{N \leq K'' \leq F(X) \\ [F(X):K''] = s}} K'' \right) \pi.$$

Hence, we can compute generators for K_0 (as words in X) by first computing s (use Todd-Coxeter algorithm), and then listing and intersecting (and finally projecting into F) all of the finitely many subgroups K'' of index s in $F(X)$ which contain N : the listing can be done by enumerating all saturated and folded Stallings graphs with s vertices, and then computing a basis for the corresponding subgroup (see [39]); intersecting is easy using the pull-back technique (again, see [39]); and deciding whether such a K'' contains N can be done, even without knowing explicit generators for N , by projecting K'' down to F , and using Todd-Coxeter again to verify if the image has index exactly s , or smaller. \square

A result by Gorjaga and Kirkinskii (see [19]) states the existence of a group F with an index two subgroup $K \leq F$, such that the conjugacy problem is solvable in K but unsolvable in F (two years later the same was independently proved by Collins and Miller in [12], together with the opposite situation, unsolvability for K and solvability for F). In other words, the conjugacy problem *does not* go up or down through finite index extensions. In contrast with this, the following proposition shows that the twisted conjugacy problem does (assuming the subgroup is characteristic).

Proposition 4.7. *Let F be a group given by a finite presentation $\langle X \mid R \rangle$, and suppose we are given a set of words $\{w_1, \dots, w_r\}$ on X such that $K = \langle w_1, \dots, w_r \rangle \leq F$ is a finite index subgroup.*

- (i) *Suppose K is characteristic in F ; if $TCP(K)$ is solvable then $TCP(F)$ is also solvable.*
- (ii) *Suppose K is normal in F ; if $TCP(K)$ is solvable then $CP(F)$ is solvable.*

Proof. By Todd-Coxeter algorithm, we can compute a set $\{1 = y_1, y_2, \dots, y_s\}$ of left coset representatives of K in F , i.e. $F = K \sqcup y_2 K \sqcup \dots \sqcup y_s K$, and use them to write any $u \in F$ (given as word in X) in the form $y_i k$ for some $i = 1, \dots, s$, and some $k \in K$ (expressed as a word in the w_i 's); in particular, $MP(K, F)$ is solvable (see Theorem 4.5).

Suppose now that K is characteristic in F , and $TCP(K)$ is solvable. Fix $\varphi \in Aut(F)$, and let $u, v \in F$ be two elements in F . With the previous procedure, we can write them as $u = y_i z$ and $v = y_j z'$, for some $i, j = 1, \dots, s$ and $z, z' \in K$. Deciding whether or not u and v are φ -twisted

conjugated in F amounts to decide whether there exist $l = 1, \dots, s$ and $k \in K$ such that

$$((ky_l^{-1})\varphi)(y_iz)(y_lk^{-1}) = y_jz'$$

or, equivalently,

$$(y_l\varphi)^{-1}y_izy_l = (k\varphi)^{-1}y_jz'k = y_j[(y_j^{-1}(k\varphi)y_j)^{-1}z'k].$$

Now observe that, if K is characteristic in F , then the bracketed element $[(y_j^{-1}(k\varphi)y_j)^{-1}z'k]$ belongs to K . For every $l = 1, \dots, s$, compute $(y_l\varphi)^{-1}y_izy_l$ and check if it belongs to the coset corresponding to y_j . If non of them do, then y_iz and y_jz' are not φ -twisted conjugated. Otherwise, we have computed the non-empty list of indices l and elements $k_l \in K$ such that $(y_l\varphi)^{-1}y_izy_l = y_jk_l$. For each one of them, it remains to decide whether there exists $k \in K$ such that $k_l = (y_j^{-1}(k\varphi)y_j)^{-1}z'k = (k\varphi\gamma_{y_j})^{-1}z'k$. This is precisely deciding whether k_l and z' are $(\varphi\gamma_{y_j})$ -twisted conjugated in K (which makes sense because $\varphi\gamma_{y_j}$ restricts to an automorphism of K). This is doable by hypothesis, so we have proven (i).

The previous argument particularized to the special case $\varphi = Id$ works assuming only that K is normal in F . This shows (ii). \square

Theorem 4.8. *Let F be a group given by a finite presentation $\langle X \mid R \rangle$. If either F is (finitely generated)*

- (i) *virtually abelian, or*
- (ii) *virtually free, or*
- (iii) *virtually surface group, or*
- (iv) *virtually polycyclic,*

then $TCP(F)$ is solvable.

Proof. We first need to compute generators for a finite index subgroup K of F being abelian, or free, or surface, or polycyclic (and so, finitely generated, like F). We shall identify such a subgroup by finding a special type of presentation for it. In the first three cases, a finite presentation will be called *canonical* if the set of relations, respectively, contains all the commutators of pairs of generators, or is empty, or consists precisely on a single element being a surface relator. For the polycyclic case, let $L = \langle y_1, \dots, y_n \mid S \rangle$ be a given finite presentation and, for every automorphism $\varphi \in Aut(L)$, consider the cyclic extension $L \rtimes_{\varphi} \mathbb{Z}$, and its redundant presentation $\mathcal{CE}_{\varphi}(\langle y_1, \dots, y_n \mid S \rangle)$ given by $L \rtimes_{\varphi} \mathbb{Z} = \langle y_1, \dots, y_n, t \mid S, t^{-1}y_it = u_i, ty_it^{-1} = v_i \ (i = 1, \dots, n) \rangle$, where u_i and v_i are words on the y_i 's describing, respectively, the images and preimages of y_i under φ . The point of considering such redundant presentations is that, once a particular presentation of this type is given, one can easily verify whether $y_i \mapsto u_i$ defines an automorphism of G (with inverse given by $y_i \mapsto v_i$). Now every non-trivial, poly- \mathbb{Z} group

admits a presentation of the form $\mathcal{CE}_{\varphi_n}(\cdots(\mathcal{CE}_{\varphi_1}(\langle x \mid - \rangle))\cdots)$, which we shall call *canonical* (add to this definition that the canonical presentation of the trivial group is the empty one). Observe that, given a presentation, it can be verified whether it is canonical in any of the above four cases.

To prove the theorem, let us first enumerate the list of all subgroups of finite index in F , say K_1, K_2, \dots . This can be done by following the strategy in the proof of Lemma 4.6: enumerate all subgroups of the free group $F(X)$ of a given (and increasing) index, and project them into F . For every $i \geq 1$, while computing K_{i+1} , start a new parallel computation following Reidemeister-Schreier process (see Section 3.2 in [27]) in order to obtain a finite presentation for K_i , say $\langle Y_i \mid S_i \rangle$. Then, start applying to $\langle Y_i \mid S_i \rangle$ the list of all possible sequences of Tietze transformations of any given (and increasing) length. When one of the running processes finds a canonical presentation (of an abelian, or free, or surface, or poly- \mathbb{Z} group) then stop all of them and output this presentation. Although many of these individual processes will never end, one of them will eventually finish because we are potentially exploring *all* finite presentations of *all* finite index subgroups of F and, by hypothesis, at least one of them admits a canonical presentation (we use here the fact that every polycyclic group is virtually poly- \mathbb{Z} , see Proposition 2 in Chapter 1 of [37]). The final output of all these parallel processes is a canonical presentation for a finite index subgroup K of F being abelian, or free, or surface, or poly- \mathbb{Z} .

Now, apply Lemma 4.6 to compute generators of a finite index characteristic subgroup $K_0 \leq F$ inside K . Note that K_0 (for which we can obtain an explicit presentation by using Reidemeister-Schreier method) will again be either abelian, or free, or surface, or polycyclic. So, $\text{TCP}(K_0)$ is solvable by results above. Hence, $\text{TCP}(F)$ is also solvable by Proposition 4.7 (ii). \square

The following results are two other interesting consequences of Propositions 4.7 and 4.1.

Corollary 4.9. *There exists a finitely presented group G with $\text{CP}(G)$ solvable, but $\text{TCP}(G)$ unsolvable.*

Proof. Consider a finitely presented group F with an index two subgroup $G \leq F$, such that $\text{CP}(G)$ is solvable but $\text{CP}(F)$ is unsolvable (see, for example, Gorjaga-Kirkinskii [19] or Collins-Miller [12]). Since index two subgroups are normal, Proposition 4.7 (ii) implies that $\text{TCP}(G)$ must be unsolvable. \square

Corollary 4.10. *There exists a finitely presented group G and an automorphism $\varphi \in \text{Aut}(G)$ such that $\text{CP}(G)$ is solvable and $\text{CP}(G \rtimes_{\varphi} \mathbb{Z})$ is unsolvable. Conversely, there also exists a finitely presented group G and*

an automorphism $\varphi \in \text{Aut}(G)$ such that $\text{CP}(G \rtimes_{\varphi} \mathbb{Z})$ is solvable and $\text{CP}(G)$ is unsolvable.

Proof. For the first assertion, consider a finitely presented group G like in the previous theorem, with $\text{CP}(G)$ solvable and $\text{TCP}(G)$ unsolvable. There must exist $\varphi \in \text{Aut}(G)$ with $\text{TCP}_{\varphi}(G)$ unsolvable. Then, by Proposition 4.1, $\text{CP}(G \rtimes_{\varphi} \mathbb{Z})$ is unsolvable too.

For the second assertion, start with Collins-Miller example of an index two extension $G \leq F$ with unsolvable $\text{CP}(G)$ but solvable $\text{CP}(F)$. In this construction, it is easy to see that $F = G \rtimes_{\varphi} C_2$ for an order two automorphism $\varphi \in \text{Aut}(G)$. And solvability of $\text{CP}(G \rtimes_{\varphi} C_2)$ directly implies solvability of $\text{CP}(G \rtimes_{\varphi} \mathbb{Z})$ because the square of the stable letter belongs to the center of this last group. \square

4.2. Centralizers in hyperbolic groups. Hypotheses (ii) and (iii) in Theorem 3.1 are also satisfied by a bigger family of groups beyond free, including finitely generated torsion-free hyperbolic groups. All these groups H provide new potential applications of Theorem 3.1.

Proposition 4.11. *Let H be a finitely generated hyperbolic group given by a finite presentation, and let h be an element of H .*

- (i) *There is an algorithm to determine whether or not the centralizer $C_H(h)$ is finite and, if it is so, to list all its elements.*
- (ii) *If h has infinite order in H , then $\langle h \rangle$ has finite index in $C_H(h)$ and there is an algorithm which computes a set of coset representatives for $\langle h \rangle$ in $C_H(h)$.*

Proof. From the given presentation, it is possible to compute a hyperbolicity constant δ for H (see [32] or [16]). Now, in 1.11 of Chapter III.Γ of [9], the authors provide an algorithm to solve the conjugacy problem in H (the reader can easily check that the q.m.c. property used there, holds with constant $K = 4\delta + 1$ in our case). In fact, the same construction (with $u = v = h$) also enables us to compute a finite set of generators for $C_H(h)$ (as labels of the closed paths at vertex h in \mathcal{G}). So, we have computed a generating set $\{g_1, \dots, g_r\}$ for $C_H(h)$.

On the other hand, by [5], each finite subgroup of H is conjugate to a subgroup which is contained in the ball of radius $4\delta + 1$ around 1, $B(4\delta + 1)$. In particular, if $C_H(h)$ is finite then $C_H(h)^x \subseteq B(4\delta + 1)$ for some $x \in H$. We can apply a solution of the conjugacy search problem for H (which is solvable) to h and each one of the members of $B(4\delta + 1)$, ending up with a list of elements $\{x_1, \dots, x_s\}$ such that $\{h^{x_1}, \dots, h^{x_s}\}$ are all the conjugates of h belonging to $B(4\delta + 1)$. Now, $C_H(h)$ is finite if and only if $C_H(h)^x \subseteq B(4\delta + 1)$ for some $x \in H$, and this happens if and only

if $C_H(h)^{x_i} = \langle g_1^{x_i}, \dots, g_r^{x_i} \rangle \subseteq B(4\delta + 1)$ for some $i = 1, \dots, s$ (because if $h^y = h$ then $C_H(h)^y = C_H(h)$). Furthermore, this last condition, for a fixed i , is decidable in the following way: recursively, for $l = 1, 2, \dots$, check whether the set B_l of all products of l or less $(g_j^{x_i})^{\pm 1}$'s is contained in $B(4\delta + 1)$; the process will eventually find a value of l for which either $B_l \not\subseteq B(4\delta + 1)$ or $B_{l+1} = B_l \subseteq B(4\delta + 1)$. In the first case $C_H(h)^{x_i} = \langle g_1^{x_i}, \dots, g_r^{x_i} \rangle \not\subseteq B(4\delta + 1)$ (and if this happens for every i then $C_H(h)$ is infinite); and in the second case $C_H(h)^{x_i} = \langle g_1^{x_i}, \dots, g_r^{x_i} \rangle \subseteq B(4\delta + 1)$, which means that $C_H(h)$ is finite and, conjugating back by x_i^{-1} , we obtain the full list of its elements. This concludes the proof of (i).

By Corollary 3.10 in Chapter III.Γ of [9], the group $\langle h \rangle$ has finite index in $C_H(h)$. As explained in the proof of that corollary, different positive powers of h are not conjugate to each other. Therefore, there exists a natural number $k \leq |B(4\delta)| + 1$, such that h^k is not conjugate into the ball $B(4\delta)$. In the proof it is claimed that each element of $C_H(h)$ lies at distance at most $2|h^k| + 4\delta$ of $\langle h^k \rangle$ and hence of $\langle h \rangle$. This means that there exists a set of coset representatives for $\langle h \rangle$ in $C_H(h)$ inside the ball of radius $K = 2(|B(4\delta)| + 1)|h| + 4\delta$. List the elements of such ball, delete those not commuting with h (use $\text{WP}(H)$ here) and, among the final list of candidates z_1, \dots, z_r , it remains to decide which pairs z_i, z_j satisfy $\langle h \rangle z_i = \langle h \rangle z_j$. This is the same as $z = z_j z_i^{-1} \in \langle h \rangle$ which can be algorithmically checked in the following way. For every element w of any group, the *translation number* of w is $\tau(w) = \lim_{n \rightarrow \infty} \frac{|w^n|}{n}$, where $|\cdot|$ denotes the word length with respect to a given presentation (see Definition 3.13 in Chapter III.Γ of [9]). Obviously, $\tau(w) \leq |w|$ and, for hyperbolic groups, there exists a computable $\epsilon > 0$ such that $\tau(w) > \epsilon$ for every w of infinite order (see Proposition 3.1 in [13], or Theorem 3.17 in Chapter III.Γ of [9]). Back to our situation, if $z = h^s$ for some integer s , then $\tau(z) = |s|\tau(h)$ and so, $|s| = \frac{\tau(z)}{\tau(h)} \leq \frac{|z|}{\epsilon}$. Thus, the exponent s has finitely many possibilities and so, we can algorithmically decide whether $z \in \langle h \rangle$. \square

Theorem 4.12. *Consider a short exact sequence given by finite presentations,*

$$1 \longrightarrow F \longrightarrow G \longrightarrow H \longrightarrow 1,$$

where F is (finitely generated) virtually free, or virtually abelian, or virtually surface, or virtually polycyclic group, and H is a (finitely generated) hyperbolic group where every non-trivial element of finite order has finite centralizer. Then, G has solvable conjugacy problem if and only if the corresponding action subgroup $A_G \leq \text{Aut}(F)$ is orbit decidable.

Proof. We just need to check that hypotheses of Theorem 3.1 are satisfied. Certainly, (i) was already considered above, and (ii) is well known

(see 1.11 in Chapter III.Γ of [9]). For (iii), apply the algorithm given in Proposition 4.11 (i) to $h \neq 1$. If it answers that $C_H(h)$ is finite, then the answer comes with the full list of its elements from which, using a solution to $WP(H)$, we can extract the required list of coset representatives for $C_H(h)$ modulo $\langle h \rangle$. Otherwise, $C_H(h)$ is infinite; in this case, our hypothesis ensures that h has infinite order in H , and hence we can apply the algorithm given in Proposition 4.11 (ii) to find the required list anyway. \square

5. THE CONJUGACY PROBLEM FOR $Aut(F_2)$

The conjugacy problem for $Aut(F_n)$ is a deep open question about free groups (a possible plan to solve it has been indicated by M. Lustig in the preprint [26], and some partial results already published by the same author in this direction). Among other partial results, it is known to be solvable for rank $n = 2$ (see [3] or [4]). As an illustration of the potential applicability of Theorem 3.1, we dedicate this section to deduce from it another solution to the conjugacy problem for $Aut(F_2)$.

Consider the standard short exact sequence involving $Aut(F_2)$,

$$1 \rightarrow Inn(F_2) \rightarrow Aut(F_2) \rightarrow GL_2(\mathbb{Z}) \rightarrow 1.$$

Although $Inn(F_2)$ is isomorphic to F_2 , which has solvable twisted conjugacy problem (see [6]), Theorem 3.1 cannot be directly applied to this sequence because some centralizers in $GL_2(\mathbb{Z})$ are too large. Specifically, this group has a centre consisting of plus and minus the identity matrix. However, if we quotient out by this order two subgroup, we obtain $PGL_2(\mathbb{Z})$ which will turn out to satisfy our required hypotheses. Let us amend the short exact sequence above as follows.

Choose a basis $\{a, b\}$ for F_2 and let σ be the (order two) automorphism of F_2 sending a to a^{-1} and b to b^{-1} . Note that $w\sigma = (w^{-1})^R = (w^R)^{-1}$, where $(\cdot)^R$ denotes the *palindromic reverse* of a word on $\{a, b\}$. Also, for every $\phi \in Aut(F_2)$, $\phi^R = \sigma^{-1}\phi\sigma$ acts by sending a to $(a\phi)^R$ and b to $(b\phi)^R$. Consider the subgroup $\langle Inn(F_2), \sigma \rangle \leq Aut(F_2)$. Since $\sigma^{-1}\phi^{-1}\sigma\phi$ abelianizes to the identity, $\phi^{-1}\sigma\phi = \sigma\gamma_x$ for some $x \in F_2$. This (together with the elementary fact $\phi^{-1}\gamma_w\phi = \gamma_{w\phi}$) shows that $\langle Inn(F_2), \sigma \rangle$ is normal in $Aut(F_2)$. Moreover, it is isomorphic to $F_2 \rtimes C_2$, a split extension of F_2 by a cyclic group of order 2. This gives us another short exact sequence as follows

$$1 \rightarrow F_2 \rtimes C_2 \rightarrow Aut(F_2) \rightarrow PGL_2(\mathbb{Z}) \rightarrow 1. \quad (1)$$

From the computational point of view we can think of (1) as given by presentations of the involved groups and the corresponding morphisms among them. But it is simpler to think of it as what is literally written: formal expressions of the type $\sigma^r w$, where $r = 0, 1$ and $w \in F_2$ (in one-to-one

correspondence with $\sigma^r \gamma_w \in \langle Inn(F_2), \sigma \rangle \trianglelefteq Aut(F_2)$; automorphisms of $F_2 = \langle a, b \mid \rangle$; and two-by-two integral matrices modulo $\pm Id$; all with the obvious morphisms among them.

Let us see that (1) satisfies hypotheses (i)-(iii) of Theorem 3.1. It is straightforward to show that F_2 is characteristic in $F_2 \rtimes C_2$. Thus, by Proposition 4.7 (i) and solvability of $TCP(F_2)$ (see [6]), we deduce that $TCP(F_2 \rtimes C_2)$ is solvable. So, our short exact sequence (1) satisfies (i). On the other hand, conditions (ii) and (iii) follow both from standard computations with two by two matrices, and also from considering the presentation of $PGL_2(\mathbb{Z})$ as amalgamated product $PGL_2(\mathbb{Z}) \cong D_2 \underset{D_1}{*} D_3$ (see, for example, page 24 in [14]).

Hence, (1) is a short exact sequence satisfying the hypotheses of Theorem 3.1. This way, the solvability of the conjugacy problem for $Aut(F_2)$ is equivalent to the orbit decidability of the action subgroup $A = \{\varphi_\phi \mid \phi \in Aut(F_2)\} \leq Aut(F_2 \rtimes C_2)$. Thus, the following proposition provides a solution to $CP(Aut(F_2))$:

Proposition 5.1. *With the above notation, the action of $Aut(F_2)$ by right conjugation on its normal subgroup $\langle Inn(F_2), \sigma \rangle \cong F_2 \rtimes C_2$ is orbit decidable.*

Proof. The action of $Aut(F_2)$ on $Inn(F_2)$ is determined by the natural action on F_2 , namely $\phi^{-1} \gamma_w \phi = \gamma_{w\phi}$, for every $\phi \in Aut(F_2)$ and $w \in F_2$. So, by the classical Whitehead algorithm (see Proposition I.4.19 in [25]), it is possible to decide whether two elements in $Inn(F_2)$ lie in the same $Aut(F_2)$ -orbit (i.e. the action of $Aut(F_2)$ on $Inn(F_2)$ is orbit decidable). But, $Inn(F_2)$ has index 2 in $\langle Inn(F_2), \sigma \rangle$ so, it only remains to decide whether two given elements of the form $\sigma \gamma_u, \sigma \gamma_v \in \langle Inn(F_2), \sigma \rangle$, with $u, v \in F_2$, lie in the same $Aut(F_2)$ -orbit or not.

Suppose then that $\sigma \gamma_u$ and $\sigma \gamma_v$ are given to us, and let us algorithmically decide whether there exists $\phi \in Aut(F_2)$ such that $\phi^{-1} \sigma \gamma_u \phi = \sigma \gamma_v$.

Note that, if this is the case, then $\phi^{-1} (\sigma \gamma_u)^2 \phi = (\sigma \gamma_v)^2$, while $(\sigma \gamma_u)^2$ and $(\sigma \gamma_v)^2$ belong to $Inn(F_2)$. So, apply Whitehead algorithm to $(\sigma \gamma_u)^2$ and $(\sigma \gamma_v)^2$; if they are not in the same $Aut(F_2)$ -orbit, the same is true for $\sigma \gamma_u$ and $\sigma \gamma_v$, and we are done. Otherwise, we come up with a particular $\alpha \in Aut(F_2)$ such that $\alpha^{-1} (\sigma \gamma_u)^2 \alpha = (\sigma \gamma_v)^2$; now, replacing $\sigma \gamma_u$ by $\alpha^{-1} \sigma \gamma_u \alpha$, we can assume that $(\sigma \gamma_u)^2 = (\sigma \gamma_v)^2$. This means

$$\gamma_{(u\sigma)u} = \sigma \gamma_u \sigma \gamma_u = (\sigma \gamma_u)^2 = (\sigma \gamma_v)^2 = \sigma \gamma_v \sigma \gamma_v = \gamma_{(v\sigma)v},$$

i.e. $(u\sigma)u = (v\sigma)v$. We can compute this element of F_2 and distinguish two cases depending whether it is trivial or not:

Case 1: $(u\sigma)u = (v\sigma)v = 1$. Note that, in this case, u and v are palindromes (i.e. $u^R = u$ and $v^R = v$). We shall show that $\sigma \gamma_u$ and $\sigma \gamma_v$

are always in the same $\text{Aut}(F_2)$ -orbit. In fact, notice that if $x \in \{a, b\}^{\pm 1}$ is any letter then,

$$\gamma_x^{-1}(\sigma\gamma_u)\gamma_x = \gamma_{x^{-1}}\sigma\gamma_u\gamma_x = \sigma\gamma_{xux}.$$

Hence, we can consecutively use conjugations of this type to shorten the length of u (and v) down to 0 or 1; that is, $\sigma\gamma_u$ (and $\sigma\gamma_v$) is in the same $\text{Aut}(F_2)$ -orbit as at least one of σ , $\sigma\gamma_a$, $\sigma\gamma_{a^{-1}}$, $\sigma\gamma_b$ or $\sigma\gamma_{b^{-1}}$. But, $\gamma_a^{-1}\sigma\gamma_{a^{-1}}\gamma_a = \sigma\gamma_a$ and $\gamma_b^{-1}\sigma\gamma_{b^{-1}}\gamma_b = \sigma\gamma_b$. Also, defining $\rho \in \text{Aut}(F_2)$ as $a \mapsto b$, $b \mapsto a$, we have $\rho^{-1}\sigma\gamma_a\rho = \sigma\gamma_b$. Finally, defining $\chi \in \text{Aut}(F_2)$ as $a \mapsto ab$, $b \mapsto b$, we have $\chi^{-1}\sigma\chi = \sigma\gamma_b$. Thus, $\sigma\gamma_u$ and $\sigma\gamma_v$ are always in the same $\text{Aut}(F_2)$ -orbit in this case.

Case 2: $(u\sigma)u = (v\sigma)v \neq 1$. Let z be its root, i.e. $(u\sigma)u = (v\sigma)v = z^s$ for some $s \geq 1$, with z not being a proper power. If some $\phi \in \text{Aut}(F_2)$ satisfies $\phi^{-1}\sigma\gamma_u\phi = \sigma\gamma_v$ then such ϕ must also satisfy

$$\gamma_{z^s}\phi = \phi^{-1}\gamma_{z^s}\phi = \phi^{-1}(\sigma\gamma_u)^2\phi = (\sigma\gamma_v)^2 = \gamma_{z^s}$$

and so, $z\phi = z$. In other words, $\sigma\gamma_u$ can only be conjugated into $\sigma\gamma_v$ by automorphisms stabilizing z . Using McCool's algorithm (see Proposition I.5.7 in [25]), we can compute a set of generators for this subgroup of $\text{Aut}(F_2)$, say $\text{Stab}(z) = \langle \phi_1, \dots, \phi_k \rangle$.

It remains to analyze how those ϕ_i 's act on $\sigma\gamma_u$. We can compute $\phi_i^{-1}\sigma\gamma_u\phi_i$ and write this element of $F_2 \rtimes C_2$ in normal form, say $\sigma\gamma_{w_i}$. We claim that $w_i = (z\sigma)^{n_i}u$ for some $n_i \in \mathbb{Z}$. In fact, squaring $\phi_i^{-1}\sigma\gamma_u\phi_i = \sigma\gamma_{w_i}$ (and using that ϕ_i stabilize z) we obtain

$$\gamma_{(u\sigma)u} = (\sigma\gamma_u)^2 = \gamma_{z^s} = \phi_i^{-1}\gamma_{z^s}\phi_i = \phi_i^{-1}(\sigma\gamma_u)^2\phi_i = (\sigma\gamma_{w_i})^2 = \gamma_{(w_i\sigma)w_i}.$$

This implies $z^s = (u\sigma)u = (w_i\sigma)w_i$ and, applying σ to both sides, $z^s\sigma = u(u\sigma) = w_i(w_i\sigma)$. Now observe that

$$\begin{aligned} (w_i u^{-1})\gamma_{z^s\sigma} &= (z^s\sigma)^{-1}(w_i u^{-1})(z^s\sigma) = (w_i^{-1}\sigma)w_i^{-1}(w_i u^{-1})u(u\sigma) \\ &= (w_i^{-1}\sigma)(u\sigma) = w_i u^{-1}, \end{aligned}$$

which means that $w_i u^{-1}$ commutes with $z^s\sigma = (z\sigma)^s$. Hence, $w_i u^{-1} = (z\sigma)^{n_i}$ and $w_i = (z\sigma)^{n_i}u$ for some computable $n_i \in \mathbb{Z}$. We have shown that, for $i = 1, \dots, k$,

$$\phi_i^{-1}\sigma\gamma_u\phi_i = \sigma\gamma_{(z\sigma)^{n_i}u}.$$

Few more computations show that conjugating by ϕ_i^{-1} makes the corresponding negative effect on the exponent: from $\phi_i^{-1}\sigma\gamma_u\phi_i = \sigma\gamma_{(z\sigma)^{n_i}u} = \gamma_{z^{n_i}}\sigma\gamma_u$ we deduce

$$\phi_i(\sigma\gamma_u)\phi_i^{-1} = \phi_i(\gamma_{z^{-n_i}}\phi_i^{-1}\sigma\gamma_u\phi_i)\phi_i^{-1} = \gamma_{z^{-n_i}}\sigma\gamma_u = \sigma\gamma_{(z\sigma)^{-n_i}u}.$$

And conjugating by another ϕ_j makes an additive effect on the exponent:

$$\begin{aligned}
\phi_j^{-1}\phi_i^{-1}\sigma\gamma_u\phi_i\phi_j &= \phi_j^{-1}\sigma\gamma_{(z\sigma)^{n_i}u}\phi_j \\
&= \phi_j^{-1}\gamma_{z^{n_i}}\sigma\gamma_u\phi_j \\
&= \gamma_{z^{n_i}}\phi_j^{-1}\sigma\gamma_u\phi_j \\
&= \gamma_{z^{n_i}}\sigma\gamma_{(z\sigma)^{n_j}u} \\
&= \sigma\gamma_{(z\sigma)^{n_i+n_j}u}.
\end{aligned}$$

So, conjugating by $\phi \in \text{Stab}(z) \leq \text{Aut}(F_2)$, we can only move from $\sigma\gamma_u$ to elements of the form $\sigma\gamma_{(z\sigma)^{\lambda n_0}u}$, where $n_0 = \gcd(n_1, \dots, n_k)$, and $\lambda \in \mathbb{Z}$. Thus, $\sigma\gamma_u$ and $\sigma\gamma_v$ belong to the same $\text{Aut}(F_2)$ -orbit if and only if $v = (z\sigma)^{\lambda n_0}u$ for some $\lambda \in \mathbb{Z}$, which happens if and only if vu^{-1} is a power of $(z\sigma)^{n_0}$. This is decidable in a free group. \square

Corollary 5.2. *$\text{Aut}(F_2)$ has solvable conjugacy problem.* \square

6. POSITIVE RESULTS

In this section, positive results for the free and free abelian cases are analyzed. Along the two parallel subsections, we will give several examples of orbit decidable subgroups in $\text{Aut}(\mathbb{Z}^n) = \text{GL}_n(\mathbb{Z})$ and $\text{Aut}(F_n)$, together with the corresponding [f.g. free abelian]-by-[f.g. free] and [f.g. free]-by-[f.g. free] groups with solvable conjugacy problem (see Theorem 3.3). The reader can easily extend all these results to [f.g. free abelian]-by-[f.g. t.f. hyperbolic] and [f.g. free]-by-[f.g. t.f. hyperbolic] groups, by direct application of Theorem 4.12 (here, t.f. stands for torsion free).

As a technical preliminaries, we first need to solve the coset intersection problem for free and virtually free groups, and to see that orbit decidability goes up to finite index.

Proposition 6.1. *Let K be the free group with basis X . Then, the coset intersection problem $\text{CIP}(K)$ is solvable.*

Proof. Without loss of generality, we can assume that X is finite, since $\text{CIP}(K)$ only involves two finitely generated subgroups $A, B \leq K$ and two words $x, y \in K$. Now, using Stallings' method (see [39]), we can construct the corresponding core-graph $\Gamma_X(A)$ (resp. $\Gamma_X(B)$) and attach to it, and fold if necessary, a path labelled x (resp. y) from a new vertex v_x (resp. v_y) to the base-point 1 in $\Gamma_X(A)$ (resp. 1 in $\Gamma_X(B)$). Now, after computing the pull-back of these two finite graphs, we can easily solve the coset intersection problem: elements from $xA \cap yB$ are precisely labels of paths from (v_x, v_y) to $(1, 1)$ in the pull-back. So, $xA \cap yB \neq \emptyset$ if and only if (v_x, v_y) and $(1, 1)$ belong to the same connected component of this pull-back. \square

Proposition 6.2. *Let L be a group given by a finite presentation $\langle X \mid R \rangle$, and let K be a finite index subgroup of L generated by a given finite set of words in X . If $CIP(K)$ is solvable then $CIP(L)$ is also solvable.*

Proof. By Todd-Coxeter algorithm (see Theorem 4.5), we can compute a set of left coset representatives $\{1 = g_0, g_1, \dots, g_q\}$ of K in L (so, $L = K \sqcup g_1K \sqcup \dots \sqcup g_qK$), plus the information on how to multiply them by generators of L on the left, say $x_i g_j \in g_{d(i,j)}K$. This information can be used to algorithmically write any given $g \in L$ in the form $g = g_p k$ for some (unique) $p = 0, \dots, q$, and $k \in K$. In particular, $MP(K, L)$ is solvable.

Suppose now several elements $x, y, a_1, \dots, a_r, b_1, \dots, b_s \in L$ are given; we shall show how to decide whether $xA \cap yB$ is empty or not, where $A = \langle a_1, \dots, a_r \rangle \leq L$ and $B = \langle b_1, \dots, b_s \rangle \leq L$.

Let us compute a (finite) set of left coset representatives W of A modulo $A \cap K$ in the following way. Enumerate all formal reduced words in the alphabet $\{a_1, \dots, a_r\}^{\pm 1}$, say $\{w_1, w_2, \dots\}$, starting with the empty word, $w_1 = 1$, and in such a way that the length never decreases. Now, starting with the empty set $U = \emptyset$, recursively enlarge it by adding w_i whenever $w_i(A \cap K) \neq w_{i'}(A \cap K)$ for all $w_{i'} \in U$ (use here $MP(K, L)$). Since K has finite index in L , $A \cap K$ has finite index in A and the set U can grow only finitely many times. Stop the process at the moment when, for some l , no word of length l can be added to U . At this moment we have exhausted the search inside the Schreier graph of A modulo $A \cap K$, and the existing list $U = \{1 = u_1, \dots, u_m\}$ is a set of left coset representatives of $A \cap K$ in A , say $A = \bigsqcup_{i=1}^m u_i(A \cap K)$. Now, for every $\alpha = 1, \dots, r$ and $i = 1, \dots, m$, compute $d(\alpha, i)$ such that $a_\alpha u_i(A \cap K) = u_{d(\alpha, i)}(A \cap K)$ (again using $MP(K, L)$). By the Reidemeister-Schreier method (see Theorem 2.7 in [27]), the set $\{u_{d(\alpha, i)}^{-1} a_\alpha u_i \mid \alpha = 1, \dots, r, i = 1, \dots, m\}$ generates $A \cap K$. Analogously, we can compute a set $V = \{v_1, \dots, v_n\}$ of left coset representatives for $B \cap K$ in B , say $B = \bigsqcup_{j=1}^n v_j(B \cap K)$, together with a set of generators for $B \cap K$. Clearly,

$$\begin{aligned} xA \cap yB &= \left(\bigsqcup_{i=1}^m x u_i(A \cap K) \right) \cap \left(\bigsqcup_{j=1}^n y v_j(B \cap K) \right) \\ &= \bigsqcup_{i,j} (x u_i(A \cap K) \cap y v_j(B \cap K)). \end{aligned}$$

Hence, $xA \cap yB \neq \emptyset$ is equivalent to $x u_i(A \cap K) \cap y v_j(B \cap K) \neq \emptyset$ for some $i = 1, \dots, m$ and some $j = 1, \dots, n$. For each (i, j) , consider the element $z_{i,j} = v_j^{-1} y^{-1} x u_i$ and rewrite it in the form $g_p k$ for some (unique) $p = 0, \dots, q$ and $k \in K$. If $p \neq 0$ (i.e. if $z_{i,j}$ does not belong to K) then the intersection $x u_i(A \cap K) \cap y v_j(B \cap K)$ is empty. Otherwise, $z_{i,j} \in K$ and we

are reduced to verify whether $z_{i,j}(A \cap K) \cap (B \cap K) \neq \emptyset$. This can be done using $\text{CIP}(K)$. \square

Let us apply this result to $GL_2(\mathbb{Z})$. It is well-known that $GL_2(\mathbb{Z})$ admits the following presentation

$$\begin{aligned} GL_2(\mathbb{Z}) &\cong D_4 \underset{D_2}{*} D_6 = \\ &= \langle t_4, x_4 \mid t_4^2, x_4^4, (t_4 x_4)^2 \rangle \left\langle \begin{array}{c} * \\ t_4 = t_6 \\ x_4^2 = x_6^3 \end{array} \right\rangle \langle t_6, x_6 \mid t_6^2, x_6^6, (t_6 x_6)^2 \rangle, \quad (2) \end{aligned}$$

where $t_4 = t_6 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, $x_4 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $x_6 = \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}$ (see Example I.5.2 in [14]). Since one can algorithmically go from matrices to words in the presentation, and viceversa, both models are algorithmically equivalent (and we shall use one or the other whenever convenient).

Corollary 6.3. *The problem $\text{CIP}(GL_2(\mathbb{Z}))$ is solvable.*

Proof. There is a natural epimorphism $\varphi: GL_2(\mathbb{Z}) \rightarrow D_{12} = \langle t_{12}, x_{12} \mid t_{12}^2, x_{12}^{12}, (t_{12} x_{12})^2 \rangle$ given by $t_4 = t_6 \mapsto t_{12}$, $x_4 \mapsto x_{12}^3$, $x_6 \mapsto x_{12}^2$. With a few straightforward calculations, one can show that $K = \ker \varphi$ is free of rank 2, with basis $P = [x_6, x_4] = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$ and $Q = [x_6^2, x_4] = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$. Since K has index $|D_{12}| = 24$ in $L = GL_2(\mathbb{Z})$, Propositions 6.1 and 6.2 conclude the proof. \square

Finally we see that, in general, orbit decidability goes up to finite index.

Proposition 6.4. *Let F be a group given in an algorithmic way, and let $A \leq B \leq \text{Aut}(F)$ be two subgroups given by finite sets of generators, such that A has finite index in B and $\text{MP}(A, B)$ is solvable. If $A \leq \text{Aut}(F)$ is orbit decidable, then $B \leq \text{Aut}(F)$ is orbit decidable.*

Proof. With a coset enumeration argument similar to that in the proof of Proposition 6.2, we can compute a (finite) list, say $\{\beta_1, \dots, \beta_m\}$, of left coset representatives of A in B , i.e. $B = \beta_1 A \sqcup \dots \sqcup \beta_m A$ (we formally need $\text{MP}(A, B)$ here, because B may have infinite index in $\text{Aut}(F)$). Now, for any given $u, v \in F$, the existence of $\beta \in B$ such that $u\beta$ is conjugate to v is equivalent to the existence of $i = 1, \dots, m$ and $\alpha \in A$ such that $(u\beta_i)\alpha$ is conjugate to v . Hence, orbit decidability for $B \leq \text{Aut}(F)$ follows from orbit decidability for $A \leq \text{Aut}(F)$. \square

6.1. The free abelian case. Let us concentrate on those short exact sequences in Theorem 3.1 with F being free abelian, say $F = \mathbb{Z}^n$, and look for orbit decidable subgroups of $\text{Aut}(\mathbb{Z}^n) = GL_n(\mathbb{Z})$.

To begin, it is an elementary fact in linear algebra that two vectors $u, v \in \mathbb{Z}^n$ lie in the same $GL_n(\mathbb{Z})$ -orbit if and only if the highest common

divisor of their entries coincide and, in this case, with the help of Euclid's algorithm, one can find an invertible matrix A such that $uA = v$. In other words,

Proposition 6.5. *The full automorphism group $GL_n(\mathbb{Z})$ of a finitely generated free abelian group \mathbb{Z}^n , is orbit decidable.* \square

Corollary 6.6. *Let $A_1, \dots, A_m \in GL_n(\mathbb{Z})$. If $\langle A_1, \dots, A_m \rangle = GL_n(\mathbb{Z})$, then the \mathbb{Z}^n -by- F_m group $G = \mathbb{Z}^n \rtimes_{A_1, \dots, A_m} F_m$ has solvable conjugacy problem.* \square

It is also a straightforward exercise in linear algebra to see that cyclic subgroups of $GL_n(\mathbb{Z})$ are orbit decidable. That is, given $A \in GL_n(\mathbb{Z})$ and $u, v \in \mathbb{Z}^n$ one can algorithmically decide whether $uA^k = v$ for some integer k : in fact, think A as a complex matrix, work out its Jordan form (approximating eigenvalues with enough accuracy) and then solve explicit equations (with the appropriate accuracy). This provides a solution to the conjugacy problem for cyclic extensions of \mathbb{Z}^n .

Proposition 6.7. *Cyclic subgroups of $GL_n(\mathbb{Z})$ are orbit decidable.* \square

Corollary 6.8. *\mathbb{Z}^n -by- \mathbb{Z} groups have solvable conjugacy problem.* \square

However, this was already known via an old result due to V.N. Remeslenikov, because \mathbb{Z}^n -by- \mathbb{Z} groups are clearly polycyclic. In [36] it was proven that polycyclic groups G are conjugacy separable. As a consequence, such a group, when given by an arbitrary finite presentation, has solvable conjugacy problem (use a brute force algorithm for solving $CP^+(G)$, and another for $CP^-(G)$ enumerating all maps into finite symmetric groups (i.e. onto finite groups) and checking whether the images of the given elements are conjugated down there). But, furthermore, this result can now be used to prove a more general fact about orbit decidability in $GL_n(\mathbb{Z})$.

J. Tits [40] proved the deep and remarkable fact that every finitely generated subgroup of $GL_n(\mathbb{Z})$ is either virtually solvable or it contains a non-abelian free subgroup. The following proposition says that the first kind of subgroups are always orbit decidable, so forcing orbit undecidable subgroups of $GL_n(\mathbb{Z})$ to contain non-abelian free subgroups. This somehow means that orbit undecidability in $GL_n(\mathbb{Z})$ is intrinsically linked to free-like structures.

Proposition 6.9. *Any virtually solvable subgroup of $GL_n(\mathbb{Z})$ is orbit decidable.*

Proof. Let B be a virtually solvable subgroup of $GL_n(\mathbb{Z})$, given by a finite generating set of matrices A_1, \dots, A_m . By a Theorem of A.I. Mal'cev

(see [28], or Chapter 2 in [37]), every solvable subgroup of $GL_n(\mathbb{Z})$ is polycyclic; so, $B = \langle A_1, \dots, A_m \rangle$ has a finite index subgroup $C \leq B$ which is polycyclic and, in particular, finitely presented.

Recurrently perform the following two lists: on one hand keep enumerating *all* finite presentations of *all* polycyclic groups (use a similar strategy as that in the proof of Theorem 4.8 above, enumerating first all canonical presentations of such groups, and diagonally applying all possible Tietze transformations to all of them). On the other hand, enumerate a set of pairs $(\mathcal{C}, \mathcal{M})$, where \mathcal{C} is a finite set of generators for a finite index subgroup C of B , and $\mathcal{M} = \{M_1, \dots, M_r\}$ is a finite set of matrices such that $B = M_1 \cdot C \cup \dots \cup M_r \cdot C$, and in such a way that C eventually visits *all* finite index subgroups of B ; we can do this in a similar way as in the proof of Lemma 4.6: enumerate all saturated and folded Stallings graphs with increasingly many vertices over the alphabet $\{A_1, \dots, A_m\}$, and map the corresponding finite index subgroup and finite set of coset representatives (one for each vertex) down to B , where possible repetitions may happen (see [39]).

These two lists are infinite so the started processes will never end; but, while running, keep choosing an element in each list in all possible ways, say $C' = \langle t_1, \dots, t_p \mid R_1, \dots, R_q \rangle$ and $(\mathcal{C}, \mathcal{M})$, and check whether there is an onto map from $\{t_1, \dots, t_p\}$ to \mathcal{C} that extend to an (epi)morphism $C' \rightarrow C$ (this just involves few matrix calculations in $GL_n(\mathbb{Z})$). Stop all the computations when finding such a map (which we are sure it exists because some finite index subgroup $C \leq B$ is polycyclic, and so isomorphic to one of the presentations in the first list). When this procedure terminates, we have got a finite presentation $\langle t_1, \dots, t_p \mid R_1, \dots, R_q \rangle$ of a polycyclic group C' and a map $C' \rightarrow C$ onto a finite index subgroup $C \leq B = \langle A_1, \dots, A_m \rangle \leq GL_n(\mathbb{Z})$, for which we also know a finite set of coset representatives \mathcal{M} , with possible repetitions.

Now, write down the natural presentation of the group $G = \mathbb{Z}^n \rtimes_C C'$. Since it is clearly polycyclic, Remeslennikov's result [36] tells us that G has solvable conjugacy problem (for instance, from the computed presentation). Thus, by Theorem 3.1 (a) \Rightarrow (c), the corresponding group of actions, $C \leq GL_n(\mathbb{Z})$, is orbit decidable (note that hypothesis (iii) of Theorem 3.1 may not be satisfied in this case, but it is not used in the proof of implication (a) \Rightarrow (c)). Finally, $B \leq GL_n(\mathbb{Z})$ is orbit decidable as well: given two vectors $u, v \in \mathbb{Z}^n$, deciding whether $v = uP$ for some $P \in B$ is the same as deciding whether $v = uM_iQ$ for some $i = 1, \dots, r$ and $Q \in C$, which reduces to finitely many claims to the orbit decidability of $C \leq GL_n(\mathbb{Z})$. \square

Corollary 6.10. *Let $A_1, \dots, A_m \in GL_n(\mathbb{Z})$. If $B = \langle A_1, \dots, A_m \rangle \leq GL_n(\mathbb{Z})$ is virtually solvable, then the \mathbb{Z}^n -by- F_m group $G = \mathbb{Z}^n \rtimes_{A_1, \dots, A_m} F_m$ has solvable conjugacy problem. \square*

Let us consider now finite index subgroups of $GL_n(\mathbb{Z})$.

Proposition 6.11. *Any finite index subgroup of $GL_n(\mathbb{Z})$ (given by generators) is orbit decidable.*

Proof. Let $B \leq GL_n(\mathbb{Z})$ be a finite index subgroup generated by some given matrices. Take your favorite presentation for $GL_n(\mathbb{Z})$ (see, for example, Section 3.5 of [27]) and write them in terms of it. With a similar argument as in the proof of Lemma 4.6, we can compute generators for the subgroup

$$A = \bigcap_{P \in GL_n(\mathbb{Z})} (P^{-1} \cdot B \cdot P) = B \cap B^{P_1} \cap \dots \cap B^{P_m} \leq GL_n(\mathbb{Z}),$$

where $Id = P_0, P_1, \dots, P_m$ is a set of right coset representatives for B in $GL_n(\mathbb{Z})$ (computable by Todd-Coxeter algorithm, see Theorem 4.5). By Proposition 6.4, we are reduced to see that $A \leq GL_n(\mathbb{Z})$ is orbit decidable.

Given $u, v \in \mathbb{Z}^n$, we have to decide whether some matrix of A sends u to v . Clearly, we can assume $u, v \neq 0$ and check whether there exists $M \in GL_n(\mathbb{Z})$ such that $uM = v$ (see Proposition 6.5). Once we have such M , the set of all those matrices carrying u to v is precisely $M \cdot Stab(v)$. And it is straightforward to compute a finite generating set for the stabilizer of v (it is conjugate to $Stab(1, 0, \dots, 0) = \left\{ \begin{pmatrix} 1 & 0 & \dots & 0 \\ * & * & & * \\ & & \ddots & \\ & & & * \end{pmatrix} \right\}$). It remains to algorithmically decide whether the intersection $A \cap (M \cdot Stab(v))$ is or is not empty; or, equivalently, whether $M \in A \cdot Stab(v)$ holds or not. This is decidable because $A \cdot Stab(v)$ is a finite index subgroup of $GL_n(\mathbb{Z})$ (here is where normality of A is needed) with a computable set of generators; hence, $MP(A \cdot Stab(v), GL_n(\mathbb{Z}))$ is solvable, again by Todd-Coxeter algorithm. \square

Corollary 6.12. *Let $A_1, \dots, A_m \in GL_n(\mathbb{Z})$. If $\langle A_1, \dots, A_m \rangle$ has finite index in $GL_n(\mathbb{Z})$ then the \mathbb{Z}^n -by- F_m group $\mathbb{Z}^n \rtimes_{A_1, \dots, A_m} F_m$ has solvable conjugacy problem. \square*

Finally, let us concentrate on rank two.

Proposition 6.13. *Every finitely generated subgroup of $GL_2(\mathbb{Z})$ is orbit decidable.*

Proof. Let $A_1, \dots, A_r \in GL_n(\mathbb{Z})$ be some given matrices and consider the subgroup they generate, $\langle A_1, \dots, A_r \rangle \leq GL_n(\mathbb{Z})$. For $n = 2$, given $u, v \in \mathbb{Z}^n$, let us decide whether there exists $A \in \langle A_1, \dots, A_r \rangle$ such that $uA = v$. We can clearly assume $u, v \neq 0$.

By Proposition 6.5, we can decide whether there exists $M \in GL_n(\mathbb{Z})$ such that $uM = v$ and, in the affirmative case, find such an M . And it is straightforward to find a set of generators for the stabilizer of v , $Stab(v) = \{B \in GL_n(\mathbb{Z}) \mid vB = v\}$, say $\{B_1, \dots, B_s\}$ (in the case $n = 2$, every such stabilizer is conjugate to $Stab(1, 0) = \langle (\begin{smallmatrix} 1 & 0 \\ 1 & 1 \end{smallmatrix}), (\begin{smallmatrix} 1 & 0 \\ 0 & -1 \end{smallmatrix}) \rangle$). Then, the matrices sending u to v are precisely those contained in the coset $M\langle B_1, \dots, B_s \rangle$. So, it remains to decide whether $\langle A_1, \dots, A_r \rangle \cap M\langle B_1, \dots, B_s \rangle$ is empty or not.

In the case $n = 2$, this can be done algorithmically (see Corollary 6.3). \square

Note that the proof of Proposition 6.13 works for every dimension n except at the end, when Corollary 6.3 is used. We shall refer to this fact later.

Corollary 6.14. *All \mathbb{Z}^2 -by-[f.g. free] groups have solvable conjugacy problem.* \square

6.2. The free case. Following the same route as in the previous subsection, let us concentrate now on those short exact sequences in Theorem 3.1 with F being free, say $F = F_n$, and look for orbit decidable subgroups of $Aut(F_n)$.

To begin, classical Whitehead algorithm (see Proposition I.4.19 in [25]) decides, given $u, v \in F_n$, whether there exists an automorphism of F_n sending u to v up to conjugacy. In other words,

Theorem 6.15 (Whitehead, [41]). *The full automorphism group $Aut(F_n)$ of a finitely generated free group F_n , is orbit decidable.* \square

Corollary 6.16. *Let F_n be a finitely generated free group. If $\varphi_1, \dots, \varphi_m$ generate $Aut(F_n)$, then the F_n -by- F_m group $G = F_n \rtimes_{\varphi_1, \dots, \varphi_m} F_m$ has solvable conjugacy problem.* \square

Like in the abelian case, cyclic subgroups of $Aut(F_n)$ are orbit decidable by a result of P. Brinkmann. This is the analog of Proposition 6.7 for the free case, but here the proof is much more complicated, making strong use of the theory of train-tracks. This was already used to solve the conjugacy problem for free-by-cyclic groups:

Theorem 6.17 (Brinkmann, [10]). *Cyclic subgroups of $Aut(F_n)$ are orbit decidable.* \square

Corollary 6.18 (Bogopolski-Martino-Maslakova-Ventura, [6]). *[f.g. free]-by-cyclic groups have solvable conjugacy problem.* \square

The analog of Proposition 6.9 and Corollary 6.10 in the free setting is not known, and seems to be an interesting and much more complicated problem. See Question 5 in the last section for some comments about it, and a clear relation with Tits alternative for $Aut(F_n)$.

Let us now consider finite index subgroups of $Aut(F_n)$.

Proposition 6.19. *Let F_n be a finitely generated free group. Any finite index subgroup of $\text{Aut}(F_n)$ (given by generators) is orbit decidable.*

Proof. Let $B \leq \text{Aut}(F_n)$ be a finite index subgroup generated by some given automorphisms. Consider Nielsen's presentation of $\text{Aut}(F_n)$ (see Proposition N1 in Section 3.5 of [27]) and write them in terms of this presentation (i.e. as products of Nielsen automorphisms). Then, with a similar argument as in the proof of Lemma 4.6, we can compute generators for the subgroup

$$A = \bigcap_{\phi \in \text{Aut}(F_n)} (\phi^{-1}B\phi) = B \cap B^{\phi_1} \cap \dots \cap B^{\phi_m} \leq \text{Aut}(F_n),$$

where $Id = \phi_0, \phi_1, \dots, \phi_m$ is a set of right coset representatives for B in $\text{Aut}(F_n)$ (computable by Todd-Coxeter algorithm, see Theorem 4.5). By Proposition 6.4, we are reduced to see that $A \leq \text{Aut}(F_n)$ is orbit decidable.

Let $u, v \in F_n$. Using Whitehead's algorithm, we can check whether there exists an automorphism $\alpha \in \text{Aut}(F_n)$ carrying u to v . Once we have such α , the set of all those automorphisms carrying u to a conjugate of v is precisely $\alpha \cdot \text{Stab}(v) \cdot \text{Inn}(F_n)$. By McCool's algorithm (see Proposition I.5.7 in [25]), we can compute a finite generating set for the stabilizer of v . It remains to algorithmically decide whether the intersection $A \cap (\alpha \cdot \text{Stab}(v) \cdot \text{Inn}(F_n))$ is or is not empty; or, equivalently, whether $\alpha \in A \cdot \text{Stab}(v) \cdot \text{Inn}(F_n)$ holds or not. This is decidable because $A \cdot \text{Stab}(v) \cdot \text{Inn}(F_n)$ is a finite index subgroup of $\text{Aut}(F_n)$ (here is where normality of A is needed) with a computable set of generators; hence, $\text{MP}(A \cdot \text{Stab}(v) \cdot \text{Inn}(F_n), \text{Aut}(F_n))$ is solvable, again by Todd-Coxeter algorithm. \square

Corollary 6.20. *Let F_n be a finitely generated free group. If $\varphi_1, \dots, \varphi_m$ generate a finite index subgroup of $\text{Aut}(F_n)$, then the F_n -by- F_m group $G = F_n \rtimes_{\varphi_1, \dots, \varphi_m} F_m$ has solvable conjugacy problem.* \square

Now, let us concentrate on rank two. Like in the abelian case, we have

Proposition 6.21. *Let F_2 be the free group of rank two. Then every finitely generated subgroup of $\text{Aut}(F_2)$ is orbit decidable.*

Proof. Let A be a finitely generated subgroup of $\text{Aut}(F_n)$ and let $u, v \in F_n$ be given. For $n = 2$, we have to decide whether there exists $\varphi \in A$ such that $u\varphi$ is conjugate to v .

Mimicking the proof of Proposition 6.13, let us apply Whitehead's algorithm to u, v (see Proposition I.4.19 in [25]). If there is no automorphism in $\text{Aut}(F_n)$ sending u to v then, clearly, the answer to our problem is also negative. Otherwise, we have found $\alpha \in \text{Aut}(F_n)$ such that $u\alpha = v$. Now, the set of all such automorphisms of F_n is $\alpha \cdot \text{Stab}(v)$. And the set of all automorphisms of F_n mapping u to a conjugate of v is $\alpha \cdot \text{Stab}(v) \cdot \text{Inn}(F_n)$.

By McCool's algorithm (see Proposition I.5.7 in [25]), we can find a finite system of generators for $Stab(v) \leq Aut(F_n)$. Finally, all we need is to verify whether $A \cap (\alpha \cdot Stab(v) \cdot Inn(F_n))$ is or is not empty.

In the case $n = 2$ this can be done algorithmically: since the kernel of the canonical projection $\bar{\cdot}: Aut(F_2) \rightarrow GL_2(\mathbb{Z})$ is $Inn(F_2)$ (which is a very special fact of the rank 2 case), our goal is equivalent to verifying whether $\bar{A} \cap (\bar{\alpha} \cdot \overline{Stab(v)})$ is or is not empty in $GL_2(\mathbb{Z})$. This can be done by Corollary 6.3. \square

Note that the proof of Proposition 6.21 works for every rank n except for the last paragraph, exactly like in Proposition 6.13. We shall refer to this fact later.

Corollary 6.22. *All F_2 -by- $[f.g. free]$ groups have solvable conjugacy problem.* \square

Another nice examples of orbit decidable subgroups in $Aut(F_n)$ come from geometry. Certain mapping class groups of surfaces with boundary and punctures turn out to embed in the automorphism group of the free group of the appropriate rank. The image of these embeddings are easily seen to be orbit decidable in two special cases. From [15] we extract the following two examples.

Let $S_{g,b,n}$ be an orientable surface of genus g , with b boundary components and n punctures. It is well known that its fundamental group has presentation

$$\begin{aligned} \Sigma_{g,b,n} = \\ = \langle x_1, y_1, \dots, x_g, y_g, z_1, \dots, z_b, t_1, \dots, t_n \mid [x_1, y_1] \cdots [x_g, y_g] z_1 \cdots z_b t_1 \cdots t_n = 1 \rangle, \end{aligned}$$

and, except for $b = n = 0$, is a free group of rank $2g + b + n - 1$. In the following cases, the mapping class group of $S_{g,b,n}$ can be viewed as a subgroup of $Aut(F_{2g+b+n-1})$, see [15] for details:

Proposition 6.23. (see [15]) *Let $S_{g,b,n}$ be an orientable surface of genus g , with b boundary components, and with n punctures.*

- (i) (**Maclachlan**) *The positive pure mapping class group of $S_{g,0,n+2}$ becomes (when the basepoint is taken to be the $(n+2)$ nd puncture) the group $Aut_{g,0,1+n+1}^+$ of automorphisms of $\Sigma_{g,0,n+1} \simeq F_{2g+n}$ which fix each conjugacy class $[t_j]$, $j = 1, \dots, n+1$ (the case $g = 0$ gives the pure braid group on $n+1$ strings modulo the center, $B_{n+1}/Z(B_{n+1})$).*
- (ii) (**A'Campo**) *The positive mapping class group of $S_{g,1,n}$ becomes (when the basepoint is taken to be on the boundary) the group*

$Aut_{g,\hat{1},n}^+$ of automorphisms of $\Sigma_{g,1,n} \simeq F_{2g+n}$ which fix z and permute the set of conjugacy classes $\{[t_1], \dots, [t_n]\}$ (the case $g = 0$ gives the braid group on n strings, B_n).

A particularly interesting case is when $g = 0$ in (ii) above: $Aut_{0,\hat{1},n}^+$ is the image of classical Artin's embedding of the braid group on n strings into $Aut(F_n)$, sending generator $\sigma_i \in B_n$ ($i = 1, \dots, n-1$) to $\sigma_i: F_n \rightarrow F_n$, $t_i \mapsto t_i t_{i+1} t_i^{-1}$, $t_{i+1} \mapsto t_i$, $t_j \mapsto t_j$ for $j \neq i, i+1$. The subgroup $Aut_{0,\hat{1},n}^+ = \langle \sigma_1, \dots, \sigma_{n-1} \rangle \leq Aut(F_n)$ is then characterized as those automorphisms $\varphi \in Aut(F_n)$ for which $(t_1 t_2 \cdots t_n) \varphi = t_1 t_2 \cdots t_n$ and there exist words $w_1, \dots, w_n \in F_n$ and a permutation σ of the set of indices such that $t_i \varphi = w_i^{-1} t_{\sigma(i)} w_i$.

All these groups of automorphisms, $Aut_{g,0,1^{\pm n+1}}^+ \leq Aut(F_{2g+n})$ and $Aut_{g,\hat{1},n}^+ \leq Aut(F_{2g+n})$, are easily seen to be orbit decidable because of the following observation.

Proposition 6.24. *Let F_n be a finitely generated free group, and let $u_i, v_i \in F_n$ ($i = 0, \dots, m$) be two lists of elements. Then,*

- (i) $A = \{\varphi \mid u_0 \varphi = v_0, u_1 \varphi \sim v_1, \dots, u_m \varphi \sim v_m\} \leq Aut(F_n)$ and
- (ii) $B = \{\varphi \mid u_0 \varphi = v_0, u_1 \varphi \sim v_{\sigma(1)}, \dots, u_m \varphi \sim v_{\sigma(m)} \text{ for some } \sigma \in Sym(m)\} \leq Aut(F_n)$

are orbit decidable.

Proof. For (i), given $u, v \in F_n$, we have to decide whether there exists an automorphism $\varphi \in Aut(F_n)$ such that $u_0 \varphi = v_0$, $[u_1] \varphi = [v_1], \dots, [u_m] \varphi = [v_m]$ and $[u] \varphi = [v]$, where brackets denote conjugacy classes. This is the same as deciding whether there exists φ such that $[u_0] \varphi = [v_0]$, $[u_1] \varphi = [v_1], \dots, [u_m] \varphi = [v_m]$ and $[u] \varphi = [v]$ (and, in the affirmative case, composing φ by $\gamma_{w_0^{-1}}$, where w_0 is the first conjugator, $u_0 \varphi = w_0^{-1} v_0 w_0$). One can make this decision by applying Proposition 4.21 in [25]. Finally, (ii) can be solved by using up to $m!$ many times the solution given for (i). \square

It is worth mentioning that D. Larue in his PhD thesis [24] analyzed the $Aut_{0,\hat{1},n}^+$ -orbit of t_1 in $\Sigma_{0,1,n}$ (i.e. the B_n -orbit of t_1 in F_n) and he provided an algorithm to decide whether a given word $w \in F_n$ belongs to this orbit (note that this is not exactly a special case of $OD(B_n)$ because B_n does not contain all inner automorphisms). Although working only for the orbit of t_1 , the algorithm provided is faster and nicer than that provided in Proposition 6.24.

Corollary 6.25. *Let F_{2g+n} be a finitely generated free group. If $\varphi_1, \dots, \varphi_m \in Aut(F_{2g+n})$ generate the positive pure mapping class group $Aut_{g,0,1^{\pm n+1}}^+$, or the positive mapping class group $Aut_{g,\hat{1},n}^+$ then the F_{2g+n} -by- F_m group*

$G = F_{2g+n} \rtimes_{\varphi_1, \dots, \varphi_m} F_m$ has solvable conjugacy problem (a particular case of this being when $\varphi_1, \dots, \varphi_m$ generate the standard copy of the braid group $B_n \leq \text{Aut}(F_n)$). \square

7. NEGATIVE RESULTS

Let us construct now negative examples, namely orbit undecidable subgroups of $GL_n(\mathbb{Z})$ and $\text{Aut}(F_n)$ which, of course, will correspond to [f.g. free abelian]-by-[f.g. free] and [f.g. free]-by-[f.g. free] groups with unsolvable conjugacy problem.

As mentioned above, Miller constructed a [f.g. free]-by-[f.g. free] group with unsolvable conjugacy problem (see [30]); here, we have a first source of examples of finitely generated subgroups of the automorphism group of a free group, which are orbit undecidable. In the present section, we will generalize this construction by giving a source of orbit undecidability in $\text{Aut}(F)$ for more groups F . When taking F to be free, this will reproduce Miller's example; when taking $F = \mathbb{Z}^n$ for $n \geq 4$, we will obtain orbit undecidable subgroups in $GL_n(\mathbb{Z})$, which correspond to the first known examples of [f.g. free abelian]-by-[f.g. free] groups with unsolvable conjugacy problem.

Let us recall Miller's construction. It begins with an arbitrary finite presentation, $H = \langle s_1, \dots, s_n \mid R_1, \dots, R_m \rangle$, where the R_j 's are words on the s_i 's. Let $F_{n+1} = \langle q, s_1, \dots, s_n \mid \rangle$ and $F_{m+n} = \langle t_1, \dots, t_m, d_1, \dots, d_n \mid \rangle$ be the free groups of rank $n+1$ and $m+n$, respectively, on the listed generators. Consider now the $m+n$ automorphisms of F_{n+1} given by

$$\begin{array}{ll} \alpha_i: F_{n+1} & \rightarrow F_{n+1} \\ q & \mapsto qR_i \\ s_k & \mapsto s_k \end{array} \quad \begin{array}{ll} \beta_j: F_{n+1} & \rightarrow F_{n+1} \\ q & \mapsto s_j^{-1}q s_j, \\ s_k & \mapsto s_k \end{array}$$

for $i = 1, \dots, m$ and $j, k = 1, \dots, n$, and denote the group of automorphisms they generate by $A(H) \leq \text{Aut}(F_{n+1})$. Next, consider the F_{n+1} -by- F_{m+n} group defined by these automorphisms,

$$G(H) = F_{n+1} \rtimes_{\alpha_1, \dots, \alpha_m, \beta_1, \dots, \beta_n} F_{m+n}.$$

The following Theorem is Corollary 5 in Chapter III of [30]. Below, we shall provide an alternative proof.

Theorem 7.1 (Miller, [30]). *If H has unsolvable word problem then $G(H)$ has unsolvable conjugacy problem.*

So, applying Miller's construction to a presentation H with n generators, m relations, and with unsolvable word problem, one obtains a $(m+n)$ -generated subgroup of $\text{Aut}(F_{n+1})$, namely $A(H)$, which is orbit undecidable.

In [8], V. Borisov constructed a group presented with 4 generators, 12 relations, and having unsolvable word problem. In order to reduce the number of generators to 2 (and so have the corresponding orbit undecidable subgroup living inside $Aut(F_3)$) we can use Higman-Neumann-Neumann embedding theorem, saying that any countable group can be embedded in a group with two generators and the same number of relations (see [21]). Since solvability of the word problem clearly passes to subgroups, we obtain a group with $n = 2$ generators, $m = 12$ relations, and having unsolvable word problem. Using Miller's construction we conclude the existence of a F_3 -by- F_{14} group with unsolvable conjugacy problem. In other words,

Corollary 7.2. *There exists a 14-generated subgroup $A \leq Aut(F_3)$ which is orbit undecidable.* \square

Let us now find a more general source of orbit undecidability that will apply to more groups F other than free (and, in the free case, will coincide with Miller's example via Mihailova's result).

Let F be a group. Recall that the *stabilizer* of a given subgroup $K \leq F$, denoted $Stab(K)$, is

$$Stab(K) = \{\varphi \in Aut(F) \mid k\varphi = k \quad \forall k \in K\} \leq Aut(F).$$

For simplicity, we shall write $Stab(k)$ to denote $Stab(\langle k \rangle)$, $k \in F$. Furthermore, we define the *conjugacy stabilizer* of K , denoted $Stab^*(K)$, to be the set of automorphisms acting as conjugation on K , formally $Stab^*(K) = Stab(K) \cdot Inn(F) \leq Aut(F)$.

Proposition 7.3. *Let F be a group. Suppose we are given two subgroups $A \leq B \leq Aut(F)$ and an element $v \in F$ such that $B \cap Stab^*(v) = \{Id\}$. If $A \leq Aut(F)$ is orbit decidable then $MP(A, B)$ is solvable.*

Proof. Given $\psi \in B \leq Aut(F)$, let us decide whether $\psi \in A$ or not. Take $w = v\psi$ and observe that

$$\{\phi \in B \mid v\phi \sim w\} = B \cap (Stab^*(v) \cdot \psi) = (B \cap Stab^*(v)) \cdot \psi = \{\psi\}.$$

So, there exists $\phi \in A$ such that $v\phi$ is conjugate to w in F , if and only if $\psi \in A$. Hence, orbit decidability for $A \leq Aut(F)$ solves $MP(A, B)$. \square

One can interpret Proposition 7.3 by saying that if, for a group F , $Aut(F)$ contains a pair of subgroups $A \leq B \leq Aut(F)$ with unsolvable $MP(A, B)$ then $A \leq Aut(F)$ is orbit undecidable.

The most classical example of unsolvability of the membership problem goes back to fifty years ago. In [29] (see also Chapter III.C of [30]) Mihailova gave a nice example of unsolvability of the membership problem. The construction goes as follows.

Like before, start with an arbitrary finite presentation, $H = \langle s_1, \dots, s_n \mid R_1, \dots, R_m \rangle$, and consider the subgroup $A = \{(x, y) \in F_n \times F_n \mid x =_H y\} \leq F_n \times F_n$. It is straightforward to verify that $A = \langle (1, R_1), \dots, (1, R_m), (s_1, s_1), \dots, (s_n, s_n) \rangle$ (and so it is finitely generated), and that $\text{MP}(A, F_n \times F_n)$ is solvable if and only if $\text{WP}(H)$ is solvable.

By Higman-Neumann-Neumann embedding theorem, we can restrict our attention to 2-generated groups (take $n = 2$ in the above paragraph). From all this, we deduce the following.

Theorem 7.4. *Let F be a finitely generated group such that $F_2 \times F_2$ embeds in $\text{Aut}(F)$ in such a way that the image intersects trivially with $\text{Stab}^*(v)$, for some $v \in F$. Then, $\text{Aut}(F)$ contains an orbit undecidable subgroup; in other words, there exist F -by- $[f.g. \text{ free}]$ groups with unsolvable conjugacy problem. \square*

In the rest of the section, we shall use Theorem 7.4 to obtain explicit examples in the free abelian and free cases.

7.1. The free abelian case. It is well known that F_2 embeds in $GL_2(\mathbb{Z})$ and so, $F_2 \times F_2$ embeds in $GL_4(\mathbb{Z})$. Hence, we can deduce the following result.

Proposition 7.5. *For $n \geq 4$, $GL_n(\mathbb{Z})$ contains finitely generated orbit undecidable subgroups.*

Proof. Consider the subgroup of $GL_2(\mathbb{Z})$ generated by $P = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$ and $Q = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$, which is free and freely generated by $\{P, Q\}$ as discussed in the proof of Corollary 6.3. We claim that $\langle P, Q \rangle \cap \text{Stab}^*((1, 0)) = \langle \begin{pmatrix} 1 & 0 \\ 12 & 1 \end{pmatrix} \rangle$. In fact, it is clear that $\text{Stab}^*((1, 0)) = \text{Stab}((1, 0)) = \left\{ \begin{pmatrix} 1 & 0 \\ n & \pm 1 \end{pmatrix} \mid n \in \mathbb{Z} \right\}$ (and we can forget the negative signum because we are interested in the intersection with $\langle P, Q \rangle \leq SL_2(\mathbb{Z})$). Now, the image of $\begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} = x_6^{-1}x_4$ under φ is $x_{12}^{-2}x_{12}^3 = x_{12} \in D_{12}$ (see the proof of Corollary 6.3 for notation). So,

$$\langle P, Q \rangle \cap \text{Stab}^*((1, 0)) = \ker \varphi \cap \text{Stab}((1, 0)) = \langle (x_6^{-1}x_4)^{12} \rangle = \langle \begin{pmatrix} 1 & 0 \\ 12 & 1 \end{pmatrix} \rangle.$$

Choose now a (free) subgroup $\langle P', Q' \rangle \leq \langle P, Q \rangle$ intersecting trivially with the cyclic subgroup $\langle \begin{pmatrix} 1 & 0 \\ 12 & 1 \end{pmatrix} \rangle$ (this always exists in non-cyclic free groups). And, for $n \geq 4$, consider

$$B = \left\langle \left(\begin{array}{c|c} P' & 0 \\ \hline 0 & Id \end{array} \right), \left(\begin{array}{c|c} Q' & 0 \\ \hline 0 & Id \end{array} \right), \left(\begin{array}{c|c} Id & 0 \\ \hline 0 & P' \end{array} \right), \left(\begin{array}{c|c} Id & 0 \\ \hline 0 & Q' \end{array} \right) \right\rangle \leq \\ \leq GL_4(\mathbb{Z}) \leq GL_n(\mathbb{Z}),$$

which is clearly isomorphic to $F_2 \times F_2$. By construction, B intersects trivially with the (conjugacy) stabilizer of $v = (1, 0, 1, 0, \dots, 0) \in \mathbb{Z}^n$. Finally, using Mihaилоva's construction, find a finitely generated subgroup $A \leq B$ with

unsolvable $\text{MP}(A, B)$. By Proposition 7.3 applied to $F = \mathbb{Z}^n$, A is a finitely generated orbit undecidable subgroup of $\text{Aut}(\mathbb{Z}^n) = \text{GL}_n(\mathbb{Z})$. \square

Corollary 7.6. *There exist \mathbb{Z}^4 -by-[f.g. free] groups with unsolvable conjugacy problem.* \square

While the constructions are quite different, these groups are reminiscent of Miller's examples, but with a free abelian base group. To the best of our knowledge, they are the first known examples of [f.g. free abelian]-by-[f.g. free] groups with unsolvable conjugacy problem. As a side consequence of the previous reasoning, we also obtain the following corollary.

Corollary 7.7. *For $n \geq 4$, $\text{CIP}(\text{GL}_n(\mathbb{Z}))$ is unsolvable.*

Proof. As noted above, the proof of Proposition 6.13 works entirely for any dimension n for which $\text{CIP}(\text{GL}_n(\mathbb{Z}))$ is solvable (for example, $n = 2$). But Proposition 7.5 states the existence of finitely generated orbit undecidable subgroups of $\text{GL}_n(\mathbb{Z})$, for $n \geq 4$. Hence, $\text{CIP}(\text{GL}_n(\mathbb{Z}))$ must be unsolvable in this case. \square

7.2. The free case. In order to apply Theorem 7.4 to the free group $F_3 = \langle q, a, b \mid \ \rangle$ of rank 3, we need to identify a copy of $F_2 \times F_2$ inside $\text{Aut}(F_3)$. For every $w \in \langle a, b \rangle$, consider the automorphisms ${}_w\theta_1: F_3 \rightarrow F_3$, $q \mapsto wq$, $a \mapsto a$, $b \mapsto b$, and ${}_1\theta_w: F_3 \rightarrow F_3$, $q \mapsto qw$, $a \mapsto a$, $b \mapsto b$. Clearly, ${}_w\theta_1 {}_{w_2}\theta_1 = {}_{w_1 w_2}\theta_1$ and ${}_1\theta_{w_1} {}_1\theta_{w_2} = {}_1\theta_{w_2 w_1}$, which means that $\{ {}_w\theta_1 \mid w \in \langle a, b \rangle \} \simeq F_2$ and $\{ {}_1\theta_w \mid w \in \langle a, b \rangle \} \simeq F_2^{\text{op}} \simeq F_2$. It is also clear that ${}_w\theta_1 {}_1\theta_{w_2} = {}_{w_1}\theta_{w_2} = {}_1\theta_{w_2} {}_{w_1}\theta_1$ (with the natural definition for ${}_w\theta_{w_2}$). So, we have an embedding $F_2 \times F_2 \simeq F_2^{\text{op}} \times F_2^{\text{op}} \rightarrow \text{Aut}(F_3)$ given by $(w_1, w_2) \mapsto {}_{w_1^{-1}}\theta_{w_2}$, whose image is

$$B = \langle {}_{a^{-1}}\theta_1, {}_{b^{-1}}\theta_1, {}_1\theta_a, {}_1\theta_b \rangle = \{ {}_{w_1}\theta_{w_2} \mid w_1, w_2 \in \langle a, b \rangle \} \leq \text{Aut}(F_3).$$

In order to use Proposition 7.3, let us consider the element $v = qaqbq$. We claim that $B \cap \text{Stab}^*(v) = \{Id\}$. In fact, suppose $w_1, w_2 \in \langle a, b \rangle$ are such that $(v) {}_{w_1}\theta_{w_2} = w_1 q w_2 a w_1 q w_2 b w_1 q w_2$ is conjugate to $v = qaqbq$ in F_3 . Since both words have exactly three occurrences of q , they must agree up to cyclic reordering. That is, $q(w_2 a w_1) q (w_2 b w_1) q (w_2 w_1)$ equals either $qaqbq$, or $qbq^2 a$, or $q^2 a q b$. From this, one can straightforwardly deduce that $w_1 = w_2 = 1$ in all three cases. Thus, ${}_{w_1}\theta_{w_2} = Id$ proving the claim.

Now, let $H = \langle a, b \mid R_1, \dots, R_{12} \rangle$ be Borisov's example of a group with unsolvable word problem, embedded in a 2-generated group via Higman-Neumann-Neumann embedding (see above and [21]). By Mihailova result and Proposition 7.3, $A = \langle {}_1\theta_{R_1}, \dots, {}_1\theta_{R_{12}}, {}_{a^{-1}}\theta_a, {}_{b^{-1}}\theta_b \rangle \leq \text{Aut}(F_3)$ is orbit undecidable. Hence, by Theorem 3.3, the F_3 -by- F_{14} group determined by

the automorphisms ${}_1\theta_{R_1}, \dots, {}_1\theta_{R_{12}}, a^{-1}\theta_a, b^{-1}\theta_b \in \text{Aut}(F_3)$,

$$G = \left\langle q, a, b, t_1, \dots, t_{12}, d_1, d_2 \left| \begin{array}{l} t_i^{-1}qt_i = qR_i, d_1^{-1}qd_1 = a^{-1}qa, d_2^{-1}qd_2 = b^{-1}qb \\ t_i^{-1}at_i = a, d_1^{-1}ad_1 = a, d_2^{-1}ad_2 = a \\ t_i^{-1}bt_i = b, d_1^{-1}bd_1 = b, d_2^{-1}bd_2 = b \end{array} \right. \right\rangle,$$

has unsolvable conjugacy problem. This is precisely Miller's group $G(H)$ associated to $H = \langle a, b \mid R_1, \dots, R_{12} \rangle$ (see the beginning of the present section). Thus, the argument just given provides an alternative proof of Miller's Theorem 7.1.

Corollary 7.8. *For $n \geq 3$, $\text{CIP}(\text{Aut}(F_n))$ is unsolvable.*

Proof. As noted above, the proof of Proposition 6.21 works entirely for any rank n for which $\text{CIP}(\text{Aut}(F_n))$ is solvable. But, for $n \geq 3$, $\text{Aut}(F_n)$ contains finitely generated orbit undecidable subgroups. Hence, $\text{CIP}(\text{Aut}(F_n))$ must be unsolvable in this case. \square

8. OPEN PROBLEMS

Finally, we collect several questions suggested by the previous results.

Question 1. Apart from finitely generated abelian, free, surface and polycyclic groups, and virtually all of them (see Theorem 4.8), find more examples of groups F with solvable twisted conjugacy problem.

Commentary. As mentioned in Section 4, for every group F with solvable twisted conjugacy problem, the study of orbit decidability/undecidability among subgroups of $\text{Aut}(F)$ becomes interesting because it directly corresponds to solving the conjugacy problem for some extensions of F . \square

Question 2. Is the twisted conjugacy problem solvable for finitely generated hyperbolic groups ?

Commentary. The first step into this direction is the solvability of the twisted conjugacy problem for finitely generated free groups, proven in [6]. However, there is no hope to extend that proof for hyperbolic groups because we do not have enough control on the automorphism group of an arbitrary hyperbolic group. \square

Question 3. Let F be a group given by a finite presentation $\langle X \mid R \rangle$, and suppose we are given a set of words $\{w_1, \dots, w_r\}$ on X such that $K = \langle w_1, \dots, w_r \rangle \leq F$ is a finite index subgroup. Does solvability of $\text{TCP}(F)$ imply solvability of $\text{TCP}(K)$? Is it true with the extra assumption that K is characteristic in F ?

Commentary. The reverse implication is proved to be true in Proposition 4.7 (i), under the characteristic assumption for K . However, to go down from F to K we would have to consider the apparently more complicated problem of dealing with possible automorphism of K which do not extend to automorphisms of F . Maybe this is a strong enough reason to build a counterexample. Note that the answer to the non-twisted version of the same question is *no* by a result of Collins-Miller (see [12]). \square

Question 4. Let F be \mathbb{Z}^n or F_n , and let $A \leq B \leq \text{Aut}(F)$ be two subgroups given by finite sets of generators, such that A has finite index in B , and $\text{MP}(A, B)$ is solvable. Is it true that orbit decidability for $B \leq \text{Aut}(F)$ implies orbit decidability for $A \leq \text{Aut}(F)$?

Commentary. The reverse implication is proven in Proposition 6.4. With the first argument there, a finite list of left coset representatives of A in B can be computed, say $B = \beta_1 A \sqcup \cdots \sqcup \beta_m A$. Then, given $u, v \in F$, and assuming we got $\beta \in B$ such that $v \sim u\beta$, the set of all such β 's is $\beta \cdot \text{Stab}(v) \cdot \text{Inn}(F)$. Since generators for $\text{Stab}(v) \cdot \text{Inn}(F)$ are computable (by straightforward matrix calculations in the case $F = \mathbb{Z}^n$, and by McCool's algorithm in the case $F = F_n$), it only remains to decide whether the intersection $\beta \cdot \text{Stab}(v) \cdot \text{Inn}(F) \cap A$ is empty or not. This can be done in the case $n = 2$ because $\text{CIP}(GL_2(\mathbb{Z}))$ is solvable (see the proof of Proposition 6.21). However, this last part of the argument does not work in the other cases, because $\text{CIP}(\text{Aut}(F))$ is unsolvable for $F = \mathbb{Z}^n$ with $n \geq 4$ (see Corollary 7.7), and for $F = F_n$ with $n \geq 3$ (see Corollary 7.8).

As partial answers, note that Propositions 6.11 and 6.19 show that the answer is *yes* in the special cases where $F = \mathbb{Z}^n$ and $B = GL_n(\mathbb{Z})$, and where $F = F_n$ and $B = \text{Aut}(F_n)$, respectively.

We formulate the question for free and free abelian groups because, if one allows an arbitrary ambient F , then the answer is negative: consider Collins-Miller example of a finitely presented group G with an index two subgroup $F \leq G$ such that $\text{CP}(G)$ is solvable but $\text{CP}(F)$ is unsolvable (see [12]); furthermore, G contains an element $g_0 \in G$ of order two which acts on F as a non-inner automorphism $\gamma_{g_0} \in \text{Aut}(F)$. So, we have the short exact sequence $1 \rightarrow F \rightarrow G \rightarrow C_2 \rightarrow 1$. Since $\text{CP}(G)$ is solvable, the action subgroup $B = \{Id, \gamma_{g_0}\} \leq \text{Aut}(F)$ is orbit decidable; however, $\text{CP}(F)$ is unsolvable meaning that the trivial subgroup $A = \{Id\} \leq \text{Aut}(F)$ is orbit undecidable. \square

Question 5. Is any virtually solvable subgroup of $\text{Aut}(F_n)$ orbit decidable?

Commentary. This is the analog of Proposition 6.9 in the free setting. However, it reduces to the same question for virtually free abelian subgroups. In

fact, Bestvina-Feighn-Handel proved in [1] that every solvable subgroup of $Out(F_n)$ contains a finitely generated free abelian subgroup of index at most 3^{5n^2} (additionally, it is also known that free abelian subgroups of $Out(F_n)$ have rank at most $2n - 3$). And the same is true for $Aut(F_n)$ because one can easily embed $Aut(F_n)$ in $Out(F_{2n})$ by sending $\alpha \in Aut(F_n)$ to the outer automorphism of F_{2n} which acts as α on both the first half and the second half of the generating set. So, the situation is formally simpler than in Proposition 6.9, but the argument there does not work here because we cannot use the trick about polycyclic groups. Apart from the possible finite index step, this question asks for a multidimensional version of Brinkmann's result (Theorem 6.17). So, due to the complexity of the proof and solution for the cyclic case, it seems a quite difficult question.

It is worth remarking that Bestvina-Feighn-Handel also proved in [2] a strong version of Tits alternative for $Out(F_n)$: every subgroup of $Out(F_n)$ is either virtually solvable (and hence virtually free abelian) or contains a non-abelian free group. Since the same is true for subgroups of $Aut(F_n)$ via the above embedding, an affirmative answer to Question 5 would then force orbit undecidable subgroups of $Aut(F_n)$ to contain non-abelian free subgroups, like in the abelian context. This would intuitively confirm that, again, orbit undecidability is intrinsically linked to free-like structures. \square

Question 6. Is any finitely presented subgroup of $Aut(F_n)$ orbit decidable?

Commentary. This question contains Question 5, so it is even more difficult to be answered in the affirmative. Note that orbit undecidable subgroups of the form $A = \langle {}_1\theta_{R_1}, \dots, {}_1\theta_{R_{12}}, {}_{a^{-1}}\theta_a, {}_{b^{-1}}\theta_b \rangle \leq Aut(F_3)$ corresponding to Miller's examples (see subsection 7.2) are *not* a counterexample to this question because they are not finitely presented by Proposition B in Grunewald [20] (there, $F \times_\phi F$ corresponds to our $A \leq B \simeq F_2 \times F_2 \leq Aut(F_3)$, and H corresponds to Borisov's group with two generators and unsolvable word problem). Alternatively, A is not finitely presented because it is not the direct product of finite index subgroups of $F_2 = \{{}_w\theta_1 \mid w \in \langle a, b \rangle\}$ and $F_2 = \{{}_1\theta_w \mid w \in \langle a, b \rangle\}$ (see Short's description of finitely presented subgroups of $F_2 \times F_2$ in [38]). \square

Question 7. Are there more sources of orbit undecidability other than exploiting the unsolvability of membership problem for certain subgroups?

Commentary. In order to find new sources, one needs to relate orbit decidability with some other algorithmic problem, for which there are known unsolvable examples. \square

Question 8. Is it true that every finitely generated subgroup of $GL_3(\mathbb{Z})$ is orbit decidable ? Or conversely, is it true that there exists a \mathbb{Z}^3 -by-free group with unsolvable conjugacy problem ? In close relation with this, is $CIP(GL_3(\mathbb{Z}))$ solvable ?

Commentary. Propositions 6.13 and 7.5, and Corollaries 6.3 and 7.7 show that the cases of dimension 2 and dimension bigger than or equal to 4 behave oppositely with respect to these three questions (answers being *yes*, *no*, *yes*, and *no*, *yes*, *no*, respectively). For the case of dimension 3, we point out that $GL_3(\mathbb{Z})$ is not virtually free, so the argument given in Proposition 6.13 does not work in this case. But, on the other hand, $F_2 \times F_2$ does not embed in $GL_3(\mathbb{Z})$ either (in fact, only very simple groups G satisfy $G \times G \leq GL_3(\mathbb{Z})$), so the argument in Proposition 7.5 does not work in dimension 3 either (unless one can find other pairs of subgroups $A \leq B \leq GL_3(\mathbb{Z})$ with unsolvable $MP(A, B)$). In the free context, the situation is easier, with the difference in behavior happening between rank two and rank three (see Propositions 6.21 and 7.2).

Also, it is interesting to remark that this situation is very similar (and maybe related) to the coherence of linear groups: it is known that $GL_2(\mathbb{Z})$ is coherent, because it is virtually free, and that $GL_n(\mathbb{Z})$ is not coherent for $n \geq 4$, precisely because it contains $F_2 \times F_2$ (see for example [20] and [38]). The question is still open in dimension 3, where none of the previous arguments work. \square

Question 9. Is it true that

- (a) for every two finitely generated groups A and B (except for $A = 1$ and $|B| < \infty$), there exists a finitely presented group G such that $Aut(G)$ simultaneously contains an orbit decidable subgroup isomorphic to A , and an orbit undecidable subgroup isomorphic to B ?
- (b) for every finitely generated group A , there exists a finitely presented group G such that $Aut(G)$ contains an orbit decidable subgroup isomorphic to A ?
- (c) for every finitely generated infinite group B , there exists a finitely presented group G such that $CP(G)$ is solvable, and $Aut(G)$ contains an orbit undecidable subgroup isomorphic to B ?

Commentary. Question (b) asks for a positive decisional condition, and question (c) for the corresponding negative one, while question (a) asks whether they are compatible within the same group G . Formally, (b) and (c) are partial cases of (a) (note that $CP(G)$ is equivalent to $OD(\{Id\})$).

However, an easy construction using the direct product shows that affirmative answers for (b) and (c) would imply an affirmative answer for (a) too.

Before, note that $A = 1$ and $|B| < \infty$ is the only situation where the copy of A will necessarily be a finite index subgroup of the copy of B and so, Proposition 6.4 would then say that solvability for $\text{OD}(A)$ implies solvability for $\text{OD}(B)$. In all other cases, even if B has a finite index subgroup isomorphic to A , it is conceivable that $\text{Aut}(G)$ could contain copies of A and B apart enough to each other to fulfill the requirements of question (a).

Now, let G_1 and G_2 be two groups, let $G = G_1 \times G_2$ be its direct product, and understand any subgroup of $\text{Aut}(G_i)$ as a subgroup of $\text{Aut}(G)$ acting trivially on the other coordinate. It is easy to see that if the orbit decidability for $A \leq \text{Aut}(G_1)$, and the conjugacy problem for G_2 are solvable, then the orbit decidability for $A \leq \text{Aut}(G)$ is also solvable. And similarly, if $B \leq \text{Aut}(G_2)$ is orbit undecidable then so is $B \leq \text{Aut}(G)$. Hence, answering question (a) in the affirmative reduces to answer in the affirmative questions (b) and (c).

If A is finitely presented, has trivial center, and $\text{CP}(A)$ is solvable, then we can take $G = A$, and the copy of A in $\text{Aut}(G)$ given by conjugations is clearly orbit decidable. This answers (b) in the affirmative in this very particular case.

Finally, let B be a finitely generated, recursively presented group. By Higman's embedding theorem (see [25]), B embeds in a finitely presented group $B' \neq 1$, which then embeds in $G_1 = B' * \mathbb{Z}$. Since G_1 has trivial center, $\text{Aut}(G_1)$ contains a copy of B given by inner automorphisms, say $B_1 \leq \text{Aut}(G_1)$. Take now another finitely presented group G_2 with unsolvable conjugacy problem, and consider their free product, $G = G_1 * G_2$. Extend the morphisms in B_1 to automorphisms of G acting trivially on G_2 ; this way, we obtain $B_2 \leq \text{Aut}(G)$, again isomorphic to B . Now, two given elements $u, v \in G_2$ lie in the same $(B_2 \cdot \text{Inn}(G))$ -orbit if and only if they are conjugate to each other in G_2 ; thus, solvability of $\text{OD}(B_2)$ would imply solvability of $\text{CP}(G_2)$. Hence, $B \simeq B_2 \leq \text{Aut}(G)$ is orbit undecidable. But, unfortunately, this does not solve question (c) because, by construction, $\text{CP}(G)$ is unsolvable, like $\text{CP}(G_2)$.

Additionally, note that the recursive presentability for B in the previous paragraph, is an extra condition also satisfied in the main source of orbit undecidability presented above. Namely, all orbit undecidable subgroups coming from Theorem 7.4 are of Mikhailova's type and so recursively presented (since they have solvable word problem). At the time of writing we are not aware of any construction producing orbit undecidable subgroups which are not recursively presented. \square

ACKNOWLEDGMENTS

We thank S. Hermiller, I. Kapovich and G. Levitt for interesting comments on the subject. The first named author is partially supported by the INTAS grant N 03-51-3663 and by the grant Complex integration projects of SB RAS N 1.9. The second and third named authors gratefully acknowledge partial support from the MEC (Spain) and the EFRD (EC) through projects BFM2003-06613 and MTM2006-13544. The third one also thanks the Department of Mathematics of the University of Nebraska-Lincoln for its hospitality during the second semester of the course 2003-2004, while a preliminary version of this research was started. The three of them thanks the Centre de Recerca Matemàtica (Barcelona, Catalonia) for its hospitality during several periods of time while this research was conducted and written.

REFERENCES

- [1] M. Bestvina, M. Feighn, M. Handel, *Solvable subgroups of $Out(F_n)$ are virtually abelian*, Geom. Dedicata, **104** (2004), 71–96.
- [2] M. Bestvina, M. Feighn, M. Handel, *The Tits alternative for $Out(F_n)$ (II), A Kolchin type theorem*, Ann. of Math. (2) **161**(1) (2005), 1–59.
- [3] O. Bogopolski, *Classification of automorphisms of the free group of rank 2 by ranks of fixed-point subgroups*, J. Group Theory **3** (2000), 339–351.
- [4] O. Bogopolski, *On the conjugacy problem for automorphisms of free groups*, Algebra i Logika **28**(1) (1989), 18–28. Translation in Algebra and Logic **28**(1) (1989), 10–17.
- [5] O. Bogopolski, V. Gerasimov, *Finite subgroups of hyperbolic groups*, Algebra i Logika **34**(6) (1995), 619–622. Translation in Algebra and Logic **34**(6) (1995), 343–345.
- [6] O. Bogopolski, A. Martino, O. Maslakova, E. Ventura, *Free-by-cyclic groups have solvable conjugacy problem*, Bulletin of the London Mathematical Society **38**(5) (2006), 787–794.
- [7] W.W. Boone, *The word problem*, Annals of mathematics **70**(2) (1959), 207–265.
- [8] V.V. Borisov, *Simple examples of groups with unsolvable word problem*, Mat. Zametki **6** (1969), 521–532. English translation in Math. Notes **6** (1969), 768–775.
- [9] M.R. Bridson, A. Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Mathematischen Wissenschaften **319**, Springer-Verlag (1999).
- [10] P. Brinkmann, *Detecting automorphic orbits in free groups*, preprint. Available at http://math.sci.cny.cuny.edu/people?name=Peter_Brinkmann.
- [11] D.J. Collins, *Recursively enumerable degrees and the conjugacy problem*, Acta Math. **122** (1969) 115–160.
- [12] D.J. Collins, C.F. Miller III, *The conjugacy problem and subgroups of finite index*, Proc. London Math. Soc. **34**(3) (1977), 535–556.
- [13] T. Delzant, *Sous-groupes distingués et quotients des groupes hyperboliques*, Duke Math. J. **83**(3) (1996), 661–682.
- [14] W. Dicks, M. Dunwoody, *Groups acting on graphs*, Cambridge Studies in Advanced Mathematics **17**, Cambridge University Press (1989).

- [15] W. Dicks, E. Formanek, *Algebraic mapping-class groups of orientable surfaces with boundaries*, Progress in Mathematics (Birkhäuser) **248** (2005), 57–116.
- [16] D.B.A. Epstein, D. Holt, *Computation in word-hyperbolic groups*, Internat. J. Algebra Comput. **11** (2001), 467–487.
- [17] A.L. Fel’shtyn, *The Reidemeister number of any automorphism of a Gromov hyperbolic group is infinite*, Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) **279** (2001), Geom. i Topol. 6, 229–240, 250. Translation in J. Math. Sci. **119**(1) (2004), 117–123.
- [18] A. Fel’shtyn, E. Troitsky, *Twisted conjugacy separable groups*, preprint available at <http://arxiv.org/abs/math/0606764>.
- [19] A.V. Gorjaga, A.S. Kirkinskii, *The decidability of the conjugacy problem cannot be transferred to finite extensions of groups*, Algebra i Logika **14** (1975), 393–406.
- [20] F.J. Grunewald, *On some groups which cannot be finitely presented*, J. London Math. Soc. **17**(2) (1978), 427–436.
- [21] G. Higman, B.H. Neumann, H. Neumann, *Embedding theorem for groups*, J. Lond. Math. Soc. **24** (1950), 247–254.
- [22] D.L. Johnson, *Presentations of groups*, London Math. Soc. Student texts **15**, Cambridge University Press, 1990.
- [23] *The Kourovka notebook: Unsolved problems in group theory*, sixteenth edition. Edited by V. D. Mazurov and E.I. Khukhro, Russian Academy of Sciences, Siberian Division, Institute of Mathematics, Novosibirsk, (2006).
- [24] D.M. Larue, *Left-distributive and left-distributive idempotent algebras*, PhD thesis (University of Colorado, Boulder) (1994), 138 pages; available at http://www.mines.edu/fs_home/dlarue/papers/dml.pdf.
- [25] R. Lyndon and P. Schupp, *Combinatorial group theory*, Springer-Verlag, (1977).
- [26] M. Lustig, *Structure and conjugacy for automorphisms of free groups*, preprint, 1999 (available at <http://junon.u-3mrs.fr/lustig/>).
- [27] W. Magnus, A. Karrass, D. Solitar, *Combinatorial Group Theory*, Interscience Publishers, New York, (1966).
- [28] A.I. Mal’cev, *On certain classes of infinite solvable groups*, Mat. Sbornik **28** (1951), 567–588. Translation in Amer. Math. Soc. Transl. **2**(2) (1956), 1–21.
- [29] K. A. Mihailova, *The occurrence problem for direct products of groups*, Dokl. Acad. Nauk SSSR **119** (1958), 1103–1105.
- [30] C.F. Miller III, *On group-theoretic decision problems and their classification*, Annals of Math. Studies **68**, (1971).
- [31] P.S. Novikov *On the algorithmic unsolvability of the word problem in group theory*, Trudy Mat. Inst. Steklov. **44** (1955), 143 pages. Translation in Amer. Math. Soc. Transl. **9**(2) (1958), 1–122.
- [32] P. Papasoglu, *An algorithm detecting hyperbolicity*, Geometric and computational perspectives on infinite groups DIMACS Ser. Discrete Math. Theoret. Comput. Sci., 25, Amer. Math. Soc. (1996), 193–200.
- [33] J.P. Preaux, *Conjugacy problem in groups of oriented geometrizable 3-manifolds.*, Topology **45**(1) (2006), 171–208.
- [34] J.P. Preaux, *Conjugacy problem in groups of non-oriented geometrizable 3-manifolds*, preprint (available at <http://arxiv.org/pdf/math.GR/0512484.pdf>).
- [35] K. Reidemeister, *Automorphismen von homotopiekettenringen*, Math. Ann. **112** (1936), 586–593.
- [36] V.N. Remeslennikov, *Conjugacy in polycyclic groups*, Algebra i Logika **8** (1969), 712–725. Translation in Algebra and Logic **8** (1969), 404–411.

- [37] D. Segal, *Polycyclic groups*, Cambridge Tracts in Mathematics **82**, Cambridge University Press, (1983).
- [38] H. Short, *Finitely presented subgroups of a product of two free groups*, Quart. J. of Math. **52** (2001), 127–131.
- [39] J.R. Stallings, *Topology of finite graphs*, Invent. Math. **71** (1983), 551-565.
- [40] J. Tits, *Free subgroups in linear groups*, J. Algebra **20** (1972), 250–270.
- [41] J.H.C. Whitehead, *On equivalent sets of elements in a free group*, Ann. of Math. **37** (1936), 782-800.

INSTITUTE OF MATHEMATICS OF SBIRAS, NOVOSIBIRSK, RUSSIA AND UNIVERSITÄT DORTMUND, FAKULTÄT FÜR MATHEMATIK, LEHRSTUHL VI (ALGEBRA), VOGELPOTHSWEG 87 D-44221 DORTMUND, GERMANY.

E-mail address: `groups@math.nsc.ru`

DEPARTAMENT DE MATEMÀTICA APLICADA IV, UNIVERSITAT POLITÈCNICA DE CATALUNYA, BARCELONA, SPAIN.

E-mail address: `Armando.Martino@upc.edu`

DEPARTAMENT DE MATEMÀTICA APLICADA III, UNIVERSITAT POLITÈCNICA DE CATALUNYA, AND CENTRE DE RECERCA MATEMÀTICA, BARCELONA, SPAIN.

E-mail address: `enric.ventura@upc.edu`