

ELEMENTS OF ALGEBRAIC GEOMETRY AND THE POSITIVE THEORY OF PARTIALLY COMMUTATIVE GROUPS

MONTSERRAT CASALS-RUIZ AND ILYA V. KAZACHKOV

ABSTRACT. The first main result of the paper is a criterion for a partially commutative group \mathbb{G} to be a domain. It allows us to reduce the study of algebraic sets over \mathbb{G} to the study of irreducible algebraic sets, and reduce the elementary theory of \mathbb{G} (of a coordinate group over \mathbb{G}) to the elementary theories of the direct factors of \mathbb{G} (to the elementary theory of coordinate groups of irreducible algebraic sets).

Then we establish normal forms for quantifier-free formulas over a non-abelian directly indecomposable partially commutative group \mathbb{H} . Analogously to the case of free groups, we introduce the notion of a generalised equation and prove that the positive theory of \mathbb{H} has quantifier elimination and that arbitrary first-order formulas lift from \mathbb{H} to $\mathbb{H} * F$, where F is a free group of finite rank. As a consequence, the positive theory of an arbitrary partially commutative group is decidable.

1. INTRODUCTION

This paper can be considered as a part of a project the aim of which is to construct algebraic (diophantine) geometry over partially commutative groups, and, more generally, to study the elementary theory of partially commutative groups.

Classical algebraic geometry is concerned with the study of the geometry of sets of solutions of systems of equations, i.e. the geometry of algebraic sets. Taking the collection of all algebraic sets as a pre-base of closed sets one gets a topology, known as the Zariski topology. In the Zariski topology, every closed set is a union (maybe infinite) of algebraic sets. In the case that the ring of coefficients is Noetherian, the Zariski topology is Noetherian, every closed set Y is a finite union of algebraic sets $Y = Y_1 \cup \dots \cup Y_k$. In the

1991 *Mathematics Subject Classification.* 20F10, 03C10, 20F06.

The first author is supported by Programa de Formación de Investigadores del Departamento de Educación, Universidades e Investigación del Gobierno Vasco.

The second author was supported by Richard. H. Tomlinson Fellowship.

case that $Y_i \not\subseteq Y_j$, $i \neq j$, and Y_i can not be non-trivially presented as a union of algebraic sets, this decomposition is unique and the sets Y_1, \dots, Y_k are referred to as the irreducible components of Y . In general, however, a finite union of algebraic sets is not necessarily again an algebraic set. In classical algebraic geometry, it suffices to require that the ring of coefficients be a domain. Under these assumptions there exists a one-to-one correspondence between algebraic sets and closed sets. Thus, the study of algebraic sets reduces completely to the study of irreducible algebraic sets.

In [1] G. Baumslag, A. Miasnikov and V. Remeslennikov lay down the foundations of algebraic geometry over groups and introduce group-theoretic counterparts of basic notions from algebraic geometry over fields. The counterpart to the notion of a Noetherian ring is the notion of an equationally Noetherian group: a group G is called *equationally Noetherian* if every system $S(X) = 1$ with coefficients from G is equivalent to a finite subsystem $S_0 = 1$, where $S_0 \subset S$, i.e. the algebraic set defined by S coincides with the one defined by S_0 . The notion of a domain carries over from rings to groups as follows: a group G is called a *domain* if for any $x, y \neq 1$ there exists $g \in G$ such that $[x, y^g] \neq 1$.

The notions of equationally Noetherian group and domain, play an analogous role (to their ring-theoretic counterparts) in algebraic geometry over groups (see [1]):

- a group G is equationally Noetherian if and only if the Zariski topology is Noetherian, in particular every closed set is a finite union of algebraic sets;
- if a group G is a domain, then the collection of all algebraic sets is a base for the Zariski topology.

Our main interest in this paper is algebraic geometry over (free) partially commutative groups. Partially commutative groups are widely studied in different branches of mathematics and computer science, which explains the variety of names they were given: *graph groups*, *right-angled Artin groups*, *semifree groups*, etc. Without trying to give an account of the literature and results in the field we refer the reader to a recent survey [4] and the introduction and references in [14].

Partially commutative groups are linear, see [19], hence, equationally Noetherian, see [1]. In [1] the authors give several sufficient conditions for a group to be a domain. In particular, any CSA group is a domain and various group-theoretic constructions preserve the property of being a domain. However, none of the criteria obtained in [1] apply to the case of partially commutative groups. The major obstacle here is that a partially commutative group may contain a direct product of two free groups.

In Section 4.2 we give a criterion for a partially commutative group to be a domain:

Theorem 4.16. *Let \mathbb{G} be a partially commutative group. Then \mathbb{G} is a domain if and only if \mathbb{G} is non-abelian and directly indecomposable.*

Note that even if a partially commutative group is directly indecomposable, it still may contain a direct product of free groups.

The proof of this theorem is given in Section 4. It makes use of the technique of van Kampen diagrams over partially commutative groups, which we present in Section 3 and the description of centralisers in partially commutative groups (see Theorem 2.3).

The remaining part of the paper has a model-theoretic flavor. In Section 5, using results from [18], we prove that the elementary theory of \mathbb{G} (of a coordinate group over \mathbb{G}) reduces to the elementary theories of the direct factors of \mathbb{G} (to the elementary theory of coordinate groups of irreducible algebraic sets):

Corollary 5.2. *Let \mathbb{G} be a non-abelian directly indecomposable partially commutative group.*

- (i) *If $Y = Y_1 \cup \dots \cup Y_k$ is an algebraic set over \mathbb{G} , where Y_1, \dots, Y_k are the irreducible components of Y , then the elementary theory of the coordinate group $\Gamma(Y)$ of Y is decidable if and only if the elementary theory of $\Gamma(Y_i)$ is decidable for all $i = 1, \dots, k$.*
- (ii) *If $Y = Y_1 \cup \dots \cup Y_k$ and $Z = Z_1 \cup \dots \cup Z_l$ are two irreducible algebraic sets, where Y_1, \dots, Y_k and Z_1, \dots, Z_l are the irreducible components of Y and Z , respectively, then $\Gamma(Y)$ is elementary equivalent to $\Gamma(Z)$ if and only if $k = l$ and, after a certain re-enumeration, $\Gamma(Y_i)$ is elementary equivalent to $\Gamma(Z_i)$ for all $i = 1, \dots, k$.*

It is known that coordinate groups of algebraic sets over \mathbb{G} are separated by \mathbb{G} (are residually \mathbb{G}), see [25]. If a coordinate group Γ is a coordinate group of an irreducible set, then Γ is discriminated by \mathbb{G} (is fully residually \mathbb{G}), or equivalently, is universally equivalent to \mathbb{G} . Hence, the class of coordinate groups of irreducible algebraic sets is much narrower and admits a convenient logical description.

In his seminal work [21], Makanin introduced the notion of a generalised equation. In [22] this notion is used in order to show that the existential theory (the compatibility problem) of free groups and monoids is decidable. Since then this result has been generalised in various ways. In [28] Schulz generalised Makanin's result to the case of systems of equations over a free monoid with regular constraints, and in [6] Diekert, Gutierrez and Hagenah showed the decidability of the compatibility problem for systems of equations over a free group with rational constraints. Using the latter result,

Diekert and Lohrey show in [7] that the existential theory of a certain class of graph products of groups is decidable. Furthermore in [8], the authors show the decidability of the existential theory for an even wider class of groups. A common feature of the results mentioned above is that they reduce the problem to the one for free groups with rational constraints.

One of the main applications of the decidability of the compatibility problem for free groups is the decidability of the positive theory of the respective group. In the case of free groups this is a very well known result. In his paper [24], Merzlyakov performs quantifier elimination for positive formulas over free groups by describing the Skolem functions. Then using the result of Makanin, [22], one gets the decidability of the positive theory.

The aim of Sections 6 and 7 is to carry over the approach of Merzlyakov and Makanin to the case of partially commutative groups.

In Section 6, we prove that any positive quantifier-free formula over a non-abelian directly indecomposable partially commutative groups is equivalent to a single equation. In order to do so we prove that

- (i) for any finite system of equations $S_1(X) = 1, \dots, S_k(X) = 1$ one can effectively find a single equation $S(X) = 1$ such that the algebraic set defined by the equations S_1, \dots, S_k and by S coincide for any non-abelian directly indecomposable partially commutative group \mathbb{G} ,
- (ii) for any finite set of equations $S_1(X) = 1, \dots, S_k(X) = 1$ one can effectively find a single equation $S(X) = 1$ such that the union of algebraic sets defined by the equations S_1, \dots, S_k coincides with the algebraic set defined by S for any non-abelian directly indecomposable partially commutative group \mathbb{G} .

In the case of free groups, the first result is due to Malcev, see [23], and in [22] Makanin attributes the second result to Gurevich. These results hold in fact in a much more general setting (for groups that satisfy certain first-order formulas), for example in [17] it is proven that this is the case for torsion-free, non-abelian, CSA groups that satisfy the Vaught's conjecture, in particular, for all non-abelian fully residually free groups and torsion-free hyperbolic groups. Note, that a non-abelian directly indecomposable partially commutative group is almost never a CSA group. We generalise the results of Malcev and Gurevich to the case of partially commutative groups. The exposition in this section as well as in Section 7 is based on [17]. As an immediate consequence of these results we get a normal form for first order formulas over partially commutative groups (in fact, over a much wider class of groups).

In Section 7, we use the normal form for Van Kampen diagrams obtained in Lemma 3.2 to describe the finite number of all possible cancellation schemes for a given equation. This allows us to introduce the notion of a generalised equation for partially commutative groups. Then we introduce an analogue of the, so called, Merzlyakov words and perform quantifier elimination for positive formulas over non-abelian directly indecomposable partially commutative groups.

Theorem 7.8. *If*

$$\mathbb{G} \models \forall x_1 \exists y_1 \dots \forall x_k \exists y_k (S(X, Y, A) = 1),$$

then there exist words (with constants from \mathbb{G}) $q_1(x_1), \dots, q_k(x_1, \dots, x_k) \in \mathbb{G}[X]$, such that

$$\mathbb{G}[X] \models S(x_1, q_1(x_1), \dots, x_k, q_k(x_1, \dots, x_k), A) = 1,$$

i.e. the equation

$$S(x_1, y_1, \dots, x_k, y_k, A) = 1$$

(in variables Y) has a solution in the group $\mathbb{G}[X]$.

Our approach, therefore, is a natural analog of the classical approach of Merzlyakov and Makanin to the positive theory of free groups and avoids the technically involved language of constraints.

In particular, quantifier elimination gives a reduction of the decidability of the positive theory of non-abelian directly indecomposable partially commutative groups to the decidability of the compatibility problem of an equation, which is known to be decidable, see [9].

Finally, in order to prove that the positive theory of any partially commutative group is decidable, we need to study the positive theory of the direct product of groups. In folklore, it is known that if $G = H_1 \times \dots \times H_k$, then the positive theory of G is decidable if the positive theories of H_1, \dots, H_k are decidable. However, we were unable to find a reference till (when this paper was already written) M. Lohrey pointed out that in [7], the authors give a proof of this result. We present another proof of this fact in the Appendix. The proof is purely model-theoretic and makes use of the ideas of the proof of Theorem 8.13 which is due to Feferman and Vaught, see [16].

2. PRELIMINARIES

2.1. Partially commutative groups. We begin with the basic notions of the theory of free partially commutative groups. Recall that a (free) *partially commutative* group is defined as follows. Let Γ be a finite, undirected, simple

graph. Let $A = V(\Gamma) = \{a_1, \dots, a_n\}$ be the set of vertices of Γ and let $F(A)$ be the free group on A . Let

$$R = \{[a_i, a_j] \in F(A) \mid a_i, a_j \in A \text{ and there is an edge of } \Gamma \text{ joining } a_i \text{ to } a_j\}.$$

The partially commutative group corresponding to the (commutation) graph Γ is the group $\mathbb{G}(\Gamma)$ with presentation $\langle A \mid R \rangle$. This means that the only relations imposed on the generators are commutation of some of the generators. When the underlying graph is clear from the context we write simply \mathbb{G} .

From now on $A = \{a_1, \dots, a_r\}$ always stands for a finite alphabet, its elements being called *letters*. We reserve the term *occurrence* to denote an occurrence of a letter in a word. In a more formal way, an occurrence is a pair (letter, its placeholder in the word).

For a given word w denote $\alpha(w)$ the set of letters occurring in w . For a word $w \in \mathbb{G}$, we denote by \bar{w} a geodesic of w . For a word $w \in \mathbb{G}$ define $\mathbb{A}(w)$ to be the subgroup of \mathbb{G} generated by all letters that do not occur in \bar{w} and commute with w . The subgroup $\mathbb{A}(w)$ is well-defined (independent of the choice of a geodesic \bar{w}), see [15]. An element $w \in \mathbb{G}$ is called *cyclically reduced* if the length of \bar{w}^2 is twice the length of \bar{w} .

For a partially commutative group \mathbb{G} consider its non-commutation graph Δ . The vertex set V of Δ is a set of generators A of \mathbb{G} . There is an edge connecting a_i and a_j if and only if $[a_i, a_j] \neq 1$. The graph Δ is a union of its connected components I_1, \dots, I_k . Then

$$(1) \quad \mathbb{G} = \mathbb{G}(I_1) \times \dots \times \mathbb{G}(I_k).$$

Consider $w \in \mathbb{G}$ and the set $\alpha(w)$. For this set, just as above, consider the graph $\Delta(\alpha(w))$ (it is a full subgraph of Δ). This graph can be either connected or not. If it is connected we will call w a *block*. If $\Delta(\alpha(w))$ is not connected, then we can split w into the product of commuting words

$$(2) \quad w = w_{j_1} \cdot w_{j_2} \cdots w_{j_t}; \quad j_1, \dots, j_t \in J,$$

where $|J|$ is the number of connected components of $\Delta(\alpha(w))$ and the word w_{j_i} involves letters from the j_i -th connected component. Clearly, the words $\{w_{j_1}, \dots, w_{j_t}\}$ pairwise commute. Each word w_{j_i} , $i \in 1, \dots, t$ is a block and so we refer to presentation (2) as the block decomposition of w .

An element $w \in \mathbb{G}$ is called a least root (or simply, root) of $v \in \mathbb{G}$ if there exists an integer $0 \neq m \in \mathbb{Z}$ such that $v = w^m$ and there does not exist $w' \in \mathbb{G}$ and $0 \neq m' \in \mathbb{Z}$ such that $w = w'^{m'}$. In this case we write $w = \sqrt{v}$. By [11], partially commutative groups have least roots, that is the root element of v is defined uniquely.

The following proposition reduces the conjugacy problem for arbitrary elements of a partially commutative group to the one for block elements.

Proposition 2.1 (Proposition 5.7 of [15]). *Let $w = w_1 \cdot w_2 \cdots w_t$ and $v = v_1 \cdot v_2 \cdots v_s$ be cyclically reduced elements decomposed into the product of blocks. Then v and w are conjugate if and only if $s = t$ and, after some certain index re-enumeration, w_i is conjugate to v_i , $i = 1, \dots, t$.*

Corollary 2.2. *Let $w = w_1^{r_1} \cdot w_2^{r_2} \cdots w_t^{r_t}$ and $v = v_1^{l_1} \cdot v_2^{l_2} \cdots v_s^{l_s}$ be cyclically reduced elements decomposed into the product of blocks, where w_i and v_j are root elements, $l_i, r_j \in \mathbb{Z}$, $i = 1, \dots, t$, $j = 1, \dots, s$. Then w and v are conjugate if and only if $s = t$ and, after some certain index re-enumeration, $r_i = l_i$ and w_i is conjugate to v_i , $i = 1, \dots, t$.*

The next result describes centralisers of elements in partially commutative groups. As the definition of “being a domain” relies on the structure of centralisers, we shall make substantial use of the following theorem.

Theorem 2.3 (Centraliser Theorem, Theorem 3.10, [11]). *Let $w \in \mathbb{G}$ be a cyclically reduced word, $w = v_1 \dots v_k$ be its block decomposition. Then, the centraliser of w is the following subgroup of \mathbb{G} :*

$$C(w) = \langle \sqrt{v_1} \rangle \times \cdots \times \langle \sqrt{v_k} \rangle \times \mathbb{A}(w).$$

Corollary 2.4. *For any $w \in \mathbb{G}$ the centraliser $C(w)$ of w is an isolated subgroup of G , i.e. $C(w) = C(\sqrt{w})$.*

2.2. Algebraic Geometry over Groups. In this section we recall basic notions of algebraic geometry over groups, see [1] for details.

Let G be a group generated by a finite set A , $F(X)$ be a free group with basis $X = \{x_1, x_2, \dots, x_n\}$, $G[X] = G * F(X)$ be the free product of G and $F(X)$. A subset $S \subset G[X]$ is called a *system of equations* over G . As an element of the free product, the left side of every equation in $S = 1$ can be written as a product of some elements from $X \cup X^{-1}$ (which are called *variables*) and some elements from $A \subset G$ (*constants*).

A *solution* of the system $S(X) = 1$ over a group G is a tuple of elements $g_1, \dots, g_n \in G$ such that every equation from S vanishes at (g_1, \dots, g_n) , i.e. $S(g_1, \dots, g_n) = 1$ in G . Equivalently, a solution of the system $S = 1$ over G is a G -homomorphism $\phi : G[X] \rightarrow G$ such that $S \subseteq \ker(\phi)$. Denote by $\text{ncl}(S)$ the normal closure of S in $G[X]$, and by G_S the quotient group $G[X]_{/\text{ncl}(S)}$. Then every solution of $S(X) = 1$ in G gives rise to a G -homomorphism $G_S \rightarrow G$, and vice versa. By $V_G(S)$ we denote the set of all solutions in G of the system $S = 1$ and call it the *algebraic set defined by S* .

Normal subgroup of $G[X]$ of the form

$$R(S) = \{T(X) \in G[X] \mid \forall A \in G^n (S(A) = 1 \rightarrow T(A) = 1)\}$$

is called the *radical of the system* S . Note that $S \subseteq R(S)$. There exists a one-to-one correspondence between algebraic sets $V_G(S)$ of systems of equations in $G[X]$ and radical subgroups.

The quotient group

$$G_{R(S)} = G[X]/R(S)$$

is called the *coordinate group* of the algebraic set $V_G(S)$, and every solution of $S(X) = 1$ in G is a G -homomorphism $G_{R(S)} \rightarrow G$.

A G -group H is called *G -equationally Noetherian* if every system $S(X) = 1$ with coefficients from G is equivalent over G to a finite subsystem $S_0 = 1$, where $S_0 \subset S$, i.e. the systems S and S_0 define the same algebraic set. If G is G -equationally Noetherian, then we say that G is equationally Noetherian. If a G -group H is equationally Noetherian every algebraic set V in G^n is a finite union of *irreducible components* of V .

Let H and K be G -groups. We say that a family of G -homomorphisms $\mathcal{F} \subset \text{Hom}_G(H, K)$ *G -separates* (*G -discriminates*) H into K if for every non-trivial element $h \in H$ (every finite set of non-trivial elements $H_0 \subset H$) there exists $\phi \in \mathcal{F}$ such that $h^\phi \neq 1$ ($h^\phi \neq 1$ for every $h \in H_0$). In this case we say that H is *G -separated* (*G -discriminated*) by K . In the case that $G = 1$, we simply say that H is separated (discriminated) by K .

A G -group H is called a *G -domain* if for any $x, y \neq 1$ there exists $g \in G$ such that $[x, y^g] \neq 1$. In the case that G is G -domain, we say that G is a domain.

3. VAN KAMPEN DIAGRAMS

In this section we present some preliminary results on Van Kampen diagrams. We refer the reader to [3] and [27] for a more detailed account on van Kampen diagrams. Our aim here is to review some basic notions and techniques and apply them to the particular case of partially commutative groups.

3.1. Van Kampen Diagrams in Partially Commutative Groups. By van Kampen's Lemma (see [3]) the word w represents the trivial element in a fixed group G given by the presentation $\langle A \mid R_A \rangle$ if and only if there exists a finite connected, oriented, based, labeled, planar graph \mathcal{D} where each oriented edge is labeled by a letter in $A^{\pm 1}$, each bounded region (cell) of $\mathbb{R}^2 \setminus \mathcal{D}$ is labeled by a word in R_A (up to shifting cyclically or taking inverses) and w can be read on the boundary of the unbounded region of $\mathbb{R}^2 \setminus \mathcal{D}$ from the base vertex. Then we say that \mathcal{D} is a *van Kampen diagram* for the boundary word w over the presentation $\langle A \mid R_A \rangle$. If $w = uv^{-1} =_{\mathbb{G}} 1$ we say that \mathcal{D} is a *van Kampen diagram* realising the equality $u = v$. In

the event that a van Kampen diagram \mathcal{D} realises the equality $w = \bar{w}$ we say that \mathcal{D} is a *geodesic van Kampen diagram* for w .

Any van Kampen diagram can also be viewed as a 2-complex, with a 2-cell attached for each bounded region (see Figure 1).

We shall further restrict our considerations to the case when G is a partially commutative group.

Following monograph [27], if we complete the set of defining relations adding the trivial relations $1 \cdot a = a \cdot 1$ for all $a \in A$, then every van Kampen diagram can be transformed so that its boundary is a simple curve. In other words, as a 2-complex the van Kampen diagram is homeomorphic to a disc tiled by cells which are also homeomorphic to a disc (see Figure 1). We further assume that all van Kampen diagrams are of this form.




FIGURE 1. van Kampen diagram and non-singular van Kampen diagram for $w = caba^{-1}b^{-1}c^{-1}$ over $\langle a, b, c | [a, b] = 1 \rangle$.

Let \mathcal{D} be a van Kampen diagram for the boundary word w . Given an occurrence a in w , there is a cell C in the 2-complex \mathcal{D} attached to a . Since every cell in a van Kampen diagram is either labelled by a relation of the form $a^{-1}b^{-1}ab$ or is a so-called 0-cell, i.e. a cell labelled by $1 \cdot a = a \cdot 1$, there is just one occurrence of a and one occurrence of a^{-1} on the boundary of C .

Since \mathcal{D} is homeomorphic to a disc, if the occurrence of a^{-1} on the boundary of C is not on the boundary of \mathcal{D} , there exists a unique cell $C' \neq C$ attached to this occurrence of a^{-1} in \mathcal{D} . Repeating this process, we obtain a unique *band* in \mathcal{D} .

Because of the structure of the cells and the fact that \mathcal{D} is homeomorphic to a disc, a band never self-intersects; indeed, since \mathcal{D} is homeomorphic to a disc, the only way a band can self-intersect is shown in Figure 2. But then, the cell corresponding to the self-intersection of the band is labelled by the word $aaa^{-1}a^{-1}$.




FIGURE 2. Bands do not self-intersect

Then, since the number of cells in \mathcal{D} is finite, in a finite number of steps the band will again meet the boundary in an occurrence of a^{-1} in w (see Figure 3).




FIGURE 3. Bands in a van Kampen diagram

We will use the notation L_a to indicate that a band begins (and thus ends) in an occurrence of a letter $a \in A^{\pm 1}$.

Remark 1.

- If two bands L_a and L_b cross then the intersection cell realises the equality $a^{-1}b^{-1}ab = 1$ and so $a \neq b$ and $[a, b] = 1$ (see Figure 3).
- Every band L_a gives a decomposition of w in the following form $w = w_1aw_2a^{-1}w_3$.

Lemma 3.1. *Let \mathbb{G} be a partially commutative group. A word w in \mathbb{G} is not geodesic if and only if w contains a subword aBa^{-1} such that $[a, \alpha(B)] = 1$, $a \in A^{\pm 1}$ if and only if there exists a geodesic van Kampen diagram for w that contains a band L_a with both ends in w .*

A proof of this lemma can be found in [29].

It is known (see [15]) that if a word w represents the trivial element in \mathbb{G} , it can be reduced to the empty word using commutation relations of letters and free cancellation. Moreover this reduction process of w to the empty word is independent of the order in which the letters are freely cancelled.

Lemma 3.1 reflects a consequence of a deeper fact: there exists a one-to-one correspondence between van Kampen diagrams for w and procedures of reductions of w to the empty word. Indeed, let \mathcal{D} be a van Kampen diagram for the boundary word w . Every band L_a gives a decomposition of the form $w = w_{1,a}aw_{2,a}a^{-1}w_{3,a}$. Let L_a be a band such that the length of $w_{2,a}$ is minimal. Hence, every band L_b with an end in an occurrence b in $w_{2,a}$ can not have the other end in an occurrence b^{-1} in $w_{2,a}$. Thus for every occurrence b in $w_{2,a}$ the band L_b crosses the band L_a and hence $[a, \alpha(w_{2,a})] = 1$. This implies that $w = w_{1,a}aw_{2,a}a^{-1}w_{3,a} = w_{1,a}w_{2,aa}a^{-1}w_{3,a}$ and thus there exists a process of reduction of w to the empty word in which the occurrence a is cancelled with the occurrence a^{-1} . Collapsing the band L_a in \mathcal{D} we get a van Kampen diagram \mathcal{D}' for the boundary word $w' = w_{1,a}w_{2,aa}w_{3,a}$, note that the number of cells in \mathcal{D}' is lower than the number of cells in \mathcal{D} . The statement follows by induction.

Conversely, if w represents the trivial element in \mathbb{G} , w can be written in the form $w = w_1aw_2a^{-1}w_3$ where $a \in A^{\pm 1}$ and $[a, \alpha(w_2)] = 1$. Construct a $|w|$ -polygon, designate a point, and orient and label its edges so that starting from the designated point and reading clockwise (or, counterclockwise) one reads w . To every edge labelled by an occurrence w_{2i} from w_2 we attach a cell labelled by $aw_{2i}a^{-1}w_{2i}^{-1}$. Identifying, as appropriate, the edges labelled by $a^{\pm 1}$ we get a band L_a with ends in a and a^{-1} , see Figure 4. We thereby get a $(|w| - 2)$ -polygon with the boundary word $w' = w_1w_2w_3$ and thus, by induction, the van Kampen diagram is constructed.

If either in a geodesic van Kampen diagram for w both ends of a band L_a lie in w or equivalently, if the occurrences a and a^{-1} freely cancel each other in a reduction process of the word w to the empty word, we say that




FIGURE 4. Constructing a van Kampen diagram by a process of reduction

the occurrence a cancels with a^{-1} . Otherwise, if one of the ends of the band L_a is in an occurrence of a in w and the other is in an occurrence of a^{-1} in \bar{w} , we say that a does not cancel.

3.2. Cancellation in a Product of Elements. We now consider in detail van Kampen diagrams corresponding to a product of k geodesic words $w_1 \cdots w_k = 1$.

By Lemma 3.1 for any van Kampen diagram \mathcal{D} of $w_1 \cdots w_k = 1$ every band with an end in w_i has its other end in w_j , $j \neq i$, $i, j = 1, \dots, k$.

Since every occurrence in w_1 cancels, there is a band with an end in a given occurrence a of w_1 and another end in w_i , $1 < i \leq k$. Then for any occurrence b in w_1 such that

- b is to the right of a , i.e. $w_1 = w'_1 a w''_1 b w'''_1$ and
- the band L_b with an end in the occurrence b has its other end in w_j , $j > i$,

the bands L_a and L_b cross and thus $[a, b] = 1$ in \mathbb{G} , see Figure 3.2. Therefore the word w_1 equals the following geodesic word $w_1 = w_1^k \cdots w_1^2$, where the band with an end in any occurrence of w_1^i has its other end in w_i .


A similar argument for w_l shows that w_l admits the following geodesic presentation (see Figure 3.2):

$$w_l = w_l^{l-1} \cdots w_l^1 w_l^k \cdots w_l^{l+1},$$

where the band with an end in any occurrence of w_l^i has its other end in w_i .

We summarise the above discussion in the following lemma

Lemma 3.2. *Let \mathbb{G} be a partially commutative group, let w_1, \dots, w_k be geodesic words in \mathbb{G} such that $w_1 \cdots w_k = 1$. Then there exist geodesic words*



w_i^j , $1 \leq i, j \leq k$ such that for any $1 \leq l \leq k$ there exists the following geodesic presentation for w_l :

$$w_l = w_l^{l-1} \cdots w_l^1 w_l^k \cdots w_l^{l+1},$$

where $w_l^i = w_i^{l-1}$.

Corollary 3.3. *Let \mathbb{G} be a partially commutative group, let w_1, \dots, w_k, v be geodesic words in \mathbb{G} such that $w_1 \cdots w_k = v$. Then there exists geodesic words v_m, w_i^j , $1 \leq i, j, m \leq k$ such that for any $1 \leq l \leq k$ there exists the following geodesic presentation for w_l :*

$$w_l = w_l^{l-1} \cdots w_l^1 v_l w_l^k \cdots w_l^{l+1},$$

where $w_l^i = w_i^{l-1}$ and $v_1 \cdots v_k = v$.

4. PARTIALLY COMMUTATIVE GROUPS AND DOMAINS

It is well-known that free groups are domains. The key point of the proof (which relies on the fact that free groups are CSA) is that for $a, x, y \in F$, $x \neq 1$:

$$\text{if } [x, y] = 1, [x, y^a] = 1, \text{ then } y \in C(a).$$

Therefore, to see that free groups are domains it suffices to apply the above argument for two elements a and b such that $C(a) \cap C(b) = 1$.

Although, directly indecomposable partially commutative groups are not CSA, using the description of centralisers, in Section 4.2 we prove that for $a, x, y \in \mathbb{G}$, such that $x \neq 1$ and $C(a)$ is cyclic:

$$\text{if } [x, y] = 1, [x, y^a] = 1, \text{ then either } y \in C(a) \text{ or } x \in \mathbb{A}(y^a).$$

The aim of Section 4.1 below is to find an element $A \in \mathbb{G}$ with cyclic centraliser for which $\mathbb{A}(y^A) = 1$. More precisely, we prove that for any $a \in \mathbb{G}$, such that $C(a)$ is cyclic, the element $A = a^{2 \cdot \text{cdim}(\mathbb{G}) + 2}$ possesses this property. Hence, for $a, x, y \in \mathbb{G}$, such that $x \neq 1$ and $C(a)$ is cyclic:

$$\text{if } [x, y] = 1, [x, y^A] = 1, \text{ then } y \in C(A).$$

4.1. Cancellation and Conjugation.

Definition 4.1. We treat the graph Δ as a metric space with the metric d being the path metric. Let y be a vertex of Δ , define $\text{adj}(y)$ to be $\{v \in \Delta \mid d(v, y) \leq 1\}$, i.e. the closed ball of radius 1 centered at y . For a subset $Y \subseteq A$, set $\text{adj}(Y) = \{v \in \Delta \mid d(v, y) \leq 1 \text{ for some } y \in Y\}$.

We set

$$\text{adj}^n(y) = \underbrace{\text{adj}(\text{adj}(\dots \text{adj}(y) \dots))}_{n \text{ times}},$$

thus $\text{adj}^n(y) = \{v \in \Delta \mid d(v, y) \leq n\}$ is the closed ball of radius n centered at y . Similarly $\text{adj}^n(Y)$, $Y \subseteq A$ is just an n -neighbourhood of Y in Δ , $\text{adj}^n(Y) = \{v \in \Delta \mid d(v, y) \leq n \text{ for some } y \in Y\}$.

Let Δ_1 be a subgraph of Δ . Then by $\text{adj}(Y)_{\Delta_1}$ we denote the following set $\text{adj}(Y)_{\Delta_1} = \text{adj}(Y) \cap \Delta_1$.

We shall further use the notion of centraliser dimension $\text{cdim}(G)$ of a group G (see Definition 4.2 below), an interested reader may consult [12, 13, 26] and references there for a detailed discussion of this notion.

Definition 4.2. If there exists an integer d such that the group G has a strictly descending chain of centralisers

$$C_0 > C_1 > \cdots > C_d$$

of length d and no centraliser chain of length greater than d then G is said to have *centraliser dimension* $\text{cdim}(G) = d$. If no such integer d exists we say that the centraliser dimension of G is infinite, $\text{cdim}(G) = \infty$.

All partially commutative groups have finite centraliser dimension, [13].

Lemma 4.3. *Let \mathbb{G} be a directly indecomposable partially commutative group, $y \in A$ then*

$$\text{adj}^{\text{cdim}(\mathbb{G})}(y) = A,$$

i.e. the diameter $\text{diam}(\Delta(\mathbb{G}))$ of $\Delta(\mathbb{G})$ is less or equals $\text{cdim}(\mathbb{G})$.

Proof. The group \mathbb{G} is directly indecomposable, hence the non-commutation graph $\Delta(\mathbb{G})$ is connected. Therefore for any pair of vertices $g, h \in V(\Delta)$ there exists a path p of minimal length connecting them. We claim that the length of p is less or equals $\text{cdim}(\mathbb{G})$.

Let $p = (g_0 = g, g_1, \dots, g_r = h)$. The path p gives rise to a strictly descending chain of centralisers of length r :

$$\mathbb{G} > C(g_0) > C(g_0, g_1) > \cdots > C(g_0, \dots, g_{r-2}) > C(g_0, \dots, g_{r-1}).$$

Indeed, to see that each of the inclusions above is strict we use the minimality of the path p . Suppose $C(g_0, \dots, g_{i-1}) = C(g_0, \dots, g_i)$ for some $1 \leq i \leq r-1$, then since $g_{i+1} \notin C(g_0, \dots, g_i)$ we also have $g_{i+1} \notin C(g_0, \dots, g_{i-1})$. So there exists $0 \leq j \leq i-1$, such that g_j does not commute with g_{i+1} , thus the distance between them is 1. Then $(g_0, g_1, \dots, g_j, g_{i+1}, \dots, g_r)$ is a shorter path from g to h , contradicting the minimality of the path p .

As the length r of any strictly descending chain of centralisers is bounded by $\text{cdim}(\mathbb{G})$, so is the distance between any two points in Δ , so $\text{adj}^{\text{cdim}(\mathbb{G})}(g) = A$. \square

Remark 2. Note that the equality $\text{diam}(\Delta(\mathbb{G})) = \text{cdim}(\mathbb{G})$ can be attained. Set \mathbb{G} to be, for example, the partially commutative group whose non-commutation graph is a path with an odd number of vertices.

Given two geodesic words $w, v \in \mathbb{G}$, if the product wv is again geodesic we write $w \circ v$. Let $g \in \mathbb{G}$ be a geodesic word. We refer to the decomposition $g = g_1 \circ g_2 \circ g_1^{-1}$, where g_2 is cyclically reduced as the *cyclic decomposition* of g .

Given a word g^n we write $g^{(i)}$ when we refer to the i -th factor g in the product $g^n = g \cdots g$. Similarly, given an occurrence a in g we write $a^{(i)}$, $1 \leq i \leq n$ to indicate that the occurrence a is in $g^{(i)}$.

Lemma 4.4. *Let $g, z \in \mathbb{G}$ be geodesic and $g = g_1 a g_2$, $a \in A^{\pm 1}$. Let \mathcal{D} be a geodesic van Kampen diagram for gz . If the occurrence a does not cancel, neither does any occurrence b in g_1 that belongs to $\text{adj}(a)$.*

Proof. If a does not cancel, the band L_a with an end in a has the other end in \overline{gz} . Then, for any $b \in g_1$ that cancels, the correspondent band L_b has one end in b and the other in an occurrence b^{-1} in z . Hence the band L_b crosses the band L_a . By Remark 1, $b \neq a$ and $[b, a] = 1$, so $b \notin \text{adj}(a)$. \square

Corollary 4.5. *Let $g, z \in \mathbb{G}$ be geodesic and let $g = g_1 a g_2$ be a cyclically reduced block, $a \in A^{\pm 1}$. Let \mathcal{D} be a geodesic van Kampen diagram for $z^g = gzg^{-1}$. If the occurrences a in g and the corresponding occurrence a^{-1} in g^{-1} do not cancel, then neither does any occurrence b in g_1 , $b \in \text{adj}(a)$, and the corresponding occurrence b^{-1} in g_1^{-1} .*

Corollary 4.6. *Let $g, z \in \mathbb{G}$ be geodesic and let g be a cyclically reduced block. Let \mathcal{D} be a geodesic van Kampen diagram for gz . If there exists an occurrence a in g that does not cancel, then $g^{\text{cdim}(\mathbb{G})+1} z = g \circ z'$, i.e. no occurrence in $g^{(1)}$ cancels.*

Proof. Let a be an occurrence of g that does not cancel in gz . Since $g^{\text{cdim}(\mathbb{G})+1}$ is geodesic, the occurrence $a^{(\text{cdim}(\mathbb{G})+1)}$ in $g^{(\text{cdim}(\mathbb{G})+1)}$ does not cancel.

Since g is a block, by definition, the graph $\Delta(\alpha(g))$ is connected, i.e. the subgroup generated by $\alpha(g)$ is a directly indecomposable partially commutative group. Thus, applying Lemma 4.3 to this subgroup and using the fact that the centraliser dimension of $\langle \alpha(g) \rangle$ is less or equals the centraliser dimension of \mathbb{G} (see [12]) we get that $\text{adj}(a)_{\Delta(\alpha(g))}^{\text{cdim}(\mathbb{G})} = \alpha(g)$.

Recursively applying Lemma 4.4, we get that no occurrence in $g^{(i)}$ that belongs to $\text{adj}_{\Delta(\alpha(g))}^{((\text{cdim}(\mathbb{G})+1)-i)}(a)$ cancels. Therefore, no occurrence from $g^{(1)}$ cancels in $g^{\text{cdim}(\mathbb{G})+1} z$. \square

Corollary 4.7. *Let $g, z \in \mathbb{G}$ be geodesic and let g be a cyclically reduced block. Let \mathcal{D} be a geodesic van Kampen diagram for $z^g = gzg^{-1}$. If there exists an occurrence a in g , such that a and the corresponding occurrence a^{-1} in g^{-1} do not cancel, then $z^{(g^{\dim(\mathbb{G})+1})} = g \circ z' \circ g^{-1}$.*

Proof. The proof is analogous to the one of Corollary 4.6 □

Definition 4.8. Let $z \in \mathbb{G}$ be a cyclically reduced word and $g \in \mathbb{G}$ be so that $z = g^{-1} \circ z_1$ (in the terminology of [15], g^{-1} is called a left-divisor of z). We say that the word $gzg^{-1} = z_1g^{-1}$ is a cyclic permutation of z .

Conjugating a cyclically reduced word z one gets a conjugation of a cyclic permutation of z . In particular, all letters of z appear in a geodesic $\overline{z^g}$ for any $g \in \mathbb{G}$. A more precise description is given in the following lemma.

Lemma 4.9. *Let $z, g \in \mathbb{G}$ and let z be cyclically reduced. Then there exist decompositions:*

$$g = g'z_2z_1^{-1}, \quad z = z_1z_3z_2, \quad \text{where } [z_1, z_2] = 1,$$

such that $\overline{z^g} = h \circ z_2z_3z_1 \circ h^{-1}$, h is a left-divisor of g' (perhaps trivial) and $z_2z_3z_1$ is a cyclic permutation of z .

Proof. Let z_1 be the maximal (where maximum is taken over all geodesic words representing g and z) common initial subword of z and g^{-1} , i.e. $z = z_1z'_1$, $g = g_1z_1^{-1}$. Note that such z_1 exists and is well-defined, in [15] the authors call it the left greatest common divisor of z and g^{-1} . Similarly, let z_2 be the right greatest common divisor of z and g , i.e. $z = z'_2z_2$, $g = g_2z_2$. Then z_1^{-1} and z_2 are both right divisors of g . Let d be the right greatest common divisor of z_1^{-1} and z_2 . By Proposition 3.18 of [15], $z_1^{-1} = d_1d$, $z_2 = d_2d$ and $g = g'd_2d_1^{-1}d$, where $\alpha(d_1) \cap \alpha(d_2) = \emptyset$ and $[\alpha(d_1), \alpha(d_2)] = 1$. Since z is cyclically reduced, it follows that $d = 1$. Hence, $z_1^{-1} = d_1$, $z_2^{-1} = d_2$ and the lemma follows. □

In general, let $z = w_1w_2w_1^{-1}$ be the cyclic decomposition of $z \in \mathbb{G}$. Then all the letters from w_2 appear in the geodesic $\overline{z^g}$ for any $g \in \mathbb{G}$. Indeed, we think of z^g as conjugating a cyclically reduced element w_2 by an element $g'' \in \mathbb{G}$.

Lemma 4.10. *Let $z, g \in \mathbb{G}$, let z be cyclically reduced, g be a block and let $[z, g] \neq 1$. Furthermore, suppose that g^{-1} does not left-divide z and z^{-1} . Then $z^g = gzg^{-1} = g_1 \circ z' \circ g_1^{-1}$, where $g_1 \neq 1$ is a left-divisor of g , i. e. there exist occurrences l in g and, correspondingly, l^{-1} in g^{-1} that do not cancel.*

Proof. Assume the contrary, then, in the notation of Lemma 4.9, $z_1 z_3 z_2^g = z_2 z_3 z_1$. By and in the notation of Corollary 3.3, we can write g , z and g^{-1} as follows: $g = v_1 w_1^3 w_1^2$, $z = w_2^1 v_2 w_2^3$, $g^{-1} = w_3^2 w_3^1 v_3$. By construction of z_1 , z_2 , z_3 and g_1 (see proof of Lemma 4.9) it follows that $z_2 = w_2^1$, $z_3 = v_2$, $z_1 = w_3^2$ and $g = z_2 g_1 z_1^{-1}$, $z = z_1 z_3 z_2$, $g^{-1} = z_2^{-1} g_1^{-1} z_1$. If $g_1 = 1$, then $g = z_1 z_2$ is not a block element by Lemma 4.9. We now assume that $g_1 \neq 1$. Equating the two expressions for g we get

$$z_2 g_1 z_1^{-1} = (z_2^{-1} g_1^{-1} z_1)^{-1} = z_1^{-1} g_1 z_2,$$

hence $[g_1, z_1 z_2] = 1$. Since $z_1 z_2 = z_1 \circ z_2$ and $[z_1, z_2] = 1$, by Theorem 2.3 we get $[g_1, z_1] = [g_1, z_2] = 1$. Then g is not a block element – a contradiction. \square

We now record some basic properties of $\mathbb{A}(w)$ which we shall use later. Given $x, y \in \mathbb{G}$ the following hold:

- (A) $x \in \mathbb{A}(y)$ if and only if $y \in \mathbb{A}(x)$;
- (B) if $\alpha(x) \subset \alpha(y)$ then $\mathbb{A}(y) < \mathbb{A}(x)$;
- (C) if the centraliser of x is cyclic then $\mathbb{A}(x) = 1$.

Lemma 4.11. *Let $g \in \mathbb{G}$ be a cyclically reduced block and let $z = z_1 z_2 z_1^{-1}$ be the cyclic decomposition of an element $z \in \mathbb{G}$. Suppose that g^{-1} does not left-divide z , z^{-1} , z_2 and z_2^{-1} , and $[g, z] \neq 1$. Then one has $z^{g^{2 \cdot \text{cdim}(\mathbb{G})+2}} = g \circ z^{g^{2 \cdot \text{cdim}(\mathbb{G})+1}} \circ g^{-1}$.*

Proof. Consider the product $g^{(\text{cdim}(\mathbb{G})+1)} z_1$, then no occurrence in $g^{(1)}$ cancels. Indeed, since g does not left-divide z^{-1} , there is an occurrence in g that does not cancel in $g z_1$. Applying Corollary 4.6, we get that no occurrence in $g^{(1)}$ cancels in $g^{(\text{cdim}(\mathbb{G})+1)} z_1$. We thereby get $g^{(\text{cdim}(\mathbb{G})+1)} z_1 = g \circ g' \circ z'_1$, where z'_1 is a right-divisor (may be trivial) of z_1 and g' is a left-divisor of $g^{(\text{cdim}(\mathbb{G}))}$.

Notice that since $\alpha(g') \subset \alpha(g)$ and g is a block, we get on the one hand that $g g'$ is a block and on the other that for any occurrence a in g' there exists an occurrence b in g such that $b \in \text{adj}(a)$.

If $[z'_1, g] \neq 1$ (or $[z'_1, g'] \neq 1$) then there exists an occurrence a in g (or in g') that belongs to $\text{adj}(\alpha(z'_1))$. Since no occurrence in z'_1 and in z_1^{-1} cancels in $z_2^{g g' z'_1}$, by Corollary 4.5 neither does the occurrence a in g (or in g'), and, correspondingly a^{-1} in g^{-1} (or in g'^{-1}). If a is an occurrence in g' , then there exists an occurrence b in g that belongs to $\text{adj}(a)$, and so by Corollary 4.5 this occurrence b in g and the corresponding occurrence b^{-1} in g^{-1} do not cancel. Hence, in any case, there exists an occurrence in g

that does not cancel in $\left((z_2)^{g'z'_1}\right)^g$. Therefore, by Corollary 4.7 we get that

$$\left((z_2)^{g'z'_1}\right)^{(g^{\text{cdim}(\mathbb{G})+1})} = g \circ z'_2 \circ g^{-1}$$

and thus

$$z^{g^{(2\text{cdim}(\mathbb{G})+1)}} = \left((z_2)^{g^{(\text{cdim}(\mathbb{G})+1)}z'_1}\right)^{(g^{\text{cdim}(\mathbb{G})})} = g \circ z^{g^{2\text{cdim}(\mathbb{G})}} \circ g^{-1}.$$

Assume now that $[z'_1, g] = 1$ and $[z'_1, g'] = 1$. If $[gg', z_2] \neq 1$ or $[g^2g', z_2] \neq 1$, since gg' (correspondingly, g^2g') is a block, by Lemma 4.10, there exists an occurrence a in gg' (in g^2g' , correspondingly), such that a and the occurrence a^{-1} in $(gg')^{-1}$ (in $(g^2g')^{-1}$) do not cancel in $z_2^{gg'}$ (in $z_2^{(g^2g')}$). If a is an occurrence in g' , then there exists an occurrence b in g (in g^2) that belongs to $\text{adj}(a)$, and thus by Corollary 4.5 this occurrence b in g (in g^2) and the corresponding occurrence b^{-1} in g^{-1} (in g^{-2}) do not cancel.

Now, by Corollary 4.7 we get

$$\left(z_2^{g'}\right)^{(g^{\text{cdim}(\mathbb{G})+2})} = g \circ z'_2 \circ g^{-1}$$

and thus

$$\begin{aligned} z^{g^{(2\text{cdim}(\mathbb{G})+2)}} &= \left((z_2)^{g^{(\text{cdim}(\mathbb{G})+2)}z'_1}\right)^{(g^{\text{cdim}(\mathbb{G})})} = \left((z_2)^{g^2g'z'_1}\right)^{(g^{\text{cdim}(\mathbb{G})})} = \\ &= \left(\left((z_2)^{g'}\right)^{(g^{\text{cdim}(\mathbb{G})+2})}\right)^{z'_1} = g \circ z^{g^{2\text{cdim}(\mathbb{G})+1}} \circ g^{-1}. \end{aligned}$$

Finally, suppose that $[z'_1, g] = 1$, $[z'_1, g'] = 1$, $[gg', z_2] = 1$ and $[g^2g', z_2] = 1$. We have $g^{\text{cdim}(\mathbb{G})+2} = g^2 \circ g' \circ d$, $z_1 = d^{-1} \circ z'_1$. If $g' \in \langle \sqrt{g} \rangle$, then $d \in \langle \sqrt{g} \rangle$ and so $[d, z'_1] = [d, z_2] = 1$. This derives a contradiction with the assumption that $z = z_2^{d^{-1}z'_1}$ is geodesic. Thus, we may assume that $g' \notin \langle \sqrt{g} \rangle$. In this case since $\alpha(g') \subseteq \alpha(g)$, $[g, g'] \neq 1$ and g is a block, by Theorem 2.3, from $[g, z'_1] = [g', z'_1] = 1$ follows that $\alpha(g) \subset \mathbb{A}(z'_1)$ and from $[gg', z_2] = [g^2g', z_2] = 1$ follows that $\alpha(g) \subset \mathbb{A}(z_2)$. Since $\alpha(d) \subseteq \alpha(g)$, we have $[d, z'_1] = [d, z_2] = 1$ – a contradiction with the assumption that $z = z_2^{d^{-1}z'_1}$ is geodesic. \square

Remark 3. A more subtle argument shows that the exponent $2\text{cdim}(\mathbb{G}) + 2$ in Lemma 4.11 can be replaced by $\text{cdim}(\mathbb{G})$.

Corollary 4.12. *Let $g \in \mathbb{G}$ have cyclic centraliser. Then for any element $z \in \mathbb{G}$ such that g^{-1} does not left-divide z , z^{-1} , z_2 and z_2^{-1} , one has $\mathbb{A}(z^{(g^{2\text{cdim}(\mathbb{G})+2})}) = 1$.*

Proof. Let $g = g_1 g_2 g_1^{-1}$ be the cyclic decomposition of g . We prove that no occurrence of $g_2^{(1)}$ cancels in $z^{(g^{2 \operatorname{cdim}(\mathbb{G})+2})}$, therefore, by properties (B) and (C) of \mathbb{A} , we get $\mathbb{A}(z^{(g^{2 \operatorname{cdim}(\mathbb{G})+2})}) \subseteq \mathbb{A}(g_2) = 1$.

By Lemma 4.11, no occurrence in $g_2^{(1)}$ cancels in $(z^{g_1^{-1}})^{(g_2^{2 \operatorname{cdim}(\mathbb{G})+2})}$. Then, since

$$z^{(g^{2 \operatorname{cdim}(\mathbb{G})+2})} = \left(\left(z^{g_1^{-1}} \right)^{g_2^{2 \operatorname{cdim}(\mathbb{G})+2}} \right)^{g_1},$$

no occurrence in $g_2^{(1)}$ cancels. \square

Corollary 4.13. *Let $g \in \mathbb{G}$ have cyclic centraliser. Then for any element $z \in \mathbb{G}$ such that $\mathbb{A}(z) \neq 1$ one has $\mathbb{A}(z^{(g^{2 \operatorname{cdim}(\mathbb{G})+2})}) = 1$.*

Proof. If g^{-1} left-divides $z^{\pm 1}$ or $z_2^{\pm 1}$, by property (B) of \mathbb{A} , we get that $\mathbb{A}(z) \subset \mathbb{A}(g) = 1$. \square

Remark 4. In Lemma 4.11, Corollary 4.12 and Corollary 4.13 we impose the condition that g^{-1} does not left-divide z , z^{-1} , z_2 and z_2^{-1} , because we seek the bound $2 \operatorname{cdim}(\mathbb{G}) + 2$ on the number of times one has to conjugate z by g . The reason for this is that the notion of centraliser dimension is axiomatisable using (existential) first-order formulas, [12] (we refer the reader to Section 6 for consequences of this result). If one does not impose this condition, Lemma 4.11 could be rephrased as follows.

Lemma. *Let $g \in \mathbb{G}$ be a cyclically reduced block, then for any element $z \in \mathbb{G}$ there exists $N \in \mathbb{N}$ such that $z^{(g^N)} = g \circ z^{(g^{N-1})} \circ g^{-1}$.*

4.2. Criterion to be a Domain.

Proposition 4.14. *Let \mathbb{G} be a non-abelian directly indecomposable partially commutative group. Let $g \in \mathbb{G}$ have cyclic centraliser, $x, y \in \mathbb{G}$, $x, y \neq 1$ be such that $[x, y] = 1$ and $[x, y^{(g^{2 \operatorname{cdim}(\mathbb{G})+2})}] = 1$. Then $C(x) = C(y) = C(g)$.*

Proof. Let $x = w x_1^{r_1} \dots x_k^{r_k} w^{-1}$, where x_1, \dots, x_k are cyclically reduced root elements such that $x_1^{r_1}, \dots, x_k^{r_k}$ are the blocks of $x^{w^{-1}}$ and $r_1, \dots, r_k \in \mathbb{Z}$. Since $[x, y] = 1$, by Theorem 2.3, after a certain re-enumeration of indices, we may assume

$$(3) \quad y = w x_1^{s_1} \dots x_l^{s_l} z w^{-1},$$

where $z \in \mathbb{A}(x_1 \dots x_k)$, $0 \leq l \leq k$ and $s_1, \dots, s_l \in \mathbb{Z}$. Thus,

$$(4) \quad y^{(g^{2 \operatorname{cdim}(\mathbb{G})+2})} = g^{2 \operatorname{cdim}(\mathbb{G})+2} w x_1^{s_1} \dots x_l^{s_l} z w^{-1} g^{-(2 \operatorname{cdim}(\mathbb{G})+2)}.$$

Since $[x, y^{(g^{2 \operatorname{cdim}(\mathbb{G})+2})}] = 1$ applying Theorem 2.3 once again, we get

$$(5) \quad y^{(g^{2 \operatorname{cdim}(\mathbb{G})+2})} = wx_{i_1}^{t_1} \dots x_{i_m}^{t_m} z' w^{-1},$$

where $z' \in \mathbb{A}(x_1 \dots x_k)$, $0 \leq m \leq k$ and $t_1, \dots, t_m \in \mathbb{Z}$. Equating (4) and (5), we get

$$(6) \quad (x_1^{s_1} \dots x_l^{s_l} z)^{w^{-1} g^{2 \operatorname{cdim}(\mathbb{G})+2} w} = x_{i_1}^{t_1} \dots x_{i_m}^{t_m} z'.$$

Suppose that $l \geq 1$. Then by Corollary 2.2, $l = m$ and for any $q \in \{1, \dots, l\}$ there exists $j \in \{1, \dots, m\}$ such that $\alpha(x_q) = \alpha(x_{i_j})$, $s_q = t_j$ and x_q is conjugated to x_{i_j} by $w^{-1} g^{2 \operatorname{cdim}(\mathbb{G})+2} w$.

Since $\alpha(x_q) = \alpha(x_{i_j})$, and since x_q and x_{i_j} are cyclically reduced root elements whose powers $x_q^{s_q}$ and $x_{i_j}^{t_j}$ are blocks of the same word $x^{(w^{-1})}$, we get that $x_q = x_{i_j}$ for all $1 \leq q \leq l$, i.e. $y^{w^{-1}}$ and $(y^{(g^{2 \operatorname{cdim}(\mathbb{G})+2})})^{w^{-1}}$ have the same blocks.

From the above it follows that

$$x_q^{w^{-1} g^{2 \operatorname{cdim}(\mathbb{G})+2} w} = x_{i_j} = x_q,$$

i. e. x_q commutes with $w^{-1} g^{2 \operatorname{cdim}(\mathbb{G})+2} w$. Since the centraliser of g is cyclic, so is the centraliser of $w^{-1} g^{2 \operatorname{cdim}(\mathbb{G})+2} w$ and thus, so is the centraliser of x_q .

More precisely, $C(x_q) = C(g^{2 \operatorname{cdim}(\mathbb{G})+2})^{w^{-1}} = C(g)^{w^{-1}}$.

Since x_q has cyclic centraliser, $x^{w^{-1}}$ and $y^{w^{-1}}$ both have a unique block; furthermore, since $z \in \mathbb{A}(x_q)$ by property (A) of \mathbb{A} , z is trivial. Therefore, $x = y = (x_q^{r_q})^w$ and so $C(x) = C(y) = C(x_q)^w = C(g)$.

Suppose next that $l = 0$. We prove then that x is trivial contradicting the assumption. Equations (3) and (6) rewrite as follows

$$y = zwz^{-1} \text{ and } z' = z^{w^{-1} g^{2 \operatorname{cdim}(\mathbb{G})+2} w}.$$

Notice that since $z, z' \in \mathbb{A}(x_1 \dots x_k)$ by property (C) of \mathbb{A} we get that $x_1, \dots, x_k \in \mathbb{A}(z) \cap \mathbb{A}(z')$. Therefore, if either $\mathbb{A}(z)$ or $\mathbb{A}(z')$ is trivial, so is x . Assume $\mathbb{A}(z)$ is non-trivial. Since the centraliser of $w^{-1} g w$ is cyclic, Corollary 4.13 applies to $z' = z^{((w^{-1} g w)^{2 \operatorname{cdim}(\mathbb{G})+2})}$, thus $\mathbb{A}(z') = 1$ and so $x = 1$. \square

Corollary 4.15. *Let \mathbb{G} be a non-abelian directly indecomposable partially commutative group. Let $a, b \in \mathbb{G}$ be elements with cyclic centralisers and such that $C(a) \cap C(b) = 1$. Then for any solution $x, y \in \mathbb{G}$ of the system*

$$[x, y] = 1, [x, y^{(a^{2 \operatorname{cdim}(\mathbb{G})+2})}] = 1, [x, y^{(b^{2 \operatorname{cdim}(\mathbb{G})+2})}] = 1,$$

either $x = 1$ or $y = 1$.

Proof. Applying Proposition 4.14 for the triples x, y, a and x, y, b we get that if $x \neq 1$ and $y \neq 1$, then $C(x) = C(y) = C(a)$ and $C(x) = C(y) = C(b)$ – a contradiction with $C(a) \cap C(b) = 1$. Note, that the elements a and b satisfying the assumption of the corollary exist (it suffices to take two distinct block elements such that $[a, b] \neq 1$ and $\alpha(a), \alpha(b) = A$). \square

Theorem 4.16 (Criterion for a partially commutative group to be a domain).

A partially commutative group \mathbb{G} is a domain if and if \mathbb{G} is non-abelian and directly indecomposable.

Proof. Since the direct product of two non-abelian groups is never a domain, see [1, 18], the result follows immediately from Corollary 4.15. \square

Note that Corollary 4.15 shows in fact that any non-abelian directly indecomposable partially commutative group is a domain with respect to only two elements $a^{2 \operatorname{cdim}(\mathbb{G})+2}$ and $b^{2 \operatorname{cdim}(\mathbb{G})+2}$ which are independent of the choice of x and y (in the notation of the definition of domain).

Theorem 4.17. *Let \mathbb{G} be a non-abelian directly indecomposable partially commutative group. Then $\mathbb{G}[X]$ is \mathbb{G} -discriminated by \mathbb{G} .*

Proof. The group $\mathbb{G}[X]$ is a non-abelian directly indecomposable partially commutative \mathbb{G} -group, thus by Theorem 4.16, $\mathbb{G}[X]$ is a domain. By Theorem C1 from [2], it suffices to prove that $\mathbb{G}[X]$ is \mathbb{G} -separated by \mathbb{G} .

Without loss of generality, we may assume that $X = \{x\}$. Take an element $w \in \mathbb{G}[X]$, $w = g_1 x^{k_1} g_2 \cdots g_l x^{k_l} g_{l+1}$, where $g_i \in \mathbb{G}$, $g_2, \dots, g_l \neq 1$. Take $a \in \mathbb{G}$ such that a satisfies the assumptions of Lemma 4.11. Consider the homomorphism $\varphi_a : \mathbb{G}[X] \rightarrow \mathbb{G}$, defined by $x \mapsto a^{4 \operatorname{cdim}(\mathbb{G})+5}$. By Lemma 4.11,

$$\varphi_a(x^{k_i} g_{i+1} x^{k_{i+1}}) = a^{k_i \cdot (2 \operatorname{cdim}(\mathbb{G})+3)} \circ g'_{i+1} \circ a^{k_{i+1} \cdot (2 \operatorname{cdim}(\mathbb{G})+3)}.$$

Therefore it follows (by induction on the length of w) that $\varphi_a(w) \neq 1$. \square

Corollary 4.18. *Let \mathbb{G} be a non-abelian directly indecomposable partially commutative group. Then the group $\mathbb{G}[X]$ is universally equivalent to \mathbb{G} (both in the language of groups and in the language $L_{\mathbb{G}}$ enriched by constants from \mathbb{G}).*

Proof. Follows from Theorem C2 in [1]. \square

Corollary 4.19. *Let \mathbb{G} be a non-abelian directly indecomposable partially commutative group. Then*

$$\mathbb{G} \models \forall X (U(X) = 1) \Leftrightarrow \mathbb{G}[X] \models U(X) = 1,$$

i.e. only the trivial equation has the whole set \mathbb{G}^n as its solution.

Proof. Since $\mathbb{G}[X]$ is \mathbb{G} -discriminated by \mathbb{G} , if the word $U(X)$ is a non-trivial element of $\mathbb{G}[X]$, then there exists a \mathbb{G} -homomorphism $\phi : \mathbb{G}[X] \rightarrow \mathbb{G}$ such that $U^\phi \neq 1$. Then $U(X^\phi) \neq 1$ in \mathbb{G} – a contradiction. \square

5. APPLICATIONS TO ALGEBRAIC GEOMETRY

The results and exposition of this section rely on paper [18]. We recall here some necessary definitions and restate some results in the case of partially commutative groups. We refer the reader to [18] for details and omitted proofs.

A *group code* C is a set of formulas

$$(7) \quad C = \{U(X, P), E(X, Y, P), \text{Mult}(X, Y, Z, P), \text{Inv}(X, Y, P)\}$$

where X, Y, Z, P are tuples of variables with $|X| = |Y| = |Z|$. If $P = \emptyset$ then C is called an *absolute code* or 0-code.

Let C be a group code, H be a group, and B be an $|P|$ -tuple of elements in H . We say that C (with parameters B) *interprets* a group $C(H, B)$ in H if the following conditions hold:

- 1) the truth set $U(H, B)$ in H of the formula $U(X, B)$ (with parameters B) is non-empty;
- 2) the truth set of the formula $E(X, Y, B)$ (with parameters B) defines an equivalence relation \sim_B on $U(H, B)$;
- 3) the formulas $\text{Mult}(X, Y, Z, B)$ and $\text{Inv}(X, Y, B)$ define, correspondingly, a binary operation ($Z = Z(X, Y)$) and a unary operation ($Y = Y(X)$) on the set $U(H, B)$ compatible with the equivalence relation \sim_B ;
- 4) the set of equivalence classes $U(H, B)/\sim_B$ forms a group with respect to the operations defined by $\text{Mult}(X, Y, Z, B)$ and $\text{Inv}(X, Y, B)$. We denote this group by $C(H, B)$.

We say that a group G is *interpretable* (or *definable*) in a group H if there exists a group code C and a set of parameters $B \subset H$ such that $G \simeq C(H, B)$. If C is 0-code then G is *absolutely* or *0-interpretable* in H . The following two types of interpretations are crucial. Let G be a definable subgroup of a group H , i.e., there exists a formula $U(x, P)$ and a set of parameters $B \subset H$ such that

$$G = \{g \in H \mid H \models U(g, B)\}.$$

Then G is interpretable in H by the code

$$C_G = \{U(x, P), x = y, xy = z, y = x^{-1}\}$$

with parameters B . If in addition G is a normal subgroup of H then the code

$$C_{H/G} = \{x = x, \exists v(x = yv \wedge U(v, P)), z = xy, y = x^{-1}\}$$

interprets the factor-group H/G in H with parameters B . Every group code (7) determines a *translation* T_C which is a map from the set of all formulas \mathcal{F}_L in the language L into itself. We define T_C by induction as follows:

- 1) $T_C(x = y) = E(X, Y, P)$;
- 2) $T_C(xy = z) = \text{Mult}(X, Y, Z, P)$ and $T_C(x^{-1} = y) = \text{Inv}(X, Y, P)$;
- 3) if $\phi, \psi \in \mathcal{F}_L$ and $\circ \in \{\wedge, \vee, \rightarrow\}$ then

$$T_C(\phi \circ \psi) = T_C(\phi) \circ T_C(\psi) \quad \text{and} \quad T_C(\neg\phi) = \neg T_C(\phi);$$

- 4) if $\phi \in \mathcal{F}_L$ then

$$T_C(\exists x\phi(x)) = \exists X(U(X, P) \wedge T_C(\phi)),$$

$$T_C(\forall x\phi(x)) = \forall X(U(X, P) \rightarrow T_C(\phi)).$$

Observe, that the formula $T_C(\phi)$ can be constructed effectively from ϕ .

Any partially commutative group is a direct product of finitely many non-abelian directly indecomposable partially commutative groups and its centre $Z(\mathbb{G})$, $Z(\mathbb{G}) \simeq \mathbb{Z}^k$, $k \in \mathbb{N}$. This decomposition is unique up to a permutation of factors. We refer to them as (direct) components of \mathbb{G} .

The centre $Z(\mathbb{G})$ is a normal subgroup and a definable subset of \mathbb{G} . It is the truth set of the following formula

$$\Phi_Z(x) : \forall y[x, y] = 1,$$

thus $Z(\mathbb{G})$ is 0-interpretable in \mathbb{G} . Consequently, as shown above, the quotient $\mathbb{G}/Z(\mathbb{G})$ is interpretable in \mathbb{G} .

Therefore, to work with partially commutative groups from model-theoretic viewpoint, it suffices to consider free partially commutative groups with the trivial centre.

Let \mathbb{G} be a partially commutative group without centre. As mentioned above, in this event \mathbb{G} is a direct product of directly indecomposable partially-commutative groups, which, in turn are domains by Theorem 4.16. Thus Theorem A and Corollary A of [18] apply and can be restated as follows.

Theorem A (cf. [18]). *Let \mathbb{G} be a partially commutative group with trivial centre. Then for each component \mathbb{G}_i of \mathbb{G} its elementary theory $\text{Th}(\mathbb{G}_i)$ is interpretable in the group \mathbb{G} .*

Corollary A (cf. [18]). *Let \mathbb{G} be a partially commutative group and let $\mathbb{G} = \mathbb{G}_1 \times \cdots \times \mathbb{G}_n \times \mathbb{Z}^r$, where \mathbb{G}_i is a non-abelian directly indecomposable partially commutative group, $i = 1, \dots, n$. Then the following hold:*

- 1) If $\mathbb{G} \equiv H$ then H is also a finite direct product of domains and the centre $Z(H)$, and if
- $$\mathbb{G} = \mathbb{G}_1 \times \dots \times \mathbb{G}_n \times \mathbb{Z}^r, \quad H = H_1 \times \dots \times H_m \times Z(H);$$
- are the corresponding decompositions for \mathbb{G} and H , then $n = m$, $Z(H) \equiv \mathbb{Z}^r$ and $\mathbb{G}_i \equiv H_i$ (after suitable re-ordering of the factors);
- 2) $\text{Th}(\mathbb{G})$ is decidable if and only if $\text{Th}(\mathbb{G}_i)$ is decidable for every $i = 1, \dots, n$.

Theorem B ([18]). *Let H be a minimal subdirect product of domains. Then the elementary theory of each component of H is interpretable in the group H .*

Corollary B ([18]). *Let H be a minimal subdirect product of k domains and*

$$H \hookrightarrow G_1 \times \dots \times G_k$$

be its minimal component decomposition. Then the following hold:

- 1) *if $\text{Th}(H)$ is decidable then $\text{Th}(G_i)$ is decidable for every $i = 1, \dots, k$;*
- 2) *if $\text{Th}(H)$ is λ -stable then $\text{Th}(G_i)$ is λ -stable for every $i = 1, \dots, k$.*

Theorem 5.1. *Let \mathbb{G} be a directly indecomposable partially commutative group, and Y be an algebraic set over \mathbb{G} . Then the following conditions hold:*

- 1) *the coordinate group $\Gamma(Y_i)$ of each irreducible component Y_i of Y is interpretable in the group $\Gamma(Y)$;*
- 2) *the elementary theory $\text{Th}(\Gamma(Y_i))$ of each irreducible component Y_i of Y is interpretable in the group $\Gamma(Y)$.*

Proof. Partially commutative groups are linear (see [20, 19]), thus equationally Noetherian (see [1]). We, therefore, can decompose Y as a finite union of irreducible algebraic sets, $Y = Y_1 \cup \dots \cup Y_k$, see Corollary 12 in [1]. By Proposition 12, [1] the coordinate group $\Gamma(Y)$ is a minimal subdirect product of the coordinate groups $\Gamma(Y_1), \dots, \Gamma(Y_k)$. Every group $\Gamma(Y_i)$, being a coordinate group of an irreducible algebraic set over a domain is again a domain by Theorem D2 in [1]. Now 1), 2) follow from Theorem B. \square

Corollary 5.2. *Let \mathbb{G} be a directly indecomposable partially commutative group.*

- (i) *If $Y = Y_1 \cup \dots \cup Y_k$ is an algebraic set over \mathbb{G} , where Y_1, \dots, Y_k are the irreducible components of Y , then the elementary theory of $\Gamma(Y)$ is decidable if and only if the elementary theory of $\Gamma(Y_i)$ is decidable for all $i = 1, \dots, k$.*

- (ii) If $Y = Y_1 \cup \dots \cup Y_k$ and $Z = Z_1 \cup \dots \cup Z_l$ are two irreducible algebraic sets, where Y_1, \dots, Y_k and Z_1, \dots, Z_l are the irreducible components of Y and Z , respectively, then $\Gamma(Y)$ is elementary equivalent to $\Gamma(Z)$ if and only if $k = l$ and, after a certain re-enumeration, $\Gamma(Y_i)$ is elementary equivalent to $\Gamma(Z_i)$ for all $i = 1, \dots, k$.

6. NORMAL FORMS OF FIRST-ORDER FORMULAS

6.1. Conjunctions of positive formulas. Let $L_{a,b}$ be the language of groups enriched by two constants and let \mathcal{S} be the class of all groups G satisfying the following universal sentences:

- (I) $\forall x (([x, a] = 1 \wedge [x, b] = 1) \rightarrow x = 1)$;
- (II) $\forall x \forall y \forall z (x^2 y^2 z^2 = 1 \rightarrow [x, y] = 1 \ \& \ [x, z] = 1 \ \& \ [y, z] = 1)$;
- (III) $\forall x \forall y (x^2 = y^2 \rightarrow x = y)$;
- (IV) $\forall x ([x^2, a] = 1 \rightarrow [x, a] = 1)$.

Let GROUPS be a set of axioms of group theory. Denote by $A_{\mathcal{S}}$ the union of axioms (I), (II), (III), (IV) and GROUPS. Notice that the axiom (II) is equivalent modulo GROUPS to the following quasi-identity

$$\forall x \forall y \forall z (x^2 y^2 z^2 = 1 \rightarrow [x, y] = 1).$$

It follows that all axioms in $A_{\mathcal{S}}$ are quasi-identities.

Lemma 6.1. *The class \mathcal{S} contains all partially commutative groups with trivial centre.*

Proof. We first prove that in any partially commutative group \mathbb{G} with trivial centre there exist two elements a and b such that $C(a) \cap C(b) = 1$. Indeed, let $\mathbb{G} = \mathbb{G}_1 \times \dots \times \mathbb{G}_k$, be the decomposition of \mathbb{G} in the form (1). Since $Z(\mathbb{G}) = 1$, each \mathbb{G}_i , $i = 1, \dots, k$ is a non-abelian directly indecomposable partially commutative group. For each i choose a pair of block elements $a_i, b_i \in \mathbb{G}_i$ such that $C_{\mathbb{G}_i}(a_i) \cap C_{\mathbb{G}_i}(b_i) = 1$. By Theorem 2.3, it follows that $C_{\mathbb{G}}(a_1 \dots a_k) = \langle \sqrt{a_1} \rangle \times \dots \times \langle \sqrt{a_k} \rangle$, $C_{\mathbb{G}}(b_1 \dots b_k) = \langle \sqrt{b_1} \rangle \times \dots \times \langle \sqrt{b_k} \rangle$ and so $C_{\mathbb{G}}(a_1 \dots a_k) \cap C_{\mathbb{G}}(b_1 \dots b_k) = 1$. This proves that \mathbb{G} satisfies Axiom (I).

In [5] Crisp and Wiest prove the following theorem.

Theorem 6.2 (J. Crisp, B. Wiest, [5]). *Let G be a partially commutative group. Then the equation $x^2 y^2 z^2 = 1$ has only commutative solutions.*

So \mathbb{G} satisfies Axiom (II).

By [11], partially commutative groups have least roots, and thus \mathbb{G} satisfies Axiom (III).

By Corollary 2.4, \mathbb{G} satisfies Axiom (IV). □

Lemma 6.3. *Let $G \in \mathcal{S}$. Then the equation*

$$(8) \quad x^2 a x^2 a^{-1} (y b y b^{-1})^{-2} = 1$$

has only the trivial solution $x = 1$ and $y = 1$ in G .

Proof. Let x, y be a solution of Equation (8) in G . Then we can rewrite (8) as follows

$$(9) \quad (x^2 a)^2 a^{-2} = ((y b)^2 b^{-2})^2.$$

Since G satisfies (II), from (9) we deduce that $[x^2 a, a^{-1}] = 1$, hence $[x^2, a^{-1}] = 1$. Since G satisfies (IV), it follows that $[x, a] = 1$. Now, we can rewrite (9) in the form

$$(x^2)^2 = ((y b)^2 b^{-2})^2,$$

and then, since G satisfies (III) we get

$$(10) \quad x^2 = (y b)^2 b^{-2}.$$

Again, since G satisfies (II) it follows that $[x, b] = 1$ and $[y, b] = 1$. Henceforth, $[x, a] = 1$ and $[x, b] = 1$ therefore, applying (I), we get $x = 1$. In this event, (10) reduces to $y^2 = 1$, so $y = 1$, as desired. \square

Corollary 6.4. *For any finite system of equations $S_1(X) = 1, \dots, S_k(X) = 1$ over G one can effectively find a single equation $S(X) = 1$ such that given a group $G \in \mathcal{S}$, the following holds:*

$$V_G(S_1, \dots, S_n) = V_G(S).$$

Proof. By induction it suffices to prove the result for $k = 2$. In this case, by the lemma above, the following equation

$$S_1(X)^2 a S_1(X)^2 a^{-1} (S_2(X) b S_2(X) b^{-1})^{-2} = 1$$

can be chosen as the equation $S(X) = 1$. \square

Corollary 6.5. *For any finite system of atomic formulas*

$$S_1(X) = 1, \dots, S_k(X) = 1$$

in $L_{a,b}$, one can effectively find a atomic formula $S(X) = 1$ in $L_{a,b}$ such that $(\bigwedge_{i=1}^k S_i(X) = 1)$ is $A_{\mathcal{S}}$ -equivalent to $S(X) = 1$,

$$\left(\bigwedge_{i=1}^k S_i(X) = 1 \right) \sim_{A_{\mathcal{S}}} S(X) = 1.$$

6.2. Disjunctions of positive formulas. Our next aim is to be able to rewrite finite disjunctions of equations into conjunctions of equations.

Let \mathcal{T} be the elementary theory (in the language $L_{a,b}$ of groups enriched by two constants) of non-abelian directly indecomposable partially commutative groups, i.e. the set of all first order sentences in the language of groups enriched by two constants a and b which are true in all non-abelian directly indecomposable partially commutative groups, together with the following two formulas:

- The intersection of centralisers of a and b is trivial:

$$\forall x([x, a] = 1 \wedge [x, b] = 1) \rightarrow x = 1$$

- The centralisers of a and b are cyclic. An interested reader may verify that this condition can indeed be written using the first order language.

Remark 5. Note that any model of \mathcal{T} lies in \mathcal{S} .

Proposition 6.6. *Let G be a model of \mathcal{T} . Let $a, b \in G$ be elements with cyclic centralisers and such that $C(a) \cap C(b) = 1$. Then for any solution $x, y \in G$ of the system*

$$[x, y] = 1, [x, y^{(a^{2 \operatorname{cdim}(G)+2})}] = 1, [x, y^{(b^{2 \operatorname{cdim}(G)+2})}] = 1,$$

either $x = 1$ or $y = 1$.

Proof. Indeed, since G is a model of \mathcal{T} , since the centraliser dimension of a group is a model-theoretic notion (see [12]) and since, by Corollary 4.15, any directly indecomposable partially commutative group satisfies the following sentence in $L_{a,b}$:

$$\forall x \forall y ([x, y] = 1 \wedge [x, y^{(a^{2 \operatorname{cdim}(G)+2})}] = 1 \wedge [x, y^{(b^{2 \operatorname{cdim}(G)+2})}] = 1) \rightarrow (x = 1 \vee y = 1),$$

the statement follows. \square

Combining Corollary 4.15 and Lemmas 6.1 and 6.3 yields an algorithm to encode an arbitrary finite disjunction of equations into a single equation.

Corollary 6.7. *For any finite set of equations $S_1(X) = 1, \dots, S_k(X) = 1$ one can effectively find a single equation $S(X) = 1$ such that given any model G of \mathcal{T} , the following holds:*

$$V_G(S_1) \cup \dots \cup V_G(S_k) = V_G(S).$$

Corollary 6.8. *For any finite set of atomic formulas $S_1(X) = 1, \dots, S_k(X) = 1$ one can effectively find a single atomic formula $S(X) = 1$ such that*

$$\left(\bigvee_{i=1}^k S_i(X) = 1 \right) \sim_{\mathcal{T}} S(X) = 1.$$

Corollary 6.9. *Every positive quantifier-free formula $\Phi(X)$ is equivalent modulo \mathcal{T} to a single equation $S(X) = 1$.*

6.3. Conjunctions and Disjunctions of Inequalities. The next result shows that one can effectively encode finite conjunctions and finite disjunctions of *inequations* (negations of atomic formulas) into a single inequation modulo \mathcal{T} .

Lemma 6.10. *For any finite set of inequations*

$$S_1(X) \neq 1, \dots, S_k(X) \neq 1,$$

one can effectively find an inequation $R(X) \neq 1$ and an inequation $T(X) \neq 1$ such that

$$\left(\bigwedge_{i=1}^k S_i(X) \neq 1 \right) \sim_{\mathcal{T}} R(X) \neq 1$$

and

$$\left(\bigvee_{i=1}^k S_i(X) \neq 1 \right) \sim_{\mathcal{T}} T(X) \neq 1.$$

Proof. Similar to Lemma 5 in [17]. □

Corollary 6.11. *For every quantifier-free formula $\Phi(X)$, one can effectively find a formula*

$$\Psi(X) = \bigvee_{i=1}^n (S_i(X) = 1 \wedge T_i(X) \neq 1)$$

which is equivalent to $\Phi(X)$ modulo \mathcal{T} . In particular, if \mathbb{G} is a model of \mathcal{T} , then every quantifier-free formula $\Phi(X)$ is equivalent over \mathbb{G} to a formula $\Psi(X)$ as above.

7. POSITIVE THEORY OF PARTIALLY COMMUTATIVE GROUPS

In this section we present a procedure of quantifier elimination for positive formulas over partially commutative groups (an analog of Merzlyakov's Theorem for free groups). Our approach to the positive theory of partially commutative groups is based on the proof of Merzlyakov's Theorem given in [17].

7.1. Generalised equations. Let $A = \{a_1, \dots, a_m\}$ be a set of constants and $X = \{x_1, \dots, x_n\}$ be a set of variables. Set $\mathbb{G} = G(A)$ to be a partially commutative group generated by A and $\mathbb{G}[X] = \mathbb{G} * F(X)$.

Definition 7.1. A combinatorial generalised equation Ω (with constants from $A^{\pm 1}$) consists of the following objects:

- (i) A finite set of *bases* $BS = BS(\Omega)$. Every base is either a constant base or a variable base. Each constant base is associated with exactly one letter from $A^{\pm 1}$. The set of variable bases \mathcal{M} consists of $2n$ elements $\mathcal{M} = \{\mu_1, \dots, \mu_{2n}\}$. The set \mathcal{M} comes equipped with two functions: a function $\varepsilon : \mathcal{M} \rightarrow \{1, -1\}$ and an involution $\Delta : \mathcal{M} \rightarrow \mathcal{M}$ (i.e., Δ is a bijection such that Δ^2 is an identity on \mathcal{M}). Bases μ and $\Delta(\mu)$ (or $\bar{\mu}$) are called *dual bases*. We denote variable bases by μ, λ, \dots
- (ii) A set of *boundaries* $BD = BD(\Omega)$. BD is a finite initial segment of the set of positive integers $BD = \{1, 2, \dots, \rho + 1\}$. We use letters i, j, \dots for boundaries.
- (iii) Two functions $\alpha : BS \rightarrow BD$ and $\beta : BS \rightarrow BD$. We call $\alpha(\mu)$ and $\beta(\mu)$ the initial and terminal boundaries of the base μ (or endpoints of μ). These functions satisfy the following conditions: $\alpha(b) < \beta(b)$ for every base $b \in BS$; if b is a constant base then $\beta(b) = \alpha(b) + 1$.

For a combinatorial generalised equation Ω , one can associate a system of equations in *variables* h_1, \dots, h_ρ over $G(A)$ (variables h_i are sometimes called *items*). This system is called a *generalised equation*, and, abusing the notation, we denote it by the same symbol Ω . The generalised equation Ω consists of the following three types of equations.

- (i) Each pair of dual variable bases $(\lambda, \Delta(\lambda))$ provides an equation over a partially commutative group \mathbb{G}

$$[h_{\alpha(\lambda)} h_{\alpha(\lambda)+1} \cdots h_{\beta(\lambda)-1}]^{\varepsilon(\lambda)} = [h_{\alpha(\Delta(\lambda))} h_{\alpha(\Delta(\lambda))+1} \cdots h_{\beta(\Delta(\lambda))-1}]^{\varepsilon(\Delta(\lambda))}.$$

These equations are called *basic equations*. In the case when $\beta(\lambda) = \alpha(\lambda) + 1$ and $\beta(\Delta(\lambda)) = \alpha(\Delta(\lambda)) + 1$, i.e. the corresponding basic equation takes the form:

$$[h_{\alpha(\lambda)}]^{\varepsilon(\lambda)} = [h_{\alpha(\Delta(\lambda))}]^{\varepsilon(\Delta(\lambda))},$$

without loss of generality, we shall assume that the equality above is graphical.

- (ii) For each constant base b we write down a *coefficient equation*

$$h_{\alpha(b)} = a,$$

where $a \in A^{\pm 1}$ is the constant associated with b .

Remark 6. We assume that every generalised equation comes associated with a combinatorial one;

Let A be finite alphabet then the monoid given by the presentation

$$\mathbb{T} = \langle A \mid R_A \rangle \text{ for } R_A \text{ a subset of } \{[a_i, a_j] \mid a_i, a_j \in A\}$$

is called *partially commutative monoid*. Partially commutative monoids, are also known as *trace monoids* and are extensively studied, see [10] and references there.

Definition 7.2. Let $\Omega(h) = \{L_1(h) = R_1(h), \dots, L_s(h) = R_s(h)\}$ be a generalised equation in variables $h = (h_1, \dots, h_\rho)$ with constants from $A^{\pm 1}$. A sequence of reduced nonempty words $U = (U_1(A), \dots, U_\rho(A))$ in the alphabet $A^{\pm 1}$ is a *solution* of Ω if:

- 1) all words $L_i(U), R_i(U)$ are geodesic (treated as elements of \mathbb{G}) as written;
- 2) $L_i(U) = R_i(U)$, $i = 1, \dots, s$ in the partially commutative monoid $\mathbb{T}(A^{\pm 1})$ with involution.

The notation (Ω, U) means that U is a solution of the generalised equation Ω .

Remark 7. Notice that a solution U of a generalised equation Ω can be viewed as a solution of Ω in the partially commutative monoid $\mathbb{T}(A^{\pm 1})$ (i.e., $L_i(U) = R_i(U)$ modulo commutation) which satisfies an additional condition: $U \in \mathbb{T}(A^{\pm 1})^\rho$ and U is a tuple of geodesic words in \mathbb{G} .

Obviously, each solution U of Ω gives rise to a solution of Ω in the partially commutative group $G(A)$. The converse does not hold in general, i.e. it might happen that U is a solution of Ω in $G(A)$ but not in $\mathbb{T}(A^{\pm 1})$, i.e. some equalities $L_i(U) = R_i(U)$ hold only after a reduction in \mathbb{G} . We introduce the following notation which will allow us to distinguish in which structure ($\mathbb{T}(A^{\pm 1})$ or $G(A)$) we are resolving Ω .

If

$$S = \{L_1(h) = R_1(h), \dots, L_s(h) = R_s(h)\}$$

is an arbitrary system of equations with constants from $A^{\pm 1}$, then by S^* we denote the system of equations

$$S^* = \{L_1(h)R_1(h)^{-1} = 1, \dots, L_s(h)R_s(h)^{-1} = 1\}$$

over the group $G(A)$.

Definition 7.3. A generalised equation Ω is called *formally consistent* if it satisfies the following conditions.

- 1) If $\varepsilon(\mu) = -\varepsilon(\Delta(\mu))$, then the bases μ and $\Delta(\mu)$ do not intersect, i.e. none of the items $h_{\alpha(\mu)}, h_{\beta(\mu)-1}$ is contained in $\Delta(\mu)$.
- 2) A variable cannot occur in two distinct coefficient equations, i.e., any two constant bases with the same left end-point are labeled by the same letter from $A^{\pm 1}$.

Lemma 7.4.

- (i) *If a generalised equation Ω has a solution then Ω is formally consistent;*
- (ii) *There is an algorithm to verify whether a given generalised equation is formally consistent or not.*

Remark 8. We further consider only formally consistent generalised equations.

7.2. Reduction to generalised equations. Similarly to the case of free groups, we now show how for a given finite system of equations $S(X, A) = 1$ over a partially commutative group \mathbb{G} one can associate a finite collection of generalised equations $\mathcal{GE}(S)$ with constants from $A^{\pm 1}$. The collection $\mathcal{GE}(S)$ to some extent describes all solutions of the system $S(X, A) = 1$.

Write $\{S(X, A) = 1\} = \{S_1 = 1, \dots, S_m = 1\}$ in the form

$$(11) \quad \begin{aligned} r_{11}r_{12} \dots r_{1l_1} &= 1, \\ r_{21}r_{22} \dots r_{2l_2} &= 1, \\ &\dots \\ r_{m1}r_{m2} \dots r_{ml_m} &= 1, \end{aligned}$$

where r_{ij} are letters of the alphabet $X^{\pm 1} \cup A^{\pm 1}$.

A *partition table* $T = (V, \Gamma)$ of the system above is a pair (a set of geodesic words, a partially commutative group)

$$V = \{V_{ij}(z_1, \dots, z_p)\} \quad (1 \leq i \leq m, 1 \leq j \leq l_i), \quad \Gamma = G(A \cup Z),$$

where $V_{ij} \in \mathbb{G} * F(Z) = \mathbb{G}[Z]$, in $\Gamma = G(A \cup Z)$ some of the letters in $Z = \{z_1, \dots, z_p\}$ commute, and the following conditions are satisfied:

- 1) The equality $V_{i1}V_{i2} \dots V_{il_i} = 1, 1 \leq i \leq m$, holds in Γ ;
- 2) $|V_{ij}| \leq l_i - 1$;
- 3) if $r_{ij} = a \in A^{\pm 1}$, then $V_{ij} = a$.

Since $|V_{ij}| \leq l_i - 1$ then at most $|S| = \sum_{i=1}^m (l_i - 1)l_i$ different letters z_i can occur in a partition table of $S(X, A) = 1$. Therefore we will always assume that $p \leq |S|$.

Each partition table encodes a particular type of cancelation that happens when one substitutes a particular solution $W(A) \in G(A)$ into $S(X, A) = 1$ and then reduces (in a certain way) the words in $S(W(A), A)$ into the empty word.

Lemma 7.5. *Let $S(X, A) = 1$ be a finite system of equations over $F(A)$. Then*

- (i) *the set $PT(S)$ of all partition tables of $S(X, A) = 1$ is finite, and its cardinality is bounded by a number which depends only on $|S(X, A)|$;*
- (ii) *one can effectively enumerate the set $PT(S)$.*

Proof. Since the words V_{ij} have bounded length, one can effectively enumerate the finite set of all collections of words $\{V_{ij}\}$ in $\mathbb{G}[Z]$ which satisfy the conditions 2), 3) above. Now for each such collection $\{V_{ij}\}$, one can effectively check whether the equalities $V_{i_1}V_{i_2}\dots V_{i_l} = 1, 1 \leq i \leq m$ hold in one of the finitely many (since $|Z| < \infty$) partially commutative groups Γ or not. This allows one to list effectively all partition tables for $S(X, A) = 1$. \square

To each partition table $T = \{V_{ij}\}$ one can assign a generalised equation Ω_T in the following way (below we use ‘ \doteq ’ for graphical equality). Consider the following word V in $M(A^{\pm 1} \cup Z^{\pm 1})$:

$$V \doteq V_{11}V_{12}\dots V_{1l_1}\dots V_{m1}V_{m2}\dots V_{ml_m} = y_1 \dots y_\rho,$$

where $y_i \in A^{\pm 1} \cup Z^{\pm 1}$ and $\rho = l(V)$ is the length of V . Then the generalised equation $\Omega_T = \Omega_T(h)$ has $\rho + 1$ boundaries and ρ variables h_1, \dots, h_ρ which are denoted by $h = (h_1, \dots, h_\rho)$.

Now we define bases of Ω_T and the functions $\alpha, \beta, \varepsilon$.

Let $z \in Z$. For any two distinct occurrences of z in V :

$$y_i = z^{\varepsilon_i}, \quad y_j = z^{\varepsilon_j} \quad (\varepsilon_i, \varepsilon_j \in \{1, -1\})$$

we introduce a pair of dual variable bases $\mu_{z,i}, \mu_{z,j}$ such that $\Delta(\mu_{z,i}) = \mu_{z,j}$ (say, if $i < j$). Put

$$\alpha(\mu_{z,i}) = i, \quad \beta(\mu_{z,i}) = i + 1, \quad \varepsilon(\mu_{z,i}) = \varepsilon_i.$$

The basic equation that corresponds to this pair of dual bases is $h_i^{\varepsilon_i} \doteq h_j^{\varepsilon_j}$.

Let $x \in X$. For any two distinct occurrences of x in $S(X, A) = 1$:

$$r_{i,j} = x^{\varepsilon_{ij}}, \quad r_{s,t} = x^{\varepsilon_{st}} \quad (\varepsilon_{ij}, \varepsilon_{st} \in \{1, -1\})$$

we introduce a pair of dual bases $\mu_{x,i,j}$ and $\mu_{x,s,t}$ such that $\Delta(\mu_{x,i,j}) = \mu_{x,s,t}$ (say, if $(i, j) < (s, t)$ in the left lexicographic order). Now let V_{ij} occurs in the word V as a subword

$$V_{ij} = y_c \dots y_d.$$

Then we put

$$\alpha(\mu_{x,i,j}) = c, \quad \beta(\mu_{x,i,j}) = d + 1, \quad \varepsilon(\mu_{x,i,j}) = \varepsilon_{ij}.$$

The basic equation which corresponds to these dual bases can be written in the form

$$[h_{\alpha(\mu_{x,i,j})} \dots h_{\beta(\mu_{x,i,j})-1}]^{\varepsilon_{ij}} =_{\mathbb{G}} [h_{\alpha(\mu_{x,s,t})} \dots h_{\beta(\mu_{x,s,t})-1}]^{\varepsilon_{st}}.$$

Let $r_{ij} = a \in A^{\pm 1}$. In this case we introduce a constant base μ_{ij} with the label a . If V_{ij} occurs in V as $V_{ij} = y_c$, then we put

$$\alpha(\mu_{ij}) = c, \quad \beta(\mu_{ij}) = c + 1.$$

The corresponding coefficient equation is written as $h_c = a$.

This defines the generalised equation Ω_T . Put

$$\mathcal{GE}(S) = \{\Omega_T \mid T \text{ is a partition table for } S(X, A) = 1\}.$$

Then $\mathcal{GE}(S)$ is a finite collection of generalised equations which can be effectively constructed for a given $S(X, A) = 1$.

For a generalised equation Ω we can also consider the same system of equations in a partially commutative group (not in the monoid). We denote this system by Ω^* . By $\mathbb{G}_{R(\Omega)}$ we denote the coordinate group of Ω^* . Now we explain relations between the coordinate groups of $S(X, A) = 1$ and of Ω_T^* .

For a letter x in X we choose an arbitrary occurrence of x in $S(X, A) = 1$ as

$$r_{ij} = x^{\varepsilon_{ij}}.$$

Let $\mu = \mu_{x,i,j}$ be the base that corresponds to this occurrence of x . Then V_{ij} occurs in V as the subword

$$V_{ij} = y_{\alpha(\mu)} \cdots y_{\beta(\mu)-1}.$$

Define a word $P_x(h) \in \mathbb{G}[h]$ (where $h = \{h_1, \dots, h_\rho\}$) as follows

$$P_x(h, A) = h_{\alpha(\mu)} \cdots h_{\beta(\mu)-1}^{\varepsilon_{ij}},$$

and put

$$P(h) = (P_{x_1}, \dots, P_{x_n}).$$

The tuple of words $P(h)$ depends on the choice of occurrences of letters from X in V . It follows from the construction above that the map $X \rightarrow \mathbb{G}[h]$ defined by $x \rightarrow P_x(h, A)$ gives rise to a \mathbb{G} -homomorphism

$$\pi : \mathbb{G}_{R(S)} \rightarrow \mathbb{G}_{R(\Omega_T)}.$$

Observe that the image $\pi(x)$ in $\mathbb{G}_{R(\Omega_T)}$ does not depend on a particular choice of the occurrence of x in $S(X, A)$ (the basic equations of Ω_T make these images equal). Hence π depends only on Ω_T . Thus, every solution of a generalised equation gives rise to a solution of Ω^* .

To relate solutions of $S(X, A) = 1$ to solutions of generalised equations from $\mathcal{GE}(S)$ we need the technique developed in Section 3.2.

Let $W(A)$ be a solution of $S(X, A) = 1$ in $G(A)$. If in the system (11) we make the substitution $\sigma : X \rightarrow W(A)$, then

$$(r_{i1}r_{i2} \cdots r_{il_i})^\sigma = r_{i1}^\sigma r_{i2}^\sigma \cdots r_{il_i}^\sigma = 1$$

in $G(A)$ for every $i = 1, \dots, m$.

Since every product $R_i = r_{i1}^\sigma r_{i2}^\sigma \cdots r_{il_i}^\sigma$ is trivial, we can choose a van Kampen diagram \mathcal{D}_{R_i} for R_i . Denote by $\tilde{z}_1, \dots, \tilde{z}_p$ the subwords w_j^k , $1 \leq$

$j < k \leq l_i$ of r_{ij}^σ , where here w_j^k are defined as in Lemma 3.2. Since, by Lemma 3.2 $w_j^k = w_k^{j-1}$, the word r_{ij} can be written as a word in $\tilde{z}_1, \dots, \tilde{z}_p$:

$$r_{ij}^\sigma = V_{ij}(\tilde{z}_1, \dots, \tilde{z}_p)$$

for some freely reduced words $V_{ij}(Z)$ in variables $Z = \{z_1, \dots, z_p\}$. Observe that if $r_{ij} = a \in A^{\pm 1}$ then $r_{ij}^\sigma = a$ and we have $V_{ij} = a$. By Lemma 3.2, r_{ij}^σ is a product of at most $l_i - 1$ words w_j^k , we have $l(V_{ij}) \leq l_i - 1$. Take a partially commutative group $\Gamma = G(A \cup Z)$ whose underlying commutation graph is defined as follows:

- two elements a_i, a_j in $A^{\pm 1}$ commute whenever they commute in \mathbb{G} ;
- an element $a \in A^{\pm 1}$ commutes with z_i if and only if a commutes with the word w_j^k corresponding to z_i ;
- two elements $z_i, z_j \in Z$ commute whenever the corresponding words w_j^k do.

In the above notation, the set $T = \{V_{ij}\}$ along with the group Γ is a partition table for $S(X, A) = 1$. Obviously,

$$U(A) = (\tilde{z}_1, \dots, \tilde{z}_p)$$

is the solution of the generalised equation Ω_T , which is induced by $W(A)$. From the construction of the map $P(h)$ we deduce that $W(A) = P(U(A))$.

The converse is also true: if $U(A)$ is an arbitrary solution of the generalised equation Ω_T , then $P(U(A))$ is a solution of $S(X, A) = 1$.

We summarize the discussion above in the following lemma.

Lemma 7.6. *For a given system of equations $S(X, A) = 1$ over \mathbb{G} , one can effectively construct a finite set*

$$\mathcal{GE}(S) = \{\Omega_T \mid T \text{ is a partition table for } S(X, A) = 1\}$$

of generalised equations such that

- (i) if the set $\mathcal{GE}(S)$ is empty, then $S(X, A) = 1$ has no solutions in \mathbb{G} ;
- (ii) for each $\Omega(h) \in \mathcal{GE}(S)$ and for each $x \in X$ one can effectively find a word $P_x(h, A) \in \mathbb{G}[h]$ of length at most $|h|$ such that the map $x \mapsto P_x(h, A)$ ($x \in X$) gives rise to a \mathbb{G} -homomorphism $\pi_\Omega : \mathbb{G}_{R(S)} \rightarrow \mathbb{G}_{R(\Omega)}$;
- (iii) for any solution $W(A) \in \mathbb{G}^n$ of the system $S(X, A) = 1$ there exists $\Omega(h) \in \mathcal{GE}(S)$ and a solution $U(A)$ of $\Omega(h)$ such that $W(A) = P(U(A))$, where $P(h) = (P_{x_1}, \dots, P_{x_n})$, and this equality holds in the partially commutative monoid $\mathbb{T}(A^{\pm 1})$;
- (iv) for any \mathbb{G} -group $\tilde{\mathbb{G}}$, if a generalised equation $\Omega(h) \in \mathcal{GE}(S)$ has a solution \tilde{U} in $\tilde{\mathbb{G}}$, then $P(\tilde{U})$ is a solution of $S(X, A) = 1$ in $\tilde{\mathbb{G}}$.

Corollary 7.7. *In the notation of Lemma 7.6 for any solution $W(A) \in \mathbb{G}^n = G(A)^n$ of the system $S(X, A) = 1$ there exists $\Omega(h) \in \mathcal{GE}(S)$ and a solution $U(A)$ of $\Omega(h)$ such that the following diagram commutes*

$$\begin{array}{ccc} \mathbb{G}_{R(S)} & \xrightarrow{\pi} & \mathbb{G}_{R(\Omega)} \\ \pi_W \downarrow & & \swarrow \pi_U \\ \mathbb{G} & & \end{array}$$

7.3. Positive theory of partially commutative groups and direct product of groups. In this section we first prove a result on elimination of quantifiers for positive sentences over a non-abelian directly indecomposable partially commutative group $\mathbb{G} = G(A)$. This proof is based on the notion of a generalised equation. Combining this result with a theorem of V. Diekert and A. Muscholl on decidability of equations over partially commutative groups, see [9], we get that the positive theory of free partially commutative groups in the language of group theory L and the language $L_{\mathbb{G}}$ enriched by constants is decidable. Note that V. Diekert and M. Lohrey, using a different method, prove a similar result in [7]. Furthermore, we apply the techniques developed for the proof of quantifier elimination to obtain a result on lifting arbitrary formulas from \mathbb{G} to $\mathbb{G} * F$, where F is a free group of finite rank (see Theorem 7.12). In order to prove that the positive theory of any partially commutative group is decidable, we need to study the positive theory of the direct product of groups. In the appendix of the paper we prove that if $G = H_1 \times \cdots \times H_k$, then the positive theory of G in both languages L and L_G is decidable if the positive theories of H_1, \dots, H_k are decidable.

Recall that every positive formula $\Psi(Z)$ in the language L_A is equivalent modulo $A_{\mathcal{T}}$ to a formula of the type

$$\forall x_1 \exists y_1 \dots \forall x_k \exists y_k (S(X, Y, Z, A) = 1),$$

where $S(X, Y, Z, A) = 1$ is an equation with constants from $A^{\pm 1}$, $X = (x_1, \dots, x_k)$, $Y = (y_1, \dots, y_k)$, $Z = (z_1, \dots, z_m)$. Indeed, one can insert auxiliary quantifiers to ensure the direct alteration of quantifiers in the prefix. In particular, every positive sentence in L_A is equivalent modulo $A_{\mathcal{T}}$ to a formula of the type

$$\forall x_1 \exists y_1 \dots \forall x_k \exists y_k (S(X, Y, A) = 1).$$

Theorem 7.8 (Elimination of Quantifiers). *If*

$$\mathbb{G} \models \forall x_1 \exists y_1 \dots \forall x_k \exists y_k (S(X, Y, A) = 1),$$

then there exist words (with constants from \mathbb{G}) $q_1(x_1), \dots, q_k(x_1, \dots, x_k) \in \mathbb{G}[X]$, such that

$$\mathbb{G}[X] \models S(x_1, q_1(x_1), \dots, x_k, q_k(x_1, \dots, x_k, A)) = 1,$$

i.e. the equation

$$S(x_1, y_1, \dots, x_k, y_k, A) = 1$$

(in variables Y) has a solution in the group $\mathbb{G}[X]$.

Proof. Let $\mathcal{GE}(u) = \{\Omega_1(Z_1), \dots, \Omega_r(Z_r)\}$ be generalised equations associated with the equation $S(X, Y, A) = 1$ in Lemma 7.6. Denote by $\rho_i = |Z_i|$ the number of variables in Ω_i .

Since the group \mathbb{G} is directly indecomposable, there exists a path p in the non-commutation graph Δ of \mathbb{G} beginning in a vertex b_1 which goes through every vertex of Δ at least once. Denote by $b_1 \cdots b_n$ the label of the path p . Set

$$b = b_1 b_2 \cdots b_{n-1} b_n b_{n-1} \cdots b_1, \quad a = b_2 b b_2 = b_2 b_1 b_2 \cdots b_{n-1} b_n b_{n-1} \cdots b_2 b_1 b_2,$$

and

$$g_1 = b^m a^{m_{1,1}} b^m a^{m_{1,2}} b \dots a^{m_{1,n_1}} b^m,$$

where $0 < m_{1,1} < m_{1,2} < \dots < m_{1,n_1}$, $\max\{\rho_1, \dots, \rho_r\} |S(X, A)| < n_1$ and $m \in \mathbb{N}$ is a constant which depends only on the generalised equation. Then there exists h_1 such that

$$\mathbb{G} \models \forall x_2 \exists y_2 \dots \forall x_k \exists y_k (S(g_1, h_1, x_2, y_2, \dots, x_k, y_k) = 1).$$

Suppose now that elements $g_1, h_1, \dots, g_{i-1}, h_{i-1} \in \mathbb{G}$ are given. We define

$$(12) \quad g_i = b^m a^{m_{i1}} b^m a^{m_{i2}} b^m \dots a^{m_{in_i}} b^m$$

such that

- 1) $0 < m_{i1} < m_{i2} < \dots < m_{in_i}$;
- 2) $\max\{\rho_1, \dots, \rho_r\} |S(X, A)| < n_i$;
- 3) no subword of the type $b^m a^{m_{ij}} b^m$ occurs in any of the words g_l , $l < i$ and in any of the (finitely many) words h'_l such that $h'_l = h_l$ in the partially commutative monoid, $l < i$.

Then there exists an element $h_i \in \mathbb{G}$ such that

$$\mathbb{G} \models \forall x_{i+1} \exists y_{i+1} \dots \forall x_k \exists y_k (S(g_1, h_1, \dots, g_i, h_i, x_{i+1}, y_{i+1}, \dots, x_k, y_k) = 1).$$

By induction we have constructed elements $g_1, h_1, \dots, g_k, h_k \in \mathbb{G}$ such that

$$S(g_1, h_1, \dots, g_k, h_k) = 1$$

and each g_i has the form (12) and satisfies conditions 1), 2), 3).

By Lemma 7.6 there exists a generalised equation $\Omega(Z) \in \mathcal{GE}(S)$, words $P_i(Z, A), Q_i(Z, A) \in \mathbb{G}[Z]$ ($i = 1, \dots, k$) of length lower than $\rho = |Z|$, and a solution $U = (u_1, \dots, u_\rho)$ of $\Omega(Z)$ in \mathbb{G} such that the following words are equal in the partially commutative monoid:

$$g_i = P_i(U), \quad h_i = Q_i(U) \quad (i = 1, \dots, k).$$

Notice that from the definition of a and b it follows that no two consecutive letters in a and b , and thus in g_i commute. Therefore, the equality $g_i = P_i(U)$ is graphical.

Since $n_i > \rho|S(X, A)|$ (by condition 2)) and $P_i(U) = y_1 \dots y_q$ with $y_i \in U^{\pm 1}$, $q \leq \rho$, the graphical equalities

$$(13) \quad g_i = ba^{m_{i1}}ba^{m_{i2}}b \dots a^{m_{in_i}}b = P_i(U) \quad (i = 1, \dots, k)$$

show that there exists a subword $v_i = b^m a^{m_{ij}} b^m$ of g_i such that every occurrence of this subword in (13) is an occurrence inside some $u_j^{\pm 1}$. For each i fix such a subword $v_i = b^m a^{m_{ij}} b^m$ in g_i . In view of condition 3) the word v_i does not occur in any of the words g_j ($j \neq i$), h_s ($s < i$), moreover, in g_i it occurs precisely once. Denote by $j(i)$ the unique index such that v_i occurs inside $u_{j(i)}^{\pm 1}$ in $P_i(U)$ from (13) (and v_i occurs in it precisely once).

The argument above shows that the variable $z_{j(i)}$ does not occur in words $P_t(Z, A)$ ($t \neq i$), $Q_s(Z, A)$ ($s < i$). Moreover, in $P_i(Z)$ it occurs precisely once. It follows that the variable $z_{j(i)}$ in the generalised equation $\Omega(Z)$ does not occur neither in coefficient equations nor in basic equations corresponding to the dual bases related to x_t ($t \neq i$), y_s ($s < i$).

We “mark” (or select) the unique occurrence of v_i (as $v_i^{\pm 1}$) in $u_{j(i)}$ $i = 1, \dots, k$. Now we are going to mark some other occurrences of v_i in the words u_1, \dots, u_ρ as follows. Suppose that some u_d has a marked occurrence of some v_i . If Ω contains an equation of the type $z_d^\varepsilon = z_r^\delta$, then $u_d^\varepsilon = u_r^\delta$ graphically. Hence u_r has an occurrence of the subword $v_i^{\pm 1}$ which corresponds to the marked occurrence of $v_i^{\pm 1}$ in u_d . We mark this occurrence of $v_i^{\pm 1}$ in u_r .

Suppose Ω contains an equation of the type

$$[h_{\alpha_1} \dots h_{\beta_1-1}]^{\varepsilon_1} = [h_{\alpha_2} \dots h_{\beta_2-1}]^{\varepsilon_2}$$

such that z_d occurs in it, say in the left-hand side of the equality. Then

$$[u_{\alpha_1} \dots u_{\beta_1-1}]^{\varepsilon_1} = [u_{\alpha_2} \dots u_{\beta_2-1}]^{\varepsilon_2}$$

in the partially commutative monoid $\mathbb{T}(A^{\pm 1})$. Since $v_i^{\pm 1}$ is a subword of u_d , a subword $v_{i,1} = b^{m-1} a^{m_{i,j}} b^{m-1}$ occurs also in the right-hand side of the above equality, say in u_r . Indeed, let $w_1 b^m a^{m_{i,j}} b^m w_2 = w$ in the monoid $\mathbb{T}(A^{\pm 1})$. Since for any letter l in w_k , $k = 1, 2$ there exists a letter l' in b such that $[l, l'] \neq 1$ and since, by the definition of a and b , no two consecutive occurrences in $b^{m-1} a^{m_{i,j}} b^{m-1}$ commute, the statement follows. We mark this occurrence of $v_{i,1}^{\pm 1}$ in u_r and in all the previously marked occurrences of $v_i^{\pm 1} = b v_{i,1} b$. We continue the marking process, but now, instead of v_i we mark the occurrences of $v_{i,1}$. The marking process stops in finitely many steps and all the occurrences of the subword $v_{i,k} = b^{m-k} a^{m_{i,j}} b^{m-k}$

are marked. For the above argument, it suffices to choose $m > k$, which depends on the generalised equation only.

Now in all words u_1, \dots, u_ρ we replace every marked occurrence of $v_{i,k} = b^{m-k} a^{m_{ij}} b^{m-k}$ with a new word $b^{m-k} a^{m_{ij}} x_i b^{m-k}$ from the group $\mathbb{G}[X]$. Denote the resulting words from $\mathbb{G}[X]$ by $\tilde{u}_1, \dots, \tilde{u}_\rho$. It follows from description of the marking process that the tuple $\tilde{U} = (\tilde{u}_1, \dots, \tilde{u}_\rho)$ is a solution of the generalised equation Ω in $\mathbb{G}[X]$. Indeed, all the equations in Ω hold in the partially commutative monoid if we substitute z_i by u_i . Hence the substitution $u_i \rightarrow \tilde{u}_i$ still makes them equal. Now by Lemma 7.6, $X = P(\tilde{U}), Y = Q(\tilde{U})$ is a solution of the equation $S(X, A) = 1$ over $\mathbb{G}[X]$ as desired. \square

Corollary 7.9. *There is an algorithm which for a given positive sentence*

$$\forall x_1 \exists y_1 \dots \forall x_k \exists y_k (S(X, Y, A) = 1)$$

in L_A determines whether or not this formula holds in \mathbb{G} , and if it does, the algorithm finds words

$$q_1(x_1), \dots, q_k(x_1, \dots, x_k) \in \mathbb{G}[X]$$

such that

$$\mathbb{G}[X] \models u(x_1, q_1(x_1), \dots, x_k, q_k(x_1, \dots, x_k)) = 1,$$

i.e. the positive theory of any directly-indecomposable partially commutative group is decidable.

Proof. The proof follows from Theorem 7.8 and decidability of equations over free partially commutative groups. Indeed, since the compatibility problem for a system of equations over a partially commutative group \mathbb{G} reduces to the compatibility of a system of equations S' over a free group with rational constraints C , see [9]. In order to prove the Corollary it suffices to check the compatibility of S' over a free group with constraints $C \cup \{y_i \in F[X_i]\}$, where $X_i = \{x_1, \dots, x_i\}$. \square

The next result follows directly from Corollary 8.14.

Corollary 7.10. *Let \mathbb{G} be an arbitrary partially commutative group. Then the positive theory of \mathbb{G} is decidable.*

Definition 7.11. Let ϕ be a sentence in the language L_A written in the standard form

$$\phi = \forall x_1 \exists y_1 \dots \forall x_k \exists y_k \phi_0(x_1, y_1, \dots, x_k, y_k),$$

where ϕ_0 is a quantifier-free formula in L_A . We say that \mathbb{G} *freely lifts* ϕ if there exist words (with constants from \mathbb{G}) $q_1(x_1), \dots, q_k(x_1, \dots, x_k) \in \mathbb{G}[X]$,

such that

$$\mathbb{G}[X] \models \phi_0(x_1, q_1(x_1), \dots, x_k, q_k(x_1, \dots, x_k, A)) = 1.$$

Theorem 7.12. *Let \mathbb{G} be a non-abelian directly indecomposable partially commutative group. Then \mathbb{G} freely lifts every sentence in L_A that is true in \mathbb{G} .*

Proof. Suppose a sentence

$$(14) \quad \phi = \forall x_1 \exists y_1 \dots \forall x_k \exists y_k (U(x_1, y_1, \dots, x_k, y_k) = 1 \wedge V(x_1, y_1, \dots, x_k, y_k) \neq 1),$$

is true in \mathbb{G} . We choose $x_1 = g_1, y_1 = h_1, \dots, x_k = g_k, y_k = h_k$ precisely as in Theorem 7.8. Then the formula

$$U(g_1, h_1, \dots, g_k, h_k) = 1 \wedge V(g_1, h_1, \dots, g_k, h_k) \neq 1$$

holds in \mathbb{G} . In particular, $U(g_1, h_1, \dots, g_k, h_k) = 1$ in \mathbb{G} . It follows from Corollary 7.9 that there are words $q_1(x_1) \in \mathbb{G}[x_1], \dots, q_k(x_1, \dots, x_k) \in \mathbb{G}[x_1, \dots, x_k]$ such that

$$\mathbb{G}[X] \models U(x_1, q_1(x_1, \dots, x_k), \dots, x_k, q_k(x_1, \dots, x_k)) = 1.$$

Moreover, it follows from the construction that $h_1 = q_1(g_1), \dots, h_k = q_k(g_1, \dots, g_k)$. We claim that

$$\mathbb{G}[X] \models V(x_1, q_1(x_1, \dots, x_k), \dots, x_k, q_k(x_1, \dots, x_k)) \neq 1.$$

Indeed, if

$$V(x_1, q_1(x_1, \dots, x_k), \dots, x_k, q_k(x_1, \dots, x_k)) = 1$$

in $\mathbb{G}[X]$, then its image in \mathbb{G} under any specialization $X \rightarrow \mathbb{G}$ is also trivial, but this is not the case for the specialization $x_1 \rightarrow g_1, \dots, x_k \rightarrow g_k$ — a contradiction. This proves the theorem for sentences ϕ of the form (14). A similar argument works for formulas of the type

$$\phi = \forall x_1 \exists y_1 \dots \forall x_k \exists y_k \bigvee_{i=1}^n (U_i(x_1, y_1, \dots, x_k, y_k) = 1 \wedge V_i(x_1, y_1, \dots, x_k, y_k) \neq 1),$$

which is, actually, the general case by Corollary 6.11. \square

APPENDIX: POSITIVE THEORY OF THE DIRECT PRODUCT OF GROUPS

In this section we prove that if $G = H_1 \times \dots \times H_k$, then the positive theory of G in the language L_G (and in L) is decidable if the positive theories of H_1, \dots, H_k are decidable. Perhaps, this result is known, nevertheless, we were not able to find a reference.

The following theorem is due to Feferman and Vaught, see [16].

Theorem 8.13. *Let $G = H_1 \times \cdots \times H_k$. Then the elementary theory of G in the language L_G is decidable, provided that the elementary theory of H_i is decidable, $i = 1, \dots, k$.*

Proof. Without loss of generality we may assume that $G = A \times B$.

We use induction on the complexity of the formula to prove the following statement. Given a formula $\varphi(x_1, \dots, x_n)$ in the language L_G , one can effectively construct a finite family of formulas $\langle \phi \rangle = \{(\psi_i(y_1, \dots, y_n), \psi'_i(z_1, \dots, z_n)) \mid i \in I\}$ such that for all $a_1, \dots, a_n, b_1, \dots, b_n$ we have

$$A \times B \models \varphi((a_1, b_1), \dots, (a_n, b_n))$$

if and only if there exists $i \in I$ such that

$$A \models \psi_i(a_1, \dots, a_n) \text{ and } B \models \psi'_i(b_1, \dots, b_n).$$

- Let $\varphi = (x_i = x_j)$, set $\langle \varphi \rangle = \{(y_i = y_j, z_i = z_j)\}$.
- Let $\varphi = (x_i = c)$, where $c \in G$, $c = (c_1, c_2)$, set $\langle \varphi \rangle = \{(y_i = c_1, z_i = c_2)\}$.
- Let $\varphi = P(x_1, \dots, x_n)$, where P is a predicate. Set $\langle \varphi \rangle = \{(P(y_1, \dots, y_n), P(z_1, \dots, z_n))\}$.
- Let $\varphi = \varphi_1 \vee \varphi_2$ and set $\langle \varphi \rangle = \langle \varphi_1 \rangle \cup \langle \varphi_2 \rangle$.
- Let $\varphi = \neg \varphi_0$ and set

$$\langle \varphi \rangle = \left\{ \left(\bigwedge_{j \in J} \neg \psi_j, \bigwedge_{i \in I \setminus J} \neg \psi'_i \right) \middle| J \in \mathcal{P}(I) \right\},$$

where $\mathcal{P}(I)$ is the power set of I and $\langle \phi_0 \rangle = \{(\psi_i, \psi'_i) \mid i \in I\}$.

- Let $\varphi = \exists x_0 \varphi_0(x_0, x_1, \dots, x_n)$ and set

$$\langle \varphi \rangle = \{(\exists y_0 \psi_i(y_0, y_1, \dots, y_n), \exists z_0 \psi'_i(z_0, z_1, \dots, z_n)) \mid i \in I\},$$

where $\langle \phi_0 \rangle = \{(\psi_i, \psi'_i) \mid i \in I\}$.

□

Corollary 8.14. *Let $G = H_1 \times \cdots \times H_k$. Then the positive theory of G in the language L_G (in the language L) is decidable, provided that the positive theories of H_1, \dots, H_k are decidable.*

Proof. In the notation of Theorem 8.13, we are left to show that if φ is a positive formula in L_G then for all $i \in I$ the formulas ψ_i and ψ'_i are also positive.

By construction of $\langle \varphi \rangle$ it follows that ψ_i and ψ'_i are positive when $\varphi = (x_i = x_j)$, $\varphi = (x_i = c)$, $\varphi = P(x_1, \dots, x_n)$, $\varphi = \varphi_1 \vee \varphi_2$, and $\varphi = \exists x_0 \varphi_0(x_0, x_1, \dots, x_n)$. We are left to consider the two following cases: $\varphi = \varphi_1 \wedge \varphi_2$ and $\varphi = \forall x_0 \varphi_0(x_0, x_1, \dots, x_n)$.

Let $\varphi = \forall x_0 \varphi_0$. Then φ is equivalent to $\neg(\exists x_0 \neg \varphi_0(x_0, x_1, \dots, x_n))$. Thus,

$$\begin{aligned} \langle \varphi \rangle &= \neg \left\{ \left(\exists x_0 \bigwedge_{j \in J} \neg \psi_j, \exists x_0 \bigwedge_{i \in I \setminus J} \neg \psi'_i \right) \middle| J \in \mathcal{P}(I) \right\} = \\ & \left\{ \left(\bigwedge_{J \in J'} \neg \left(\exists x_0 \bigwedge_{j \in J} \neg \psi_j \right), \bigwedge_{I \in \mathcal{P} \setminus J'} \neg \left(\exists x_0 \bigwedge_{i \in I} \neg \psi'_i \right) \right) \middle| J' \in \mathcal{P}(\mathcal{P}(I)) \right\} = \\ & \left\{ \left(\bigwedge_{J \in J'} \forall x_0 \bigvee_{j \in J} \psi_j, \bigwedge_{I \in \mathcal{P} \setminus J'} \forall x_0 \bigvee_{i \in I} \psi'_i \right) \middle| J' \in \mathcal{P}(\mathcal{P}(I)) \right\}. \end{aligned}$$

Let now $\varphi = \varphi_1 \wedge \varphi_2$ and $\langle \varphi_l \rangle = \{(\psi_{l,i}, \psi'_{l,i}) \mid i \in I_l\}$, $l = 1, 2$. Then φ is equivalent to $\neg(\neg \varphi_1 \vee \neg \varphi_2)$. Thus,

$$\begin{aligned} \langle \varphi \rangle &= \neg \left\{ \left\{ \left(\bigwedge_{j \in J} \neg \psi_{1,j}, \bigwedge_{i \in I_1 \setminus J} \neg \psi'_{1,i} \right) \middle| J \in \mathcal{P}(I_1) \right\} \cup \right. \\ & \quad \left. \left\{ \left(\bigwedge_{j \in J} \neg \psi_{2,j}, \bigwedge_{i \in I_2 \setminus J} \neg \psi'_{2,i} \right) \middle| J \in \mathcal{P}(I_2) \right\} \right\} = \\ & \left\{ \neg \left\{ \left(\bigwedge_{j \in J} \neg \psi_{1,j}, \bigwedge_{i \in I_1 \setminus J} \neg \psi'_{1,i} \right) \middle| J \in \mathcal{P}(I_1) \right\} \cap \right. \\ & \quad \left. \neg \left\{ \left(\bigwedge_{j \in J} \neg \psi_{2,j}, \bigwedge_{i \in I_2 \setminus J} \neg \psi'_{2,i} \right) \middle| J \in \mathcal{P}(I_2) \right\} \right\} = \\ & \left\{ \left(\bigwedge_{J \in J'} \neg \left(\bigwedge_{j \in J} \neg \psi_{1,j} \right), \bigwedge_{I \in \mathcal{P}(I_1) \setminus J} \neg \left(\bigwedge_{i \in I_1 \setminus J} \neg \psi'_{1,i} \right) \right) \middle| J' \in \mathcal{P}(\mathcal{P}(I_1)) \right\} \cap \\ & \left\{ \left(\bigwedge_{J \in J'} \neg \left(\bigwedge_{j \in J} \neg \psi_{2,j} \right), \bigwedge_{I \in \mathcal{P}(I_2) \setminus J} \neg \left(\bigwedge_{i \in I_2 \setminus J} \neg \psi'_{2,i} \right) \right) \middle| J' \in \mathcal{P}(\mathcal{P}(I_2)) \right\} = \\ & \left\{ \left(\bigwedge_{J \in J'} \bigvee_{j \in J} \psi_{1,j}, \bigwedge_{I \in \mathcal{P}(I_1) \setminus J} \bigvee_{i \in I_1 \setminus J} \psi'_{1,i} \right) \middle| J' \in \mathcal{P}(\mathcal{P}(I_1)) \right\} \cap \end{aligned}$$

$$\left\{ \left(\bigwedge_{J \in J'} \bigvee_{j \in J} \psi_{2,j}, \bigwedge_{I \in \mathcal{P}(I_2) \setminus J} \bigvee_{i \in I_2 \setminus J} \psi'_{2,i} \right) \middle| J' \in \mathcal{P}(\mathcal{P}(I_2)) \right\}$$

□

REFERENCES

- [1] G. Baumslag, A. G. Myasnikov, V. N. Remeslennikov, *Algebraic geometry over groups I. Algebraic sets and Ideal Theory*. J. Algebra 219 (1999), 16-79.
- [2] G. Baumslag, A. G. Myasnikov, V. N. Remeslennikov, *Discriminating Completions of Hyperbolic Groups*. Geom. Dedicata 92 (2002), 115-143.
- [3] N. Brady, H. Short, T. Riley *The Geometry of the Word Problem for Finitely Generated Groups (Advanced Courses in Mathematics - CRM Barcelona)* Birkhauser; 2007.
- [4] R. Charney, *An introduction to right-angled Artin groups*, arXiv:math/0610668.
- [5] J. Crisp and B. Wiest, *Embeddings of graph braid groups and surface groups in right-angled Artin groups and braid groups*, Algebraic and Geometric Topology 4 (2004) 439-472.
- [6] V. Diekert, C. Gutierrez, C. Hagenah *The existential theory of equations with rational constraints in free groups is PSPACE-complete*, Information and Computation 202(2005), 105-140.
- [7] V. Diekert and M. Lohrey, *Existential and Positive Theories of Equations in Graph Products*, Theory of Computing Systems **37** (2004), 133-156.
- [8] V. Diekert and M. Lohrey, *Word equations over graph products*, FSTTCS 2003, LNCS 2914, 156-167.
- [9] V. Diekert and A. Muscholl *Solvability of Equations in Free Partially Commutative Groups is Decidable*, Internat. J. Algebra Comput. **16** (2006), 1047-1070.
- [10] V. Diekert, G. Rozenberg, ed. *The Book of Traces*; World Scientific, 1995.
- [11] G. Duchamp and D. Krob, *Partially Commutative Magnus Transformations*, Internat. J. Algebra Comput. **3** (1993) 15-41.
- [12] A. J. Duncan, I. V. Kazachkov, V. N. Remeslennikov, *Centraliser dimension and universal classes of groups*, SEMR 3(2006), 197-215.
- [13] A. J. Duncan, I. V. Kazachkov, V. N. Remeslennikov, *Centraliser Dimension of Partially Commutative Groups*, Geom. Dedicata, 120 (2006), p. 73-97.
- [14] A. J. Duncan, I. V. Kazachkov, V. N. Remeslennikov, *Parabolic and Quasiparabolic Subgroups of Free Partially Commutative Groups*, J. Algebra, to appear.
- [15] E. S. Esyp, I. V. Kazachkov and V. N. Remeslennikov, *Divisibility Theory and Complexity of Algorithms for Free Partially Commutative Groups*, In: Groups, Languages, Algorithms. Contemporary Mathematics **378** (2005), 319-348.
- [16] S. Feferman, L. Vaught, *The first order properties of products of algebraic systems*, Fund. Math., 47 (1959), 57-103.
- [17] O. Kharlampovich, A. Myasnikov, *Implicit function theorem over free groups*, J. Algebra, vol 290 (2005), 1-203.
- [18] A. Kvaschuk, A. G. Myasnikov, V. N. Remeslennikov, *Algebraic geometry over groups III. Elements of Model Theory*. J. of Algebra 288 (2005), 7898.
- [19] T. Hsu and D. Wise, *On linear and residual properties of graph products*, Mich. Math. J. 46 (1999), 251-259.
- [20] S. Humphries, *On representations of Artin groups and the Tits conjecture*, J. Algebra 169 (1994) 847-862.

- [21] G.S. Makanin. *Equations in a free group* (Russian). *Izv. Akad. Nauk SSSR, Ser. Mat.* 46, 1199-1273, 1982. transl. in *Math. USSR Izv.*, V. 21, 1983.
- [22] G.S. Makanin. *Decidability of the universal and positive theories of a free group* (Russian). *Izv. Akad. Nauk SSSR, Ser. Mat.* 48(1), 735-749, 1985. transl. in *Math. USSR Izv.*, V. 25, 1985.
- [23] A. I. Mal'cev. *On the equation $zxyx^{-1}y^{-1}z^{-1} = aba^{-1}b^{-1}$ in a free group.* (Russian), *Algebra i Logika Sem.* 1, 1962 no. 5, 45-50.
- [24] Ju. I. Merzlyakov. *Positive formulae on free groups.* *Algebra i Logika*, 5(1966), 25-42.
- [25] A. G. Myasnikov, V. N. Remeslennikov, *Algebraic geometry over groups II. Logical Foundations.* *J. of Algebra* 234(2000), 225-276.
- [26] A. Myasnikov and P. Shumyatsky, *Discriminating groups and c-dimension*, *J. Group Theory* 7(1), p. 135142.
- [27] A. Yu. Ol'shanskii, *The Geometry of Defining Relations in Groups [in Russian]*, Nauka, Moscow (1989).
- [28] K. Schulz, *Makanin's Algorithm for Word Equations - Two Improvements and a Generalization*, *Proceedings of the First International Workshop on Word Equations and Related Topics*, p.85-150, 1990.
- [29] S. L. Shestakov, *The equation $[x, y] = g$ in partially commutative groups*, *Siberian Math. J.*, V. 46 (2005), p. 364-372.

MONTSERRAT CASALS-RUIZ, DEPARTMENT OF MATHEMATICS AND STATISTICS, MCGILL UNIVERSITY, 805 SHERBROOKE ST. WEST, MONTREAL, QUEBEC H3A 2K6, CANADA

ILYA V. KAZACHKOV, DEPARTMENT OF MATHEMATICS AND STATISTICS, MCGILL UNIVERSITY, 805 SHERBROOKE ST. WEST, MONTREAL, QUEBEC H3A 2K6, CANADA